NEFNESS OF ADJOINT BUNDLES FOR AMPLE VECTOR BUNDLES OF CORANK 3

ANDREA LUIGI TIRONI

ABSTRACT. Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on a smooth complex projective variety X of dimension n. The aim of this paper is to describe the structure of pairs (X, \mathcal{E}) as above whose adjoint bundles $K_X + \det \mathcal{E}$ are not nef for r = n - 3. Furthermore, we give some immediate consequences of this result in adjunction theory.

1. INTRODUCTION

Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on a smooth complex projective variety X of dimension n. We say that a Cartier divisor D on X is numerically effective (nef) if it has non-negative intersection number with any curve C on X.

In the late 1980s, Ye and Zhang started to study the nefness of the adjoint bundle $K_X + \det \mathcal{E}$, obtaining in [40] complete results for $r \ge n-1$.

Subsequently, Zhang investigated in [41] also the case r = n - 2, showing that in this situation $K_X + \det \mathcal{E}$ is nef except in a few cases.

Some years later, by the adjunction theory and the remark that a key result of [37] is still valid without the extra assumption of spannedness for \mathcal{E} , Maeda gave in [17] an exhaustive description of all the ample vector bundles \mathcal{E} for the case r = n - 2.

Let us note that a weaker version of the main result of [41], in which \mathcal{E} is assumed to be also generated by global sections, was already proved by Wiśniewski in [37] and that all the above results are a generalization of those due to Fujita [10] and Ionescu [13].

Thus the purpose of this paper is to describe the structure of pairs (X, \mathcal{E}) as above whose adjoint bundles $K_X + \det \mathcal{E}$ are not nef in the next case r = n - 3.

The final statement which we obtain in Section 3 is as follows:

Theorem 1.1. Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on a smooth complex projective variety X of dimension n. If r = n - 3, then $K_X + \det \mathcal{E}$ is nef except when (X, \mathcal{E}) is one of the following:

- (1) $(\mathbb{P}^n, \bigoplus_{i=1}^{n-3} \mathcal{O}_{\mathbb{P}^n}(a_i))$, where all the a_i 's are positive integers such that $\sum_{i=1}^{n-3} a_i \leq n$;
- (2) $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-3-a} \oplus \mathcal{V})$, where \mathcal{V} is an indecomposable Fano vector bundle on \mathbb{P}^n of rank a such that $3 \leq a \leq n-3$, $c_1(\mathcal{V}) = a+3$ and its generic splitting types are (3, 2, 1, 1, ..., 1) or (2, 2, 2, 1, ..., 1); moreover, $\mathcal{V}(-1) := \mathcal{V} \otimes \mathcal{O}(-1)$ is nef for $3 \leq a \leq n-6$, but it is not globally generated for $n \geq 6$;

Date: December 7, 2017.

- (3) $(\mathbb{Q}^n, \bigoplus_{j=1}^{n-3} \mathcal{O}_{\mathbb{Q}^n}(b_j))$, where all the b_j 's are positive integers such that $\sum_{j=1}^{n-3} b_j \leq n-1$;
- (4) $(\mathbb{Q}^n, \mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-3-b} \oplus \mathcal{V})$, where \mathcal{V} is an indecomposable Fano vector bundle on \mathbb{Q}^n of rank b such that $3 \leq b \leq n-3$, $c_1(\mathcal{V}) = b+2$ and its generic splitting type is (2, 2, 1, ..., 1); moreover, $\mathcal{V}(-1)$ is nef on \mathbb{Q}^n for $3 \leq b \leq n-5$, but it is not globally generated for $n \geq 7$;
- (5) X is a Fano n-fold of index n-1 with $\operatorname{Pic}(X)$ generated by an ample line bundle \mathcal{H} and either (α) $\mathcal{E} \cong \mathcal{H}^{\oplus n-3}$, or (β) $\mathcal{E}|_l \cong \mathcal{H}_l^{\oplus n-4} \oplus \mathcal{H}_l^{\otimes 2}$ for every line l of (X, \mathcal{H}) ;
- (6) X is a Fano n-fold of index n-2 with $\operatorname{Pic}(X)$ generated by an ample line bundle \mathcal{L} and $\mathcal{E} \cong \mathcal{L}^{\oplus n-3}$;
- (7) $(\mathbb{P}^3 \times \mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(1, 1)^{\oplus 3});$
- (8) $(\mathbb{P}^2 \times \mathbb{Q}^3, \mathcal{O}_{\mathbb{P}^2 \times \mathbb{Q}^3}(1, 1)^{\oplus 2});$
- (9) $(\mathbb{P}_{\mathbb{P}^3}(T), [\xi_T]^{\oplus 2})$, where T is the tangent bundle of \mathbb{P}^3 and ξ_T is the tautological line bundle;
- (10) $(\mathbb{P}_{\mathbb{P}^3}(\mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus 2}), [\xi]^{\oplus 2}), \xi$ being the tautological line bundle;
- (11) there is a vector bundle \mathcal{V} on a smooth curve C such that $X \cong \mathbb{P}_C(\mathcal{V})$; moreover, for any fibre $F \cong \mathbb{P}^{n-1}$ of $X \to C$, we have $\mathcal{E}|_F$ is isomorphic to one of the following: (a) $\mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3}$; (b) $\mathcal{O}_{\mathbb{P}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}$; (c) $\mathcal{O}_{\mathbb{P}^{n-1}}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-5}$; (d) $\mathcal{O}_{\mathbb{P}^{n-1}}(3) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}$;
- (12) X is a section of a divisor of relative degree two in a projective space $\mathbb{P}_C(\mathcal{G})$, where \mathcal{G} is a vector bundle of rank n + 1 on a smooth curve C; moreover, for any smooth fibre $F \cong \mathbb{Q}^{n-1}$ of $X \to C$, where \mathbb{Q}^{n-1} is a smooth quadric hypersurface of \mathbb{P}^n , we have $(F, \mathcal{E}|_F)$ is one of the following pairs:
 - (i) $(\mathbb{Q}^{n-1}, \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus n-3});$
 - (ii) $(\mathbb{Q}^{n-1}, \mathcal{O}_{\mathbb{Q}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus n-4});$
 - (iii) $(\mathbb{Q}^4, \mathcal{S} \otimes \mathcal{O}_{\mathbb{Q}^4}(2))$, where \mathcal{S} is the spinor bundle on $\mathbb{Q}^4 \subset \mathbb{P}^5$;
- (13) the map $\Phi: X \to C$ associated to the linear system $|(n-3)K_X + (n-2)\det \mathcal{E}|$ makes X a Del Pezzo fibration over a smooth curve C; moreover, any general smooth fibre F of Φ is either a Del Pezzo (n-1)-fold with $\operatorname{Pic}(F) \cong \mathbb{Z}[\mathcal{O}_F(1)]$ and such that $\mathcal{E}|_F \cong \mathcal{O}_F(1)^{\oplus n-3}$, or $\mathbb{P}^2 \times \mathbb{P}^2$ with $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^2}(1, 1)^{\oplus 2}$;
- (14) there is a vector bundle \mathcal{V} on a smooth surface S such that $X \cong \mathbb{P}_{S}(\mathcal{V})$; moreover, for any fibre $F \cong \mathbb{P}^{n-2}$ of $X \to S$, we have $\mathcal{E}|_{F} \cong \mathcal{O}_{\mathbb{P}^{n-2}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus n-4}$;
- (15) there exists a \mathbb{P}^{n-2} -fibration $\pi : X \to S$, locally trivial in the complex (or étale) topology, over a smooth surface S such that $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus n-3}$ for any closed fiber $F \cong \mathbb{P}^{n-2}$ of the map π ;
- (16) the map $\psi : X \to S$ associated to the linear system $|(n-3)K_X + (n-2)\det \mathcal{E}|$ makes X a quadric fibration over a smooth surface S; moreover, for any general fibre $F \cong \mathbb{Q}^{n-2}$ we have $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{Q}^{n-2}}(1)^{\oplus n-3}$, where \mathbb{Q}^{n-2} is a smooth quadric hypersurface of \mathbb{P}^{n-1} ;
- (17) there is a vector bundle \mathcal{F} on a smooth 3-fold V such that $X \cong \mathbb{P}_V(\mathcal{F})$; moreover, for any fibre $F \cong \mathbb{P}^{n-3}$ of $X \to V$, we have $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{n-3}}(1)^{\oplus n-3}$;

 $\mathbf{2}$

- (18) there exists a smooth variety X' and a morphism $\varphi : X \to X'$ expressing X as the blowing-up of X' at a finite set of points B and an ample vector bundle \mathcal{E}' on X' such that $\mathcal{E} \otimes ([\varphi^{-1}(B)]) \cong \varphi^* \mathcal{E}'$ and $K_{X'} + \tau' \det \mathcal{E}'$ is nef, where $\tau' < \frac{n-1}{n-3}$ is the nefvalue of the pair (X', det \mathcal{E}'). Moreover, $\mathcal{E}|_E \cong \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3}$ for any irreducible component E of the exceptional locus of φ ;
- (19) the map $\psi: X \to X'$ associated to the linear system $|(n-2)K_X + (n-1)\det \mathcal{E}|$ is a birational morphism which contracts an extremal face spanned by extremal rays R_i for some *i* in a finite set of index. Let $\psi_i: X \to X_i$ be the contraction associated to R_i . Then each ψ_i is birational and of divisorial type; moreover, if E_i is an exceptional divisor of ψ_i , then $(E_i, [E_i]_{E_i}, \mathcal{E}|_{E_i}) \cong (\mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(-2), \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3});$
- (20) the map $\phi : X \to X'$ associated to the linear system $|(n-3)K_X + (n-2)\det \mathcal{E}|$ is a birational morphism which contracts an extremal face. Let R_i be the extremal rays spanning this face for some *i* in a finite set of index. Call $\rho_i : X \to W_i$ the contraction associated to one of the R_i . Then we have each ρ_i is birational and of divisorial type; if D_i is one of the exceptional divisors and $Z_i = \rho_i(D_i)$, we have $\dim Z_i \leq 1$ and one of the following possibilities can occur:
 - (j) dim $Z_i = 0, D_i \cong \mathbb{P}^{n-1}$ and $([D_i]_{D_i}, \mathcal{E}|_{D_i}) \cong (\mathcal{O}_{\mathbb{P}^{n-1}}(-1), \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4} \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(2));$
 - (jj) dim $Z_i = 0, D_i$ is a (possible singular) quadric $\mathbb{Q}^{n-1} \subset \mathbb{P}^n$ and $[D_i]_{D_i} = \mathcal{O}_{\mathbb{Q}^{n-1}}(-1)$; moreover, $\mathcal{E}|_{D_i} \cong \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus n-3}$;
 - (jjj) dim $Z_i = 1$, W_i and Z_i are smooth projective varieties and ρ_i is the blow-up of W_i along Z_i ; moreover, $\mathcal{E}|_{F_i} \cong \mathcal{O}_{F_i}(1)^{\oplus n-3}$ and $\mathcal{O}_{D_i}(D_i)|_{F_i} = \mathcal{O}_{\mathbb{P}^{n-2}}(-1)$ for any fibre $F_i \cong \mathbb{P}^{n-2}$ of $D_i \to Z_i$.

Moreover, the map ϕ is a composition of disjoint extremal contractions as in (j), (jj) and (jjj).

The Theorem 1.1 is a generalization of a result due to Fujita [10, theorem 4] and its proof makes use principally of [2] and all the above papers [17], [40] and [41], together with some recent results ([3] and [26]) about the nef value of pairs $(X, \det \mathcal{E})$.

Note that Theorem 1.1 is complete for n = 5 (or $\operatorname{Pic}(X) \neq \mathbb{Z}$). When $n \geq 6$ and X is a Fano manifold of index $i(X) \geq n-2$ with $\operatorname{Pic}(X) = \mathbb{Z}[\mathcal{O}_X(1)]$, a finest description of \mathcal{E} is equivalent to have in this setting a complete classification of ample vector bundles $\mathcal{F} := \mathcal{E} \oplus \mathcal{O}_X(1)$ of rank $n-2 \geq 4$ such that either $c_1(\mathcal{F}) = -K_X$ (see also [2, theorem 5.1 (2)(i)]), or $\mathbb{P}_X(\mathcal{F}) := W$ is a Fano manifold of index $i(W) = \frac{\dim W - 1}{2} \geq 4$ (the case i(W) = 3 implies n = 5; see [23, theorem 1.3] for the case i(W) = 2). However, a more satisfactory and effective description of \mathcal{E} when $\operatorname{Pic}(X) = \mathbb{Z}[\mathcal{O}_X(1)]$ is given in Corollary 3.1 under the extra assumption that $\mathcal{E} \otimes \mathcal{O}_X(-1)$ is globally generated on X.

In the last section, we consider two immediate applications of Theorem 1.1. More precisely, in §4.1 we extend [15, theorem 3], by giving a description of ample vector bundles \mathcal{E} of rank $r \geq 2$ on a smooth projective variety X of dimension n which admit a global section $s \in \Gamma(\mathcal{E})$ whose zero locus $Z := (s)_0 \subset X$ is a smooth subvariety of the expected dimension $n - r \geq 3$ and such that $K_Z + (\dim Z - 3)H_Z$ is not nef (Propositions 4.1 and

4.2), where H_Z denotes the restriction to Z of an ample line bundle H on X. These results partially overlap with those obtained in [4, §§4, 5, 6 and 7], but they are proved in a simple and different way, and together with [15, theorems 1, 2 and 3], they describe the structure of all pairs (X, \mathcal{E}) as above such that (Z, H_Z) is a special polarized variety until the second reduction map in the sense of the adjunction theory (e.g., see [6, chapter 7] for a complete classification of such pairs (Z, H_Z)). Finally, in §4.2 we study classical scrolls over manifolds of dimension five and Picard number one (Corollary 4.2 and Remark 4.1), extending some results of [7] and [36, §3] about classical scrolls which are not adjunction-theoretic scrolls (we refer to §1 and the first part of §4.2 for the difference between the two definitions of scrolls).

2. NOTATION AND BACKGROUND MATERIAL.

In this note varieties are always assumed to be defined over the complex number field \mathbb{C} . We use the standard notation from algebraic geometry. The words "vector bundles" and "locally free sheaves" are used interchangeably. Let X be a smooth irreducible projective variety of dimension n (for simplicity, n-fold). The group of line bundles on X is denoted by Pic(X). Moreover, we denote by $\rho(X)$ the Picard number of X. The tensor products of line bundles are denoted additively. The pull-back $j^*\mathcal{E}$ of a vector bundle \mathcal{E} on X by an embedding $j: V \to X$ is denoted by $\mathcal{E}|_V$ (or, for simplicity, by \mathcal{E}_V when no confusion arises). The canonical bundle of an n-fold X is denoted by K_X . An n-fold X is said to be a Fano manifold if its anticanonical bundle $-K_X$ is ample. For Fano manifolds X, the largest integer i(X) which divides $-K_X$ in Pic(X) is called the *index* of X.

A polarized n-fold is a pair (X, L) consisting of an n-fold X and an ample line bundle L on X. A polarized n-fold (X, L) is said to be a classical scroll over a smooth variety Y if $(X, L) \cong (\mathbb{P}_Y(\mathcal{V}), \xi_{\mathcal{V}})$ for some ample vector bundle \mathcal{V} on Y, where $\xi_{\mathcal{V}}$ is the tautological line bundle on the projective space $\mathbb{P}_Y(\mathcal{V})$ associated to \mathcal{V} . We say that (X, L) is an adjunction-theoretic scroll (respectively a quadric fibration, respectively a Del Pezzo fibration, respectively a Mukai fibration) over a normal variety Y of dimension m if there exists a surjective morphism with connected fibres $p: X \to Y$ and an ample line bundle H on Y, such that $K_X + (n-m+1)L \simeq p^*H$ (respectively $K_X + (n-m)L \simeq p^*H$, respectively $K_X + (n-m-1)L \simeq p^*H$, respectively $K_X + (n-m-2)L \simeq p^*H$). We say that (X, L)is a Del Pezzo variety (respectively a Mukai variety) if $K_X \simeq -(n-1)L$ (respectively $K_X \simeq -(n-2)L$). Moreover, for general results about the adjunction theory, we refer to [6] and [34].

A part of Mori's theory of extremal rays is to be used throughout the paper. So, let $Z_1(X)$ be the free abelian group generated by integral curves on an *n*-fold X. The intersection pairing gives a bilinear map $\operatorname{Pic}(X) \times Z_1(X) \to \mathbb{Z}$ and the *numerical equivalence* \equiv is defined so that the pairing

$$((\operatorname{Pic}(X)/\equiv)\otimes\mathbb{Q})\times((Z_1(X)/\equiv)\otimes\mathbb{Q})\to\mathbb{Q}$$

is non-degenerate. The closed cone of curves $\overline{NE}(X)$ is the closed convex cone generated by effective 1-cycles in the \mathbb{R} -vector space $(Z_1(X)/\equiv)\otimes\mathbb{R}$. We say that $L \in \operatorname{Pic}(X)$ is nef if the numerical class of L in $(\operatorname{Pic}(X)/\equiv)\otimes\mathbb{R}$ gives a non-negative function on $\overline{NE}(X) - \{0\}$. Let Z be a 1-cycle on X. We denote by [Z] the numerical class of Z in $(Z_1(X)/\equiv)\otimes\mathbb{R}$. A half line $R = \mathbb{R}_+[Z]$ in $\overline{NE}(X)$ is called an *extremal ray* if

(1)
$$K_X \cdot Z < 0$$
, and

(2) if $z_1, z_2 \in \overline{NE}(X)$ satisfy $z_1 + z_2 \in R$, then $z_1, z_2 \in R$.

A rational (possibly singular) reduced and irreducible curve C on X is called an *extremal* rational curve if $\mathbb{R}_+[C]$ is an extremal ray and $(-K_X) \cdot C \leq n+1$. Let

$$\overline{NE}(X)^+ = \{ z \in \overline{NE}(X) \mid K_X \cdot z \ge 0 \}.$$

Then we have the following basic theorem in the Mori theory.

Theorem 2.1 (Cone Theorem). Let X be a smooth projective variety. Then NE(X) is the smallest closed convex cone containing $\overline{NE}(X)^+$ and all the extremal rays:

$$\overline{NE}(X) = \overline{NE}(X)^+ + \sum_i R_i,$$

where the R_i are extremal rays of $\overline{NE}(X)$ for X. For any open convex cone V containing $\overline{NE}(X)^+ - \{0\}$ there exist only a finite number of extremal rays that do not lie in $V \cup \{0\}$. Furthermore, every extremal ray is spanned by a numerical class of an extremal rational curve.

For the proof of the above result, we refer to [21, theorem 1.5] and [22, (1.2)]. In the next section, we will use also the following well-known

Lemma 2.1. Let \mathcal{E} be an ample vector bundle of rank r on a rational curve C. Then $\det \mathcal{E} \cdot C \geq r$.

Now, let us give here some technical results about vector bundles on some Fano *n*-folds X with $\rho(X) = 1$.

Lemma 2.2. Let \mathcal{V} be an ample vector bundle of rank r on a Fano n-fold X with $\rho(X) = 1$. Assume that $c_1(\mathcal{V}) \leq i(X) - 1$, where i(X) is the index of X. If $\frac{n-r+1}{i(X)-c_1(\mathcal{V})} < 2$ then $\mathcal{V} \otimes \mathcal{O}_X(-1)$ is a nef vector bundle on X.

Proof. By adjunction we have $-K_{\mathbb{P}(\mathcal{V})} \simeq r\xi_{\mathcal{V}} + \pi^*\mathcal{O}_X(i(X) - c_1(\mathcal{V}))$, where $\xi_{\mathcal{V}}$ is the tautological line bundle on $\mathbb{P}(\mathcal{V})$ and $\pi : \mathbb{P}(\mathcal{V}) \to X$ is the projection map. Since $i(X) - c_1(\mathcal{V}) > 0$ and $\xi_{\mathcal{V}}$ is ample, we see that $\mathbb{P}(\mathcal{V})$ is a Fano (n+r-1)-fold. Let R be an extremal ray of $\mathbb{P}(\mathcal{V})$ different from a line in a fiber of π . Then we get

$$n+1 \ge -K_{\mathbb{P}(\mathcal{V})} \cdot R \ge r + [i(X) - c_1(\mathcal{V})]\pi^*\mathcal{O}_X(1) \cdot R.$$

This gives $2 > \frac{n-r+1}{i(X)-c_1(\mathcal{V})} \ge \pi^* \mathcal{O}_X(1) \cdot R$, showing that $0 \le \pi^* \mathcal{O}_X(1) \cdot R' \le 1$ for any extremal ray R' on $\mathbb{P}(\mathcal{V})$. So, let C be an effective irreducible curve on $\mathbb{P}(\mathcal{V})$. From Theorem 2.1 it

follows that $C \equiv \sum_i \alpha_i R_i$ with $\alpha_i > 0$, where each R_i is an extremal ray of $\mathbb{P}(\mathcal{V})$. Therefore we deduce that

$$[\xi_{\mathcal{V}} - \pi^* \mathcal{O}_X(1)] \cdot C = \sum_i \alpha_i [\xi_{\mathcal{V}} - \pi^* \mathcal{O}_X(1)] \cdot R_i \ge 0.$$

So by definition $\xi_{\mathcal{V}} - \pi^* \mathcal{O}_X(1)$ is nef on $\mathbb{P}(\mathcal{V})$, i.e. $\mathcal{V} \otimes \mathcal{O}_X(-1)$ is nef on X. \Box

Lemma 2.3. Let \mathcal{V} be a globally generated vector bundle of rank r < n on \mathbb{P}^n with $c_1 :=$ $c_1(\mathcal{V}) \leq \frac{n}{2}$. If $r \geq c_1 + 1$, then one of the following possibilities can occurs:

- (1) $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}(c_1) \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r-1};$ (2) $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1} \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r-c_1};$ (3) $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}(2) \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1-2} \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r-c_1+1};$

- (4) $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}(3) \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1 3} \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r c_1 + 2};$ (5) $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1 4} \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r c_1 + 2};$
- (6) $c_1 \geq 5$ and there exist the following exact sequences:

$$0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V} \to \mathcal{V}^{r-1} \to 0, \ \dots, 0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V}^{k+1} \to \mathcal{V}^k \to 0,$$

where $2 \leq k \leq c_1 - 3$, $\operatorname{rk} \mathcal{V}^i = i$, $c_1(\mathcal{V}^i) = c_1$, all the \mathcal{V}^i 's are globally generated and $(s_k)_0 \neq \emptyset$ for a generic section s_k of \mathcal{V}^k .

Proof. Let $Z_r = (s_r)_0$ be the zero locus of a generic section s_r of \mathcal{V} . If $Z_r \neq \emptyset$, then Z_r is smooth and of dimension n-r. By adjunction $K_{Z_r} = \mathcal{O}_{\mathbb{P}^n}(-n-1+c_1)|_{Z_r}$. Thus $n+1-c_1 \leq \dim Z_r+1 = n-r+1$, i.e. $r \leq c_1$, but this is a contradiction. Hence $Z_r = \emptyset$ and by induction (see, e.g., [27, (4.3.2), p.83]) we obtain the following exact sequences:

 $0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V} \to \mathcal{V}^{r-1} \to 0, \ \dots, 0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V}^{k+1} \to \mathcal{V}^k \to 0.$

where $k \leq c_1$, $\operatorname{rk}\mathcal{V}^i = i$, $c_1(\mathcal{V}^i) = c_1$, all the \mathcal{V}^i 's are globally generated and $Z_k = (s_k)_0 \neq \emptyset$ for a generic section s_k of \mathcal{V}^k . If k = 1 then $\mathcal{V}^1 = \mathcal{O}_{\mathbb{P}^n}(c_1)$ and by induction we deduce that $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}^{\oplus r-1} \oplus \mathcal{O}_{\mathbb{P}^n}(c_1)$. So $2 \leq k \leq c_1$. Assume that $k = c_1 \geq 2$. Since $Z_{c_1} = Z_k \neq \emptyset$, Z_{c_1} is smooth and of dimension $n - c_1$. Moreover, since $2(n - c_1) - n = n - 2c_1 \ge 0$, we obtain that Z_{c_1} is also irreducible. Thus by the Kobayashi-Ochiai Theorem we have $Z_{c_1} = \mathbb{P}^{n-c_1}$. Note that $\mathcal{N}_{Z_{c_1}/\mathbb{P}^n} \cong \mathcal{O}_{Z_{c_1}}(1)^{\oplus c_1}$. Since

$$\mathcal{V}_{|Z_{c_1}}^{c_1} \cong \mathcal{N}_{Z_{c_1}/\mathbb{P}^n} \cong \mathcal{O}_{\mathbb{P}^{n-c_1}}(1)^{\oplus c_1}$$

and dim $Z_{c_1} = n - c_1 \ge c_1 \ge 2$, by [27, Ch.I (2.3.2)] we conclude that \mathcal{V}^{c_1} splits, and then $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}^{r-c_1} \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1}$. Suppose now that $k = c_1 - 1 \geq 2$. By arguing as above, we have $Z_{c_1-1} \neq \emptyset$ is smooth, irreducible and of dimension $n - c_1 + 1$. Therefore by the Kobayashi-Ochiai Theorem we know that $Z_{c_1-1} \cong \mathbb{Q}^{n-c_1+1} \subset \mathbb{P}^n$. Hence $\mathcal{N}_{Z_{c_1-1}/\mathbb{P}^n} \cong$ $\mathcal{O}_{\mathbb{P}^n}(2)_{|Z_{c_1-1}} \oplus \mathcal{O}_{\mathbb{P}^n}(1)_{|Z_{c_1-1}}^{c_1-2} \cong \mathcal{V}_{|Z_{c_1-1}}^{c_1-1}$. Since dim $Z_{c_1-1} = n - c_1 + 1 \ge c_1 + 1 = k + 2 \ge 4$, we see that there exists a linear $\mathbb{P}^2 \subset Z_{c_1-1}$ such that $\mathcal{V}_{\mathbb{P}^2}^{c_1-1}$ splits. This shows that $\mathcal{V} \cong \mathcal{O}_{\mathbb{P}^n}^{r-c_1+1} \oplus \mathcal{O}_{\mathbb{P}^n}(2) \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1-2}$. Therefore $2 \leq k \leq c_1-2$. Finally, suppose that $k = c_1 - 2 \ge 2$. By arguing as in the above cases, we obtain that $Z_{c_1-2} \neq \emptyset$ is a smooth

irreducible Del Pezzo $(n-c_1+2)$)-fold with index $i(Z_{c_1-2}) = n-c_1+1 \ge \frac{n}{2}+1 > \frac{\dim Z_{c_1-2}}{2}+1$. Then by [38] we know that $\rho(Z_{c_1-2}) = 1$. Furthermore, since $\dim Z_{c_1-2} = n-c_1+2 \ge \frac{n}{2}+2 \ge c_1+2 \ge 6$, by the classification of Del Pezzo manifolds (see, e.g., [12]), we conclude that Z_{c_1-2} is one of the following manifolds:

- (a) a cubic hypersurface in \mathbb{P}^{n-c_1+3} ;
- (b) a complete intersection of two quadric hypersurfaces $\mathbb{Q}_i \subset \mathbb{P}^{n-c_1+4}$;
- (c) $\mathbb{G}(1,4) \subset \mathbb{P}^9$.

Note that Case (c) does not occur, since in this situation $n = c_1 + 4 \leq \frac{n}{2} + 4$, i.e. $\mathbb{G}(1,4) \subset \mathbb{P}^n$ with $n \leq 8$, a contradiction. On the other hand, in Case (a) we get

$$\mathcal{V}_{|Z_{c_1-2}}^{c_1-2} \cong \mathcal{N}_{Z_{c_1-2}/\mathbb{P}^n} \cong \mathcal{O}_{\mathbb{P}^n}(3)_{|Z_{c_1-2}} \oplus \mathcal{O}_{\mathbb{P}^n}(1)_{|Z_{c_1-2}}^{\oplus c_1-3}$$

while in Case (b) we have

$$\mathcal{V}_{|Z_{c_1-2}}^{c_1-2} \cong \mathcal{N}_{Z_{c_1-2}/\mathbb{P}^n} \cong \mathcal{O}_{\mathbb{P}^n}(2)_{|Z_{c_1-2}}^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^n}(1)_{|Z_{c_1-2}}^{\oplus c_1-4}$$

By [33] we see that in both cases there exists a linear $\mathbb{P}^2 \subset Z_{c_1-2}$. So \mathcal{V}^{c_1-2} splits and by induction we obtain that \mathcal{V}^r is either $\mathcal{O}_{\mathbb{P}^n}(3) \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1-3} \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r-c_1+2}$, or $\mathcal{O}_{\mathbb{P}^n}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus c_1-4} \oplus \mathcal{O}_{\mathbb{P}^n}^{\oplus r-c_1+2}$. \Box

Corollary 2.1. Let \mathcal{V} be a globally generated vector bundle of rank r < n on \mathbb{P}^n with $c_1(\mathcal{V}) = 3$. If $n \ge 6$ then \mathcal{V} splits.

Proof. If $r \ge 4$ then we conclude by Lemma 2.3. Since $\mathcal{V}(1)$ is ample and $c_1(\mathcal{V}(1)) = c_1(\mathcal{V}) + r = 3 + r$, if $r \le 2$ then by [1, theorem (9.1)] we know that $\mathcal{V}(1)$ splits, i.e. \mathcal{V} splits. So let r = 3. Let s_3 be a general section of \mathcal{V} and put $Z_3 = (s_3)_0$. If $Z_3 = \emptyset$ then we obtain the following exact sequence

$$0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V} \to \mathcal{V}^2 \to 0,$$

where \mathcal{V}^2 is a rank-2 vector bundle on \mathbb{P}^n . Since $\mathrm{rk}\mathcal{V}^2 = 2$ and $n \geq 6$, from [1, (9.1)] we deduce by a similar argument as above that \mathcal{V}^2 splits, i.e. \mathcal{V} splits. Therefore we can assume that $Z_3 \neq \emptyset$. Then Z_3 is smooth and of dimension n-3. Since $2(n-3)-n = n-6 \geq 0$, we see that Z_3 is irreducible and by adjunction $K_{Z_3} = (-n-1+c_1)\mathcal{O}_{\mathbb{P}^n}(1)|_{Z_3} = (-n+2)\mathcal{O}_{\mathbb{P}^n}(1)|_{Z_3}$. Since $n-2 = \dim Z_3+1$, by the Kobayashi-Ochiai Theorem we obtain that $Z_3 = \mathbb{P}^{n-3} \subset \mathbb{P}^n$. Hence

$$\mathcal{V}|_{Z_3} \cong \mathcal{N}_{Z_3/\mathbb{P}^n} \cong \mathcal{O}_{\mathbb{P}^{n-3}}(1)^{\oplus 3}$$

with dim $Z_3 = n - 3 \ge 3$, and this shows that \mathcal{V} splits. \Box

Corollary 2.2. Let \mathcal{V} be a globally generated vector bundle of rank r < n on \mathbb{P}^n with $c_1(\mathcal{V}) = 4$. If $n \geq 8$ then \mathcal{V} splits.

Proof. If $r \ge 5$ then we obtain the statement by Lemma 2.3. Let $2 \le r \le 4$. Therefore we have either (i) $Z_r := (s_r)_0 \ne \emptyset$ for a generic section s_r of \mathcal{V} , or by induction (ii) there exist the following exact sequences

$$0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V} \to \mathcal{V}^{r-1} \to 0, \ \dots, 0 \to \mathcal{O}_{\mathbb{P}^n} \to \mathcal{V}^{k+1} \to \mathcal{V}^k \to 0,$$

where $1 \leq k \leq r-1 \leq 3$, $\operatorname{rk}\mathcal{V}^i = i$, $c_1(\mathcal{V}^i) = c_1$, all the \mathcal{V}^i 's are globally generated and $Z_k := (s_k)_0 \neq \emptyset$ for a generic section s_k of \mathcal{V}^k . In Case (i), $Z_r \neq \emptyset$ is a smooth irreducible (n-r)-fold with $K_{Z_r} + (n-3)\mathcal{O}_{\mathbb{P}^n}(1)|_{Z_r} = \mathcal{O}_{\mathbb{P}^n}$ and $n-3 = \dim Z_r + r - 3$. Since $2 \leq r \leq 4$, we get one of the following possibilities:

(α) r = 4 and $Z_4 = \mathbb{P}^{n-4}$; (β) r = 3 and $Z_3 = \mathbb{Q}^{n-3} \subset \mathbb{P}^{n-2}$; (γ) r = 2.

In Case (γ) , since $\mathcal{V}(1)$ is ample and $c_1(\mathcal{V}(1)) = c_1(\mathcal{V}) + r = 6$, from [1, (9.1)] it follows that \mathcal{V} splits. Moreover, by similar arguments as in the proof of Corollary 2.1, we see that \mathcal{V} splits also in Cases (α) and (β) .

Consider now Case (ii). Note that k = 3, since otherwise $k \leq 2$ and we could deduce by [1, (9.1)] that \mathcal{V}^k splits, i.e. \mathcal{V} splits. Thus $Z_3 \neq \emptyset$ is a smooth irreducible (n-3)-fold. Moreover, by adjunction $K_{Z_3} = \mathcal{O}_{\mathbb{P}^n}(n-3)|_{Z_3}$ and from the Kobayashi-Ochiai Theorem we get $Z_3 = \mathbb{Q}^{n-3} \subset \mathbb{P}^n$. Thus $\mathcal{V}|_{Z_3} \cong \mathcal{N}_{Z_3/\mathbb{P}^n} \cong \mathcal{O}_{\mathbb{Q}^{n-3}}(2) \oplus \mathcal{O}_{\mathbb{Q}^{n-3}}(1)^{\oplus 2}$ and since dim $Z_3 = n-3 \geq 5$, we conclude that there exists a $\mathbb{P}^2 \subset Z_3$, i.e. \mathcal{V} splits. \Box

Finally, let us give here also some useful results about globally generated vector bundles on a smooth quadric hypersurface $\mathbb{Q}^n \subset \mathbb{P}^{n+1}$.

Lemma 2.4. Let \mathcal{V} be a globally generated vector bundle of rank r < n on $\mathbb{Q}^n \subset \mathbb{P}^{n+1}$ with $c_1 := c_1(\mathcal{V}) \leq \frac{n-3}{2}$. If $r \geq c_1 + 2$, then either \mathcal{V} splits, or $c_1 \geq 2$ and there exist exact sequences

 $0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V} \to \mathcal{V}^{r-1} \to 0, \ \dots, 0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V}^{k+1} \to \mathcal{V}^k \to 0,$

where $2 \leq k \leq c_1$, $\operatorname{rk} \mathcal{V}^i = i$, $c_1(\mathcal{V}^i) = c_1$, all the \mathcal{V}^i 's are globally generated and $(s_k)_0 \neq \emptyset$ is an irreducible smooth (n-k)-fold for a generic section s_k of \mathcal{V}^k .

Proof. Let Z_r be the zero locus of a general section s_r of \mathcal{V} . If $Z_r \neq \emptyset$ then Z_r is smooth and of dimension n-r. Since by adjunction $K_{Z_r} + (n-c_1)\mathcal{O}_{\mathbb{Q}^n}(1)|_{Z_r} = \mathcal{O}_{Z_r}$, we deduce that $n-c_1 \leq \dim Z_r + 1 = n-r+1$, i.e. $r \leq c_1 + 1$, a contradiction. Thus $Z_r = \emptyset$ and by induction there exist the following exact sequences:

$$0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V} \to \mathcal{V}^{r-1} \to 0, \ \dots, 0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V}^{k+1} \to \mathcal{V}^k \to 0,$$

where $1 \leq k \leq c_1+1$, $\operatorname{rk}\mathcal{V}^i = i$, $c_1(\mathcal{V}^i) = c_1$, all the \mathcal{V}^i 's are globally generated and $(s_k)_0 \neq \emptyset$ is smooth and of dimension (n-k) for a generic section s_k of \mathcal{V}^k . Note that \mathcal{V} splits for k = 1. Thus suppose that $2 \leq k \leq c_1+1$. Moreover, if $c_1 \leq 1$, then from [32] we know that \mathcal{V} splits again. So we can assume also that $c_1 \geq 2$. If $k = c_1 + 1 \geq 3$, then $Z_{c_1+1} = Z_k \neq \emptyset$ is smooth and of dimension $n - c_1 - 1$. Since

$$2(n-k) - n - 1 = n - 2k - 1 = n - 2(c_1 + 1) - 1 = n - 2c_1 - 3 \ge 0,$$

we see that Z_{c_1+1} is also irreducible. Furthermore, by adjunction we get

$$K_{Z_{c_1+1}} + (n-c_1)\mathcal{O}_{\mathbb{Q}^n}(1)|_{Z_{c_1+1}} \simeq \mathcal{O}_{Z_{c_1+1}}$$

where $n - c_1 = \dim Z_{c_1+1} + 1$. From the Kobayashi-Ochiai Theorem it follows that $Z_{c_1+1} = \mathbb{P}^{n-c_1-1} \subset \mathbb{Q}^n$. Hence $n - c_1 - 1 \leq \frac{n}{2}$, i.e. $n \leq 2c_1 + 2$, but this is absurd. \Box

Corollary 2.3. Let \mathcal{V} be a globally generated vector bundle of rank r < n on $\mathbb{Q}^n \subset \mathbb{P}^{n+1}$ with $c_1(\mathcal{V}) = 2$. If $n \geq 7$ then \mathcal{V} splits.

Proof. Note that $\mathcal{V}' := \mathcal{V} \otimes \mathcal{O}_{\mathbb{Q}^n}(1)$ is ample with $c_1(\mathcal{V}') = c_1(\mathcal{V}) + \mathrm{rk}\mathcal{V} = 2 + \mathrm{rk}\mathcal{V}$. Since $n \geq 7$, if r = 2 then \mathcal{V}' is a Fano bundle of rank-2 on \mathbb{Q}^n and by [1, (2.4)(2)] we can conclude that \mathcal{V}' splits, i.e. \mathcal{V} splits. Let $r \geq 3$. If $r \geq 4$, then from Lemma 2.4 we deduce that either \mathcal{V} splits, or there exist exact sequences

$$0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V} \to \mathcal{V}^{r-1} \to 0, \ \dots, 0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V}^3 \to \mathcal{V}^2 \to 0,$$

where $\operatorname{rk} \mathcal{V}^i = i$, $c_1(\mathcal{V}^i) = 2$, all the \mathcal{V}^i 's are globally generated and $(s_2)_0 \neq \emptyset$ is smooth and of dimension (n-2) for a general section s_2 of \mathcal{V}^2 . Since $\mathcal{V}^2(1)$ is a Fano bundle of rank 2 on \mathbb{Q}^n with $n \geq 7$, by [1, main theorem (2.4)(2)] we see that $\mathcal{V}^2(1)$ splits, i.e. \mathcal{V}^2 splits. Therefore, from the above exact sequences we deduce that also \mathcal{V} splits.

Finally, suppose that r = 3. Let $Z_3 = (s_3)_0$ be the zero locus of a general section s_3 of \mathcal{V} . If $Z_3 = \emptyset$ then there exists an exact sequence $0 \to \mathcal{O}_{\mathbb{Q}^n} \to \mathcal{V} \to \mathcal{V}^2 \to 0$, where \mathcal{V}^2 splits since $n \ge 7$, that is, \mathcal{V} splits. Hence $Z_3 \neq \emptyset$ is smooth and of dimension n-3. Since $2(n-3) - n - 1 = n - 7 \ge 0$, we see that Z_3 is also irreducible. By adjunction we obtain that $K_{Z_3} + (n-2)\mathcal{O}_{\mathbb{Q}^n}(1)|_{Z_3} = \mathcal{O}_{Z_3}$ with $n-2 = \dim Z_3 + 1$. Therefore $Z_3 = \mathbb{P}^{n-3} \subset \mathbb{Q}^n$ and this implies that $n-3 \le \frac{n}{2}$, i.e. $n \le 6$, but this gives a contradiction. \Box

Let X be an n-fold. If R is an extremal ray, then its length l(R) is defined as

 $l(R) := \min\{(-K_X) \cdot C \mid C \text{ is a rational curve such that } [C] \in R\}.$

Note that $0 < l(R) \le n + 1$ from Theorem 2.1 and the definition of an extremal rational curve. Let \mathcal{E} be an ample vector bundle of rank $r \ge 2$ on X and let $\Omega(X, \mathcal{E})$ be the set of extremal rays R such that $(K_X + \det \mathcal{E}) \cdot R < 0$. Then it follows from Theorem 2.1 that the set $\Omega(X, \mathcal{E})$ is finite. For any extremal ray R in $\Omega(X, \mathcal{E})$ we define a positive integer

$$\Lambda(X, \mathcal{E}, R) := (-K_X - \det \mathcal{E}) \cdot C,$$

where C is an extremal rational curve such that $-K_X \cdot C = l(R)$.

3. Proof of Theorem 1.1.

Suppose that $K_X + \det \mathcal{E}$ is not nef. By Theorem 2.1 we can find an extremal ray R with $(K_X + \det \mathcal{E}) \cdot R < 0$, and so $\Omega(X, \mathcal{E}) \neq \emptyset$. Since the set $\Omega(X, \mathcal{E})$ is finite, define the positive integer

$$\Lambda(X,\mathcal{E}) := \max\{\Lambda(X,\mathcal{E},R) \mid R \in \Omega(X,\mathcal{E})\}.$$

Therefore we have only the following three possibilities:

(1) $\Lambda(X, \mathcal{E}) \ge 3;$ (2) $\Lambda(X, \mathcal{E}) = 2;$ (3) $\Lambda(X, \mathcal{E}) = 1.$

From now on, we proceed with a case-by-case analysis.

Case (1). Note that, by definition of $\Lambda(X, \mathcal{E})$, there exists an extremal rational curve $C \in R$ such that $-(K_X + \det \mathcal{E}) \cdot C \geq 3$ and $l(R) = -K_X \cdot C$. So, by Lemma 2.1 we get

$$-K_X \cdot C \ge \det \mathcal{E} \cdot C + 3 \ge (n-3) + 3 = n,$$

i.e. $l(R) \ge n$. Let $f_R : X \to Y$ be the extremal ray contraction associated to R. From [6, lemma 6.3.12], we deduce that f_R is of fibre type with dim $Y \le 1$. Let F be a smooth general fibre of f_R . Thus we have the following two cases:

(a)
$$\dim F = n - 1$$
, $\dim Y = 1$; (b) $\dim F = n$, $\dim Y = 0$.

In Case (a), by [6, lemma 6.3.12(2)] we know that Y is a smooth irreducible curve and $\rho(F) = 1$. Note that $\mathcal{E}|_F$ is an ample vector bundle on F of rank dimF - 2.

Claim. $K_F + \det \mathcal{E}|_F$ is not nef and $\Lambda(F, \mathcal{E}|_F) \ge 3$ for a general fibre F of $f_R : X \to Y$.

Since $\rho(F) = 1$, we have $\overline{NE}(F) \cong \langle C' \rangle$ with $[C'] \in R = \mathbb{R}_+[C]$, i.e. $C' \equiv \delta C$ for some $\delta > 0$. Thus for any effective curve $\gamma \subset F$, we know that $\gamma \equiv \delta' C'$ for some $\delta' > 0$ and so

$$(K_F + \det \mathcal{E}|_F) \cdot \gamma = (K_X + \det \mathcal{E})_F \cdot (\delta'C') = \delta'(K_X + \det \mathcal{E}) \cdot (\delta C) = \delta\delta'(K_X + \det \mathcal{E}) \cdot C < 0.$$

This shows that $K_F + \det \mathcal{E}|_F$ cannot be nef and that $-K_F$ is ample on F. Then by Theorem 2.1, we see that $\overline{NE}(F) \cong \mathbb{R}_+[C']$, where C' is an extremal rational curve on F. Moreover, since

$$\delta(-K_X \cdot C) = -K_X \cdot C' \ge l(R) = -K_X \cdot C > 0,$$

we deduce that $\delta \geq 1$. Thus we get for $C' \subset F$

$$(-K_F - \det \mathcal{E}|_F) \cdot C' = (-K_X - \det \mathcal{E}) \cdot C' = (-K_X - \det \mathcal{E}) \cdot \delta C \ge 3\delta \ge 3,$$

and this gives $\Lambda(F, \mathcal{E}|_F) \ge 3$ for a general fibre F of $f_R : X \to Y$. Q.E.D.

By the *Claim* and [41, Proposition 1.1'], we obtain that $(F, \mathcal{E}|_F)$ is one of the following pairs:

$$(a_{1}) \ (\mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3}); (a_{2}) \ (\mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}); (a_{3}) \ (\mathbb{Q}^{n-1}, \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus n-3}).$$

In Case (a_2) , we get $-(K_F + \det \mathcal{E}|_F) = \mathcal{O}_{\mathbb{P}^{n-1}}(2)$. Take a line $l \subset F$. Since $l \in R = \mathbb{R}_+[C]$ and $l(R) = -K_X \cdot C$, by arguing as in the proof of *Claim 1*, we deduce that $l \equiv \rho C$ for some $\rho \geq 1$, but this gives the following numerical contradiction

$$2 = -(K_F + \det \mathcal{E}|_F) \cdot l = -(K_X + \det \mathcal{E}) \cdot \rho C \ge 3\rho \ge 3.$$

By a similar argument, we see that also Case (a_3) cannot occur. Consider now Case (a_1) and let $c := \det \mathcal{E} \cdot C$. As in Case (a_2) , taking a line $l \subset F$, we deduce that $3 = (-K_F - \det \mathcal{E}|_F) \cdot l \ge -(K_X + \det \mathcal{E}) \cdot C$, i.e. $(K_X + \det \mathcal{E}) \cdot C = -3$. Therefore

$$[cK_X + (c+3)\det\mathcal{E}] \cdot C = c[K_X + \det\mathcal{E}] \cdot C + 3c = 0$$

and from [6, theorem 4.3.1] it follows that

$$cK_X + (c+3) \det \mathcal{E} \simeq f_R^* D,$$

for some $D \in Pic(Y)$. Then we have

(1)
$$cK_X + (c+3)(\det \mathcal{E} + (n-3)f_R^*A) \simeq f_R^*H$$

for a suitable very ample line bundle A on Y such that H := D + (c+3)(n-3)A is ample on Y. Restricting (1) to a general fibre F of f_R , we get

$$\mathcal{O}_F \simeq cK_F + (c+3)\det \mathcal{E}|_F = \mathcal{O}_F(-nc + (c+3)(n-3)).$$

Hence c = n - 3 and (1) becomes

(2)
$$(n-3)K_X + n(\det \mathcal{E} + (n-3)f_R^*A) \simeq f_R^*H.$$

Put $\mathcal{E}' := \mathcal{E} \otimes f_R^* A$ and note that \mathcal{E}' is ample and such that $\det \mathcal{E}' = \det \mathcal{E} + (n-3)f_R^* A$. Thus we get

$$(n-3)K_X + n(\det \mathcal{E}') \simeq f_R^* H$$

Moreover, since

$$\left[K_X + \left(\frac{n}{n-3}\right)\det \mathcal{E}'\right] \cdot l = \left[K_F + \left(\frac{n}{n-3}\right)\det \mathcal{E}|_F\right] \cdot l = 0,$$

for any line $l \subset F \cong \mathbb{P}^{n-1}$, we see that the nef value $\tau(X, \mathcal{E}')$ of the pair (X, \mathcal{E}') is such that $\tau(X, \mathcal{E}') \geq \frac{n}{n-3} = \frac{n}{\mathrm{rk}\mathcal{E}'} > 1$. Thus by [3, proposition 4(3)] we have $X \cong \mathbb{P}_Y(\mathcal{V})$ for a suitable rank *n* vector bundle \mathcal{V} on *Y* and from [26, theorem 1.3(5)] we conclude that $\mathcal{E}' \cong \xi_{\mathcal{V}} \otimes f_R^* \mathcal{G}$, where \mathcal{G} is an ample vector bundle of rank n-3 on *Y*.

Finally, in Case (b), since X is a Fano *n*-fold with $\rho(X) = 1$, for any effective rational curve Γ on X we see that $\Gamma \in R$ and $-K_X \cdot \Gamma \geq l(R) \geq n$. Thus by [19], [20] and [9] we can conclude that X is either \mathbb{P}^n or \mathbb{Q}^n . In the former case, since $\Lambda(X, \mathcal{E}) \geq 3$, by Lemma 2.1 we have

$$n-3 \le c_1(\mathcal{E}) \cdot l \le -K_{\mathbb{P}^n} \cdot l - 3 = n-2,$$

where l is a line in \mathbb{P}^n . Therefore by [27, theorems 3.2.1 and 3.2.3], \mathcal{E} is isomorphic to either $\mathcal{O}(1)^{\oplus n-3}$, or $\mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus n-4}$. In the latter case, since $X \cong \mathbb{Q}^n$ and $\Lambda(X, \mathcal{E}) \ge 3$, we see that

$$n-3 \le c_1(\mathcal{E}) \cdot l \le -K_{\mathbb{Q}^n} \cdot l - 3 = n-3,$$

i.e. $c_1(\mathcal{E}) \cdot l = n - 3$, and then $\mathcal{E} \cong \mathcal{O}(1)^{\oplus n-3}$ by [37, lemma 3.6.1].

Summing up the above discussion, we obtain the following

Proposition 3.1. If \mathcal{E} is an ample vector bundle of rank $n-3 \geq 2$ such that $K_X + \det \mathcal{E}$ is not nef and $\Lambda(X, \mathcal{E}) \geq 3$, then the pair (X, \mathcal{E}) is one of the following:

- (1') $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-3});$ (2') $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2) \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-4});$ (3') $(\mathbb{Q}^n, \mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-3});$
- (4) there is a vector bundle \mathcal{V} on a smooth curve C such that $X \cong \mathbb{P}_C(\mathcal{V})$ and $\mathcal{E} \cong$ $\xi_{\mathcal{V}} \otimes \pi^* \mathcal{G}'$, where $\pi : X \to C$ is the projection map, $\xi_{\mathcal{V}}$ is the tautological line bundle of $\mathbb{P}(\mathcal{V})$ and \mathcal{G}' is a suitable vector bundle of rank n-3 on C; in particular, $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3}$ for any fibre $F \cong \mathbb{P}^{n-1}$ of the map π .

Case (2). In this situation, we know that there exists an extremal rational curve $C \in R$ such that $-(K_X + \det \mathcal{E}) \cdot C = 2$ and $-K_X \cdot C = l(R)$. Moreover, note that

$$\left[K_X + \left(1 + \frac{2}{\det \mathcal{E} \cdot C}\right) \det \mathcal{E}\right] \cdot C = 0,$$

i.e. the nef value $\tau(X, \mathcal{E})$ of the pair $(X, \det \mathcal{E})$ is such that

(3)
$$\tau(X,\mathcal{E}) \ge 1 + \frac{2}{\det \mathcal{E} \cdot C}.$$

Since $-K_X \cdot C \leq n+1$, by Lemma 2.1 we get det $\mathcal{E} \cdot C = l(R) - 2$ and

$$n-3 \leq \det \mathcal{E} \cdot C = -K_X \cdot C - 2 \leq n-1.$$

So we have only the following three possibilities:

- (I) det $\mathcal{E} \cdot C = n 1$, $\tau(X, \mathcal{E}) \ge \frac{n+1}{n-1} = \frac{l(R)}{n-1}$; (II) det $\mathcal{E} \cdot C = n 2$, $\tau(X, \mathcal{E}) \ge \frac{n}{n-2} = \frac{l(R)}{n-2}$; (III) det $\mathcal{E} \cdot C = n 3$, $\tau(X, \mathcal{E}) \ge \frac{n-1}{n-3} = \frac{l(R)}{n-3}$.

In Case (I), we see that $-K_X \cdot C = n+1$, i.e. l(R) = n+1. From the Ionescu-Wiśniewski inequality, [6, lemma 6.3.12] and [9], it follows that $X \cong \mathbb{P}^n$. Since by (3) we have

$$\tau(X,\mathcal{E}) \ge \frac{n+1}{n-1} \ge \frac{n-2}{n-3} = \frac{n-2}{\operatorname{rk} \mathcal{E}} > 1$$

and det $\mathcal{E} \cdot C - (n-3) = 2$, we obtain by [3, proposition 6(a)(1)] and [26] that \mathcal{E} is a decomposable vector bundle with $c_1(\mathcal{E}) = \mathcal{O}_{\mathbb{P}^n}(n-1)$. Since \mathcal{E} is ample, this gives $\mathcal{E} \cong \mathcal{O}(3) \oplus \mathcal{O}(1)^{\oplus n-4} \text{ or } \mathcal{O}(2)^{\oplus 2} \oplus \mathcal{O}(1)^{\oplus n-5}.$

In Case (II), note that

$$\tau(X,\mathcal{E}) \ge \frac{n}{n-2} > \frac{n-2}{n-3} = \frac{n-2}{\operatorname{rk}\mathcal{E}} > 1$$

and det $\mathcal{E} \cdot C - (n-3) = 1$. Moreover, $l(R) = -K_X \cdot C = n$ and from [6, lemma 6.3.12] we have only the following two cases:

- (A) X is a Fano n-fold with $\rho(X) = 1$;
- (B) $\rho(X) = 2$ and the contraction $f_R: X \to Y$ associated to the extremal ray $R = \mathbb{R}_+[C]$ is a morphism onto a smooth curve Y, whose general fibre F is a smooth Fano (n-1)-fold with $\rho(F) = 1$.

In Case (A), by [19] and [20] we obtain that $X \cong \mathbb{Q}^n \subset \mathbb{P}^{n+1}$ with $c_1(\mathcal{E}) = \mathcal{O}_{\mathbb{Q}^n}(n-2)$. Thus from [26, theorem 1.3(15)] we get $\mathcal{E} \cong \mathcal{O}_{\mathbb{Q}^n}(2) \oplus \mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-4}$. In Case (B), we conclude by [26, theorem 1.3(10)].

Finally, in Case (III) by [3, theorem 1] we have

$$\tau(X, \mathcal{E}) = \frac{n-1}{n-3} = \frac{l(R)}{\operatorname{rk} \mathcal{E}}$$

Since $l(R) = -K_X \cdot C = n-1$ and $n = r+3 \ge 5$, we can conclude by [3, propositions 3, 4 and 5] and [26, theorem 1.3(6), (8), (11) and (12)(a)]. Moreover, if $\rho(X) = 1$ then from [3, theorem 1(2)] it follows that $\mathcal{E} \cong \mathcal{O}_X(1)^{\oplus n-3}$, which leads to Case (5)(α) of Theorem 1.1.

The above discussion can be summed up in the following

Proposition 3.2. If \mathcal{E} is an ample vector bundle of rank $n-3 \ge 2$ such that $K_X + \det \mathcal{E}$ is not nef and $\Lambda(X, \mathcal{E}) = 2$, then the pair (X, \mathcal{E}) is one of the following:

- $(1'') (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(3) \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-4});$
- $(2'') (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-5});$
- (3") $(\mathbb{Q}^n, \mathcal{O}_{\mathbb{Q}^n}(2) \oplus \mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-4});$
- (4") there exist a vector bundle \mathcal{V} on a smooth curve C such that $X \cong \mathbb{P}_C(\mathcal{V})$ and an exact sequence

$$0 \to \pi^* \mathcal{L} \otimes \xi_{\mathcal{V}}^{\otimes 2} \to \mathcal{E} \to \pi^* \mathcal{G} \otimes \xi_{\mathcal{V}} \to 0$$

for some line bundle \mathcal{L} on C and some vector bundle \mathcal{G} of rank $n-4 \geq 1$ on C, where $\pi : \mathbb{P}(\mathcal{V}) \to C$ is the projection map and $\xi_{\mathcal{V}}$ is the tautological line bundle on $\mathbb{P}_{C}(\mathcal{V})$; in particular, we have $\mathcal{E}|_{F} \cong \mathcal{O}_{\mathbb{P}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}$ for any fibre $F \cong \mathbb{P}^{n-1}$ of the map π ;

- (5") X is a Del Pezzo n-fold with $\rho(X) = 1$ and $\mathcal{E} \cong L^{\oplus n-3}$, where L is the ample generator of $\operatorname{Pic}(X)$;
- (6") X is a quadric fibration, $q: X \to C$, of the relative Picard number one over a smooth curve C (i.e. X is a section of a divisor of relative degree two in a projective space $\mathbb{P}_C(\mathcal{F})$, where \mathcal{F} is a vector bundle of rank n + 1 on C) and there exist a q-ample line bundle $\mathcal{O}_X(1)$ on X and an ample vector bundle \mathcal{G} of rank n - 3 on C such that $\mathcal{E} \cong \mathcal{O}_X(1) \otimes q^*\mathcal{G}$, where $\mathcal{O}_X(1)|_F \simeq \mathcal{O}_{\mathbb{Q}^{n-1}}(1)$ for any fibre $F \cong \mathbb{Q}^{n-1} \subset \mathbb{P}^n$ of the map q;
- (7") there exists a \mathbb{P}^{n-2} -fibration $\pi : X \to S$, locally trivial in the complex (or étale) topology, over a smooth surface S such that $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus n-3}$ for any closed fiber $F \cong \mathbb{P}^{n-2}$ of the map π ;
- (8") there exists a smooth variety X' and a morphism $\varphi : X \to X'$ expressing X as the blowing-up of X' at a finite set of points B and an ample vector bundle \mathcal{E}' on X' such that $\mathcal{E} \otimes ([\varphi^{-1}(B)]) \cong \varphi^* \mathcal{E}'$ and $K_{X'} + \tau' \det \mathcal{E}'$ is nef, where $\tau' < \frac{n-1}{n-3}$ is the nefvalue of the pair (X', det \mathcal{E}'). Moreover, $\mathcal{E}|_E \cong \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3}$, where E is any irreducible component of the exceptional locus of the map φ .

Case (3). By arguing as in [17, p.77] and [37, proposition 3.5], note that in fact we get $L \cdot C = 1$, where $L := (m-1)K_X + m \det \mathcal{E}$ is ample on X and $m \ge n-3 \ge 2$ with

 $m := \min\{(\det \mathcal{E}) \cdot C \mid C \text{ extremal rational curve s.t. } (K_X + \det \mathcal{E}) \cdot C = -1\}.$

So, define $\mathcal{V} := \mathcal{E} \oplus L$. Then \mathcal{V} is an ample vector bundle of rank n-2 on X such that

$$(K_X + \det \mathcal{V}) \cdot C = (K_X + \det \mathcal{E}) \cdot C + L \cdot C = -1 + 1 = 0.$$

Therefore $K_X + \det \mathcal{V}$ cannot be ample on X. Note also that $K_X + \det \mathcal{V}$ must be nef. Otherwise, by Theorem 2.1 we could find an extremal ray R on X such that

$$-1 \ge (K_X + \det \mathcal{V}) \cdot R \ge (K_X + \det \mathcal{E}) \cdot R + 1,$$

i.e. $(K_X + \det \mathcal{E}) \cdot R \leq -2$, but this would contradict $\Lambda(X, \mathcal{E}) = 1$.

Thus assume that $K_X + \det \mathcal{V}$ is nef but not big. From [2, theorem 5.1(2)] we know that there exists a morphism $\Phi : X \to W$ onto a normal variety W supported by (a large multiple of) $K_X + \det \mathcal{V}$ and $\dim W \leq 3$. Moreover, let F be a general fibre of Φ . We have the following according to $s := \dim W$:

- (*i*₁) if s = 0 then X is a Fano *n*-fold with $K_X + \det \mathcal{V} = \mathcal{O}_X$. If $n \ge 6$, then $b_2(X) = 1$ except if $X \cong \mathbb{P}^3 \times \mathbb{P}^3$ and $\mathcal{E} \cong \mathcal{O}(1, 1)^{\oplus 4}$;
- (*i*₂) if s = 1 then W is a smooth curve and Φ is a flat (equidimensional) map. If $n \ge 6$ then Φ is an elementary contraction. Moreover, by [24] and [31], $(F, \mathcal{V}|_F)$ is one of the following pairs:
 - (a) $(\mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4});$
 - (b) $(\mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(3) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3});$
 - (c) $(\mathbb{Q}^{n-1}, \mathcal{O}_{\mathbb{Q}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus n-3});$
 - (d) $(\mathbb{Q}^4, \mathcal{S}(2) \oplus \mathcal{O}_{\mathbb{Q}^{n-1}}(1))$, where \mathcal{S} is the spinor bundle on $\mathbb{Q}^4 \subset \mathbb{P}^5$;
 - (e) F is a Del Pezzo (n-1)-fold with $\rho(F) = 1, -K_F \simeq (n-2)H_F$, $\operatorname{Pic}(F) \cong \mathbb{Z}[H_F]$ and $\mathcal{V}|_F \cong H_F^{\oplus n-2}$;
 - (f) $(\mathbb{P}^2 \times \mathbb{P}^2, \mathcal{O}(1, 1)^{\oplus 3}).$

If the general fibre is \mathbb{P}^{n-1} then X is a classical scroll, while if the general fibre is $\mathbb{Q}^{n-1} \subset \mathbb{P}^n$ then X is a quadric bundle;

- (*i*₃) if s = 2 then W is a smooth surface, Φ is a flat map and by [11] and [30], $(F, \mathcal{V}|_F)$ is one of the following pairs:
 - (α) ($\mathbb{P}^{n-2}, \mathcal{O}_{\mathbb{P}^{n-2}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus n-3}$);
 - (β) ($\mathbb{P}^{n-2}, T_{\mathbb{P}^{n-2}}$), where $T_{\mathbb{P}^{n-2}}$ is the tangent bundle on \mathbb{P}^{n-2} ;

$$(\gamma) \ (\mathbb{Q}^{n-2}, \mathcal{O}_{\mathbb{Q}^{n-2}}(1)^{\oplus n-2}).$$

In particular, if the general fibre is \mathbb{P}^{n-2} then all the fibres are \mathbb{P}^{n-2} ;

 (i_4) if s = 3 then W is a smooth 3-fold and all the fibres of Φ are isomorphic to \mathbb{P}^{n-3} .

Furthermore, since $(K_X + \det \mathcal{V}) \cdot C = 0$, note that $L_F \cdot C = L \cdot C = 1$, C being an extremal rational curve of X which belongs into a fibre of Φ . Moreover, from $(K_X + \det \mathcal{E}) \cdot C = -1$, we deduce that

$$-K_X \cdot C = \det \mathcal{E} \cdot C + 1 \ge m + 1 = (m+1)L \cdot C,$$

and then the nef value $\tau(X, L)$ of the polarized pair (X, L) is greater than or equal to $m+1 \ge n-2$.

Case (i₁). Observe that X is a Fano *n*-fold with $-K_X \simeq \eta L$ and

$$\eta = \tau(X, L) \ge m + 1 \ge n - 2 \ge 3.$$

If $\rho(X) = 1$, then by [6, chapter 7] we know that X is either \mathbb{P}^n , \mathbb{Q}^n , a Del Pezzo *n*-fold with $-K_X \simeq (n-1)L$, or a Mukai *n*-fold with $-K_X \simeq (n-2)L$. Since det $\mathcal{E} \cdot C = -K_X \cdot C - 1 \ge \eta - 1$, note that $m \ge \eta - 1$, that is, $\eta \le m + 1$. Thus $\eta = \tau(X, L) = m + 1$ and so we get

$$L = (\eta - 2)K_X + (\eta - 1)\det \mathcal{E} = -\eta(\eta - 2)L + (\eta - 1)\det \mathcal{E},$$

i.e. det $\mathcal{E} = (\eta - 1)L$, where $\operatorname{Pic}(X) \cong \mathbb{Z}[L]$. Let us proceed with a case-by-case analysis.

First of all, assume that $X \cong \mathbb{P}^n$. Note that $L = \mathcal{O}_{\mathbb{P}^n}(1)$ and $c_1(\mathcal{E}) = \mathcal{O}_{\mathbb{P}^n}(n)$. Without loss of generality, we can write $\mathcal{E} \cong V_1 \oplus V_2$, where V_1 is an indecomposable vector bundle on \mathbb{P}^n . Put $r_i = \operatorname{rk} V_i$ and note that $r_1 + r_2 = n - 3$ with $1 \leq r_1 \leq n - 3$. Furthermore, if $\deg V_i \leq r_i + 1$, then V_i is an uniform vector bundle on \mathbb{P}^n of rank $r_i \leq n - 3$, i.e. V_i is a sum of line bundles on \mathbb{P}^n . Moreover, since V_1 is an ample vector bundle on \mathbb{P}^n such that $\deg V_1 \leq \deg \mathcal{E} = n$, if $r_1 = 2$ then by [1, theorem (9.1)] we see that V_1 splits. Thus we can assume that $r_1 \geq 3$. Then we get the following three possibilities:

- (a) $r_1 = n 3 \ge 3$ and deg $V_1 = n$;
- (b) $3 \le r_1 \le n 4$, deg $V_1 = r_1 + 2$ and deg $V_2 = r_2 + 1$;
- (c) $3 \le r_1 \le n-4$, deg $V_1 = r_1 + 3$ and deg $V_2 = r_2$.

By Lemma 2.2, we have $V_1(-1)$ is a nef vector bundle on \mathbb{P}^n either in Case (b), or in Case (c) for $3 \leq r_1 \leq n-6$. Note that Case (b) does not occur. Indeed, since $V_1(-1)$ is nef and $c_1(V_1(-1)) = 2$, by [32, Lemma 6 (1)] we know that $V_1(-1)$ is decomposable, but this leads to a contradiction. In Case (c), since V_2 is ample, we get that $V_2 \cong \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus r_2}$, that is, $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus r_2} \oplus V_1$. Finally, assume that $V_1(-1)$ is globally generated on \mathbb{P}^n . Since $n \geq 6$, from Corollary 2.1 we know that V_1 splits, a contradiction. This completes Case (2) of Theorem 1.1.

Suppose now that $X \cong \mathbb{Q}^n \subset \mathbb{P}^{n+1}$. Note that $L = \mathcal{O}_{\mathbb{Q}^n}(1)$ and $c_1(\mathcal{E}) = \mathcal{O}_{\mathbb{Q}^n}(n-1)$. As above, write $\mathcal{E} \cong V_1 \oplus V_2$, where V_1 is an indecomposable vector bundle on \mathbb{Q}^n . Put $r_i = \operatorname{rk} V_i$ and note that $r_1 + r_2 = n-3$ with $1 \leq r_1 \leq n-3$. Furthermore, if deg $V_i \leq r_i + 1$, then V_i is an uniform vector bundle on \mathbb{Q}^n of rank $r_i \leq n-3$ and by [14] we know that V_i is a sum of line bundles on \mathbb{Q}^n . Thus deg $V_1 \geq r_1 + 2$ and then

$$\deg V_2 = n - 1 - \deg V_1 \le n - 3 - r_1 = r_2.$$

This shows that deg $V_2 = r_2$ and deg $V_1 = r_1 + 2$, that is, $\mathcal{E} \cong \mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-3-r_1} \oplus V_1$. Since V_1 is an ample vector bundle on \mathbb{Q}^n such that deg $V_1 \leq \deg \mathcal{E} = n - 1$, we deduce that V_1 is a Fano bundle on \mathbb{Q}^n . Thus if $r_1 = 2$, then by [1, theorem (2.4)(2)] we see that either V_1 splits, or n = 5 and $V_1 = \mathcal{C}(a)$, where \mathcal{C} is a Cayley bundle on \mathbb{Q}^5 such that $c_1(\mathcal{C}) = -1$.

Since $4 = c_1(V_1) = -1 + 2a$, we see that a is not an integer, but this is not possible. Hence V_1 splits for $r_1 = 2$. On the other hand, if $r_1 \ge 3$ then by restricting V_1 to any line on \mathbb{Q}^n , we have the only possible splitting types are (3, 1, 1, ..., 1) or (2, 2, 1, ..., 1). So either V_1 is a uniform vector bundle of splitting type (3, 1, 1, ..., 1) and by [14] V_1 is a direct sum of line bundles on \mathbb{Q}^n , or V_1 has the generic splitting type (2, 2, 1, ..., 1). Finally, by Lemma 2.2 and Corollary 2.3, we know that $V_1(-1)$ is nef for $3 \le r_1 \le n - 6$, and it is not globally generated for $n \ge 7$. This gives Case (4) of Theorem 1.1.

Assume that X is a Fano n-fold of index n - 1. Since $c_1(\mathcal{E}) = \mathcal{O}_X(n-2)$, we see that for every line l of (X, H), where H is the ample generator of $\operatorname{Pic}(X)$, it happens that $\mathcal{E}|_l \cong H|_l^{\oplus n-4} \oplus 2H|_l$, i.e. \mathcal{E} is a uniform vector bundle on X with splitting type (2, 1, 1, ..., 1). This leads to Case $(5)(\beta)$ of Theorem 1.1.

Finally, suppose that X is a Fano *n*-fold of index n-2. From [18, theorem 2.10], it follows by hyperplane sections that there exists at least a line l of (X, L), where $\text{Pic}(X) = \mathbb{Z}[L]$. Recall that $\mathcal{V} = \mathcal{E} \oplus L$. Since $\rho(X) = 1$ and

$$\tau(X, \mathcal{V}) \ge 1 = \frac{n-2}{\mathrm{rk}\mathcal{V}} = \frac{-K_X \cdot l}{\mathrm{rk}\mathcal{V}},$$

by [5, proposition 1.2] and [3, theorem 1(2)], we obtain that $\tau(X, \mathcal{V}) = \frac{-K_X \cdot l}{\mathrm{rk}\mathcal{V}}$ and so $\mathcal{V} \cong L^{\oplus n-2}$, i.e. $\mathcal{E} \cong L^{\oplus n-3}$. This gives Case (6) in Theorem 1.1.

Assume now that $\rho(X) \ge 2$. Since $\eta \ge n-2 \ge \frac{n+1}{2}$, from [25], [38] and [39], it follows that $\eta \le \frac{n+2}{2}$, i.e. $5 \le n \le 6$, and that (X, \mathcal{V}, L) is one of the following triplets:

- $(\mathbb{P}^3 \times \mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(1, 1)^{\oplus 4}, \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(1, 1));$
- $(\mathbb{P}^2 \times \mathbb{Q}^3, \mathcal{O}_{\mathbb{P}^2 \times \mathbb{Q}^3}(1, 1)^{\oplus 3}, \mathcal{O}_{\mathbb{P}^2 \times \mathbb{Q}^3}(1, 1));$
- $(\mathbb{P}_{\mathbb{P}^3}(T), [\xi_T]^{\oplus 3}, \xi_T)$, where T is the tangent bundle of \mathbb{P}^3 and ξ_T is the tautological line bundle;
- $(\mathbb{P}_{\mathbb{P}^3}(\mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus 2}), [\xi]^{\oplus 3}, \xi)$, where ξ is the tautological line bundle.

Since $\mathcal{V} = \mathcal{E} \oplus L$, we obtain Cases (7) to (10) of Theorem 1.1.

Case (i₂). Note that $L|_F = -(K_X + \det \mathcal{E})|_F = -K_F - \det \mathcal{E}|_F$. Then

$$K_F + mL|_F = K_F + m[-K_F - \det \mathcal{E}|_F] = -[(m-1)K_F + m\det \mathcal{E}|_F] = -L|_F,$$

since $L := (m-1)K_X + m \det \mathcal{E}$. This gives

$$L|_F = -K_F - \det \mathcal{E}|_F = (m+1)L|_F - \det \mathcal{E}|_F,$$

i.e. det $\mathcal{E}|_F = mL|_F$. Moreover, observe also that the extremal rational curve C such that $L \cdot C = 1$ is contained in a fibre of Φ , since $(K_X + \det \mathcal{V}) \cdot C = 0$. Keeping in mind that $\mathcal{V} = \mathcal{E} \oplus L$ and that $L = (m-1)K_X + m \det \mathcal{E}$ with $m \ge n-3$, by all the above remarks we can easily obtain Cases (11)(c)(d), (12)(ii)(iii), and (13) of Theorem 1.1.

Case (i₃). Note that Case (β) cannot occur, since $\mathcal{V}|_F = \mathcal{E}|_F \oplus L|_F$ is a decomposable vector bundle on \mathbb{P}^{n-2} with $n-2 \geq 3$. Moreover, since the extremal rational curve C such

that $L \cdot C = 1$ is contained in a fibre of Φ , we obtain immediately Cases (14) and (16) of Theorem 1.1.

Case (i_4) . Since $K_F + \det \mathcal{V}|_F = \mathcal{O}_F$, where $\mathcal{V}|_F$ is an ample vector bundle of rank dimF+1, from [29] we deduce that $(F, \mathcal{V}|_F) \cong (\mathbb{P}^{n-3}, \mathcal{O}_{\mathbb{P}^{n-3}}(1)^{\oplus n-2})$ and then $L|_F = \mathcal{O}_{\mathbb{P}^{n-3}}(1)$ for any fibre $F \cong \mathbb{P}^{n-3}$ of Φ . Moreover, as observed in *Case* (i_2) , the extremal rational curve Csuch that $L \cdot C = 1$ is contained in a fibre of Φ . Thus by [10, (2.12)] we get immediately Case (17) of Theorem 1.1.

Finally, suppose that $K_X + \det \mathcal{V}$ is nef and big, but not ample. Then from [2, theorem 5.1(3)] we deduce that a high multiple of $K_X + \det \mathcal{V}$ defines a birational map, $\varphi : X \to X'$, which contracts an extremal face. Let $R_i = \mathbb{R}_+[C_i]$ be the extremal rays spanning this face for some *i* in a finite set of index. Call $\rho_i : X \to W_i$ the contraction associated to one of the R_i . Then we have each ρ_i is birational and of divisorial type. If D_i is one of the exceptional divisors and $Z_i = \rho_i(D_i)$, we know that $\dim Z_i \leq 1$ and one of the following possibilities can occur:

- $(j_1) \dim Z_i = 0, D_i \cong \mathbb{P}^{n-1} \text{ and } ([D_i]_{D_i}, \mathcal{V}|_{D_i}) \text{ is either } (\alpha) (\mathcal{O}(-2), \mathcal{O}(1)^{\oplus n-2}), \text{ or } (\beta)$ $(\mathcal{O}(-1), \mathcal{O}(1)^{\oplus n-3} \oplus \mathcal{O}(2));$
- $(j_2) \dim Z_i = 0, D_i$ is a (possible singular) quadric $\mathbb{Q}^{n-1} \subset \mathbb{P}^n$ and $[D_i]_{D_i} = \mathcal{O}(-1)$; moreover, $\mathcal{V}|_{D_i} \cong \mathcal{O}(1)^{\oplus n-2}$;
- $(j_3) \dim Z_i = 1, W_i \text{ and } Z_i \text{ are smooth projective varieties and } \rho_i \text{ is the blow-up of } W_i$ along Z_i ; moreover, $\mathcal{V}|_{F_i} \cong \mathcal{O}(1)^{\oplus n-2}$ for any fibre $F_i \cong \mathbb{P}^{n-2}$ of $D_i \to Z_i$.

Furthermore, we know also that φ is a composition of disjoint extremal contractions as in $(j_1), (j_2)$ and (j_3) . Note that the extremal rational curves C_i such that $R_i = \mathbb{R}_+[C_i]$ and $L \cdot C_i = 1$ are contained in D_i , since $(K_X + \det \mathcal{V}) \cdot C_i = 0$. Moreover

det
$$\mathcal{E} \cdot C_i = -K_X \cdot C_i - 1 = \begin{cases} n-2 & \text{in Case } (j_1)(\alpha) \\ n-3 & \text{otherwise} \end{cases}$$

This shows that if some ρ_i is a contraction associated to an extremal ray as in Case $(j_1)(\alpha)$, then $\varphi: X \to X'$ contracts an extremal face spanned by only extremal rays as in $(j_1)(\alpha)$. So, keeping in mind that $\mathcal{V} = \mathcal{E} \oplus L, L \cdot C_i = 1$ and $L = (m-1)K_X + m \det \mathcal{E}$ with $m \ge n-3 \ge 2$, we get Cases (19) and (20) of Theorem 1.1.

Taking into account the above discussion and the Propositions 3.1 and 3.2, we obtain the statement of Theorem 1.1. $\hfill \Box$

Finally, under the extra assumption that $\rho(X) = 1$ and $\mathcal{E}(-1) := \mathcal{E} \otimes \mathcal{O}_X(-1)$ is globally generated on X, we have the following

Corollary 3.1. Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on a smooth complex projective variety X of dimension n with $\rho(X) = 1$. Assume that $\mathcal{E}(-1)$ is globally generated on X. If r = n-3, then $K_X + \det \mathcal{E}$ is nef except when (X, \mathcal{E}) is one of the following pairs:

(a) $(\mathbb{P}^n, \bigoplus_{i=1}^{n-3} \mathcal{O}_{\mathbb{P}^n}(a_i))$, where all the a_i 's are positive integers such that $\sum_{i=1}^{n-3} a_i \leq n$;

- (b) $(\mathbb{Q}^n, \bigoplus_{j=1}^{n-3} \mathcal{O}_{\mathbb{Q}^n}(b_j))$, where all the b_j 's are positive integers such that $\sum_{j=1}^{n-3} b_j \leq n-1$;
- (c) $X \cong \mathbb{Q}^6$ and \mathcal{E} is an indecomposable Fano vector bundle on \mathbb{Q}^6 of rank 3 such that $c_1(\mathcal{E}) = 5$ and its generic splitting type is (2, 2, 1);
- (d) X is a Fano n-fold of index n-1 with $\operatorname{Pic}(X)$ generated by an ample line bundle \mathcal{H} and either (α) $\mathcal{E} \cong \mathcal{H}^{\oplus n-3}$, or (β) $\mathcal{E}|_{l} \cong \mathcal{H}_{l}^{\oplus n-4} \oplus \mathcal{H}_{l}^{\otimes 2}$ for every line l of (X, \mathcal{H}) ;
- (e) X is a Fano n-fold of index n-2 with $\operatorname{Pic}(X)$ generated by an ample line bundle \mathcal{L} and $\mathcal{E} \cong \mathcal{L}^{\oplus n-3}$;

Let us show that in the above result Case (c) really exists.

Example. Let $\mathbb{Q}^6 \subset \mathbb{P}^7$ be the 6-dimensional quadric hypersurface endowed with a spinor bundle S of rank 4. Note that $c_4(S^{\nu}) = c_4(S) = 0$ and that S^{ν} is globally generated on \mathbb{Q}^6 , where S^{ν} is the dual bundle of S (see [28, (2.8)(ii), (2.9)]). Consider now the following exact sequence

$$0 \to \mathcal{O}_{\mathbb{Q}^6} \to \mathcal{S}^\nu \to \mathcal{F} \to 0$$

given by a section of \mathcal{S}^{ν} . Then \mathcal{F} is a globally generated vector bundle on \mathbb{Q}^{6} of rank 3 such that $c_{1}(\mathcal{F}) = -c_{1}(\mathcal{S}) = 2$ ([28, (2.9)]). Put $\mathcal{E} := \mathcal{F} \otimes \mathcal{O}_{\mathbb{Q}^{6}}(1)$. This gives an example of an ample indecomposable vector bundle \mathcal{E} of rank 3 on \mathbb{Q}^{6} with $c_{1}(\mathcal{E}) = 5$ and such that $\mathcal{E} \otimes \mathcal{O}_{\mathbb{Q}^{6}}(-1) \cong \mathcal{S}^{\nu}/\mathcal{O}_{\mathbb{Q}^{6}}$ is globally generated on \mathbb{Q}^{6} .

4. Two applications.

Within the adjunction theory, let us give here two easy and immediate consequences of Theorem 1.1.

4.1. Ample vector bundles and special varieties in adjunction theory. Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on an *n*-fold X. Given a smooth projective variety Z, the classification of such varieties X containing Z as an ample divisor occupies an important position in the theory of polarized varieties. Moreover, it is well-known that the structure of Z imposes severe restrictions on that of X. Inspired by this philosophy, we generalize some results on ample divisors to ample vector bundles. In particular, by using a peculiarity of adjoint bundles, Lanteri and Maeda [15] investigated when $K_Z + tH_Z$ is not nef for $t \geq \dim Z - 2$, where H is an ample line bundle on X and $H_Z := H|_Z$. So, following here this idea, we deal with the next case $t = \dim Z - 3$. More precisely, we give a detailed description of triplets (X, \mathcal{E}, H) as above under the assumption that $K_Z + (\dim Z - 3)H_Z$ is not nef for dim $Z \geq 3$.

In particular, for dim Z = 3, we obtain the following

Proposition 4.1. Let X be an n-fold. Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on X such that there exists a global section $s \in \Gamma(\mathcal{E})$ whose zero locus $Z = (s)_0 \subset X$ is a smooth subvariety of dimension n - r = 3. Then K_Z is nef if and only if the pair (X, \mathcal{E}) is not as in Theorem 1.1 with $n \geq 5$.

18

Proof. To prove the 'if' part, assume that K_Z is not nef. Recalling that $K_Z = (K_X + \det \mathcal{E})_Z$, we deduce that $K_X + \det \mathcal{E}$ is not nef. Therefore we have (X, \mathcal{E}) is as in Theorem 1.1.

Conversely, to prove the 'only if' part of the statement, assume that the pairs (X, \mathcal{E}) are as in Theorem 1.1. Therefore, if (X, \mathcal{E}) is as in cases (1) to (10), it is easy to see that $-K_Z$ is ample, and so K_Z cannot be nef.

Suppose that (X, \mathcal{E}) is as in case (11). Let π be the projection map $X \to C$ and let s_F denote the restriction of the global section s to a general fiber F of π . Then $s_F \in \Gamma(\mathcal{E}|_F)$. Putting $D := (s_F)_0 = Z \cap F$, we have dim $D \ge 2$. On the other hand, since Z is irreducible, we have dim $D < \dim Z = 3$, i.e. dim D = 2. Moreover, note that $[D]_D \cong \mathcal{O}_D$. Thus by the adjunction formula we have $(K_Z)_D \cong K_D - [D]_D \cong K_D$. Since $-K_D$ is ample, this implies that K_Z is not nef.

In cases (12) and (13), by similar arguments as in (11), we see that K_Z cannot be nef.

Consider now case (14). Let π be the projection map $X \to S$. Since $s_F \in \Gamma(\mathcal{E}|_F)$ for a general fibre $F \cong \mathbb{P}^{n-2}$ of π , putting $D := Z \cap F$ we have dim $D \ge n-2-r = \dim Z-2 = 1$. Since $D \subset F, D \subset Z$, we deduce that

$$(K_Z)_D \cong (K_X + \det \mathcal{E})|_D \cong (K_F + \det \mathcal{E}|_F)|_D = \mathcal{O}_F(-1)|_D$$

This shows that K_Z cannot be nef.

In cases (15) and (16), by arguing as in case (14), we deduce that K_Z is not nef.

Assume now that (X, \mathcal{E}) is as in case (17). In this situation, from section 2 we know that $L := (n-4)K_X + (n-3) \det \mathcal{E}$ is an ample line bundle on X such that $L_F = \mathcal{O}_{\mathbb{P}^{n-3}}(1)$ for any fibre $F \cong \mathbb{P}^{n-3}$ of $\pi : X \to V$. Since $X \cong \mathbb{P}_V(\mathcal{F})$, we see that there exists an ample vector bundle \mathcal{V} of rank n-2 on V such that $(X, L) \cong (\mathbb{P}(\mathcal{V}), \xi_{\mathcal{V}})$, where $\xi_{\mathcal{V}}$ is the tautological line bundle on $\mathbb{P}(\mathcal{V})$. Thus

(4)
$$K_X + \det \mathcal{E} \cong -(n-2)\xi_{\mathcal{V}} + \pi^*(K_V + \det \mathcal{V}) + \det \mathcal{E}.$$

By $\left[\det \mathcal{E} - (n-3)\xi_{\mathcal{V}}\right]|_F = \mathcal{O}_F$, the relation (4) becomes

 $K_X + \det \mathcal{E} \cong -\xi_{\mathcal{V}} + \pi^* (K_V + \det \mathcal{V} + \mathcal{H}),$

for some $\mathcal{H} \in \operatorname{Pic}(V)$. Choose a very ample line bundle H on V such that $H' := K_V + \det \mathcal{V} + \mathcal{H} + H$ is ample on V. Put $L' := L + \pi^* H$. Note that also L' is ample and

(5)
$$K_X + \det \mathcal{E} + L' \cong \pi^*(H')$$

Since dim Z = 3, [16, theorem 1.1] tells us that $\rho(X) = \rho(Z)$. Obviously, $\rho(X) = \rho(V) + 1$. Hence $\rho(Z) = \rho(V) + 1$ and this shows that Z cannot be isomorphic to V via π . Moreover, since $s_F \in \Gamma(\mathcal{O}_{\mathbb{P}^{n-3}}(1)^{\oplus n-3})$ for any fibre $F \cong \mathbb{P}^{n-3}$ of π , $D := (s_F)_0 = Z \cap F$ is a linear subspace of F, i.e. D has degree one in $F \cong \mathbb{P}^{n-3}$. Thus $\pi_Z : Z \to V$ is neither an isomorphism nor a finite-to-one map onto V. So, there exists a curve $\gamma \subset Z$ such that $\pi_Z(\gamma)$ is a point of V. Restricting (5) to Z, we get $K_Z + L'_Z \cong \pi_Z^* H'$, and this gives

 $(K_Z + L'_Z) \cdot \gamma = 0$, i.e. $K_Z \cdot \gamma = -L'_Z \cdot \gamma < 0$ by the ampleness of L'_Z . This implies that K_Z is not nef.

Now, consider case (18). Let s_E denote the restriction of s to E. Therefore $s_E \in \Gamma(\mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-3})$. Thus $D := (s_E)_0 = Z \cap E$ is a linear subspace of E and we have

$$\dim D \ge n - 1 - (n - 3) = 2 = \dim Z - 1.$$

Obviously, $D \subseteq Z$. If D = Z, then $Z \cong \mathbb{P}^3$, but this gives the contradiction $\rho(X) = \rho(Z) = 1$ by [16, theorem 1.1]. Therefore $D \subsetneq Z$, and the irreducibility of Z implies that $\dim D < \dim Z$, i.e. $D \cong \mathbb{P}^2$. Furthermore,

$$\mathcal{O}_D(D) \cong \mathcal{O}_Z(Z \cap E)|_D \cong \mathcal{O}_X(E)|_D \cong \mathcal{O}_E(E)|_D \cong \mathcal{O}_{\mathbb{P}^2}(-1).$$

Thus by the adjunction formula applied to $D \subset Z$, we get

$$(K_Z)_D \cong K_D - \mathcal{O}_D(D) = \mathcal{O}_{\mathbb{P}^2}(-3) + \mathcal{O}_{\mathbb{P}^2}(1) = \mathcal{O}_{\mathbb{P}^2}(-2).$$

This shows that K_Z is not nef.

Suppose that (X, \mathcal{E}) is as in case (19). Let s_{E_i} denote the restriction of s to E_i . Then $s_{E_i} \in \Gamma(\mathcal{E}|_{E_i})$. So $D := (s_{E_i})_0 = Z \cap E_i$ is a linear subspace of E_i and

$$\dim D \ge n - 1 - (n - 3) = 2 = \dim Z - 1.$$

By [16, theorem 1.1] note that $D \neq Z$ since dim Z = 3. Thus $D \subsetneq Z$ and the irreducibility of Z implies that D is \mathbb{P}^2 . Furthermore,

$$\mathcal{O}_D(D) \cong \mathcal{O}_Z(Z \cap E_i)|_D \cong \mathcal{O}_{E_i}(E_i)|_D \cong \mathcal{O}_{\mathbb{P}^2}(-2)$$

Thus by the adjunction formula applied to $D \subset Z$, we get $(K_Z)|_D \cong K_D - \mathcal{O}_D(D) = \mathcal{O}_D(-1)$, i.e. K_Z cannot be nef.

Finally, assume that we are in case (20). Suppose that D_i is as in cases (j) and (jj). Let s_{D_i} denote the restriction of s to D_i . Then $s_{D_i} \in \Gamma(\mathcal{E}|_{D_i})$. So $D := (s_{D_i})_0 = Z \cap D_i$ is a quadric hypersurface contained in D_i and

$$\dim D \ge n - 1 - (n - 3) = 2 = \dim Z - 1.$$

By [16, theorem 1.1] note that $D \neq Z$ since dim Z = 3. Thus $D \subsetneq Z$ and the irreducibility of Z implies that D is a (possible singular) quadric $\mathbb{Q}^2 \subset \mathbb{P}^3$. Furthermore,

$$\mathcal{O}_D(D) \cong \mathcal{O}_Z(Z \cap D_i)|_D \cong \mathcal{O}_{D_i}(D_i)|_D \cong \mathcal{O}_{\mathbb{Q}^2}(-1)$$

Thus by the adjunction formula applied to $D \subset Z$, we have $(K_Z)|_D \cong K_D - \mathcal{O}_D(D) = \mathcal{O}_D(-1)$. Therefore K_Z is not nef.

Suppose now that D_i is as in case (jjj). Let s_{F_i} be the restriction of s to any fiber $F_i \cong \mathbb{P}^{n-2}$ of $D_i \to Z_i$. Then $s_{F_i} \in \Gamma(\mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus n-3})$. So $D := (s_{F_i})_0 = Z \cap F_i$ is a linear subspace of F_i and

$$\dim D \ge n - 2 - (n - 3) = 1 = \dim Z - 2.$$

20

Since $\rho(X) \neq 1$, by [16, theorem 1.1] we see that $D \neq Z$. Thus the irreducibility of Z implies that $1 \leq \dim D \leq 2$. Since $D \subset Z$ and $D \subset F_i \subset D_i$, we obtain that

$$K_Z|_D \cong (K_X + \det \mathcal{E})|_D \cong [(K_X + \det \mathcal{E})_{D_i}]|_D \cong [K_{D_i} - \mathcal{O}_{D_i}(D_i) + \det \mathcal{E}_{D_i}]|_D \cong$$
$$\cong [K_F - \mathcal{O}_{D_i}(D_i)|_F + \det \mathcal{E}_F]|_D \cong \mathcal{O}_F(-1)|_D.$$

This shows that K_Z cannot be nef. \Box

When dim $Z \ge 4$, having in mind [15, theorems 1, 2 and 3], we get the following

Proposition 4.2. Let X be an n-fold. Let \mathcal{E} be an ample vector bundle of rank $r \geq 2$ on X such that there exists a global section $s \in \Gamma(\mathcal{E})$ whose zero locus $Z = (s)_0 \subset X$ is a smooth subvariety of dimension $n - r \geq 4$ and let H be an ample line bundle on X. Suppose that $K_Z + (\dim Z - 2)H_Z$ is nef. Then $K_Z + (\dim Z - 3)H_Z$ is nef if and only if (X, \mathcal{E}, H) is not any of the following triplets:

- $\begin{array}{ll} (1') \ X \cong \mathbb{P}^n \ and \ (\mathcal{E}, H) \ is \ one \ of \ the \ following \ pairs: \\ (i) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{P}^n}(4)); \ (ii) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{P}^n}(3)); \\ (iii) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-5} \oplus \mathcal{O}_{\mathbb{P}^n}(2), \mathcal{O}_{\mathbb{P}^n}(3)); \ (iv) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-5} \oplus \mathcal{O}_{\mathbb{P}^n}(2), \mathcal{O}_{\mathbb{P}^n}(2)); \\ (v) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-6} \oplus \mathcal{O}_{\mathbb{P}^n}(2)^{\oplus 2}, \mathcal{O}_{\mathbb{P}^n}(2)); \ (vi) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-5} \oplus \mathcal{O}_{\mathbb{P}^n}(3), \mathcal{O}_{\mathbb{P}^n}(2)); \\ (vii) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-6}, \mathcal{O}_{\mathbb{P}^n}(2)); \ (viii) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-6} \oplus \mathcal{O}_{\mathbb{P}^n}(2), \mathcal{O}_{\mathbb{P}^n}(2)); \\ (ix) \ (\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n-6}, \mathcal{O}_{\mathbb{P}^n}(2)); \ (x) \ c_1(\mathcal{E}) = r+3 \ and \ H = \mathcal{O}_{\mathbb{P}^n}(1); \\ (2') \ X \cong \mathbb{Q}^n \ and \ (\mathcal{E}, H) \ is \ one \ of \ the \ following \ pairs: \\ (1) \ (\mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{Q}^n}(3)); \\ (11) \ (\mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-5} \oplus \mathcal{O}_{\mathbb{Q}^n}(2), \mathcal{O}_{\mathbb{Q}^n}(2)); \\ (111) \ (\mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-5}, \mathcal{O}_{\mathbb{Q}^n}(2), \mathcal{O}_{\mathbb{Q}^n}(2)); \\ (1v) \ (\mathcal{O}_{\mathbb{Q}^n}(1)^{\oplus n-5}, \mathcal{O}_{\mathbb{Q}^n}(2)); \end{array} \right)$
 - (v) $c_1(\mathcal{E}) = \mathcal{O}_{\mathbb{Q}^n}(r+2)$ and $H = \mathcal{O}_{\mathbb{Q}^n}(1)$;
- (3') X is a Del Pezzo n-fold with $\operatorname{Pic}(X) \cong \mathbb{Z}[\mathcal{O}_X(1)]$ and one of the following possibilities can occur:
 - (α) $\mathcal{E} \cong \mathcal{O}_X(1)^{\oplus r}$ and $H = \mathcal{O}_X(1);$
 - (β) for any line l of $(X, \mathcal{O}_X(1))$ either $\mathcal{E}|_l \cong \mathcal{O}_l(1)^{\oplus n-4}$ and $H|_l = \mathcal{O}_l(2)$, or $\mathcal{E}|_l \cong \mathcal{O}_l(1)^{\oplus r-1} \oplus \mathcal{O}_l(2)$ and $H|_l = \mathcal{O}_l(1)$;
- (4') X is a Mukai n-fold with $\operatorname{Pic}(X) \cong \mathbb{Z}[\mathcal{O}_X(1)]$ and $\mathcal{E} \cong \mathcal{O}_X(1)^{\oplus r}$;
- (5') $(\mathbb{P}^3 \times \mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(1, 1)^{\oplus 2}, \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(1, 1));$
- (6') there exists a vector bundle \mathcal{V} on a smooth curve C such that $X \cong \mathbb{P}_C(\mathcal{V})$; moreover, for any fibre $F \cong \mathbb{P}^{n-1}$ of $X \to C$, we have $(\mathcal{E}|_F, H_F)$ is isomorphic to one of the following pairs:
 - (a) $(\mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{P}^{n-1}}(2));$
 - (b) $(\mathcal{O}_{\mathbb{P}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-5}, \mathcal{O}_{\mathbb{P}^{n-1}}(2));$
 - (c) $(\mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-5}, \mathcal{O}_{\mathbb{P}^{n-1}}(2));$
 - (d) $(\mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{P}^{n-1}}(3));$
 - (e) $(\mathcal{O}_{\mathbb{P}^{n-1}}(3) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus r-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(1));$
 - (f) $(\mathcal{O}_{\mathbb{P}^{n-1}}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus r-2}, \mathcal{O}_{\mathbb{P}^{n-1}}(1));$

- (7') X is a section of a divisor of relative degree two in a projective space $\mathbb{P}_C(\mathcal{G})$, where \mathcal{G} is a vector bundle of rank n + 1 on a smooth curve C; moreover, for any smooth fibre $F \cong \mathbb{Q}^{n-1}$ of $X \to C$, where \mathbb{Q}^{n-1} is a smooth quadric hypersurface of \mathbb{P}^n , we have $(\mathcal{E}|_F, H_F)$ is isomorphic to either $(\mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{Q}^{n-1}}(2))$, or $(\mathcal{O}_{\mathbb{Q}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus r-1}, \mathcal{O}_{\mathbb{Q}^{n-1}}(1));$
- (8') the map $\Phi: X \to C$ associated to the linear system $|(n-3)K_X + (n-2)\det \mathcal{E}|$ makes X a Del Pezzo fibration over a smooth curve C; moreover, any general smooth fibre F of Φ is a Del Pezzo (n-1)-fold with $\operatorname{Pic}(F) \cong \mathbb{Z}[\mathcal{O}_F(1)]$ such that $\mathcal{E}|_F \cong \mathcal{O}_F(1)^{\oplus r}$ and $H_F \simeq \mathcal{O}_F(1)$;
- (9') there exists a vector bundle \mathcal{V} on a smooth surface S such that $X \cong \mathbb{P}_{S}(\mathcal{V})$; moreover, for any fibre $F \cong \mathbb{P}^{n-2}$ of $X \to S$, we have $(\mathcal{E}|_{F}, H_{F})$ is isomorphic to either $(\mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{P}^{n-2}}(2))$, or $(\mathcal{O}_{\mathbb{P}^{n-2}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus r-1}, \mathcal{O}_{\mathbb{P}^{n-2}}(1))$;
- (10') the map $\psi : X \to S$ associated to the linear system $|(n-3)K_X + (n-2)\det \mathcal{E}|$ makes X a quadric fibration over a smooth surface S; moreover, for any general fibre $F \cong \mathbb{Q}^{n-2}$ we have $(\mathcal{E}|_F, H_F) \cong (\mathcal{O}_{\mathbb{Q}^{n-2}}(1)^{\oplus r}, \mathcal{O}_{\mathbb{Q}^{n-2}}(1))$, where \mathbb{Q}^{n-2} is a smooth quadric hypersurface of \mathbb{P}^{n-1} ;
- (11') there is a vector bundle \mathcal{F} on a smooth 3-fold V such that $X \cong \mathbb{P}_V(\mathcal{F})$; moreover, for any fibre $F \cong \mathbb{P}^{n-3}$ of $X \to V$, we get $(\mathcal{E}|_F, H_F) \cong (\mathcal{O}_{\mathbb{P}^{n-3}}(1)^{\oplus r}, \mathcal{O}_{\mathbb{P}^{n-3}}(1))$;
- (12') the map $\psi: X \to X'$ associated to the linear system $|(n-2)K_X + (n-1)\det \mathcal{E}|$ is a birational morphism which contracts an extremal face spanned by extremal rays R_i for some *i* in a finite set of index. Let $\psi_i: X \to X_i$ be the contraction associated to R_i . Then each ψ_i is birational and of divisorial type; moreover, if E_i is an exceptional divisor of ψ_i , then $E_i \cong \mathbb{P}^{n-1}$ and $([E_i]_{E_i}, \mathcal{E}|_{E_i}, H_{E_i}) \cong$ $(\mathcal{O}_{\mathbb{P}^{n-1}}(-2), \mathcal{O}_{\mathbb{P}^{n-1}}(1))^{\oplus r}, \mathcal{O}_{\mathbb{P}^{n-1}}(1));$
- (13') the map $\phi : X \to X'$ associated to the linear system $|(n-3)K_X + (n-2)\det \mathcal{E}|$ is a birational morphism which contracts an extremal face. Let R_i be the extremal rays spanning this face for some *i* in a finite set of index. Call $\rho_i : X \to W_i$ the contraction associated to one of the R_i . Then we have each ρ_i is birational and of divisorial type; if D_i is one of the exceptional divisors and $Z_i = \rho_i(D_i)$, we have $\dim Z_i \leq 1$ and one of the following possibilities can occur:
 - (j) dim $Z_i = 0, D_i \cong \mathbb{P}^{n-1}$ and $([D_i]_{D_i}, \mathcal{E}|_{D_i}, H_{D_i})$ is either $(j_1) (\mathcal{O}_{\mathbb{P}^{n-1}}(-1), \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus n-4}, \mathcal{O}_{\mathbb{P}^{n-1}}(2)); or$ $(j_2) (\mathcal{O}_{\mathbb{P}^{n-1}}(-1), \mathcal{O}_{\mathbb{P}^{n-1}}(2) \oplus \mathcal{O}_{\mathbb{P}^{n-1}}(1)^{\oplus r-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(1));$
 - (jj) dim $Z_i = 0, D_i$ is a (possible singular) quadric hypersurface $\mathbb{Q}^{n-1} \subset \mathbb{P}^n$ and $([D_i]_{D_i}, \mathcal{E}|_{D_i}, H_{D_i}) \cong (\mathcal{O}_{\mathbb{Q}^{n-1}}(-1), \mathcal{O}_{\mathbb{Q}^{n-1}}(1)^{\oplus r}, \mathcal{O}_{\mathbb{Q}^{n-1}}(1));$
 - (jjj) dim $Z_i = 1, W_i$ and Z_i are smooth projective varieties and ρ_i is the blow-up of W_i along Z_i ; furthermore, the triplets $(\mathcal{O}_{D_i}(D_i)|_{F_i}, \mathcal{E}|_{F_i}, H_{F_i})$ are isomorphic to $(\mathcal{O}_{\mathbb{P}^{n-2}}(-1), \mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus r}, \mathcal{O}_{\mathbb{P}^{n-2}}(1))$ for any fibre $F_i \cong \mathbb{P}^{n-2}$ of the restriction map $\rho_i|_{D_i}: D_i \to Z_i$.

Moreover, the map ϕ is a composition of disjoint extremal contractions as in (j), (jj) and (jjj).

Proof. To prove the 'if' part, suppose that $K_Z + (\dim Z - 3)H_Z$ is not nef. Set $\mathcal{E}' := \mathcal{E} \oplus H^{\oplus n-r-3}$. Then \mathcal{E}' is an ample vector bundle of rank n-3 on X. Now, we get

$$(K_X + \det \mathcal{E}')|_Z = [K_X + \det \mathcal{E} + (\dim Z - 3)H]|_Z \cong K_Z + (\dim Z - 3)H_Z.$$

So, $K_X + \det \mathcal{E}'$ is not nef on X. Thus we deduce that (X, \mathcal{E}') is as in Theorem 1.1. Keeping in mind that $\mathcal{E}' = \mathcal{E} \oplus H^{\oplus n-r-3}$ with $n \ge r+4 \ge 6$, by [15, theorems 1, 2 and 3] we can easily see that cases (8), (9), (10), (11)(a), (12)(i)(iii), (18) of Theorem 1.1 cannot occur. Moreover, as to case (15), we have $H|_F \cong \mathcal{O}_{\mathbb{P}^{n-2}}(1)$ for any closed fiber $F \cong \mathbb{P}^{n-2}$ of $\pi : X \to S$. This shows that (X, H) is an adjunction-theoretic scroll over S and by [6, proposition 3.2.1 and theorem 14.1.1] and [10, (2.12)] we can conclude that there exists a suitable vector bundle \mathcal{V} of rank n-1 on S such that $(X, H) \cong (\mathbb{P}_S(\mathcal{V}), \xi_{\mathcal{V}})$, where $\xi_{\mathcal{V}}$ is the tautological line bundle on $\mathbb{P}_S(\mathcal{V})$. From [15, theorem 3(14)] it follows that $K_Z + (\dim Z - 2)H_Z$ cannot be nef, a contradiction. In all the other cases, by [15, theorems 1, 2 and 3] and easy computations we are done.

Conversely, if (X, \mathcal{E}, H) is as in cases (1') to (5'), by the adjunction formula we see that $K_Z + (\dim Z - 3)H_Z$ is not nef.

Suppose we are in case (6'). Let π be the projection map $X \to C$ and let s_F denote the restriction of the global section s to a general fibre $F \cong \mathbb{P}^{n-1}$ of π . Then $s_F \in \Gamma(\mathcal{E}|F)$ and putting $D := (s_F)_0 = Z \cap F$, we have dim $D \ge n-r-1 \ge 3$. On the other hand, since Z is irreducible, we have dim $D < \dim Z = n-r$, i.e. dim $D = \dim Z - 1 = n-r-1$. Moreover, note that $[D]_D \cong \mathcal{O}_D$. Thus $(K_Z)_D \cong K_D - [D]_D \cong K_D$. Hence $-[K_Z + (\dim Z - 3)H_Z]|_D = -[K_D + (\dim D - 2)H_D]$ is ample on $D \subset Z$ and this shows that $K_Z + (\dim Z - 3)H_Z$ cannot be nef.

In cases (7') and (8'), by a similar argument as in (5'), we can see that $K_Z + (\dim Z - 3)H_Z$ is not nef.

Consider now case (9'). Let π be the projection map $X \to S$. Since $s_F \in \Gamma(\mathcal{E}|F)$ for a general fibre $F \cong \mathbb{P}^{n-2}$ of π , putting $D := Z \cap F = (s_F)_0$ we get dim $D \ge n - r - 2 = \dim Z - 2 \ge 2$. Since $D \subset F$ and $D \subset Z$, we deduce in both situations that

$$(K_Z + (\dim Z - 3)H_Z)|_D = (K_X + \det \mathcal{E} + (\dim Z - 3)H)|_D = [K_F + \det \mathcal{E}|_F + (\dim Z - 3)H_F]|_D \cong \mathcal{O}_{\mathbb{P}^{n-2}}(-1)|_D.$$

This implies that $K_Z + (\dim Z - 3)H_Z$ cannot be nef.

By arguing as in (9'), we can see that also in case (10') the line bundle $K_Z + (\dim Z - 3)H_Z$ is not nef.

Assume now that (X, \mathcal{E}, H) is as in (11'). Let π be the projection map $\mathbb{P}_V(\mathcal{V}) \to V$. Since $s_F \in \Gamma(\mathcal{E}|_F)$ for any fibre $F \cong \mathbb{P}^{n-3}$ of π , $D := (s_F)_0 = Z \cap F$ is not empty for any fibre F, i.e. the restriction map $\pi_Z : Z \to V$ of π to Z is surjective. On the other hand,

we have dim $Z > \dim D \ge n - r - 3 = \dim Z - 3$, because of the irreducibility of Z. So dim $Z - 3 \le \dim D \le \dim Z - 1$. Furthermore, since $K_X \cong -(n-2)\xi_{\mathcal{V}} + \pi^*(K_V + \det \mathcal{V})$, where $\xi_{\mathcal{V}}$ is the tautological line bundle on $\mathbb{P}_V(\mathcal{V})$, we get

(6)
$$K_X + \det \mathcal{E} + (n - r - 3)H = [\det \mathcal{E} + (n - r - 3)H - (n - 3)\xi_{\mathcal{V}}] + \pi^*(K_V + \det \mathcal{V}) - \xi_{\mathcal{V}}.$$

Since $[\det \mathcal{E} + (n - r - 3)H - (n - 3)\xi_{\mathcal{V}}]|_F \cong \mathcal{O}_F$, it follows that

$$\det \mathcal{E} + (n - r - 3)H - (n - 3)\xi_{\mathcal{V}} \cong \pi^* \mathcal{D},$$

for some $\mathcal{D} \in \operatorname{Pic}(V)$. By (6) we obtain that

$$[K_{Z} + (\dim Z - 3)H_{Z}]|_{D} = ([K_{X} + \det \mathcal{E} + (n - r - 3)H]_{Z})|_{D} \cong$$
$$\cong ([K_{X} + \det \mathcal{E} + (n - r - 3)H]_{F})|_{D} = -(\xi_{\mathcal{V}F})|_{D} \cong \mathcal{O}_{\mathbb{P}^{n-3}}(-1)|_{D},$$

for a general fibre $D = F|_Z$ of π_Z . This implies that $K_Z + (\dim Z - 3)H_Z$ is not nef.

Suppose we are in case (12'). Let s_{E_i} denote the restriction of the section s to E_i . Then $s_{E_i} \in \Gamma(\mathcal{E}|_{E_i})$ and so $D := (s_{E_i})_0 = E_i \cap Z$ is a linear subspace of $E_i \cong \mathbb{P}^{n-1}$. Moreover, dim $D \ge n - r - 1 = \dim Z - 1 \ge 3$. Obviously, $D \subseteq Z$. If D = Z, then $Z \subset E_i \subset X$ and we have the canonical surjection $\mathcal{N}_{Z/X} \to \mathcal{N}_{E_i/X}|_Z$ between normal bundles. Since $\mathcal{N}_{Z/X} \cong \mathcal{E}|_Z$ is ample, we deduce that $\mathcal{N}_{E_i/X}|_Z$ is ample, but this gives a contradiction since E_i is an exceptional divisor on X. Thus $D \subsetneq Z$ and the irreducibility of Z implies that dim $D < \dim Z$, i.e. dim $D = \dim Z - 1 = n - r - 1$. Therefore D is isomorphic to \mathbb{P}^{n-r-1} and

$$H_Z = H_D = (H_{E_i})|_D = \mathcal{O}_{\mathbb{P}^{n-r}}(1)$$

Note that

$$\mathcal{O}_D(D) \cong \mathcal{O}_Z(Z \cap E_i)|_D \cong \mathcal{O}_X(E_i)|_D \cong \mathcal{O}_{E_i}(E_i)|_D \cong \mathcal{O}_{\mathbb{P}^{n-r-1}}(-2)$$

and

$$(D, H_D, \mathcal{O}_D(D)) \cong (\mathbb{P}^{n-r-1}, \mathcal{O}_{\mathbb{P}^{n-r-1}}(1), \mathcal{O}_{\mathbb{P}^{n-r-1}}(-2)).$$

Thus by the adjunction formula applied to $D \subset Z$, we obtain that

$$(K_Z)|_D \cong K_D - \mathcal{O}_D(D) \cong \mathcal{O}_{\mathbb{P}^{n-r-1}}(-n+r+2).$$

Hence $(K_Z + (\dim Z - 3)H_Z)|_D \cong \mathcal{O}_D(-1)$ and this shows that $K_Z + (\dim Z - 3)H_Z$ cannot be nef.

Consider now case (13')(j). Let s_{D_i} denote the restriction of the section s to D_i . Then $s_{D_i} \in \Gamma(\mathcal{E}|_{D_i})$ and so $D := (s_{D_i})_0 = D_i \cap Z$ is either a linear subspace of $D_i \cong \mathbb{P}^{n-1}$, or a (possible singular) quadric hypersurface on D_i . Moreover, dim $D \ge n - r - 1 = \dim Z - 1 \ge 3$. Obviously, $D \subseteq Z$. If D = Z, then $Z \subset D_i \subset X$ and we have the canonical surjection $\mathcal{N}_{Z/X} \to \mathcal{N}_{D_i/X}|_Z$ between normal bundles. Since $\mathcal{N}_{Z/X} \cong \mathcal{E}|_Z$ is ample, we deduce that $\mathcal{N}_{D_i/X}|_Z$ is ample, but this gives a contradiction since D_i is an exceptional divisor on X. Thus $D \subsetneq Z$ and the irreducibility of Z implies that dim $D < \dim Z$,

i.e. dim $D = \dim Z - 1 = n - r - 1$. Furthermore, D is isomorphic to either \mathbb{P}^{n-r-1} , or $\mathbb{Q}^{n-r-1} \subset \mathbb{P}^{n-r}$ and

(7)
$$H_Z = H_D = (H_{D_i})|_D = \begin{cases} \mathcal{O}_{\mathbb{P}^4}(2) & \text{in case } (j_1) \\ \mathcal{O}_{\mathbb{Q}^{n-r}}(1) & \text{in case } (j_2) \end{cases}$$

Note that

$$\mathcal{O}_D(D) \cong \mathcal{O}_Z(Z \cap D_i)|_D \cong \mathcal{O}_X(D_i)|_D \cong \mathcal{O}_{D_i}(D_i)|_D \cong \begin{cases} \mathcal{O}_{\mathbb{P}^3}(-1) & \text{in case } (j_1) \\ \mathcal{O}_{\mathbb{Q}^{n-r-1}}(-1) & \text{in case } (j_2) \end{cases}$$

Thus

$$(D, H_D, \mathcal{O}_D(D)) \cong \begin{cases} (\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(2), \mathcal{O}_{\mathbb{P}^3}(-1)) & \text{in case } (j_1) \\ (\mathbb{Q}^{n-r-1}, \mathcal{O}_{\mathbb{Q}^{n-r-1}}(1), \mathcal{O}_{\mathbb{Q}^{n-r-1}}(-1)) & \text{in case } (j_2) \end{cases}$$

and by the adjunction formula applied to $D \subset Z$, we obtain that

$$(K_Z)|_D \cong K_D - \mathcal{O}_D(D) \cong \begin{cases} \mathcal{O}_{\mathbb{P}^3}(-3) & \text{in case } (j_1) \\ \mathcal{O}_{\mathbb{Q}^{n-r-1}}(-n+r+2) & \text{in case } (j_2) \end{cases}$$

Hence $(K_Z + (\dim Z - 3)H_Z)|_D \cong \mathcal{O}_D(-1)$ and this shows that $K_Z + (\dim Z - 3)H_Z$ is not nef.

In case (13')(jj), by arguing as in (11')(j), we see that $D := Z \cap D_i \subsetneq Z$ and dim D = n-r-1. Hence $D \cong \mathbb{Q}^{n-r-1} \subset D_i$ and $H_Z|_D \cong (H_{D_i})|_D \cong \mathcal{O}_D(1)$. Moreover, $\mathcal{O}_D(D) \cong \mathcal{O}_Z(Z \cap D_i)|_D \cong \mathcal{O}_{D_i}(D_i)|_D \cong \mathcal{O}_D(-1)$ and so $K_Z|_D \cong K_D - \mathcal{O}_D(D) \cong \mathcal{O}_{\mathbb{Q}^{n-r-1}}(-n+r+2)$. This gives $(K_Z + (\dim Z - 3)H_Z)|_D \cong \mathcal{O}_D(-1)$, i.e. $K_Z + (\dim Z - 3)H_Z$ cannot be nef.

Finally, let us consider case (13')(jjj). Let s_F denote the restriction of the section sto a fibre $F \cong \mathbb{P}^{n-2}$ of $D_i \to Z_i$. Then $s_F \in \Gamma(\mathcal{O}_{\mathbb{P}^{n-2}}(1)^{\oplus r})$ with $r \leq n-4$. Thus $D := (s_F)_0 = Z \cap F$ is a linear subspace of $F \cong \mathbb{P}^{n-2}$ and dim $D \geq n-r-2 = \dim Z-2 \geq 2$. Obviously, $D \subseteq Z$. If D = Z, then $Z \cong \mathbb{P}^{n-r}$ and $H_Z = H_D = (H_F)|_D \cong \mathcal{O}_{\mathbb{P}^{n-r}}(1)$. Thus $K_Z + (\dim Z - 2)H_Z \cong \mathcal{O}_{\mathbb{P}^{n-r}}(-3)$, but this is a contradiction. Therefore, $D \subsetneq Z$ and the irreducibility of Z implies that $n - r - 2 \leq \dim D \leq n - r - 1$. Then

$$(K_Z + (\dim Z - 3)H_Z)|_D \cong [(K_X + \det \mathcal{E} + (n - r - 3)H)|_F]|_D =$$

$$= [(K_X + \det \mathcal{E} + (n - r - 3)H)|_{D_i}]|_D = [(K_{D_i} - [D_i]_{D_i} + \det \mathcal{E}|_{D_i} + (n - r - 3)H_{D_i})|_F]|_D =$$

$$= [(K_F - \mathcal{O}_{D_i}(D_i)|_F + \det \mathcal{E}|_F + (n - r - 3)H_F)]|_D = \mathcal{O}_{\mathbb{P}^{n-r}}(-1)|_D.$$

This shows that $K_Z + (\dim Z - 3)H_Z$ cannot be nef again. \Box

4.2. Some remarks on classical scrolls over fivefolds. In very classical times scrolls naturally occurred very often. Let us recall that by a *classical scroll* we mean a \mathbb{P}^k -bundle X over a variety Y together with an ample line bundle L such that $L|_F \simeq \mathcal{O}_{\mathbb{P}^k}(1)$ for any fibre $F \cong \mathbb{P}^k$ with $k = \dim X - \dim Y$. This is equivalent to saying that $(X, L) \cong (\mathbb{P}(\mathcal{E}), \xi_{\mathcal{E}})$, where $\mathcal{E} = p_*L$ is an ample vector bundle of rank k + 1 on Y and $p : X \to Y$ is the projection map. In this situation, the canonical bundle formula gives

(8)
$$K_X + (\dim X - \dim Y + 1)L = K_X + (\operatorname{rk} \mathcal{E})L \simeq p^*(K_Y + \det \mathcal{E}).$$

From the adjunction theoretic point of view the correct definition of scroll (see [35]) is that of *adjunction-theoretic scroll* over a normal variety Y. This means that there exists a morphism with connected fibres, $p: X \to Y$, such that

$$K_X + (\dim X - \dim Y + 1)L \simeq p^*H,$$

for some ample line bundle H on Y. The general fibre is $(\mathbb{P}^k, \mathcal{O}_{\mathbb{P}^k}(1))$ with $k = \dim X - \dim Y$, but the special fibres can vary quite a lot.

Note that if L is further very ample and (X, L) is an adjunction theoretic scroll over a normal variety Y of dimension $m \leq 4$, then Y is smooth and (X, L) is a \mathbb{P}^k -bundle over Y with $k = \dim X - m$. This follows from a general result due to Sommese ([35, theorem 3.3]) for $m \leq 2$ (and L merely spanned), from [8, proposition 3.2.3] and [36, proposition 2.1] for m = 3 (and L ample and spanned), and from [36, theorem 2.2] for m = 4.

The following results are concerned with the other direction. In the stable range dim $X \ge 2\dim Y - 1$, a classical scroll is also an adjunction theoretic scroll with a few easy exceptions. These results depend on Mori theory [21] and [11]. In the unstable range dim $X \le 2\dim Y - 2$ and dim $Y \le 4$, classical scrolls that are not adjunction theoretic scrolls in the modern sense have been classified in [7] and [36, §3].

As to the cases $n = 2m - 3 \ge 7$, i.e. $k = m - 3 \ge 2$ and rk $\mathcal{E} = m - 2$, let us give here this immediate consequence of [2, theorem 5.1], [17] and [24, theorem 1], as noted in [36, remark 3.3].

Proposition 4.3. Let X be an n-fold and let L be an ample line bundle on X. Assume that $(X, L) \cong (\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))$ is a \mathbb{P}^{n-m} -bundle, $\pi : X \to Y$, over a smooth variety Y of dimension m with $\mathcal{E} = \pi_* L$. If $n = 2m - 3 \ge 7$, then (X, L) is an adjunction-theoretic scroll over Y under π unless either:

- (1) $Y \cong \mathbb{P}^m$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^m}(1)^{\oplus m-2}$, $\mathcal{O}_{\mathbb{P}^m}(2) \oplus \mathcal{O}_{\mathbb{P}^m}(1)^{\oplus m-3}$, $\mathcal{O}_{\mathbb{P}^m}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^m}(1)^{\oplus m-4}$, $\mathcal{O}_{\mathbb{P}^m}(3) \oplus \mathcal{O}_{\mathbb{P}^m}(1)^{\oplus m-3}$;
- (2) $Y \cong \mathbb{Q}^m$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{Q}^m}(1)^{\oplus m-2}$, $\mathcal{O}_{\mathbb{Q}^m}(2) \oplus \mathcal{O}_{\mathbb{Q}^m}(1)^{\oplus m-3}$, where \mathbb{Q}^m is a smooth quadric hypersurface of \mathbb{P}^{m+1} ;
- (3) Y is a Del Pezzo m-fold with $b_2(Y) = 1$, i.e. $\operatorname{Pic}(Y)$ is generated by an ample line bundle $\mathcal{O}_Y(1)$ such that $-K_Y \cong (m-1)\mathcal{O}_Y(1)$ and $\mathcal{E} \cong \mathcal{O}_Y(1)^{\oplus m-2}$;
- (4) there is a vector bundle \mathcal{V} on a smooth curve C such that $Y \cong \mathbb{P}(\mathcal{V})$ and $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{m-1}}(1)^{\oplus m-2}$ for any fibre $F \cong \mathbb{P}^{m-1}$ of $Y \to C$;

- (5) there is a vector bundle \mathcal{V} on a smooth curve C such that $Y \cong \mathbb{P}(\mathcal{V})$ and $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{m-1}}(2) \oplus \mathcal{O}_{\mathbb{P}^{m-1}}(1)^{\oplus m-3}$ for any fibre $F \cong \mathbb{P}^{m-1}$ of $Y \to C$;
- (6) there is a surjective morphism $q: Y \to C$ onto a smooth curve C such that any general fibre F of q is a smooth quadric hypersurface $\mathbb{Q}^{m-1} \subset \mathbb{P}^m$ with $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{Q}^{m-1}}(1)^{\oplus m-2}$;
- (7) there is a vector bundle \mathcal{V} on a smooth surface S such that $Y \cong \mathbb{P}(\mathcal{V})$ and $\mathcal{E}|_F \cong \mathcal{O}_{\mathbb{P}^{m-2}}(1)^{\oplus m-2}$ for any fibre $F \cong \mathbb{P}^{m-2}$ of $Y \to S$;
- (8) Y is a Fano m-fold with $-K_Y \simeq \det \mathcal{E}$; moreover, if $m \ge 6$ then $b_2(Y) = 1$ except for $Y \cong \mathbb{P}^3 \times \mathbb{P}^3$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(1, 1)^{\oplus 4}$;
- (9) there exists a morphism $\Phi: Y \to W$ onto a normal variety W supported by (a large multiple of) $K_Y + \det \mathcal{E}$ and $\dim W \leq 3$; if F is a general fibre of Φ , then we have the following possibilities:
 - (a) W is a smooth curve and Φ is flat (equidimensional) map; moreover, the pair $(F, \mathcal{E}|_F)$ is one of the following:
 - (a1) $F \cong \mathbb{P}^{m-1}$ and $\mathcal{E}|_F \cong \mathcal{O}(2)^{\oplus 2} \oplus \mathcal{O}(1)^{m-4}, \mathcal{O}(3) \oplus \mathcal{O}(1)^{m-3};$
 - (a2) $F \cong \mathbb{Q}^{m-1}$ and $\mathcal{E}|_F \cong \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus m-3}$, where \mathbb{Q}^{m-1} is the quadric hypersurface of \mathbb{P}^m ;
 - (a3) $F \cong \mathbb{Q}^4$ and $\mathcal{E}|_F \cong E(2) \oplus \mathcal{O}(1)$, where E is a spinor bundle on the quadric hypersurface \mathbb{Q}^4 of \mathbb{P}^5 ;
 - (a4) F is a Del Pezzo (m-1)-fold with $b_2(F) = 1$, i.e. $\operatorname{Pic}(F)$ is generated by an ample line bundle $\mathcal{O}_F(1)$ such that $-K_F \simeq (m-2)\mathcal{O}_F(1)$ and $\mathcal{E}|_F \cong \mathcal{O}_F(1)^{\oplus m-2}$;
 - (a5) $F \cong \mathbb{P}^2 \times \mathbb{P}^2$ and $\mathcal{E}|_F \cong \mathcal{O}(1,1)^{\oplus 3}$.

In particular, if $m \ge 6$ then Φ is an elementary contraction. Furthermore, if $F \cong \mathbb{P}^{m-1}$ then Y is a classical scroll, while if $F \cong \mathbb{Q}^{m-1} \subset \mathbb{P}^m$ then Y is a quadric bundle;

- (b) W is a smooth surface, Φ is flat and for a general fibre F of Φ the pair $(F, \mathcal{E}|_F)$ is one of the following:
 - (b1) $(\mathbb{P}^{m-2}, \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus m-3});$
 - (b2) $(\mathbb{P}^{m-2}, T_{\mathbb{P}^{m-2}})$, where $T_{\mathbb{P}^{m-2}}$ is the tangent bundle on \mathbb{P}^{m-2} ;

(b3) $(\mathbb{Q}^{m-2}, \mathcal{O}(1)^{\oplus m-2})$, where \mathbb{Q}^{m-2} is a smooth quadric hypersurface in \mathbb{P}^{m-1} . In particular, if $F \cong \mathbb{P}^{m-2}$ then all the fibres of Φ are \mathbb{P}^{m-2} ;

- (c) W is a 3-fold and $\mathcal{E}|_F \cong \mathcal{O}(1)^{\oplus m-2}$ for all the fibres $F \cong \mathbb{P}^{m-3}$ of Φ ;
- (10) there exist an m-fold Y', a morphism $\psi: Y \to Y'$ expressing Y as blown up at a finite set B of points and an ample vector bundle \mathcal{E}' on Y' such that $\mathcal{E} = \pi^* \mathcal{E}' \otimes [-\psi^{-1}(B)]$; or
- (11) a high multiple of $K_Y + \det \mathcal{E}$ defines a birational map, $\varphi : Y \to \widehat{Y}$, which contracts an extremal face. Let R_i , for *i* in a finite set of index, the extremal rays spanning this face; call $\rho_i : Y \to \widehat{Y}_i$ the contraction associated to one of the R_i . Then we

have each ρ_i is birational and divisorial; if D_i is one of the exceptional divisors and $Z_i = \rho_i(D_i)$, we have dim $Z_i \leq 1$ and the following possibilities can occur:

- (i) dim $Z_i = 0, D_i \cong \mathbb{P}^{m-1}$ and $[D_i]_{D_i} \simeq \mathcal{O}(-2)$, or $\mathcal{O}(-1)$; moreover, $\mathcal{E}|_{D_i} \cong \mathcal{O}(1)^{\oplus m-2}$, or $\mathcal{E}|_{D_i} \cong \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus m-3}$;
- (ii) dim $Z_i = 0, D_i$ is a (possible singular) quadric hypersurface \mathbb{Q}^{m-1} and $[D_i]_{D_i} \simeq \mathcal{O}(-1)$ with $\mathcal{E}|_{D_i} \cong \mathcal{O}(1)^{m-2}$;
- (iii) dim $Z_i = 1$, Z_i and \widehat{Y}_i are smooth projective varieties and ρ_i is the blow-up of \widehat{Y}_i along Z_i ; moreover, $\mathcal{E}|_f \cong \mathcal{O}(1)^{\oplus m-2}$ for any fibre $f \cong \mathbb{P}^{m-2}$ of $\rho_i|_{D_i} : D_i \to Z_i$.

Specializing the above result to the case m = 5, we get also the following

Corollary 4.1. Let X be a smooth n-fold with $n \ge 7$ and L an ample line bundle on X. Assume that $(X, L) \cong (\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))$ is a \mathbb{P}^k -bundle, $\pi : X \to Y$, over a smooth 5-fold Y with $\mathcal{E} = \pi_* L$. Then (X, L) is an adjunction scroll over Y under π unless either

- (1') $n = 10, Y \cong \mathbb{P}^5$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 6}$;
- (2') $n = 9, Y \cong \mathbb{P}^5$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 5}, \mathcal{O}_{\mathbb{P}^5}(2) \oplus \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 4}$ or $T_{\mathbb{P}^5}$, where $T_{\mathbb{P}^5}$ is the tangent bundle of \mathbb{P}^5 ;
- (3') $n = 9, Y \cong \mathbb{Q}^5$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{Q}^5}(1)^{\oplus 5}$, where \mathbb{Q}^5 is a smooth quadric hypersurface of \mathbb{P}^6 ;
- (4') n = 9, there is a rank 5 vector bundle \mathcal{V} over a smooth curve C such that $Y \cong \mathbb{P}(\mathcal{V})$ and $\mathcal{E}|_f \cong \mathcal{O}_f(1)^{\oplus 5}$ for any fibre $f \cong \mathbb{P}^4$ of $Y \to C$;
- (5') n = 8, $Y \cong \mathbb{P}^5$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 4}$, $\mathcal{O}_{\mathbb{P}^5}(2) \oplus \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 3}$, $\mathcal{O}_{\mathbb{P}^5}(2)^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 2}$, $\mathcal{O}_{\mathbb{P}^5}(3) \oplus \mathcal{O}_{\mathbb{P}^5}(1)^{\oplus 3}$;
- (6') n = 8, $Y \cong \mathbb{Q}^5$ and \mathcal{E} is either $\mathcal{O}_{\mathbb{Q}^5}(1)^{\oplus 4}$ or $\mathcal{O}_{\mathbb{Q}^5}(2) \oplus \mathcal{O}_{\mathbb{Q}^5}(1)^{\oplus 3}$, where \mathbb{Q}^5 is a smooth quadric hypersurface of \mathbb{P}^6 ;
- (7') n = 8, Y is a Del Pezzo 5-fold with $b_2(Y) = 1$, i.e. $\operatorname{Pic}(Y)$ is generated by an ample line bundle $\mathcal{O}_Y(1)$ such that $-K_Y \simeq \mathcal{O}_Y(4)$ and $\mathcal{E} \cong \mathcal{O}_Y(1)^{\oplus 4}$;
- (8') n = 8, there is a vector bundle \mathcal{V} on a smooth curve C such that $Y \cong \mathbb{P}_C(\mathcal{V})$ and $\mathcal{E}|_F \cong \mathcal{O}_F(1)^{\oplus 4}$ for any fibre $F \cong \mathbb{P}^4$ of $Y \to C$;
- (9') n = 8, there is a surjective morphism $q: Y \to \Gamma$ onto a smooth curve Γ such that any general fibre F of q is a smooth quadric hypersurface \mathbb{Q}^4 in \mathbb{P}^5 with $\mathcal{E}|_F \cong \mathcal{O}_F(1)^{\oplus 4}$;
- (10') n = 8, there is a vector bundle \mathcal{V} on a smooth surface S such that $Y \cong \mathbb{P}_S(\mathcal{V})$ and $\mathcal{E}|_F \cong \mathcal{O}_F(1)^{\oplus 4}$ for any fibre $F \cong \mathbb{P}^3$ of $Y \to S$;
- (11') n = 8, there exists a smooth projective 5-fold W and a morphism $\pi : Y \to W$ expressing Y as blown up at a finite set B of points and an ample vector bundle \mathcal{E}' on W such that $\mathcal{E} = \pi^* \mathcal{E}' \otimes [-\pi^{-1}(B)]$ and $K_W + \det \mathcal{E}'$ is ample; or
- (12') n = 7 and (Y, \mathcal{E}) is as in Proposition 4.3.

Proof. If $n \ge 11$, then $k = n - 5 \ge 6 > \dim Y$, so (X, L) is an adjunction scroll over Y by [7, (2.1.1)]. If $7 \le n \le 10$, then we conclude by [7, (2.1.2), (2.1.3)], [36, proposition 3.1] and Proposition 4.3. \Box

Finally, as to the case n = 6 and m = 5, under the extra assumption that $\operatorname{Pic}(Y) \cong \mathbb{Z}[\mathcal{O}_Y(1)]$, we can easily deduce from Theorem 1.1 the following

Corollary 4.2. Let X be a smooth 6-fold and L an ample line bundle on X. Assume that $(X, L) \cong (\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1))$ is a \mathbb{P}^1 -bundle, $\pi : X \to Y$, over a smooth 5-fold $Y, \mathcal{E} = \pi_* L$. If $\operatorname{Pic}(Y) \cong \mathbb{Z}[\mathcal{O}_Y(1)]$, then (X, L) is an adjunction-theoretic scroll over Y under π except when:

- (1") $Y \cong \mathbb{P}^5$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^5}(a_1) \oplus \mathcal{O}_{\mathbb{P}^5}(a_2)$ with $a_1 + a_2 \leq 5$ and $a_i > 0$ for i = 1, 2;
- (2") $Y \cong \mathbb{Q}^5$ and $\mathcal{E} \cong \mathcal{O}_{\mathbb{Q}^5}(a_1) \oplus \mathcal{O}_{\mathbb{Q}^5}(a_2)$, with $a_1 + a_2 \leq 4$ and $a_i > 0$ for i = 1, 2, where \mathbb{Q}^5 is a smooth quadric hypersurface of \mathbb{P}^6 ;
- (3") Y is a Del Pezzo 5-fold, i.e. $-K_Y \simeq \mathcal{O}_Y(4)$, and either (α) $\mathcal{E} \cong \mathcal{O}_Y(1)^{\oplus 2}$, or (β) $\mathcal{E}|_l \cong \mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1}(2)$ for any line $l \cong \mathbb{P}^1$ of $(Y, \mathcal{O}_Y(1))$;
- (4") Y is a Mukai 5-fold, i.e. $-K_Y \simeq \mathcal{O}_Y(3)$, and $\mathcal{E} \cong \mathcal{O}_Y(1)^{\oplus 2}$;
- (5") Y is a Fano 5-fold such that $-K_Y \simeq \det \mathcal{E}$ and X is a Fano 6-fold such that $-K_X \simeq 2L$.

Proof. Since $\operatorname{Pic}(Y) \cong \mathbb{Z}[\mathcal{O}_Y(1)]$, if (X, L) is not an adjunction-theoretic scroll over Y under π , then from the formula (8) we deduce that $K_Y + \det \mathcal{E} = \mathcal{O}_Y(a)$ for some integer $a \leq 0$. Therefore, we have either a = 0, obtaining Case (5") of the statement, or a < 0. In the latter case, since $K_Y + \det \mathcal{E}$ is not nef, we conclude by Theorem 1.1. \Box

Remark 4.1. If (X, L) and (Y, \mathcal{E}) are as in Corollary 4.2, then similar techniques and arguments used for the proofs of [7, theorem 3.1] and [36, proposition 3.4] work also in the case $\operatorname{Pic}(Y) \not\cong \mathbb{Z}[\mathcal{O}_Y(1)]$. More precisely, we can say in this situation when (X, L) is not an adjunction-theoretic scroll over Y under the projection map $\pi : X \to Y$, but the final classification result appears at present not yet complete in some of its parts. This fact is due especially to the lack of classification results of pairs (Y, \mathcal{E}) as above with K_Y + det \mathcal{E} trivial, and of properties on the third reduction of (Y, \mathcal{E}) in the same spirit of [3, remark 4]. We refer to [3, §3] and [26] for the case $m - 2 \geq 2$ and the second reduction of the pair (Y, \mathcal{E}) .

Acknowledgement. I would thank Proff. M.C. Beltrametti, A. Lanteri and A. Laface for their kind comments and some useful remarks about the final form of this paper. Finally, the author wants to thank for some funds supporting this research from the University of Milan (FIRST 2006 and 2007).

References

- V. Ancona, T. Peternell, J.A. Wiśniewski. Fano bundles and splitting theorems on projective spaces and quadrics. *Pacific Journal of Math.* 163 (1994), no. 1, 17–42.
- [2] M. Andreatta, M. Mella. Contractions on a manifold polarized by an ample vector bundle. Trans. Amer. Math. Soc. 349 (1997), no. 11, 4669–4683.

- [3] M. Andreatta, C. Novelli. Manifolds polarized by vector bundles. Ann. Mat. Pura Appl. (4) 186 (2007), no. 2, 281–288.
- [4] M. Andreatta, G. Occhetta. Ample vector bundles with sections vanishing on special varieties. *Internat. J. Math.* 10 (1999), no. 6, 677–696.
- [5] M. Andreatta, J.A. Wiśniewski. On manifolds whose tangent bundle contains an ample subbundle. *Invent. Math.* 146 (2001), no. 1, 209–217.
- [6] M.C. Beltrametti, A.J. Sommese. The Adjunction Theory of Complex Projective Varieties. Expositions in Mathematics, vol. 16, W. de Gruyter, Berlin–New York (1995).
- [7] M.C. Beltrametti, A.J. Sommese. Comparing the classical and the adjunction theoretic definition of scrolls, Geometry of complex projective varieties (Cetraro, 1990), 55–74.
- [8] M.C. Beltrametti, A.J. Sommese, J.A. Wiśniewski. Results on varieties with many lines and their applications to adjunction theory (with an appendix by M.C. Beltrametti and A.J. Sommese), in Complex Algebraic Varieties, Bayreuth 1990, ed. by K. Hulek, T. Peternell, M. Schneider, and F.-O. Schreyer, *Lecture Notes in Math.* **1507** (1992), 16–38.
- [9] K. Cho, Y. Miyaoka, N.I. Shepherd-Barron. Characterizations of projective space and applications to complex symplectic manifolds. Higher dimensional birational geometry (Kyoto, 1997), Math. Soc. Japan, Tokyo, Adv. Stud. Pure Math. 35 (2002), 1–88.
- [10] T. Fujita. On polarized manifolds whose adjoint bundles are not semipositive. Algebraic Geometry, Sendai, 1985, Adv. Stud. Pure Math. 10 (1987), 167–178.
- [11] T. Fujita. On adjoint bundles of ample vector bundles. In Proc. Alg. Geom. Conf. Bayreuth, Lect. Notes Math. 1507 (1990), 105–112.
- [12] T. Fujita. Classification Theories of Polarized Varieties, London Math. Soc. Lecture Notes Series 155, Cambridge Univ. Press, Cambridge (1990)
- [13] P. Ionescu. Generalized adjunction and applications. Math. Proc. Cambridge Philos. Soc. 99 (1986), no. 3, 457–472.
- [14] Y. Kachi, E. Sato. Segre's reflexivity and an inductive characterization of hyperquadrics. Mem. Amer. Math. Soc. 160 (2002), n. 763.
- [15] A. Lanteri, H. Maeda. Special varieties in adjunction theory and ample vector bundles. Math. Proc. Cambridge Philos. Soc. 130 (2001), 61–75.
- [16] A. Lanteri, H. Maeda. Ample vector bundle characterizations of projective bundles and quadric fibrations over curves. *Higher Dimensional Complex Varieties*, Trento, 1994, (ed. M. Andreatta and T. Peternell), (Walter de Gruyter, 1996), pp. 247–259.
- [17] H. Maeda. Nefness of adjoint bundles for ample vector bundles. Le Matematiche Vol. L, Fasc. I (1995), 73–82.
- [18] M. Mella. Existence of good divisors on Mukai varieties. J. Algebraic Geom. 8 (1999), no. 2, 197–206.
- [19] Y. Miyaoka. Numerical characterisations of hyperquadrics. Complex analysis in several variables– Memorial Conference of Kiyoshi Oka's Centennial Birthday (Tokyo) (Math. Soc. Japan, Ed.), North-Holland, Adv. Stud. Pure Math. 42 (2004), 209–235.
- [20] Y. Miyaoka. Numerical characterisations of hyperquadrics. The Fano Conference, Univ. Torino, Turin (2004), 559–562.
- [21] S. Mori. Threefolds whose canonical bundles are not numerically effective. Ann. of Math. (2) 116 (1982), no. 1, 133–176.
- [22] S. Mori. Threefolds whose canonical bundles are not numerically effective, in Algebraic Threefolds, Proceedings Varenna, 1981, ed. by A. Conte, *Lecture notes in Math.* 947 (1982), 125–189, Springer-Verlag, New York.
- [23] C. Novelli, G. Occhetta. Ruled Fano fivefolds of index two. Indiana Univ. Math. J. 56 (2007), no. 1, 207–241.

- [24] G. Occhetta. A note on the classification of Fano manifolds of middle index. Manuscripta Math. 117 (2005), no. 1, 43–49.
- [25] G. Occhetta. On some Fano manifolds of large pseudoindex. Manuscripta Math. 104 (2001), 111–121.
- [26] M. Ohno. Classification of generalized polarized manifolds by their nef values. Adv. Geom. 6 (2006), no. 4, 543–599; Addendum to: "Classification of generalized polarized manifolds by their nef values" [Adv. Geom. 6 (2006), no. 4, 543–599], Adv. Geom. 7 (2007), no. 2, 315–316.
- [27] C. Okonek, M. Schneider, H. Spindler. Vector bundles on complex projective spaces. Progress in Mathematics 3 (1980), Birkhuser, Boston, Mass.
- [28] G. Ottaviani. Spinor bundles on quadrics. Trans. Amer. Math. Soc. 307 (1988), no. 1, 301–316.
- [29] T. Peternell. A characterization of \mathbb{P}^n by vector bundles. Math. Z. 205 (1990), no. 3, 487–490.
- [30] T. Peternell. Ample vector bundles on Fano manifolds. Internat. J. Math. 2 (1991), no. 3, 311–322.
- [31] T. Peternell, M. Szurek, J.A. Wiśniewski. Fano manifolds and vector bundles. Math. Ann. 294 (1992), 151–165.
- [32] T. Peternell, M. Szurek, J.A. Wiśniewski. Numerically effective vector bundles with small Chern classes. Complex algebraic varieties (Bayreuth, 1990), *Lecture Notes in Math.* 1507, 145–156, Springer, Berlin, 1992.
- [33] A. Predonzan. Intorno agli S_k giacenti sulla variet intersezione completa di più forme. Atti Accad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Nat. (8) 5 (1948), 238–242.
- [34] A.J. Sommese. On the adjunction theoretic structure of projective varieties. In Complex Analysis and Algebraic Geometry, Proceedings Göttingen 1985, *Lecture Notes in Math.* 1194 (1986), 175–213.
- [35] A.J. Sommese. On the adjunction theoretic structure of projective varieties, Complex Analysis and Algebraic Geometry, Proceedings Göttingen 1985, Springer-Verlag, Lecture Notes in Math. 1194 (1986), 175–213.
- [36] A.L. Tironi. Scrolls over a four dimensional variety. Adv. Geom. 10 (2010), 145–159.
- [37] J.A. Wiśniewski. Length of extremal rays and generalized adjunction. Math. Z. 200 (1989), 409–427.
- [38] J.A. Wiśniewski. On a conjecture of Mukai. Manuscripta Math. 68 (1990), 135–141.
- [39] J.A. Wiśniewski. On Fano manifolds of large index. Manuscripta Math. 70 (1991), 145–152.
- [40] Y.-G. Ye, Q. Zhang. On ample vector bundles whose adjunction bundles are not numerically effective. Duke Math. J. 60 (1990), no. 3, 671–687.
- [41] Q. Zhang. A theorem on the adjoint system for vector bundles. Manuscripta Math. 70 (1991), no. 2, 189–201.

Departamento de Matemática Universidad de Concepción Casilla 160-C, Concepción (Chile)

E-mail address: atironi@udec.cl