

BERTINI THEOREM FOR NORMALITY ON LOCAL RINGS IN MIXED CHARACTERISTIC (APPLICATIONS TO CHARACTERISTIC IDEALS)

TADASHI OCHIAI AND KAZUMA SHIMOMOTO

ABSTRACT. In this article, we prove a strong version of local Bertini theorem for normality on local rings in mixed characteristic. The main result asserts that a generic hyperplane section of a normal, Cohen-Macaulay, and complete local domain of dimension at least 3 is normal. Applications include the study of characteristic ideals attached to torsion modules over normal domains, which is fundamental in the study of Euler system theory, Iwasawa's main conjectures, and the deformation theory of Galois representations.

1. INTRODUCTION

The classical Bertini theorem says that a generic hyperplane section of a smooth complex projective variety is smooth. We would like to call by a "local Bertini theorem" the following problem for a local Noetherian ring $(R, \mathfrak{m}, \mathbf{k})$:

Let \mathbf{P} be a ring-theoretic property (e.g. regular, reduced, normal, seminormal and so on). Then if R is \mathbf{P} and $x \in \mathfrak{m}$ is a non-zero divisor, then is it true that R/xR is \mathbf{P} for a generic choice of x ?

We refer the reader to Definition 2.3 for a more precise formulation of the problem. A local Bertini theorem (in a slightly weak form) was first raised by Grothendieck ([8], Exposé XIII, Conjecture 2.6) and was proved by Flenner [4] and Trivedi [19]. We use the following notation. For an ideal $I \subseteq R$, we denote by $U(I)$ the set of all primes of R which do not contain I , and by $V(I)$ the complement of $U(I)$ in $\text{Spec } R$. For a graded ring R with a graded ideal I , we use the notation $V^+(I)$ and $U^+(I)$ for a closed subset with its complement in $\text{Proj } R$. Before stating our main theorems, let us recall the following result from [4]:

Theorem 1.1 (Flenner-Trivedi). *Let (R, \mathfrak{m}) be a local Noetherian ring and let $I \subseteq \mathfrak{m}$ be an ideal. Assume that Q is a finite subset of $U(I)$. Then there exists an element $x \in I$ such that:*

- (1) $x \notin \mathfrak{p}^{(2)}$ for all $\mathfrak{p} \in U(I)$;

1991 *Mathematics Subject Classification.* 13H10, 13K05, 13N05.

Key words and phrases. Bertini-type theorem, characteristic ideal, differential module, hyperplane section.

The first-named author is partially supported by Grant-in-Aid for Challenging Exploratory Research (24654004) and Sumitomo Foundation.

The second-named author is partially supported by Grant-in-Aid for Young Scientists (B) # 25800028.

(2) $x \notin \mathfrak{p}$ for all $\mathfrak{p} \in Q$;

for $\mathfrak{p}^{(n)} := \mathfrak{p}^n R_{\mathfrak{p}} \cap R$, the n -th symbolic power ideal of \mathfrak{p} .

We make a remark on the second symbolic power of ideals. Let (R, \mathfrak{m}) be a local ring, let $x \in \mathfrak{p}$ be a non-zero divisor and let $x \notin \mathfrak{p}^{(2)}$ for a prime ideal $\mathfrak{p} \subseteq R$. Then $R_{\mathfrak{p}}$ is regular if and only if $R_{\mathfrak{p}}/xR_{\mathfrak{p}}$ is regular. Many ring-theoretic properties such as regular, normal, reduced can be verified at the localization $R_{\mathfrak{p}}$, which is the reason why we require $x \notin \mathfrak{p}^{(2)}$ (but not merely $x \notin \mathfrak{p}^2$), which is equivalent to the condition: $x \notin \mathfrak{p}^2 R_{\mathfrak{p}} \cap R$. A strong version of local Bertini theorem similar to our main theorem below was already proved for local rings containing a field in [4]. To extend it to the mixed characteristic case, we need to introduce some new ideas. Theorem 1.1 has the following implication. If (R, \mathfrak{m}) is a complete local normal domain of depth $R \geq 3$, then there is a non-zero element $x \in \mathfrak{m}$ for which R/xR is normal ([4], Korollar 3.4). However, this does not suffice for our purpose, because we would like to give a family of infinitely many such specializations R/xR which are parametrized by a certain explicit p -adic space.

Here is our notation which will be used throughout the paper. We denote by $\mathbb{P}^n(S)$ (resp. $\mathbb{A}^n(S)$) the set of S -rational points of the n -dimensional projective space (resp. affine space) over an integral domain S . In case when S is a field or a discrete valuation ring, these spaces come equipped with some topology (see discussions in § 2). For a complete discrete valuation ring (A, π_A, \mathbf{k}) , we have a *specialization map* $\text{Sp}_A : \mathbb{P}^n(A) \rightarrow \mathbb{P}^n(\mathbf{k})$ (cf. Definition 2.1).

In § 2 and § 3, we make discussions on basic tools and prove some requisite results, including the specialization map and basic elements, due to Swan.

In § 4, we prove the first main theorem (see Theorem 4.4 together with Theorem 4.3) in this article:

Main Theorem A (Local Bertini Theorem). Let $(R, \mathfrak{m}, \mathbf{k})$ be a complete local domain of mixed characteristic $p > 0$ and suppose the following conditions:

- (1) let $A \rightarrow R$ be a coefficient ring map for a complete discrete valuation ring (A, π_A) ;
- (2) let x_0, x_1, \dots, x_d be a fixed set of minimal generators of \mathfrak{m} ;
- (3) R is normal, of depth $R \geq 3$, and the residue field \mathbf{k} is infinite.

Then there exists a Zariski dense open subset $U \subseteq \mathbb{P}^d(\mathbf{k})$ satisfying the following properties. For any $a = (a_0 : \dots : a_d) \in \text{Sp}_A^{-1}(U)$, the quotient $R/x_{\bar{a}}R$ is a normal domain of mixed characteristic $p > 0$ (cf. Definition 2.3 for the notation $x_{\bar{a}}R$).

By the minimality condition of x_0, x_1, \dots, x_d , it is immediate to see that $x_{\bar{a}} \neq 0$, that is, $R/x_{\bar{a}}R$ is a non-trivial quotient of R . We will also discuss a version of the above theorem for the case when the residue field is finite at the end of § 4. The above theorem allows us to find sufficiently many local normal domains of mixed characteristic as specializations,

but it does not tell us how to find U . We will show how to find U in Example 4.9. It is worth pointing out that if $\dim R = 2$, the local Bertini theorem fails due to a simple reason. In fact, if the quotient R/yR is normal, then it is a discrete valuation ring, so R must be regular. By Cohen's structure theorem, there is a surjection $A[[z_0, \dots, z_d]] \twoheadrightarrow R$, where $d + 1$ is the number of the minimal generators of \mathfrak{m} , so that the minimal generators of \mathfrak{m} are just the image of z_0, z_1, \dots, z_d under this surjection. Related to our main result, we mention that if (R, \mathfrak{m}) is a local ring and $y \in \mathfrak{m}$ is a non-zero divisor such that R/yR is normal, then R is normal ([7] 5.12.7).

In § 5, we establish a fact that the parameter set of specializations in the local Bertini theorem is infinite (see Proposition 5.1), which is used to prove the main result on characteristic ideals.

In § 6, we prove a version of the local Bertini theorem in the case when Serre's condition (R_n) or (S_n) is satisfied.

In § 7, we define characteristic ideals of finitely generated torsion modules over a Noetherian normal domain as reflexive ideals, following the paper [18].

In § 8, the main purpose is to generalize the results proved in [14] over a regular ring to the case over a Noetherian normal domain. First, we prove some preliminary results on characteristic ideals. Then combining Main Theorem A, we prove another main theorem (see Theorem 8.8) as follows:

Main Theorem B (Control Theorem for Characteristic Ideals). Let $(R, \mathfrak{m}, \mathbf{k})$ be a complete local domain of mixed characteristic $p > 0$ satisfying the conditions (1), (2) and (3) of Main Theorem A, except that we now assume the residue field to be finite. Let M and N be finitely generated torsion R -modules. Then for certain infinite subsets $\mathcal{L}_{W(\overline{\mathbb{F}})}(M_{W(\overline{\mathbb{F}})})$ and $\mathcal{L}_{W(\overline{\mathbb{F}})}(N_{W(\overline{\mathbb{F}})})$ of the projective space $\mathbb{P}^d(W(\overline{\mathbb{F}}))$ defined in a natural way (cf. Definition 8.1), the following statements are equivalent:

- (1) $\text{char}_R(M) \subseteq \text{char}_R(N)$, where $\text{char}_R(-)$ denotes the characteristic ideal (cf. Definition 7.1).
- (2) For all but finitely many height-one primes:

$$\mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{W(\overline{\mathbb{F}})}) \cap \mathcal{L}_{W(\overline{\mathbb{F}})}(N_{W(\overline{\mathbb{F}})}),$$

we have

$$\text{char}_{R_{W(\mathbb{F}')} / \mathbf{x}_{\tilde{a}} R_{W(\mathbb{F}')}}(M_{W(\mathbb{F}')} / \mathbf{x}_{\tilde{a}} M_{W(\mathbb{F}')})) \subseteq \text{char}_{R_{W(\mathbb{F}')} / \mathbf{x}_{\tilde{a}} R_{W(\mathbb{F}')}}(N_{W(\mathbb{F}')} / \mathbf{x}_{\tilde{a}} N_{W(\mathbb{F}')})),$$

where \mathbb{F}' is any finite field extension of \mathbb{F} depending on \tilde{a} such that $\mathbf{x}_{\tilde{a}} \in R_{W(\mathbb{F}')}.$

- (3) For all but finitely many height-one primes:

$$\mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{W(\overline{\mathbb{F}})}) \cap \mathcal{L}_{W(\overline{\mathbb{F}})}(N_{W(\overline{\mathbb{F}})}),$$

we have

$$\text{char}_{R_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}}(M_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\bar{a}}M_{W(\overline{\mathbb{F}})}) \subseteq \text{char}_{R_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}}(N_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\bar{a}}N_{W(\overline{\mathbb{F}})}).$$

Main Theorem B will be crucial in a forthcoming paper [15], where we plan to compare characteristic ideals of certain torsion modules arising from Iwasawa theory as developed in [14] (for example, those torsion modules arise as the Pontryagin dual modules of Selmer groups associated with two-dimensional certain Galois representations with values in a complete local ring with finite residue field).

2. SPECIALIZATION MAP AND THE TOPOLOGY FOR BERTINI-TYPE THEOREMS

In this section, we introduce notation and make definitions. We fix a prime integer $p > 0$. First, we discuss the specialization map. Let (A, π_A, \mathbf{k}) be a complete discrete valuation ring and let us choose a point:

$$a = (a_0 : \cdots : a_n) \in \mathbb{P}^n(A).$$

Suppose that the valuation v of π_A is equal to 1 and that the i -th coordinate a_i has the minimal valuation $v_i := v(a_i)$ among the valuations of a_0, \dots, a_n . We put $a'_j = a_j / \pi_A^{v_i}$ ($0 \leq j \leq n$). Then we have a presentation $a = (a'_0 : \cdots : a'_n)$, so that $a_j \in A$ ($0 \leq j \leq n$) and that a'_i is a unit of A . So we may and we will think of the projective space as

$$\mathbb{P}^n(A) = \{\text{homogeneous coordinates } (a_0 : \cdots : a_n) \text{ in } A \mid a_i \in A^\times \text{ for some } i\} / \sim$$

where the equivalence relation \sim is a simultaneous multiplication by a unit of A .

Definition 2.1. Let (A, π_A, \mathbf{k}) be a complete discrete valuation ring and pick a point $a = (a_0 : \cdots : a_n) \in \mathbb{P}^n(A)$.

- (1) We define a *lift* $\tilde{a} = (\tilde{a}_0, \dots, \tilde{a}_n) \in \mathbb{A}^{n+1}(A)$ of $a \in \mathbb{P}^n(A)$ to be an element belonging to the set consisting of all inverse images of $a \in \mathbb{P}^n(A)$ via the projection map:

$$\mathbb{A}^{n+1}(A) \setminus \{\text{points whose coordinates are all non-units}\} \twoheadrightarrow \mathbb{P}^n(A).$$

- (2) Let $\tilde{a} \in \mathbb{A}^{n+1}(A)$ be a lift of $a \in \mathbb{P}^n(A)$. Then \tilde{a} gets mapped to a point $\bar{a} = (\bar{a}_0 : \cdots : \bar{a}_n) \in \mathbb{P}^n(\mathbf{k})$ via the surjection map $A \rightarrow \mathbf{k} := A/\pi_A A$. Thus, we construct a *specialization map*:

$$\text{Sp}_A : \mathbb{P}^n(A) \rightarrow \mathbb{P}^n(\mathbf{k})$$

by setting $\text{Sp}_A(a) = \bar{a}$. This map does not depend on the choice of the lift of a .

The set $\mathbb{P}^n(\mathbf{k})$ is endowed with the Zariski topology, while $\mathbb{P}^n(A)$ is endowed with the topology induced by the valuation on A . Hence we simply regard $\mathbb{P}^n(A)$ as a set of

points equipped with this topology. It is also straightforward from the definition that we naturally identify the set $\mathbb{P}^n(A)$ with $\mathbb{P}^n(\text{Frac}(A))$ as well as the induced topology.

We begin to pin down the suitable topology for formulating Bertini-type theorems in mixed characteristic. Let $(R, \mathfrak{m}, \mathbf{k})$ be a local Noetherian ring. Then we say that a reduced local ring $(R, \mathfrak{m}, \mathbf{k})$ is of *mixed characteristic* $p > 0$, if every component of the total ring of fractions of R is of characteristic zero and the residue field \mathbf{k} is of characteristic p .

Now we assume that $(R, \mathfrak{m}, \mathbf{k})$ is a complete reduced local ring of mixed characteristic $p > 0$ and (A, π_A, \mathbf{k}) is a complete discrete valuation ring such that $\pi_A A = pA$ and there is an injection $A \hookrightarrow R$ of rings, which induces an isomorphism on residue fields, say $\mathbf{k} = A/\pi_A A \simeq R/\mathfrak{m}$. In this situation, we call (A, π_A, \mathbf{k}) together with a map $A \hookrightarrow R$ a *coefficient ring* of R .

Example 2.2. Let $R := \mathbb{Z}_p[[x, y]]/(p - xy)$. Then R is a finite extension of $\mathbb{Z}_p[[x + y]]$ defined by the Eisenstein equation $t^2 - (x + y)t + p = 0$ and \mathbb{Z}_p is a coefficient ring of R .

In what follows, we will fix a coefficient ring (A, π_A, \mathbf{k}) . We denote by $\mathbf{Loc.alg}_{/A}$ the category of complete local A -algebras for a discrete valuation ring A . Note that the category of complete local \mathbf{k} -algebras is a full subcategory of $\mathbf{Loc.alg}_{/A}$. Let $(R, \mathfrak{m}, \mathbf{k})$ be a local Noetherian ring and let M be a finitely generated R -module. We say that a set of elements x_0, \dots, x_d of M is the *minimal generators* of M , if the following conditions hold:

$$\sum_{i=0}^d \mathbf{k} \cdot \bar{x}_i = M/\mathfrak{m}M \text{ and } \dim_{\mathbf{k}} M/\mathfrak{m}M = d + 1,$$

where \bar{x}_i denotes the image of x_i in $M/\mathfrak{m}M$.

Definition 2.3. Let $(R, \mathfrak{m}, \mathbf{k}) \in \mathbf{Loc.alg}_{/A}$ be a reduced local ring of mixed characteristic $p > 0$.

- (1) Fix a set of minimal generators $x_0, \dots, x_n \in \mathfrak{m}$ and let $a = (a_0 : \dots : a_n) \in \mathbb{P}^n(A)$. For any lift $\tilde{a} = (\tilde{a}_0, \dots, \tilde{a}_n) \in \mathbb{A}^{n+1}(A)$ of a in the sense of Definition 2.1, we put

$$\mathbf{x}_{\tilde{a}} := \sum_{i=0}^n \tilde{a}_i x_i.$$

By definition, the element $\mathbf{x}_{\tilde{a}} \in R$ depends on the choice of a lift \tilde{a} , but the ideal $\mathbf{x}_{\tilde{a}}R$ depends only on $a \in \mathbb{P}^n(A)$.

- (2) With the notation as above, assume that \mathbf{P} is a ring-theoretic property on Noetherian rings. We say that $(R, \mathfrak{m}, \mathbf{k}) \in \mathbf{Loc.alg}_{/A}$ satisfies a *local Bertini theorem* for the property \mathbf{P} if for a fixed set of minimal generators $x_0, \dots, x_n \in \mathfrak{m}$, there exists a Zariski (dense) open subset $U \subseteq \mathbb{P}^n(\mathbf{k})$ such that $R/\mathbf{x}_{\tilde{a}}R$ has \mathbf{P} for all $a = (a_0 : \dots : a_n) \in \text{Sp}_A^{-1}(U) \subseteq \mathbb{P}^n(A)$.

One may formulate the local Bertini theorem in a different way. For example, the completeness of R may be dispensed with, or a set of minimal generators of the maximal ideal of R may be replaced with a set of minimal generators of a smaller ideal. In this article, we adopt the above definition.

Remark 2.4. The naturality of the above definition is explained as follows. We endow $\mathbb{P}^n(\mathbf{k})$ with the Zariski topology. Let $\bar{f} \in \mathbf{k}[x_0, \dots, x_n]$ be a non-zero homogeneous polynomial, let $f \in A[x_0, \dots, x_n]$ be any fixed homogeneous lifting of \bar{f} , and let $U^+(\bar{f}) \subseteq \mathbb{P}^n(\mathbf{k})$ be an open subset defined by $\bar{f} \neq 0$. Then the inverse image of the open subset $U^+(\bar{f})$ under the map $\mathrm{Sp}_A : \mathbb{P}^n(A) \rightarrow \mathbb{P}^n(\mathbf{k})$ can be described as follows:

$$\mathrm{Sp}_A^{-1}(U^+(\bar{f})) = \{a = (a_0 : \dots : a_n) \in \mathbb{P}^n(A) \mid f(a) \in A^\times\}.$$

This is an admissible open subset of $\mathbb{P}^n(A)$ ([5] for this fact). Our objective is to show that this topology is suitable in formulating the local Bertini theorem in the mixed characteristic case.

The following proposition is indispensable for the proof of Theorem 4.3 and Theorem 4.4.

Proposition 2.5. *Let $U \subseteq \mathbb{A}^n(L)$ be any non-empty Zariski open subset for an infinite field L . Then U is dense. Furthermore, if $K \subseteq L$ is any subfield of L such that K is infinite, the intersection $U \cap \mathbb{A}^n(K)$ is also a Zariski dense open subset of $\mathbb{A}^n(K)$. The above assertions hold over the projective space as well.*

Proof. The first assertion about density is obvious. So we prove the second assertion. Then it suffices to prove the following statement:

- Let $V := V(I)$ be a Zariski closed subset of $\mathbb{A}_L^n := \mathrm{Spec} L[X_1, \dots, X_n]$, where I is an ideal of $L[X_1, \dots, X_n]$. If $\mathbb{A}^n(K)$ is contained in the set of K -rational points of V , then V is equal to \mathbb{A}_L^n .

Let us prove this. If $\mathbb{A}^n(K)$ is contained in the set of K -rational points of V , we have

$$I \subseteq \bigcap_{(a_1, \dots, a_n) \in K^n} (X_1 - a_1, X_2 - a_2, \dots, X_n - a_n) \subseteq L[X_1, \dots, X_n].$$

Since K is infinite, we have

$$\bigcap_{(a_1, \dots, a_n) \in K^n} (X_1 - a_1, X_2 - a_2, \dots, X_n - a_n) = 0$$

and thus $I = (0)$. This implies that $V(I) = \mathbb{A}_L^n$. □

3. DISCUSSION ON BASIC ELEMENTS

We start with the definition of basic elements.

Definition 3.1. Let R be a Noetherian ring and let M be a finitely generated R -module.

- (1) Assume that (R, \mathfrak{m}) is local. The depth of M , denoted by $\text{depth}_R M$, is defined to be the maximal length of all M -regular sequences contained in \mathfrak{m} .
- (2) Let $\mathfrak{p} \in \text{Spec } R$. Then $\mu_{\mathfrak{p}}(M)$ denotes the number of minimal generators of the $R_{\mathfrak{p}}$ -module $M_{\mathfrak{p}}$.
- (3) (Swan) Let $\mathfrak{p} \in \text{Spec } R$. We say that an element $m \in M$ is *basic* at \mathfrak{p} , if we have

$$\mu_{\mathfrak{p}}(M) - \mu_{\mathfrak{p}}(M/(R \cdot m)) = 1.$$

More generally, we say that a set of elements m_1, \dots, m_n of M is *k-fold basic* at \mathfrak{p} , if the following inequality holds:

$$\mu_{\mathfrak{p}}(M) - \mu_{\mathfrak{p}}(M/(\sum_{i=1}^n R \cdot m_i)) \geq k,$$

that is, $N = R \cdot m_1 + \dots + R \cdot m_n$ contains at least k minimal generators at \mathfrak{p} .

Remark 3.2. Under the setting of the above definition, let $M^{(r)} := M/(\sum_{i=1}^r R \cdot m_i)$ for a set of elements m_1, \dots, m_k of M and r satisfying $0 \leq r \leq k-1$ and pick a prime ideal \mathfrak{p} of R . Then

$$\mu_{\mathfrak{p}}(M^{(r)}) - \mu_{\mathfrak{p}}(M^{(r)}/(R \cdot m_{r+1})) = 1 \iff m_{r+1} \notin \mathfrak{p}M_{\mathfrak{p}}^{(r)}$$

for $0 \leq r \leq k-1$ by Nakayama's lemma. In other words, m_1, \dots, m_k form partial generators of the $k(\mathfrak{p})$ -vector space $M \otimes_R k(\mathfrak{p})$ with $k(\mathfrak{p}) := R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}$.

We shall use (finite) Kähler differentials ([10] as a reference). For a complete local ring (R, \mathfrak{m}) with its coefficient ring A , the usual module of Kähler differentials $\Omega_{R/A}$ is not a finitely generated R -module. Instead, one uses the completed module $\widehat{\Omega}_{R/A}$. This is the \mathfrak{m} -adic completion of $\Omega_{R/A}$ and it is a finitely generated R -module. It can be also defined as follows. Let I denote the kernel of the map $\mu : R \widehat{\otimes}_A R \rightarrow R$ defined by $\mu(a \otimes b) = ab$. Then $\widehat{\Omega}_{R/A} := I/I^2$. Let $d : R \rightarrow \widehat{\Omega}_{R/A}$ be the canonical derivation defined by $a \mapsto a \otimes 1 - 1 \otimes a$. The connection of Kähler differential modules with the symbolic power ideals is expressed by the following simple fact ([4], Lemma 2.2 for its proof).

Lemma 3.3. Let M be a module over a ring R , let \mathfrak{p} be a prime of R , and let $d : R \rightarrow M$ be a derivation. If for $x \in R$, $dx \in M$ is basic at \mathfrak{p} , then $x \notin \mathfrak{p}^{(2)}$.

Let $I \subseteq R$ be an ideal. We denote by $\text{Min}_R(I)$ the set of all prime ideals that are minimal over I . The authors are grateful to Prof. V. Trivedi for explaining the proof of the following lemma ([4], Lemma 1.2).

Lemma 3.4 (Flenner). *Suppose that R is a Noetherian ring, M is a finitely generated R -module, $U \subseteq \text{Spec } R$ is a Zariski open subset, and $\{m_0, \dots, m_n\}$ is a set of elements of M , which generates the submodule $N \subseteq M$. Suppose that we have $t \in \mathbb{Z}$ (which can be negative) such that*

$$\mu_{\mathfrak{p}}(M) - \mu_{\mathfrak{p}}(M/N) \geq \dim(V(\mathfrak{p}) \cap U) - t$$

for every $\mathfrak{p} \in U$. Let $(\phi^*)^{-1}(U)$ be the inverse image of U under $\phi^* : \text{Spec } R[X_0, \dots, X_n] \rightarrow \text{Spec } R$ induced by the natural injection $\phi : R \rightarrow R[X_0, \dots, X_n]$. Then there exists an ideal $(F_1, \dots, F_r) \subseteq R[X_0, \dots, X_n]$ such that

$$\dim(V(F_1, \dots, F_r) \cap (\phi^*)^{-1}(U)) \leq n + 1 + t$$

and the element

$$\sum_{i=0}^n m_i \otimes X_i \in M \otimes_R R[X_0, \dots, X_n]$$

is basic on $U(F_1, \dots, F_r) \cap (\phi^*)^{-1}(U)$.

We need the following technical lemma for the proof of the main theorem. We recall that a local domain S is *catenary*, if and only if $\text{ht } \mathfrak{p} + \dim S/\mathfrak{p} = \dim S$ for all $\mathfrak{p} \in \text{Spec } S$.

Lemma 3.5. *Let (R, \mathfrak{m}) be an excellent local domain and let $\phi : R \rightarrow R[X_0, \dots, X_d]$ be a natural injection with an ideal $(F_1, \dots, F_r) \subseteq R[X_0, \dots, X_d]$. Suppose that the following conditions hold:*

- (1) $\dim(V(F_1, \dots, F_r) \cap (\phi^*)^{-1}(U(\mathfrak{m}))) \leq d + 1$.
- (2) $(F_1, \dots, F_r) \subseteq \mathfrak{m}R[X_0, \dots, X_d]$.

Then the set of prime ideals $P \in \text{Spec } R[X_0, \dots, X_d]$ satisfying the property:

$$(F_1, \dots, F_r) \subseteq P \subseteq \mathfrak{m}R[X_0, \dots, X_d]$$

is finite.

Proof. Let $I := (F_1, \dots, F_r)$ and consider the finite set:

$$S := \{P \in \text{Min}_{R[X_0, \dots, X_d]}(I) \mid P \subseteq \mathfrak{m}R[X_0, \dots, X_d]\}.$$

We would like to show that

$$(3.1) \quad 0 \leq \text{ht}(\mathfrak{m}R[X_0, \dots, X_d]) - \text{ht } P \leq 1$$

for $P \in S$. If (3.1) is true for $P \in S$, then it is immediate that a prime ideal between P and $\mathfrak{m}R[X_0, \dots, X_d]$ is either P or $\mathfrak{m}R[X_0, \dots, X_d]$. Hence if (3.1) is proved for all $P \in S$, the lemma follows. So we will prove (3.1) for all $P \in S$.

Since R is an excellent local domain, $R[X_0, \dots, X_d]$ is a catenary domain. From this it follows that $0 \leq \text{ht}(\mathfrak{m}R[X_0, \dots, X_d]) - \text{ht } P$ for $P \in S$. If we have $\text{ht}(\mathfrak{m}R[X_0, \dots, X_d]) = \text{ht } P$, then $\mathfrak{m}R[X_0, \dots, X_d] = P$. So it suffices to establish the inequality:

$$\text{ht}(\mathfrak{m}R[X_0, \dots, X_d]) - \text{ht } P \leq 1.$$

Let T be the localization of $R[X_0, \dots, X_d]$ at the prime ideal $\mathfrak{m}R[X_0, \dots, X_d]$. Then $R \rightarrow T$ is a flat local map of local domains and $\dim R = \dim T$. Now assume that $P \neq \mathfrak{m}R[X_0, \dots, X_d]$ for $P \in S$. Since R is an excellent local domain, T is a catenary local domain and we have $\text{ht } P = \dim T - \dim T/PT$. So if we can prove

$$\dim T/PT = 1,$$

the relation (3.1) follows from this and we are done. Let

$$(3.2) \quad P_0 := P \subsetneq P_1 \subsetneq \cdots \subsetneq P_s = \mathfrak{m}R[X_0, \dots, X_d]$$

be a chain of prime ideals of maximal length. Since $\dim T/PT = s$, it suffices to prove $s = 1$. Consider a chain of prime ideals ($0 \leq k \leq s$) in $R[X_0, \dots, X_d]$:

$$(3.3) \quad P_0 \subsetneq \cdots \subsetneq P_k \subsetneq P_k + (X_0) \subsetneq P_k + (X_0, X_1) \subsetneq \cdots \subsetneq P_k + (X_0, \dots, X_d).$$

On the other hand, $\dim(V(F_1, \dots, F_r) \cap (\phi^*)^{-1}(U(\mathfrak{m})))$ is equal to the length of the chain of prime ideals of maximal length in $R[X_0, \dots, X_d]$:

$$(3.4) \quad Q_0 \subsetneq Q_1 \subsetneq \cdots \subsetneq Q_t \text{ such that } I \subseteq Q_0 \text{ and } \phi^{-1}(Q_t) \neq \mathfrak{m}.$$

First, note that $s = 0$ is impossible, since we assumed $P \neq \mathfrak{m}R[X_0, \dots, X_d]$. So in the rest of the proof, let us assume $s \geq 2$ and derive a contradiction. In this case, taking $k = 1$ in (3.3), we get a chain of prime ideals of length $d + 2$:

$$(3.5) \quad P_0 \subsetneq P_1 \subsetneq P_1 + (X_0) \subsetneq P_1 + (X_0, X_1) \subsetneq \cdots \subsetneq P_1 + (X_0, \dots, X_d).$$

Since $P_1 \subsetneq P_2 \subseteq \mathfrak{m}R[X_0, \dots, X_d]$ by $s \geq 2$, we have $\phi^{-1}(P_1 + (X_0, \dots, X_d)) \neq \mathfrak{m}$. By comparing (3.4) and (3.5), we must have $d + 2 \leq t$, because (3.4) is of maximal length. However, this is not compatible with the condition (1). Therefore, we get $s = 1$ as desired. \square

In this paper, the Noetherian induction and the lemma of generic freeness are important tools. For this, we often need the following fact and use it freely.

Lemma 3.6. *Let R be a Noetherian ring and let M be a finitely generated R -module. Let I be an ideal of R and let \mathfrak{p} be a prime ideal of R satisfying $I \subseteq \mathfrak{p}$. We denote the quotient ring R/I by \overline{R} and the image of \mathfrak{p} in \overline{R} by $\overline{\mathfrak{p}}$. Then an element $x \in M$ is basic at $\mathfrak{p} \in \text{Spec } R$ if and only if $\overline{x} \in \overline{M} := M/IM$ is basic at $\overline{\mathfrak{p}} \in \text{Spec } \overline{R}$.*

Proof. Note that there is a commutative square:

$$\begin{array}{ccc} M & \longrightarrow & \overline{M} \\ \pi \downarrow & & \overline{\pi} \downarrow \\ M \otimes_R k(\mathfrak{p}) & \xlongequal{\quad} & \overline{M} \otimes_{\overline{R}} k(\overline{\mathfrak{p}}) \end{array}$$

In this diagram, we have $\pi(x) = \overline{\pi(\overline{x})}$. By Remark 3.2, $x \in M$ (resp. $\overline{x} \in \overline{M}$) is basic at $\mathfrak{p} \in \text{Spec } R$ (resp. $\overline{\mathfrak{p}} \in \text{Spec } \overline{R}$) if and only if $\pi(x) \in M \otimes_R k(\mathfrak{p})$ (resp. $\overline{\pi(\overline{x})} \in \overline{M} \otimes_{\overline{R}} k(\overline{\mathfrak{p}})$) is not zero. This completes the proof. \square

Let R be a Noetherian ring and let M be a finitely generated R -module. Fix an element $x \in M$. Let us define

$$Z_x := \{\mathfrak{p} \in \text{Spec } R \mid x \text{ is basic at } \mathfrak{p}\}.$$

Then Z_x is not necessarily a Zariski open subset. However, we have the following result.

Proposition 3.7. *Let R be a Noetherian ring and suppose M is a finitely generated R -module. Fix an element $x \in M$ and let Z_x be as above. Then Z_x is a (possibly empty) constructible subset of $\text{Spec } R$.*

Proof. First we consider the special case where M is finitely generated and projective over R . Recall that for a finitely generated projective R -module N , $x \in N$ is basic at $\mathfrak{p} \in \text{Spec } R$ if and only if $R_{\mathfrak{p}} \cdot x$ spans a direct summand of $N_{\mathfrak{p}}$ (see [2], Lemma 1 for the proof). Note that, for any short exact sequence of R -modules, the locus on which it splits is a Zariski open subset of $\text{Spec } R$. Hence, we proved that Z_x is Zariski open when M is finitely generated and projective over R .

Then we will prove the case for general R -modules M . First, we apply Lemma 3.6 by taking I to be the nil-radical $\sqrt{0}$ of R and by taking \overline{R} to be $R_{\text{red}} := R/\sqrt{0}$. Since $R \rightarrow R_{\text{red}}$ induces an isomorphism of topological spaces $\text{Spec } R_{\text{red}} \xrightarrow{\sim} \text{Spec } R$, we may and we will assume that R is reduced from now on. Thus there exists $f \in R$ for which $M[f^{-1}]$ is a free $R[f^{-1}]$ -module. Denote by $U \subseteq \text{Spec } R[f^{-1}]$ the locus on which $x \in M[f^{-1}]$ is basic.

Continuing this process, we may find a chain of closed subsets:

$$V(f) =: V_1 \supseteq V_2 \supseteq V_3 \supseteq \cdots$$

such that $x \in M$ is basic at $\mathfrak{p} \in V_i$ if and only if $\mathfrak{p} \in Z_i := V_i \setminus V_{i-1} \subseteq V_i$. Moreover, Z_i is an open subset of V_i . Since $\text{Spec } R$ is a Noetherian space, this chain stabilizes. Thus, there is an integer $N > 0$ such that $V_N = V_{N+1} = \cdots$. We set $Z := Z_1 \cup Z_2 \cup \cdots \cup Z_N$. Then it follows that $x \in M$ is basic at each point of Z and Z is a constructible subset of $\text{Spec } R$. So $Z_x := U \cup Z$ is the sought one and constructible. \square

4. MAIN THEOREMS

In this section, we establish our main theorems.

Lemma 4.1. *Let (A, π_A, \mathbf{k}) be a discrete valuation ring and let $f \in A[y_1, \dots, y_d]$ be a non-zero (possibly constant) polynomial. Then there exists $t \in \mathbb{Z}_{\geq 0}$ such that $\pi_A^{-t} f \in A[y_1, \dots, y_d]$ and the reduction of $\pi_A^{-t} f$ modulo π_A is a non-zero (possibly constant) polynomial in $\mathbf{k}[y_1, \dots, y_d]$.*

Proof. The proof goes by induction on d . If $d = 1$, we write $f = a_m y_1^m + a_{m-1} y_1^{m-1} + \cdots + a_0$ with $a_i \in A$. Let $0 \leq h \leq m$ be such that the valuation $v(a_h)$ is the smallest in the set $\{v(a_i) \mid 0 \leq i \leq m\}$ and write $a_h = (\text{unit}) \cdot \pi_A^t$. Dividing f by π_A^t , we get the desired polynomial.

In general, write $f = b_n y_d^n + b_{n-1} y_d^{n-1} + \cdots + b_0$ for $b_i \in A[y_1, \dots, y_{d-1}]$. Applying the induction hypothesis to every b_i , we may find $t_i \in \mathbb{Z}_{\geq 0}$ such that $\pi_A^{-t_i} b_i$ has the desired property. Let $t_s := \min\{t_i \mid 0 \leq i \leq n\}$. Then the term $\pi_A^{-t_s} b_s y_d^s$ modulo π_A is non-zero and it is clear that $\pi_A^{-t_s} f$ is contained in $A[y_1, \dots, y_d]$. Hence it suffices to put $t := t_s$. \square

The following lemma will play a role in the final step of the proof of the next theorem.

Lemma 4.2. *Let $(R, \mathfrak{m}, \mathbf{k})$ be a complete local domain of mixed characteristic $p > 0$ with infinite residue field \mathbf{k} and a coefficient ring (A, π_A) . Fix a set of minimal generators x_0, x_1, \dots, x_d of \mathfrak{m} together with a prime ideal \mathfrak{p} of R with $\mathfrak{p} \neq \mathfrak{m}$. Then there exists a non-empty open subset $U \subseteq \mathbb{P}^d(\mathbf{k})$ such that*

$$\mathbf{x}_{\tilde{a}} := \sum_{i=0}^d \tilde{a}_i x_i \notin \mathfrak{p}$$

for every $a = (a_0 : \cdots : a_d) \in \text{Sp}_A^{-1}(U)$.

Proof. Consider the homogeneous polynomial:

$$F(X_0, \dots, X_d) := \sum_{i=0}^d x_i X_i \in R[X_0, \dots, X_d].$$

Then we have an equivalence of conditions:

$$\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i \notin \mathfrak{p} \iff F(\tilde{a}_0, \dots, \tilde{a}_d) \not\equiv 0 \pmod{\mathfrak{p}}.$$

We will divide the proof of the lemma into the separate cases.

Case1: Assume that $\pi_A \in \mathfrak{p}$. Let S be the localization of R at \mathfrak{p} . Then $A \rightarrow S$ is a flat local map of local rings and S is of mixed characteristic. Let \mathbf{k}' denote the residue field of S . Then we have a mapping:

$$\mathbb{P}^d(\mathbf{k}) \rightarrow \mathbb{P}^d(\mathbf{k}').$$

Since F is a homogeneous polynomial, the condition $F(\tilde{a}_0, \dots, \tilde{a}_d) \not\equiv 0 \pmod{\mathfrak{p}}$ defines an open subset $V \subseteq \mathbb{P}^d(\mathbf{k}')$. By Proposition 2.5,

$$U := V \cap \mathbb{P}^d(\mathbf{k})$$

is a dense open subset of $\mathbb{P}^d(\mathbf{k})$, since the field \mathbf{k} is infinite. Then this U is a non-empty open subset of $\mathbb{P}^d(\mathbf{k})$ with the desired property.

Case2: Assume that $\pi_A \notin \mathfrak{p}$. In this case, notation being as in **Case1**, we have a mapping:

$$\mathbb{P}^d(\text{Frac}(A)) \rightarrow \mathbb{P}^d(\mathbf{k}').$$

Again, the condition that $F(\tilde{a}_0, \dots, \tilde{a}_d) \not\equiv 0 \pmod{\mathfrak{p}}$ defines a Zariski open subset $V \subseteq \mathbb{P}^d(\mathbf{k}')$. We use an identification $\mathbb{P}^d(\text{Frac}(A)) = \mathbb{P}^d(A)$.

Proposition 2.5 implies that $V \cap \mathbb{P}^d(\text{Frac}(A))$ is a dense open subset of $\mathbb{P}^d(\text{Frac}(A))$, and it is covered by open sets of the form $U^+(f)$ for a homogeneous polynomial $f \in A[X_0, \dots, X_d]$. Let $\bar{f} \in \mathbf{k}[X_0, \dots, X_d]$ be its reduction modulo π_A and let $U^+(\bar{f}) \subseteq \mathbb{P}^d(\mathbf{k})$ be the corresponding open subset. In view of Lemma 4.1, by replacing f with $\pi_A^{-t}f$ if necessary, we may assume that $f \in A[X_0, \dots, X_d]$ and $\bar{f} \in \mathbf{k}[X_0, \dots, X_d]$ is not equal to 0. Since $f(\tilde{a}) \in A^\times$ if and only if $\bar{f}(\text{Sp}_A(a)) \neq 0$, we have

$$\text{Sp}_A^{-1}(U^+(\bar{f})) = \{a = (a_0 : \dots : a_d) \in \mathbb{P}^d(A) \mid f(\tilde{a}) \in A^\times\} \subseteq \mathbb{P}^d(A).$$

Applying this description to the open covering of $V \cap \mathbb{P}^d(\text{Frac}(A))$ consisting of $U^+(f)$, the image

$$U := \text{Sp}_A(V \cap \mathbb{P}^d(\text{Frac}(A))) \subseteq \mathbb{P}^d(\mathbf{k})$$

is a Zariski open subset. Then this U is a non-empty open subset of $\mathbb{P}^d(\mathbf{k})$ with the desired property.

Combining **Case1** and **Case2** together, we finish the proof of the lemma. \square

The following is our first main theorem.

Theorem 4.3. *Let $(R, \mathfrak{m}, \mathbf{k})$ be a complete local domain of mixed characteristic $p > 0$ and suppose the following conditions:*

- (1) *let $A \rightarrow R$ be a coefficient ring map for a complete discrete valuation ring (A, π_A) ;*
- (2) *let x_0, x_1, \dots, x_d be a fixed set of minimal generators of \mathfrak{m} ;*
- (3) *the residue field \mathbf{k} is infinite.*

Then there exists a Zariski dense open subset $U' \subseteq \mathbb{P}^d(\mathbf{k})$ such that we have

$$\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i \notin \mathfrak{p}^{(2)}$$

for every prime \mathfrak{p} of R and every $a = (a_0 : \dots : a_d) \in \text{Sp}_A^{-1}(U') \subseteq \mathbb{P}^d(A)$.

Note that x_0, x_1, \dots, x_d are not a system of parameters of R , unless R is a regular local ring.

Proof. First, the R -module $\widehat{\Omega}_{R/A}$ is generated by dx_0, \dots, dx_d , which is easily verified by considering the surjective ring map $A[[X_0, \dots, X_d]] \twoheadrightarrow R$ defined by mapping each X_i to x_i . Let us recall an important fact. If R is a complete local domain with (A, π_A) its

coefficient ring, it follows from ([4], Lemma 2.6, or [19], Lemma 2) that for any fixed element $\mathfrak{p} \in U(\mathfrak{m})$, we have

$$\mu_{\mathfrak{p}}(\widehat{\Omega}_{R/A}) \geq \dim(R/\mathfrak{p}) - 1.$$

Noting that $\dim(R/\mathfrak{p}) - 1 = \dim(V(\mathfrak{p}) \cap U(\mathfrak{m}))$ for a subscheme $V(\mathfrak{p}) \cap U(\mathfrak{m}) \subseteq \text{Spec } R$ and applying Lemma 3.4 for the R -module $\widehat{\Omega}_{R/A}$, there is an ideal $(F_1, \dots, F_r) \subseteq R[X_0, \dots, X_d]$ such that we have

$$(4.1) \quad \dim(V(F_1, \dots, F_r) \cap (\phi^*)^{-1}(U(\mathfrak{m}))) \leq d + 1,$$

where $(\phi^*)^{-1}(U(\mathfrak{m}))$ is the inverse image of $U(\mathfrak{m})$ under $\phi^* : \text{Spec } R[X_0, \dots, X_d] \rightarrow \text{Spec } R$ and

$$\sum_{k=0}^d dx_k \otimes X_k \in \widehat{\Omega}_{R/A} \otimes_R R[X_0, \dots, X_d]$$

is basic on $U(F_1, \dots, F_r) \cap (\phi^*)^{-1}(U(\mathfrak{m}))$. Let T denote the localization of $R[X_0, \dots, X_d]$ at the prime ideal $\mathfrak{m}R[X_0, \dots, X_d]$. Then we have a flat local map of catenary local domains: $(R, \mathfrak{m}) \rightarrow (T, \mathfrak{m}_T)$ and $\dim R = \dim T$. By abuse of notation, we denote by $\phi^* : \text{Spec } T \rightarrow \text{Spec } R$ the scheme map induced by $R \rightarrow R[X_0, \dots, X_d] \rightarrow T$. Then since ϕ^* is faithfully flat, it is surjective.

In what follows, it suffices to consider the case when $(F_1, \dots, F_r)T$ is a proper ideal. Let $P \subseteq T$ be any prime ideal containing $(F_1, \dots, F_r)T$. Then it follows from Lemma 3.5 together with (4.1) above, that $V := V(F_1, \dots, F_r) \subseteq \text{Spec } T$ is a finite set and

$$\sum_{k=0}^d dx_k \otimes X_k$$

is basic on $\text{Spec } T \setminus V$ (the complement of $\phi^*(\text{Spec } T \setminus V)$ in $\text{Spec } R$ is a finite set). Then it suffices to consider the proof on $U(g) \subseteq \text{Spec } T$ for some $g \in R$ such that $\widehat{\Omega}_{R/A} \otimes_R T[g^{-1}]$ is a $T[g^{-1}]$ -free module, and we reason this as follows by Noetherian induction. By applying the lemma of generic freeness to the R/\mathfrak{p} -module $\widehat{\Omega}_{R/A} \otimes_R R/\mathfrak{p}$ for each prime $\mathfrak{p} \in \text{Min}_R(gR)$ and repeating the same discussion for $R/\mathfrak{p} \rightarrow T \otimes_R R/\mathfrak{p}$ in place of $R \rightarrow T$, we see that this process stabilizes, because R is a Noetherian ring. Hence it suffices to prove the theorem on the open subset $U(g) \subseteq \text{Spec } T$.

Now there exists a (not necessarily homogeneous) polynomial

$$G \in R[X_0, \dots, X_d] \setminus \mathfrak{m}R[X_0, \dots, X_d]$$

such that

$$\sum_{k=0}^d dx_k \otimes X_k$$

is basic on $U(g \cdot G) \subseteq \text{Spec } R[X_0, \dots, X_d]$. Choose a point $a = (a_0, \dots, a_d) \in \mathbb{P}^d(A)$ together with its lift $\tilde{a} = (\tilde{a}_0, \dots, \tilde{a}_d) \in \mathbb{A}^{d+1}(A)$ such that $G(\tilde{a}) \notin \mathfrak{m}$ (such a point exists, because $\#\mathbf{k} = \infty$ and $G \notin \mathfrak{m}R[X_0, \dots, X_d]$). Then by Noetherian induction hypothesis,

$$\sum_{k=0}^d \tilde{a}_k dx_k \in \widehat{\Omega}_{R/A}$$

is basic on $\phi^*(\text{Spec } T \setminus V) \subseteq \text{Spec } R$ in view of ([4], Lemma 1.1). Now we get the following implication. Let $\tilde{a} \in \mathbb{A}^{d+1}(A)$ be such that $G(\tilde{a}) \notin \mathfrak{m}$. Then for all $\mathfrak{p} \in \phi^*(\text{Spec } T \setminus V) = \text{Spec } R \setminus \{\mathfrak{p}_1, \dots, \mathfrak{p}_r, \mathfrak{m}\}$, it follows from Lemma 3.3 that

$$\mathbf{x}_{\tilde{a}} = \sum_{k=0}^d \tilde{a}_k x_k \notin \mathfrak{p}^{(2)}.$$

So it remains to deal with the issue on a finite set $\{\mathfrak{p}_1, \dots, \mathfrak{p}_r, \mathfrak{m}\}$. First, if we have $\tilde{a}_i \in A^\times$ for some i , then $\mathbf{x}_{\tilde{a}} \notin \mathfrak{m}^2$ by the minimality of x_0, \dots, x_d . Therefore, we have verified the condition: $\mathbf{x}_{\tilde{a}} \notin \mathfrak{p}^{(2)}$ for all $\mathfrak{p} \in \text{Spec } R \setminus \{\mathfrak{p}_1, \dots, \mathfrak{p}_r\}$, whenever $G(\tilde{a}) \notin \mathfrak{m}$.

Put $\overline{U}_0 := U(\overline{G}) \subseteq \mathbb{A}^{d+1}(\mathbf{k}) \setminus \{0\}$ (the origin of $\mathbb{A}^{d+1}(\mathbf{k})$ is excluded). Then define an open subset

$$(4.2) \quad U_0 \subseteq \mathbb{P}^d(\mathbf{k})$$

as the image of \overline{U}_0 under the projection map: $\mathbb{A}^{d+1}(\mathbf{k}) \setminus \{0\} \rightarrow \mathbb{P}^d(\mathbf{k})$. To deal with the issue on $\{\mathfrak{p}_1, \dots, \mathfrak{p}_r\}$, take the homogeneous polynomial:

$$F(X_0, \dots, X_d) := \sum_{i=0}^d x_i X_i \in R[X_0, \dots, X_d].$$

Then it suffices to force

$$\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i \notin \mathfrak{p}_j$$

for all $1 \leq j \leq r$. We fix a point $a = (a_0 : \dots : a_d) \in \mathbb{P}^d(A)$. For each $1 \leq j \leq r$, we have an equivalence of conditions:

$$(4.3) \quad \mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i \notin \mathfrak{p}_j \iff F(\tilde{a}_0, \dots, \tilde{a}_d) \not\equiv 0 \pmod{\mathfrak{p}_j}.$$

Moreover, the condition (4.3) is stable under taking multiplication by elements of A^\times .

Our final goal is to identify the set of points of $\mathbb{P}^d(A)$ satisfying the condition (4.3) and describe it as the inverse image of an open subset under the map $\text{Sp}_A : \mathbb{P}^d(A) \rightarrow \mathbb{P}^d(\mathbf{k})$. Since the maximal ideal \mathfrak{m} of R is generated by x_0, \dots, x_d , it is clear that \mathfrak{m} is generated by the set $\{F(\tilde{a}_0, \dots, \tilde{a}_d) \mid (a_0 : \dots : a_d) \in \mathbb{P}^d(A)\}$. Since the union of prime ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_r$ is strictly contained in \mathfrak{m} , there exists a point $(a_0 : \dots : a_d) \in \mathbb{P}^d(A)$ for which $F(\tilde{a}_0, \dots, \tilde{a}_d) \notin \mathfrak{p}_i$ for all $1 \leq i \leq r$. In fact, more is true. Applying Lemma 4.2 to the primes ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_r$, each of which is different from \mathfrak{m} , we find non-empty

open subsets U_1, \dots, U_r of $\mathbb{P}^d(\mathbf{k})$ with each U_i attached to \mathfrak{p}_i . Hence U' is defined as the intersection $U_0 \cap U_1 \cap \dots \cap U_r \subseteq \mathbb{P}^d(\mathbf{k})$, where U_0 is as given in (4.2). This completes the proof of the theorem. \square

As the main result of this article, we obtain the following theorem.

Theorem 4.4 (Local Bertini Theorem). *Let $(R, \mathfrak{m}, \mathbf{k})$ be a complete local domain of mixed characteristic $p > 0$ and suppose the following conditions:*

- (1) *let $A \rightarrow R$ be a coefficient ring map for a complete discrete valuation ring (A, π_A) ;*
- (2) *let x_0, x_1, \dots, x_d be a fixed set of minimal generators of \mathfrak{m} ;*
- (3) *R is normal, of depth $R \geq 3$, and the residue field \mathbf{k} is infinite.*

Then there exists a Zariski dense open subset $U \subseteq \mathbb{P}^d(\mathbf{k})$ satisfying the following properties. For any $a = (a_0 : \dots : a_d) \in \mathrm{Sp}_A^{-1}(U)$, the quotient $R/\mathfrak{x}_{\tilde{a}}R$ is a normal domain of mixed characteristic $p > 0$, where we put

$$\mathfrak{x}_{\tilde{a}} := \sum_{i=0}^d \tilde{a}_i x_i.$$

Proof. The first step of the proof has been completed in Theorem 4.3. Let $\mathrm{Reg}(R)$ denote the regular locus of $\mathrm{Spec} R$. Taking $U' \subseteq \mathbb{P}^d(\mathbf{k})$ as given in the conclusion of Theorem 4.3, we have

$$(4.4) \quad \mathrm{Reg}(R) \cap V(\mathfrak{x}_{\tilde{a}}) \subseteq \mathrm{Reg}(R/\mathfrak{x}_{\tilde{a}}R)$$

for all $a = (a_0 : \dots : a_d) \in \mathrm{Sp}_A^{-1}(U')$. Let us explain basic ideas of this proof. In **Step1**, we find an open set $U'' \subseteq \mathbb{P}^d(\mathbf{k})$ to deal with finitely many bad primes, with the help of Lemma 4.2. Then in **Step2**, we show that for $a \in \mathrm{Sp}_A^{-1}(U' \cap U'')$, the quotients $R/\mathfrak{x}_{\tilde{a}}R$ satisfy the well-known conditions (R_1) and (S_2) and thus are normal. Finally in **Step3**, we find an open set $U''' \subseteq \mathbb{P}^d(\mathbf{k})$ to show that the quotient rings $R/\mathfrak{x}_{\tilde{a}}R$ with $a \in \mathrm{Sp}_A^{-1}(U''')$ are of mixed characteristic.

Step1: The goal of this step is to find a candidate of an open subset $U'' \subseteq \mathbb{P}^d(\mathbf{k})$. Since R is a complete local domain, the singular locus $\mathrm{Sing}(U(\mathfrak{m}))$ of $U(\mathfrak{m}) \subseteq \mathrm{Spec} R$ is a proper closed subset. Hence the set of minimal primes in $\mathrm{Sing}(U(\mathfrak{m}))$ is finite, and let

$$Q_1 := \{\mathfrak{p} \in U(\mathfrak{m}) \mid \mathfrak{p} \text{ is a minimal prime in } \mathrm{Sing}(U(\mathfrak{m}))\}.$$

Note that every prime in Q_1 has height ≥ 2 , due to the (R_1) condition on R . On the other hand,

$$Q_2 := \{\mathfrak{p} \in U(\mathfrak{m}) \mid \mathrm{depth} R_{\mathfrak{p}} = 2 \text{ and } \dim R_{\mathfrak{p}} > 2\}$$

is also a finite set by ([4], Lemma 3.2 and the (S_2) condition on R). Now let $Q_1 \cup Q_2 := \{\mathfrak{p}_1, \dots, \mathfrak{p}_m\}$ and let us put $F(X_0, \dots, X_d) = \sum_{i=0}^d x_i X_i \in R[X_0, \dots, X_d]$. Then for each

$1 \leq j \leq m$, it follows that

$$\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i \notin \mathfrak{p}_j \iff F(\tilde{a}_0, \dots, \tilde{a}_d) \not\equiv 0 \pmod{\mathfrak{p}_j}.$$

Then applying Lemma 4.2 to $\{\mathfrak{p}_1, \dots, \mathfrak{p}_m\}$, we obtain U_1, \dots, U_m , which are non-empty open subsets of $\mathbb{P}^d(\mathbf{k})$ with each U_i attached to \mathfrak{p}_i . Put

$$(4.5) \quad U'' := U_1 \cap \dots \cap U_m.$$

Step2: The goal of this step is to show that the quotient $R/\mathbf{x}_{\tilde{a}}R$ is a normal domain for $a \in \mathrm{Sp}_A^{-1}(U'')$, or equivalently, $\mathbf{x}_{\tilde{a}}$ is contained in no prime of $Q_1 \cup Q_2$. Pick $\mathfrak{p} \in U(\mathfrak{m}) \cap V(\mathbf{x}_{\tilde{a}})$ with $\mathrm{ht} \mathfrak{p} \geq 2$ and assume that $\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d a_i x_i$ satisfies the condition (4.4) and that $\mathbf{x}_{\tilde{a}}$ is contained in no prime of $Q_1 \cup Q_2$.

(i) If we have $\mathrm{ht} \mathfrak{p} > 2$, then since $\mathbf{x}_{\tilde{a}}$ is contained in no prime of Q_2 , it follows that $\dim(R/\mathbf{x}_{\tilde{a}}R)_{\mathfrak{p}} \geq 2$ and $\mathrm{depth}(R/\mathbf{x}_{\tilde{a}}R)_{\mathfrak{p}} \geq 2$.

(ii) If we have $\mathrm{ht} \mathfrak{p} = 2$, then since $\mathbf{x}_{\tilde{a}}$ is contained in no prime of Q_1 and the height of every prime in Q_1 is at least 2, it follows that $R_{\mathfrak{p}}$ is regular. By (4.4), one finds that $(R/\mathbf{x}_{\tilde{a}}R)_{\mathfrak{p}}$ is a discrete valuation ring. On the other hand, the hypothesis that $\mathrm{depth}(R) \geq 3$ implies that $\mathrm{depth}(R/\mathbf{x}_{\tilde{a}}R) \geq 2$ by ([1], Proposition 1.2.9). Hence $R/\mathbf{x}_{\tilde{a}}R$ is a normal domain in view of Serre's normality criterion.

Step3: In this final step, we make $R/\mathbf{x}_{\tilde{a}}R$ into a local ring of mixed characteristic $p > 0$. Let $\{q_1, \dots, q_n\}$ be a set of all height-one primes of R lying above π_A . Then again, applying Lemma 4.2 to $\{q_1, \dots, q_n\}$, we find a non-empty open subset

$$(4.6) \quad U''' \subseteq \mathbb{P}^d(\mathbf{k}),$$

such that for $a = (a_0 : \dots : a_d) \in \mathrm{Sp}_A^{-1}(U) \subseteq \mathbf{P}^d(A)$, we have $\mathbf{x}_{\tilde{a}} \notin q_1 \cup \dots \cup q_n$.

Combining (4.4), (4.5), together with (4.6), and taking a non-empty open subset

$$U := U' \cap U'' \cap U''' \subseteq \mathbb{P}^d(\mathbf{k}),$$

it turns out that $\mathrm{Sp}_A^{-1}(U) \subseteq \mathbb{P}^d(A)$ has the required property. \square

Remark 4.5. Assume that R is a Cohen-Macaulay local normal domain. Then $R/\mathbf{x}_{\tilde{a}}R$ is Cohen-Macaulay and normal. One can continue this process until $\dim R = 2$ is attained. As to Cohen-Macaulay property, the following fact is known. Assume that $S \rightarrow R$ is a torsion free module-finite extension of local domains such that S is regular. Then R is a flat S -module if and only if R is Cohen-Macaulay by the Auslander-Buchsbaum formula.

Next, let us consider the case when the residue field is finite. Let $(R, \mathfrak{m}, \mathbb{F})$ be a complete local normal domain of mixed characteristic $p > 0$ with finite residue field \mathbb{F} . In other words, R is a finite extension of $W(\mathbb{F})[[z_1, \dots, z_n]]$, where $W(\mathbb{F})$ is the ring of Witt

vectors of \mathbb{F} . Let $W(\mathbb{F})^{\text{ur}}$ be the maximal unramified extension of $W(\mathbb{F})$. Then $W(\overline{\mathbb{F}})$ is the completion of $W(\mathbb{F})^{\text{ur}}$. Put

$$R_{W(\overline{\mathbb{F}})} := R \widehat{\otimes}_{W(\mathbb{F})} W(\overline{\mathbb{F}}) \text{ (resp. } R_{W(\mathbb{F})^{\text{ur}}} := \text{strict henselization of } R).$$

Then $R_{W(\mathbb{F})^{\text{ur}}}$ is local Noetherian, but not complete. By the main result of [6], $R_{W(\overline{\mathbb{F}})}$ is the completion of $R_{W(\mathbb{F})^{\text{ur}}}$ and a local normal domain. From algebraic number theory, $W(\mathbb{F})^{\text{ur}}$ is obtained from $W(\mathbb{F})$ by adjoining all n -th roots of unity for $(n, p) = 1$. There is a structure map $W(\overline{\mathbb{F}}) \rightarrow R_{W(\overline{\mathbb{F}})}$. We define the multiplicative map (not additive)

$$[-] : \overline{\mathbb{F}} \rightarrow W(\overline{\mathbb{F}})$$

as the Teichmüller map (see Proposition 8 in Chap. II §4 in [17] for details). An element in the image of $[-]$ is called a *Teichmüller lift*. In particular, we have $q \circ [-] = \text{Id}_{\overline{\mathbb{F}}}$, where $q : W(\overline{\mathbb{F}}) \rightarrow \overline{\mathbb{F}}$ is the residue field map. There is a set-theoretic mapping:

$$(4.7) \quad \theta_{W(\overline{\mathbb{F}})} : \mathbb{P}^d(\overline{\mathbb{F}}) \rightarrow \mathbb{P}^d(W(\overline{\mathbb{F}}))$$

defined by $\theta_{W(\overline{\mathbb{F}})}(b) := ([b_0] : \cdots : [b_d])$ for $b = (b_0 : \cdots : b_d) \in \mathbb{P}^d(\overline{\mathbb{F}})$. Then $\theta_{W(\overline{\mathbb{F}})}$ is well-defined, since $[-]$ is multiplicative. We write $R_{W(\mathbb{F}')} := R \otimes_{W(\mathbb{F})} W(\mathbb{F}')$ for a finite field extension $\mathbb{F} \rightarrow \mathbb{F}'$.

Corollary 4.6 (Finite Residue Field Case). *Let the hypothesis be as in Theorem 4.4 for $(R, \mathfrak{m}, \mathbb{F})$, except that we now assume the residue field \mathbb{F} is finite. Then there exists a non-empty open subset $U \subseteq \mathbb{P}^d(\overline{\mathbb{F}})$ such that the following holds:*

Fix an element $a = (a_0 : \cdots : a_d) \in \theta_{W(\overline{\mathbb{F}})}(U)$. Then there exists a finite extension $W(\mathbb{F}) \rightarrow W(\mathbb{F}')$ such that there is a choice of a lift \tilde{a}_i of a_i for each $0 \leq i \leq d$ with $\tilde{a}_i \in W(\mathbb{F}')$, and the quotient $R_{W(\mathbb{F}')} / \mathfrak{x}_{\tilde{a}} R_{W(\mathbb{F}')}$ is a normal domain of mixed characteristic $p > 0$.

The point of the proof is to construct a multiplicative map: $[\widetilde{-}] : \overline{\mathbb{F}} \rightarrow W(\mathbb{F})^{\text{ur}}$ which extends to the map $[-] : \overline{\mathbb{F}} \rightarrow W(\overline{\mathbb{F}})$.

Proof. We keep the notation as in Theorem 4.4. First, note that a priori a choice of a lift \tilde{a}_i of each a_i is contained in $W(\overline{\mathbb{F}})$. Since x_0, x_1, \dots, x_d are the minimal generators of the maximal ideal of $R_{W(\overline{\mathbb{F}})}$, the hypotheses of Theorem 4.4 are fulfilled for the complete local domain $R_{W(\overline{\mathbb{F}})}$. Let us construct a multiplicative map $[\widetilde{-}] : \overline{\mathbb{F}} \rightarrow W(\mathbb{F})^{\text{ur}}$ which extends to the map $[-]$. Note that $W(\mathbb{F})^{\text{ur}} = \varinjlim_{\lambda \in \Lambda} W(\mathbb{F}_\lambda)$ and we have the Teichmüller mapping $\mathbb{F}_\lambda \rightarrow W(\mathbb{F}_\lambda)$. Then we have a commutative diagram:

$$\begin{array}{ccc} \mathbb{F}_{\lambda'} & \xrightarrow{\text{Teich}} & W(\mathbb{F}_{\lambda'}) \\ \uparrow & & \uparrow \\ \mathbb{F}_\lambda & \xrightarrow{\text{Teich}} & W(\mathbb{F}_\lambda) \end{array}$$

which naturally forms a direct system, so the desired map $[\widetilde{-}]$ is given by its direct limit. On the other hand, it is easy to see that the map $[-] : \overline{\mathbb{F}} \rightarrow W(\overline{\mathbb{F}})$ factors as

$$\overline{\mathbb{F}} \xrightarrow{[\widetilde{-}]} W(\mathbb{F})^{\text{ur}} \longrightarrow W(\overline{\mathbb{F}}),$$

and thus for any $a \in \theta_{W(\overline{\mathbb{F}})}(U)$, we have $\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i \in R_{W(\mathbb{F})}^{\text{ur}}$ for an appropriate choice of \tilde{a}_i for every a_i . Since the map

$$R_{W(\mathbb{F})}^{\text{ur}} / \mathbf{x}_{\tilde{a}} R_{W(\mathbb{F})}^{\text{ur}} \rightarrow R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$$

is local flat, $R_{W(\mathbb{F})}^{\text{ur}} / \mathbf{x}_{\tilde{a}} R_{W(\mathbb{F})}^{\text{ur}}$ is a local normal domain.

By what we have said above, all the coefficients of a linear form $\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i x_i$ are contained in some finite extension $W(\mathbb{F}) \rightarrow W(\mathbb{F}')$. In other words, for a finite étale extension $R \rightarrow R_{W(\mathbb{F}')}$ of normal domains, the quotient ring $R_{W(\mathbb{F}')} / \mathbf{x}_{\tilde{a}} R_{W(\mathbb{F}')}$ is normal. \square

Remark 4.7. Let $\phi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ be a flat local map of local rings. Then one might think of the relationship between R/xR and S/xS for a non-zero divisor $x \in \mathfrak{m}$. In fact, in order to use the local Bertini theorem for S in terms of R , for example, assume that R and all fibers of ϕ are normal. Then for any x such that R/xR is normal, S/xS is also normal.

It is important and necessary to answer the following question:

Question 4.8. Resume the hypothesis of Theorem 4.4 and assume that an equality holds: $\mathbf{x}_{\tilde{a}} = u \mathbf{x}_{\tilde{a}'}$ for a unit $u \in R^\times$. Then is it true that $u \in A^\times$?

This question is restated as follows: Which subset of $\mathbb{P}^d(A)$ parametrizes the set of height-one primes $\{\mathbf{x}_{\tilde{a}}R\}$? In other words, does it give a set of distinct height-one primes of R ? In the next section, we will answer the above question. In fact, we need to restrict our attention to the set of those points which are in the image of the Teichmüller mapping. This will be important in the proof of the control theorem, which will be discussed later. We end this section with an example, which applies Theorem 4.4 and its proof for a given normal domain R .

Example 4.9. In this example, we are dealing with the case where the residue field is finite, since we can find a non-empty set which does the job. However, the result is valid for any discrete valuation coefficient ring. Suppose that $p \geq 3$ and

$$R := \mathbb{Z}_p[[x_1, x_2, x_3]] / (x_1^2 + x_2^2 + x_3^2),$$

which is a three-dimensional Cohen-Macaulay local normal domain. Now let us find a non-empty open subset $U \subseteq \mathbb{P}^3(\mathbb{F}_p)$ by keeping track of the proof of Theorem 4.4.

- (i) We need to have $d\mathbf{x}_{\tilde{a}} \in \widehat{\Omega}_{R/\mathbb{Z}_p}$ basic at every $\mathfrak{p} \in \text{Spec } R$.

(ii) We need to determine two finite sets of primes Q_1 and Q_2 in Theorem 4.4.

(iii) We need to avoid Q_1 and Q_2 as above.

Then we know

$$\widehat{\Omega}_{R/\mathbb{Z}_p} \simeq \frac{Rdx_1 \oplus Rdx_2 \oplus Rdx_3}{R(x_1dx_1 + x_2dx_2 + x_3dx_3)},$$

the singular locus of R is defined by the ideal (x_1, x_2, x_3) , $Q_1 = \{(x_1, x_2, x_3)\}$ and $Q_2 = \emptyset$.

To get a normal ring of mixed characteristic, take $\mathbf{x}_{\bar{a}} := \tilde{a}_0 p + \sum_{i=1}^3 \tilde{a}_i x_i$ such that

$$\bar{a} = (\bar{a}_0 : \bar{a}_1 : \bar{a}_2 : \bar{a}_3) \in U := U^+(z_0) \cap \left(\bigcup_{i=1}^3 U^+(z_i) \right) \subseteq \mathbb{P}^3(\mathbb{F}_p)$$

for the homogeneous coordinate $(z_0 : z_1 : z_2 : z_3)$. Let us check that this U is what we want. We have $\mathbf{x}_{\bar{a}} \notin Q_1$. If we assume \tilde{a}_1 is a unit for simplicity, we see that

$$\widehat{\Omega}_{R/\mathbb{Z}_p} \left[\frac{1}{x_3} \right] \simeq R \left[\frac{1}{x_3} \right] dx_1 \oplus R \left[\frac{1}{x_3} \right] dx_2$$

is a free module, in which the image of $d\mathbf{x}_{\bar{a}}$ spans a direct summand. On the other hand, for $\bar{R} := R/x_3R$,

$$\widehat{\Omega}_{R/\mathbb{Z}_p}/x_3 \cdot \widehat{\Omega}_{R/\mathbb{Z}_p} \simeq \frac{\bar{R}dx_1 \oplus \bar{R}dx_2 \oplus \bar{R}dx_3}{\bar{R}(x_1dx_1 + x_2dx_2)}.$$

To show that the image of $d\mathbf{x}_{\bar{a}}$ is basic on $\widehat{\Omega}_{R/\mathbb{Z}_p}/x_3 \cdot \widehat{\Omega}_{R/\mathbb{Z}_p}$, keep track of the same steps as above by inverting and killing first x_2 , and then x_1 .

Remark 4.10. If one takes $R := \mathbb{Z}_p[[x]]$, then $p \notin \mathfrak{p}^{(2)}$ for every prime \mathfrak{p} of R , since p is a regular parameter. But $dp = 0$ and so the converse of Lemma 3.3 does not hold true.

5. DISTINCT HYPERPLANE SECTIONS IN THE LOCAL BERTINI THEOREM

In this section, we give an answer to Question 4.8. Assume that $(R, \mathfrak{m}, \mathbf{k})$ is a complete local (not necessarily normal) domain with perfect residue field of characteristic $p > 0$ with its coefficient ring $W(\mathbf{k})$, the ring of Witt vectors. Then as defined in (4.7), the mapping:

$$\theta_{W(\mathbf{k})} : \mathbb{P}^d(\mathbf{k}) \rightarrow \mathbb{P}^d(W(\mathbf{k}))$$

is induced by the Teichmüller mapping $\mathbf{k} \rightarrow W(\mathbf{k})$. Note that the field \mathbf{k} can be finite. The following proposition asserts that the parameter set of specializations in the local Bertini theorem may be identified with a certain open subset $U \subseteq \mathbb{P}^d(\mathbf{k})$.

Proposition 5.1. *Let the notation be as above and let x_0, \dots, x_d be a set of minimal generators of the maximal ideal of R . Assume the following conditions:*

- (1) *If $\pi_{W(\mathbf{k})} \in \mathfrak{m}^2$, we put $\mathbf{x}_{\bar{a}} = \sum_{i=0}^d \tilde{a}_i x_i$.*
- (2) *If $\pi_{W(\mathbf{k})} \notin \mathfrak{m}^2$, we put $x_0 = \pi_{W(\mathbf{k})}$ and $\mathbf{x}_{\bar{a}} = \tilde{a}_0 \pi_{W(\mathbf{k})} + \sum_{i=1}^d \tilde{a}_i x_i$.*

Suppose that $\mathbf{x}_{\tilde{a}} = u\mathbf{x}_{a'}$ for $a, a' \in \theta_{W(\mathbf{k})}(\mathbb{P}^d(\mathbf{k})) \subseteq \mathbb{P}^d(W(\mathbf{k}))$ and $u \in R^\times$. Then we have $u \in W(\mathbf{k})^\times$.

Proof. To clarify the notation, we simply write a_i in place of \tilde{a}_i .

(1) Assume that $\pi_{W(\mathbf{k})} \in \mathfrak{m}^2$ and denote for simplicity by a_i the image of $a_i \in W(\mathbf{k})$ under the surjection:

$$W(\mathbf{k})[[X_0, \dots, X_d]] \twoheadrightarrow R \quad (X_i \mapsto x_i).$$

Let P be its kernel and we prove that

$$(5.1) \quad P \subseteq \pi_{W(\mathbf{k})}W(\mathbf{k})[[X_0, \dots, X_d]] + \mathcal{I}^2.$$

Here, we put $\mathcal{I} := (X_0, \dots, X_d)$. Let $F \in P$ be a non-zero element. We prove (5.1) by contradiction and so assume that

$$(5.2) \quad F \notin \pi_{W(\mathbf{k})}W(\mathbf{k})[[X_0, \dots, X_d]] + \mathcal{I}^2.$$

Under this assumption, after reducing $W(\mathbf{k})[[X_0, \dots, X_d]]$ by $\pi_{W(\mathbf{k})}$, we find that

$$f := F \pmod{\pi_{W(\mathbf{k})}} \notin \mathcal{I}^2\mathbf{k}[[X_0, \dots, X_d]].$$

The number of the minimal generators of the maximal ideal of $R/\pi_{W(\mathbf{k})}R$ is equal to

$$\dim_{\mathbf{k}} \mathfrak{m}/(\pi_{W(\mathbf{k})}R + \mathfrak{m}^2) = \dim_{\mathbf{k}} \mathfrak{m}/\mathfrak{m}^2,$$

because we assumed $\pi_{W(\mathbf{k})} \in \mathfrak{m}^2$. Hence $\bar{x}_0, \dots, \bar{x}_d$ form the minimal generators of the maximal ideal of $R/\pi_{W(\mathbf{k})}R$.

Claim 5.2. *Under the assumption (5.2), we can choose s with $0 \leq s \leq d$ such that*

$$f = \sum_{i=0}^{\infty} h_i X_s^i \in \mathbf{k}[[X_0, \dots, X_d]],$$

$h_i \in \mathbf{k}[[X_1, \dots, X_{s-1}, X_{s+1}, \dots, X_d]]$ for all $i \geq 0$, and h_1 is a unit.

Proof of Claim 5.2. We explain how to choose h_1 as a unit element. Fix a presentation $f = \sum_{i=0}^{\infty} h_i X_s^i \in \mathbf{k}[[X_0, \dots, X_d]]$ with respect to s and assume that h_1 is not a unit. Then since $f \notin \mathcal{I}^2\mathbf{k}[[X_0, \dots, X_d]]$, the element h_0 contains a non-zero linear term after presenting h_0 as an (infinite) sum of homogeneous polynomials. So we can write

$$h_0 = \sum_{i=1, i \neq s}^d a_i X_i + (\text{terms of degree at least 2})$$

and we have $a_t \neq 0$ for some t . By replacing s with t , we can achieve the requirement that h_1 is a unit. \square

By mapping $f = \sum_{i=0}^{\infty} h_i X_s^i$ to the quotient $R/\pi_{W(\mathbf{k})}R$, we get

$$\sum_{i=1}^{\infty} \bar{h}_i \bar{x}_s^i = -\bar{h}_0 \text{ in } R/\pi_{W(\mathbf{k})}R.$$

Note that $\sum_{i=1}^{\infty} \bar{h}_i \bar{x}_s^{i-1}$ is a unit since \bar{h}_1 is a unit of $R/\pi_{W(\mathbf{k})}R$. Hence, we have:

$$\bar{x}_s \cdot (\text{unit}) = -\bar{h}_0.$$

But this gives a contradiction to the fact that $\bar{x}_0, \dots, \bar{x}_d$ are the minimal generators of the maximal ideal of $R/\pi_{W(\mathbf{k})}R$ and that $-\bar{h}_0 \in (\bar{x}_0, \dots, \bar{x}_{s-1}, \bar{x}_{s+1}, \dots, \bar{x}_d)$. Hence (5.1) is established.

In the next place, fix an arbitrary lifting $\tilde{u} \in W(\mathbf{k})[[X_0, \dots, X_d]]$ of $u \in R^\times$ under the map $W(\mathbf{k})[[X_0, \dots, X_d]] \rightarrow R$. We write:

$$\tilde{u} = \sum_{s_0, \dots, s_d} \left(\sum_{r=0}^{\infty} b_r^{(s_0, \dots, s_d)} \pi_{W(\mathbf{k})}^r \right) X_0^{s_0} \cdots X_d^{s_d},$$

where (s_0, \dots, s_d) denotes the multi-index and $b_r^{(s_0, \dots, s_d)}$ are the Teichmüller lifts. Since \tilde{u} is a unit, we have $b_0^{(0, \dots, 0)} \neq 0$. By lifting the relation $\mathbf{x}_{\tilde{u}} = u\mathbf{x}_{\tilde{u}}$ to $W(\mathbf{k})[[X_0, \dots, X_d]]$, we have

$$\sum_{i=0}^d a_i X_i \equiv \left(\sum_{s_0, \dots, s_d} \left(\sum_{r=0}^{\infty} b_r^{(s_0, \dots, s_d)} \pi_{W(\mathbf{k})}^r \right) X_0^{s_0} \cdots X_d^{s_d} \right) \left(\sum_{i=0}^d a'_i X_i \right) \pmod{P}.$$

Rewrite the above equation as:

$$\begin{aligned} \sum_{i=0}^d (a_i - a'_i b_0^{(0, \dots, 0)}) X_i &\equiv \left(\sum_{s_0, \dots, s_d} \left(\sum_{r=1}^{\infty} b_r^{(s_0, \dots, s_d)} \pi_{W(\mathbf{k})}^r \right) X_0^{s_0} \cdots X_d^{s_d} \right) \left(\sum_{i=0}^d a'_i X_i \right) \\ &+ \left(\sum_{\substack{(s_0, \dots, s_d) \\ \neq (0, \dots, 0)}} b_0^{(s_0, \dots, s_d)} X_0^{s_0} \cdots X_d^{s_d} \right) \left(\sum_{i=0}^d a'_i X_i \right) \pmod{P}. \end{aligned}$$

Then by mapping the above equation to the quotient $\mathbf{k}[[X_0, \dots, X_d]]$, comparing the degrees on both sides, and then using the relation (5.1), we find that

$$a_i = a'_i b_0^{(0, \dots, 0)} + \pi_{W(\mathbf{k})} \cdot v_i$$

for some $v_i \in W(\mathbf{k})$. However if $v_i \neq 0$, this implies that a_i is not a Teichmüller lift, since both a'_i and $b_0^{(0, \dots, 0)}$ are Teichmüller lifts. So we must have $v_i = 0$ for all $0 \leq i \leq d$ and the following relation holds:

$$\begin{aligned} &\left(\sum_{s_0, \dots, s_d} \left(\sum_{r=1}^{\infty} b_r^{(s_0, \dots, s_d)} \pi_{W(\mathbf{k})}^r \right) X_0^{s_0} \cdots X_d^{s_d} \right) \left(\sum_{i=0}^d a'_i X_i \right) \\ &+ \left(\sum_{\substack{(s_0, \dots, s_d) \\ \neq (0, \dots, 0)}} b_0^{(s_0, \dots, s_d)} X_0^{s_0} \cdots X_d^{s_d} \right) \left(\sum_{i=0}^d a'_i X_i \right) \in P. \end{aligned}$$

Since P is a prime ideal, we deduce that

$$\left(\sum_{s_0, \dots, s_d} \left(\sum_{r=1}^{\infty} b_r^{(s_0, \dots, s_d)} \pi_{W(\mathbf{k})}^r \right) X_0^{s_0} \cdots X_d^{s_d} + \sum_{\substack{(s_0, \dots, s_d) \\ \neq (0, \dots, 0)}} b_0^{(s_0, \dots, s_d)} X_0^{s_0} \cdots X_d^{s_d} \right) \in P$$

and thus $\tilde{u} \equiv b_0^{(0, \dots, 0)} \pmod{P}$ and $u \in W(\mathbf{k})$. Now we obtain $u \in W(\mathbf{k})^\times$.

(2) Assume that $\pi_{W(\mathbf{k})} \notin \mathfrak{m}^2$. Then taking $x_0 = \pi_{W(\mathbf{k})}$, we may consider the surjection $W(\mathbf{k})[[X_1, \dots, X_d]] \rightarrow R (X_i \mapsto x_i)$ and let P be its kernel. In this case, we prove that

$$(5.3) \quad P \subseteq \mathcal{M}^2,$$

where we put $\mathcal{M} := (\pi_{W(\mathbf{k})}, X_1, \dots, X_d)$. Let $F \in P$ be any non-zero element. Then we can write

$$(5.4) \quad F = \sum_{s_1, \dots, s_d} \left(\sum_{r=0}^{\infty} b_r^{(s_1, \dots, s_d)} \pi_{W(\mathbf{k})}^r \right) X_1^{s_1} \cdots X_d^{s_d},$$

for Teichmüller lifts $b_r^{(s_1, \dots, s_d)}$. Now assume that $F \notin \mathcal{M}^2$ for a contradiction. Then the equation (5.4) can be written as

$$(5.5) \quad F = b_0 \pi_{W(\mathbf{k})} + \sum_{i=1}^d b_i X_i + (\text{terms of degree at least 2})$$

for Teichmüller lifts b_i and at least one of b_0, \dots, b_d is not zero, say b_k . Then mapping (5.5) to R , F goes to 0 and we find that $x_k \in (\pi_{W(\mathbf{k})}, x_2, \dots, x_{k-1}, x_{k+1}, \dots, x_d)$, due to $b_k \neq 0$. But since $\pi_{W(\mathbf{k})}, x_1, \dots, x_d$ are the minimal generators of \mathfrak{m} , this is a contradiction. Thus, we must have $F \in \mathcal{M}^2$ and (5.3) is proved. Assume that $\mathbf{x}_{\tilde{a}} = u \mathbf{x}_{\tilde{a}}$ for some $u \in R^\times$. Then by applying the final step of (1), together with the fact that $P \subseteq \mathcal{M}^2$, we conclude that $u \in W(\mathbf{k})^\times$, as desired. \square

Now Proposition 5.1 assures us that there are sufficiently many normal hyperplane sections for a local normal domain. We start with the following lemma:

Lemma 5.3. *Let R be a Noetherian domain and let $\{P_\lambda\}_{\lambda \in \Lambda}$ be an infinite set of distinct height-one primes. Then we have*

$$\bigcap_{\lambda \in \Lambda} P_\lambda = 0.$$

Proof. Assume that there is a non-zero element $a \in \bigcap_{\lambda \in \Lambda} P_\lambda$. Let \overline{P}_λ denote the image of P_λ under the surjection $R \rightarrow R/aR$. Then since $a \in P_\lambda$ for all λ , the set $\{\overline{P}_\lambda\}_{\lambda \in \Lambda}$ gives an infinite set of minimal prime ideals of the Noetherian ring R/aR . But this is a contradiction and we must have $\bigcap_{\lambda \in \Lambda} P_\lambda = 0$, as desired. \square

Applying this lemma, we get the following.

Corollary 5.4. *In addition to the hypothesis of Proposition 5.1, assume that $S \subseteq \mathbb{P}^d(\mathbf{k})$ is an infinite subset such that the quotient $R/\mathfrak{x}_{\tilde{a}}R$ is an integral domain for every $a \in \theta_{W(\mathbf{k})}(S)$. Then we have*

$$\bigcap_{a \in \theta_{W(\mathbf{k})}(S)} \mathfrak{x}_{\tilde{a}}R = 0.$$

Proof. By Proposition 5.1 and our assumption, we see that $\{\mathfrak{x}_{\tilde{a}}R\}_{a \in \theta_{W(\mathbf{k})}(S)}$ is an infinite set of height-one primes of a Noetherian domain R . Now the corollary follows from Lemma 5.3. \square

It is not clear if the proof of Proposition 5.1 can be modified so that it holds for any residue field. Via the proof of the proposition, we obtain the following corollary, which is useful in dealing with the unramified case.

Corollary 5.5. *In addition to the hypothesis of Proposition 5.1, assume that $\pi_{W(\mathbf{k})}$ is part of minimal generators of \mathfrak{m} and $R/\mathfrak{x}_{\tilde{a}}R$ is an integral domain of mixed characteristic for*

$$\mathfrak{x}_{\tilde{a}} = \tilde{a}_0 \pi_{W(\mathbf{k})} + \sum_{i=1}^d \tilde{a}_i x_i$$

and $a = (a_0 : \dots : a_d) \in \mathbb{P}^d(W(\mathbf{k}))$. Then $\pi_{W(\mathbf{k})}$ is part of minimal generators of the maximal ideal of the quotient ring $R/\mathfrak{x}_{\tilde{a}}R$.

Note that the linear form $\mathfrak{x}_{\tilde{a}}$ is not assumed to define a normal quotient $R/\mathfrak{x}_{\tilde{a}}R$.

Proof. Let $W(\mathbf{k})[[X_1, \dots, X_d]] \twoheadrightarrow R$ be a surjective ring map with its kernel P such that the images of $\pi_{W(\mathbf{k})}, X_1, \dots, X_d$ are the minimal generators of \mathfrak{m} . Then we obtain that $P \subseteq (\pi_{W(\mathbf{k})}, X_1, \dots, X_d)^2$ as in the proof of Proposition 5.1. Let

$$\tilde{\mathfrak{x}}_{\tilde{a}} = \tilde{a}_0 \pi_{W(\mathbf{k})} + \sum_{i=1}^d \tilde{a}_i X_i \in W(\mathbf{k})[[X_1, \dots, X_d]]$$

be a lift of $\mathfrak{x}_{\tilde{a}}$. Since R and $R/\mathfrak{x}_{\tilde{a}}R$ are integral domains of mixed characteristic by assumption, $a_i \in W(\mathbf{k})$ must be a unit for some $1 \leq i \leq d$, which gives an isomorphism:

$$W(\mathbf{k})[[X_1, \dots, X_d]]/(\tilde{\mathfrak{x}}_{\tilde{a}}) \simeq W(\mathbf{k})[[X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_d]],$$

together with

$$\pi_{W(\mathbf{k})} \notin (\pi_{W(\mathbf{k})}, X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_d)^2.$$

But since $\overline{P} \subseteq (\pi_{W(\mathbf{k})}, X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_d)^2$, where \overline{P} denotes the image of P in the quotient $W(\mathbf{k})[[X_1, \dots, X_d]]/(\tilde{\mathfrak{x}}_{\tilde{a}})$, it follows that $\pi_{W(\mathbf{k})}$ forms part of minimal generators of the maximal ideal of $R/\mathfrak{x}_{\tilde{a}}R$, as required. \square

Remark 5.6. In this section, it is essential to assume that the residue field is perfect, which allows us to present an element in the ring $W(\mathbf{k})[[X_1, \dots, X_d]]$ in a unique way using the Teichmüller mapping. Note that the results in this section as well as Theorem 4.4 can be applied to a complete local normal domain $(R, \mathfrak{m}, \mathbf{k})$ with infinite perfect residue field and depth $R \geq 3$.

6. SERRE'S CONDITIONS (R_n) AND (S_n)

In this section, we prove Bertini theorems in the case when R satisfies Serre's conditions on the punctured spectra of local rings. The essential part for these cases is found in the proof of Theorem 4.4. As usual, we put

$$\mathbf{x}_{\tilde{a}} = \sum_{i=0}^d \tilde{a}_i X_i$$

for $a = (a_0, \dots, a_d) \in \mathbb{P}^d(A)$.

Corollary 6.1. *Suppose that $(R, \mathfrak{m}, \mathbf{k})$ is a complete local reduced ring of mixed characteristic $p > 0$, that conditions (1) and (2) of Theorem 4.4 hold, and that the residue field \mathbf{k} is infinite. If $R_{\mathfrak{p}}$ has Serre's (R_n) (resp. (S_n)) for all $\mathfrak{p} \in X$, then there exists a Zariski dense open subset $U \subseteq \mathbb{P}^d(\mathbf{k})$ such that for every*

$$a = (a_0 : \dots : a_d) \in \mathrm{Sp}_A^{-1}(U),$$

the quotient $R_{\mathfrak{p}}/\mathbf{x}_{\tilde{a}}R_{\mathfrak{p}}$ has (R_n) (resp. (S_n)) for all $\mathfrak{p} \in U(\mathfrak{m}) \cap V(\mathbf{x}_{\tilde{a}})$.

Proof. We briefly sketch the proof of the corollary. Since R is complete local and reduced, the non-singular locus of R is non-empty. Thus, the set

$$Q_1 := \{\mathfrak{p} \in U(\mathfrak{m}) \mid \mathfrak{p} \text{ is a minimal prime in } \mathrm{Sing}(U(\mathfrak{m}))\}$$

is finite. Let

$$Q_2 := \{\mathfrak{p} \in U(\mathfrak{m}) \mid \mathrm{depth} R_{\mathfrak{p}} = n \text{ and } \dim R_{\mathfrak{p}} > n\},$$

which is also finite by ([4], lemma 3.2). The proof is similar to that of Theorem 4.4. So it suffices to avoid the union of finite set of prime ideals in $Q_1 \cup Q_2$. Namely, for any $a = (a_0 : \dots : a_d) \in \mathrm{Sp}_A^{-1}(U)$, the localization $R_{\mathfrak{p}}/\mathbf{x}_{\tilde{a}}R_{\mathfrak{p}}$ has (R_n) (resp. (S_n)) for all $\mathfrak{p} \in U(\mathfrak{m}) \cap V(\mathbf{x}_{\tilde{a}})$ and $\mathrm{Reg}(R) \cap V(\mathbf{x}_{\tilde{a}}) \subseteq \mathrm{Reg}(R/\mathbf{x}_{\tilde{a}}R)$. \square

The above proof also shows that the Bertini theorem holds for mixed Serre's conditions. That is, if R has $(R_s) + (S_r)$, then so does $R_{\mathfrak{p}}/\mathbf{x}_{\tilde{a}}R_{\mathfrak{p}}$ for all $a = (a_0 : \dots : a_d) \in \mathrm{Sp}_A^{-1}(U)$ and all $\mathfrak{p} \in U(\mathfrak{m}) \cap V(\mathbf{x}_{\tilde{a}})$. For instance, we obtain the Bertini theorem for reduced rings, since we know that R is reduced if and only if R has $(R_0) + (S_1)$. To be precise, we have the following version of the local Bertini theorem:

Corollary 6.2. *Suppose that $(R, \mathfrak{m}, \mathbf{k})$ is a complete local normal domain of dimension 2 in mixed characteristic $p > 0$, that conditions (1) and (2) of Theorem 4.4 hold, and that the residue field \mathbf{k} is infinite. Then there exists a Zariski dense open subset $U \subseteq \mathbb{P}^d(\mathbf{k})$ such that for every*

$$a = (a_0 : \cdots : a_d) \in \mathrm{Sp}_A^{-1}(U),$$

the quotient $R/\mathfrak{x}_{\bar{a}}R$ is a reduced ring of mixed characteristic $p > 0$.

Proof. We first show that there exists a Zariski dense open subset $U' \subseteq \mathbb{P}^d(\mathbf{k})$ for which $R/\mathfrak{x}_{\bar{a}}R$ is reduced for all $a \in \mathrm{Sp}_A^{-1}(U')$. Since R is a domain, $R_{\mathfrak{p}}$ has $(R_0) + (S_1)$ for all $\mathfrak{p} \in \mathrm{Spec} R$. Then by Corollary 6.1, we find a desired $U' \subseteq \mathbb{P}^d(\mathbf{k})$ such that, for $a \in \mathrm{Sp}_A^{-1}(U')$, $R_{\mathfrak{p}}/\mathfrak{x}_{\bar{a}}R_{\mathfrak{p}}$ is reduced for all $\mathfrak{p} \in U(\mathfrak{m}) \cap V(\mathfrak{x}_{\bar{a}})$. It remains to show that the localization of $R/\mathfrak{x}_{\bar{a}}R$ at \mathfrak{m} , which is $R/\mathfrak{x}_{\bar{a}}R$ itself, has (S_1) . But since R is a local normal domain of dimension 2, it is Cohen-Macaulay and thus, $R/\mathfrak{x}_{\bar{a}}R$ has (S_1) . Hence $R/\mathfrak{x}_{\bar{a}}R$ is reduced for all $a \in \mathrm{Sp}_A^{-1}(U')$.

To make the quotient $R/\mathfrak{x}_{\bar{a}}R$ into a ring of mixed characteristic, we need to shrink U' to an open subset U . To this aim, it suffices to choose U such that π_A is not a zero divisor of $R/\mathfrak{x}_{\bar{a}}R$ for all $a \in \mathrm{Sp}_A^{-1}(U)$. In fact, we may take $\mathfrak{x}_{\bar{a}}$ so that it is contained in none of prime ideals in $\mathrm{Min}_R(\pi_A R)$, since every system of parameters of R is a regular sequence. Let $U''' \subseteq \mathbb{P}^d(\mathbf{k})$ be an open subset as in **Step3** of Theorem 4.4. Then the required open subset is defined as $U := U' \cap U'''$. \square

Remark 6.3. In place of the hypothesis of Corollary 6.2, assume that R is only an integral domain. Then can one find $\mathfrak{x}_{\bar{a}}$ such that $R/\mathfrak{x}_{\bar{a}}R$ is also an integral domain? In the mixed characteristic case, the answer to this question is not clear yet. But there is a two-dimensional complete local normal domain over \mathbb{C} without principal prime ideals at all (such an example is due to Laufer, as mentioned in [4]. However, an explicit example is not given there). In light of this, both Corollary 6.1 and Corollary 6.2 seem to be the best results.

7. CHARACTERISTIC IDEALS OF TORSION MODULES OVER NORMAL DOMAINS

Throughout this section, we assume that R is a Noetherian normal domain and M is a finitely generated torsion R -module. Then the localization of R at every height-one prime is a discrete valuation ring. We introduce an invariant of the module M . For an ideal I of R , let $M[I]$ denote the maximal submodule of M which is annihilated by I . We follow the definition of characteristic ideals by Skinner-Urban as in [18]. For more results and properties on characteristic ideals with its relation to the Fitting ideal, see § 9. For a finitely generated R -module M , we denote by

$$M^{\mathrm{rc}} := \mathrm{Hom}_R(\mathrm{Hom}_R(M, R), R),$$

the reflexive closure of M .

Definition 7.1. Let the notation be as above. Then the *characteristic ideal* is an ideal of R defined by

$$\text{char}_R(M) = \{x \in R \mid v_P(x) \geq \ell_{R_P}(M_P) \text{ for any height-one prime } P\},$$

where $v_P(-)$ is the normalized valuation of R_P ; that is, $v_P(PR_P) = 1$.

Since M is torsion, $\ell_{R_P}(M_P) = 0$ for all but finitely many height-one primes \mathfrak{p} of R and it suffices to take only height-one primes in the support of M in the definition. When M is not a torsion R -module, we put $\text{char}_R(M) = 0$.

Remark 7.2. The formation of characteristic ideals does not commute with base change in general. For example, let $R = \mathbb{Z}_p[[x, y]]$ and let $M = R/xR$. Then M is a torsion module and $\text{char}_R(M) = xA$. However, the R/xR -module M/xM is not torsion. Therefore,

$$0 = \text{char}_{R/xR}(M/xM) \subsetneq xR = \text{char}_R(M)R/xR$$

in this case. In general, even when M/xM is a torsion R/xR -module, it may happen that $\text{char}_{R/xR}(M/xM) \neq \text{char}_R(M)R/xR$, which is caused by the presence of pseudo-null submodules (see the definition below).

Definition 7.3. A finitely generated module M over a Noetherian normal domain R is *pseudo-null*, if $M_{\mathfrak{p}} = 0$ for all height-one primes $\mathfrak{p} \in \text{Spec } R$. A homomorphism of R -modules $f : M \rightarrow N$ is a *pseudo-isomorphism*, if both $\ker(f)$ and $\text{coker}(f)$ are pseudo-null modules.

The proof of the next lemma is found in ([12], Proposition 5.1.7).

Lemma 7.4 (Structure Theorem). *Let M be a finitely generated torsion module over a Noetherian normal domain R . Then there exist a finite set of height-one primes $\{P_i\}_{i \in I}$ (which is not necessarily a redundant set of height-one primes) and a set of natural numbers $\{e_i\}_{i \in I}$ such that there is a homomorphism:*

$$f : M \rightarrow \bigoplus_{i \in I} R/P_i^{e_i}$$

that is a pseudo-isomorphism. Moreover, both $\{P_i\}_{i \in I}$ and $\{e_i\}_{i \in I}$ are uniquely determined.

Henceforth, we use the notation $M \approx N$ to indicate that there is a pseudo-isomorphism between M and N . As mentioned before, the formation of characteristic ideals does not commute with base change in general, which can produce extra error terms.

Proposition 7.5. *Let M be a finitely generated torsion module over a Noetherian normal domain R . Let x be an element of R which satisfies the following conditions:*

- (1) R/xR is a normal domain (which implies that xR is a prime ideal);
- (2) x is contained in no prime ideal of height-one in the support of M (which implies that M/xM is a torsion R/xR -module).

Then we have:

$$\text{char}_{R/xR}(M/xM) = \left(\text{char}_{R/xR}(M[x]) \cdot \prod_{\substack{\text{ht } \mathfrak{p}=1, \\ \mathfrak{p} \in \text{Spec } R}} (\mathfrak{p}(R/xR))^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} \right)^{\text{rc}}.$$

Proof. Recall from Proposition 9.6 in Appendix, that if $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is a short exact sequence of finitely generated R -modules, then $\text{char}_R(M) = (\text{char}_R(L) \cdot \text{char}_R(N))^{\text{rc}}$. The condition stated in the proposition implies that all relevant modules are torsion. First off, we claim that

$$(7.1) \quad \text{char}_{R/xR} \left(\bigoplus_{\text{ht } \mathfrak{p}=1} R/(\mathfrak{p} + xR)^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} \right) = \left(\prod_{\substack{\text{ht } \mathfrak{p}=1, \\ \mathfrak{p} \in \text{Spec } R}} (\mathfrak{p}(R/xR))^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} \right)^{\text{rc}}.$$

For the proof of (7.1), let $\{P_1, \dots, P_m\}$ be a set of all height-one primes of R/xR which contain $\mathfrak{p}(R/xR)$. Then the localization $(R/xR)_{P_i}$ is a discrete valuation ring and we have

$$\mathfrak{p}(R/xR)_{P_i} = P_i^{\ell_{(R/xR)_{P_i}}((R/(\mathfrak{p}+xR))_{P_i})} (R/xR)_{P_i}$$

for all i . Then (7.1) follows from this and the definition of characteristic ideals. Thus, it suffices to prove the following equality:

$$(7.2) \quad \left(\text{char}_{R/xR}(M/xM) \cdot (\text{char}_{R/xR}(M[x]))^{-1} \right)^{\text{rc}} = \text{char}_{R/xR} \left(\bigoplus_{\substack{\text{ht } \mathfrak{p}=1, \\ \mathfrak{p} \in \text{Spec } R}} R/(\mathfrak{p} + xR)^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} \right),$$

which is regarded as an element in the group of reflexive fractional ideals of R , and $(\text{char}_{R/xR}(M[x]))^{-1}$ is the unique fractional ideal which is an inverse of $\text{char}_{R/xR}(M[x])$.

By considering all torsion R -modules satisfying the condition (2) with respect to $x \in R$, we show that both sides of (7.2), regarded as operations on such R -modules, is multiplicative on short exact sequences. Let

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

be a short exact sequence of R -modules satisfying the condition (2). Then since the function $\ell(-)$ is additive, it follows that the right hand side of (7.2) is multiplicative. On the other hand, there follows the exact sequence:

$$0 \rightarrow L[x] \rightarrow M[x] \rightarrow N[x] \rightarrow L/xL \rightarrow M/xM \rightarrow N/xN \rightarrow 0$$

by the snake lemma. If \mathfrak{p} is a height-two prime ideal of R containing $x \in R$, we may localize the above exact sequence at \mathfrak{p} , so it follows that the left hand side of (7.2) is multiplicative as well.

Hence we are reduced to the case that $M = R/\mathfrak{q}$ for a prime ideal \mathfrak{q} by the prime filtration argument. Assume first that $\text{ht } \mathfrak{q} = 1$. Then we clearly have $\text{Supp}(M) = \{\mathfrak{q}\}$, $M[x] = 0$, and $\ell_{R_{\mathfrak{q}}}(M_{\mathfrak{q}}) = 1$, because $M_{\mathfrak{q}}$ is a simple R -module. Now (7.2) is obviously true. Next assume that $\text{ht } \mathfrak{q} > 1$. Then it is easy to see that both sides of (7.2) are equal to a unit ideal, which completes the proof. \square

8. APPLICATIONS TO CHARACTERISTIC IDEALS

Our final goal is to prove Theorem 8.8. The aim of the main theorem is to establish some techniques, which enable us to study the Iwasawa's main conjecture attached to a p -adic family of modular forms (see a forthcoming paper [15]). For this reason, it is necessary to deal with local rings with finite residue field. Let $(R, \mathfrak{m}, \mathbb{F})$ be a complete local normal domain of mixed characteristic with finite residue field. In other words, R is the integral closure of $\mathbb{Z}_p[[z_1, \dots, z_n]]$ in a finite field extension of the field of fractions of $\mathbb{Z}_p[[z_1, \dots, z_n]]$. We recall the set-up of Corollary 4.6 and prove some preliminary results.

Let $R_{W(\overline{\mathbb{F}})} := R \widehat{\otimes}_{W(\mathbb{F})} W(\overline{\mathbb{F}})$ with its coefficient ring $W(\overline{\mathbb{F}})$. Then if $\text{depth } R \geq 3$, the complete local ring $R_{W(\overline{\mathbb{F}})}$ fits into the hypothesis of Theorem 4.4. We have the set-theoretic mapping: $\theta_{W(\overline{\mathbb{F}})} : \mathbb{P}^d(\overline{\mathbb{F}}) \rightarrow \mathbb{P}^d(W(\overline{\mathbb{F}}))$ as constructed in (4.7).

To establish the fundamental theorem for characteristic ideals, we need to relate torsion R -modules to torsion $R_{W(\overline{\mathbb{F}})}$ -modules and then descend to R by faithful flatness. The advantage of working with $R_{W(\overline{\mathbb{F}})}$ is that the residue field is the algebraic closure of a finite field. We introduce some notation. Denote by $\text{Fitt}_A(M)$ the Fitting ideal of an A -module M . We make free use of results and notation from Appendix.

Definition 8.1. Under the notation as above, fix a set of minimal generators x_0, \dots, x_d of the unique maximal ideal of R and let $U \subseteq \mathbb{P}^d(\overline{\mathbb{F}})$ be as given in Corollary 4.6. Then we set

$$\mathcal{L}_{W(\overline{\mathbb{F}})} := \{\mathfrak{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} \mid a = (a_0 : \dots : a_d) \in \theta_{W(\overline{\mathbb{F}})}(U)\}$$

for the mapping $\theta_{W(\overline{\mathbb{F}})} : \mathbb{P}^d(\overline{\mathbb{F}}) \rightarrow \mathbb{P}^d(W(\overline{\mathbb{F}}))$. For a finitely generated torsion $R_{W(\overline{\mathbb{F}})}$ -module M , we define a subset:

$$\mathcal{L}_{W(\overline{\mathbb{F}})}(M) \subseteq \mathcal{L}_{W(\overline{\mathbb{F}})}$$

which consists of all height-one primes $\mathfrak{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}$ such that the following conditions are satisfied:

- (A) $M/\mathfrak{x}_{\tilde{a}} M$ is a torsion $R_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$ -module.
- (B) The following equalities of ideals hold in $R_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$:

$$\text{char}_{R_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}}(M/\mathfrak{x}_{\tilde{a}} M) = \left(\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)(R_{W(\overline{\mathbb{F}})}/\mathfrak{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}}$$

$$= \left(\text{Fitt}_{R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \right) (R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}},$$

where $M \approx \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i}$ is a fundamental pseudo-isomorphism.

When the base ring R is isomorphic to a power-series ring over a discrete valuation ring, compare the above definition with the one given in ([14], Definition 3.2.). The way of interpreting the condition **(B)** in Definition 8.1 is that one wants to consider the characteristic ideals through Fitting ideal as an intermediate invariant. Note that all of three ideals appearing in **(B)** may differ in general.

We are going to prove a number of lemmas to describe an explicit structure of the set $\mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ (see Lemma 8.5 and Lemma 8.6). To this aim, we prove some preliminary results from [14] over general normal domains by making necessary modifications. The next lemma is necessary for a technical reason and it will be shown in Lemma 8.4 how to use it in a more concrete situation. The reader may skip the proof at the first reading.

Lemma 8.2. *Under the notation and the hypothesis as in Corollary 4.6, assume that M is a finitely generated torsion $R_{W(\overline{\mathbb{F}})}$ -module. For a fundamental pseudo-isomorphism $M \approx \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i}$, set*

$$I = \text{Fitt}_{R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \right)$$

and consider the natural injection $I \rightarrow I^{\text{rc}}$ with its cokernel N . Then there exists a finite set $\{Q_i\}_{1 \leq i \leq \ell}$ consisting of height-two primes of $R_{W(\overline{\mathbb{F}})}$ with the following condition:

Fix an arbitrary element

$$\mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} \in \bigcap_{1 \leq i \leq \ell} \mathcal{L}_{W(\overline{\mathbb{F}})}(R_{W(\overline{\mathbb{F}})} / Q_i)$$

and let $P \in \text{Spec } R_{W(\overline{\mathbb{F}})}$ be any prime ideal such that $\mathbf{x}_{\tilde{a}} \in P$ and $\text{ht } P \leq 2$. Then $N_P = 0$.

Proof. By definition, the module N is supported on a closed subset of codimension two in $\text{Spec } R_{W(\overline{\mathbb{F}})}$. So there are only finitely many height-two primes contained in $\text{Supp } N$. Then we show that it is sufficient to choose Q_1, \dots, Q_ℓ as those height-two primes contained in $\text{Supp } N$. Let $N_i := R_{W(\overline{\mathbb{F}})} / Q_i$. Then N_i is a pseudo-null $R_{W(\overline{\mathbb{F}})}$ -module and we have $\text{char}_{R_{W(\overline{\mathbb{F}})}}(N_i) = R_{W(\overline{\mathbb{F}})}$. Then by the condition **(B)**, we have

$$\text{char}_{R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}}(N_i / \mathbf{x}_{\tilde{a}} N_i) = R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})},$$

and this implies that $N_i / \mathbf{x}_{\tilde{a}} N_i$ is a pseudo-null $R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$ -module. In other words, $\mathbf{x}_{\tilde{a}} \notin Q_i$ for all $1 \leq i \leq \ell$. So if we choose a prime ideal P such that $\mathbf{x}_{\tilde{a}} \in P$ and $\text{ht } P \leq 2$, then we must have $P \notin \text{Supp } N$, which proves the lemma. \square

We will discuss when equalities occur between various ideals in the condition **(B)** in Definition 8.1.

Discussion 8.3. Suppose M is a finitely generated torsion module over a Noetherian normal domain A . Then we have

$$\text{Fitt}_A(M) \subseteq \text{char}_A(M) \text{ and } \text{Fitt}_A(M)^{\text{rc}} = \text{char}_A(M)$$

(see Proposition 9.6 in Appendix). Take a fundamental pseudo-isomorphism:

$$M \approx \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i}$$

for a (not necessarily redundant) finite set of height-one primes $\{P_i\}$ of $R_{W(\overline{\mathbb{F}})}$. Choose $\mathbf{x}_{\tilde{a}} \in R_{W(\overline{\mathbb{F}})}$ such that $R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$ is normal and $M / \mathbf{x}_{\tilde{a}} M$ is a torsion $R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$ -module. In particular, the multiplication map:

$$\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \xrightarrow{\mathbf{x}_{\tilde{a}}} \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i}$$

is injective. Then since $\text{Fitt}_B(M \otimes_A B) = \text{Fitt}_A(M)B$ for any Noetherian A -algebra B , letting $A = R_{W(\overline{\mathbb{F}})}$ and $B = R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$, we have

$$(8.1) \quad \left(\text{Fitt}_{R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \right) (R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}} = \left(\text{Fitt}_{R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / (\mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} + P_i^{e_i}) \right) \right)^{\text{rc}}.$$

Let $I := \text{Fitt}_{R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \right)$. Then we have a commutative square:

$$\begin{array}{ccc} I(R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) & \xrightarrow{\phi} & I^{\text{rc}}(R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \\ \parallel & & \parallel \\ \text{Fitt}_{R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \right) (R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) & \longrightarrow & \text{char}_{R_{W(\overline{\mathbb{F}})}}(M)(R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \end{array}$$

Applying Lemma 8.2 to the inclusion $I \rightarrow I^{\text{rc}}$, we have a set of height-two primes $\{Q_i\}_{1 \leq i \leq \ell}$. Therefore, the localization of ϕ at any height-one prime of $R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}$ is an isomorphism if and only if the following condition holds:

$$(8.2) \quad \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} \in \bigcap_{1 \leq i \leq \ell} \mathcal{L}_{W(\overline{\mathbb{F}})}(R_{W(\overline{\mathbb{F}})} / Q_i).$$

We conclude that if (8.2) is satisfied, then we have

$$(8.3) \quad \left(\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)(R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}} = \left(\text{Fitt}_{R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \right) (R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}}.$$

It also yields that

$$(8.4) \quad \left(\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)(R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}} = \text{char}_{R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})}} \left(\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})} / (\mathbf{x}_{\tilde{a}} R_{W(\overline{\mathbb{F}})} + P_i^{e_i}) \right).$$

We state the conclusion of the above discussion as a lemma.

Lemma 8.4. *Let the set-up be as in Discussion 8.3. Then the equation (8.1) holds true without any condition and if (8.2) is satisfied, then the equations (8.3) and (8.4) hold true.*

We prove a lemma which describes the set $\mathcal{L}_{W(\overline{\mathbb{F}})}(M)$.

Lemma 8.5. *Under the notation and the hypothesis as in Corollary 4.6, assume that M is a finitely generated torsion $R_{W(\overline{\mathbb{F}})}$ -module. Then the following assertions hold:*

(1) *The set $\mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ is identified with the intersection:*

$$\{\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})} \mid M/\mathbf{x}_{\bar{a}}M \text{ is a torsion } R_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}\text{-module}\} \cap \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{\text{null}}) \cap \bigcap_{1 \leq i \leq \ell} \mathcal{L}_{W(\overline{\mathbb{F}})}(R_{W(\overline{\mathbb{F}})}/Q_i),$$

where M_{null} is the maximal pseudo-null submodule of M and $\{Q_i\}_{1 \leq i \leq \ell}$ is a set of height-two primes attached to M as stated in Lemma 8.2.

(2) *Assume that N is a finitely generated pseudo-null $R_{W(\overline{\mathbb{F}})}$ -module and $\{Q'_i\}_{1 \leq i \leq k}$ is a set of all associated prime ideals of height two for the module N . Then we have:*

$$\mathcal{L}_{W(\overline{\mathbb{F}})}(N) = \bigcap_{1 \leq i \leq k} \mathcal{L}_{W(\overline{\mathbb{F}})}(R_{W(\overline{\mathbb{F}})}/Q'_i).$$

Proof. (1) This is taken from ([14], Lemma 3.4.), but we give its proof, as it requires some modifications. Let M_{null} be the maximal pseudo-null submodule of M . Then by Lemma 7.4, there is a fundamental pseudo-isomorphism $M \rightarrow \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})}/P_i^{e_i}$, together with the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M/M_{\text{null}} & \longrightarrow & \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})}/P_i^{e_i} & \longrightarrow & N \longrightarrow 0 \\ & & \mathbf{x}_{\bar{a}} \downarrow & & \mathbf{x}_{\bar{a}} \downarrow & & \mathbf{x}_{\bar{a}} \downarrow \\ 0 & \longrightarrow & M/M_{\text{null}} & \longrightarrow & \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})}/P_i^{e_i} & \longrightarrow & N \longrightarrow 0 \end{array}$$

for a (not necessarily redundant) set of height-one primes $\{P_i\}_{1 \leq i \leq m}$ of $R_{W(\overline{\mathbb{F}})}$ and N is a pseudo-null module.

First, we note that $\mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ is contained in the intersection as stated in the lemma. So we need to establish the other inclusion. Now assume that $\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}$ satisfies the following conditions:

- $M/\mathbf{x}_{\bar{a}}M$ is a torsion $R_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}$ -module.
- $\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})} \in \bigcap_{1 \leq i \leq \ell} \mathcal{L}_{W(\overline{\mathbb{F}})}(R_{W(\overline{\mathbb{F}})}/Q_i)$.

Then the set of all elements $\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}$ satisfying the above conditions contains $\mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ as a subset. So assuming that $\mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})}$ satisfies the above conditions, it suffices to prove the following implication:

$$(8.5) \quad \mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{\text{null}}) \Rightarrow \mathbf{x}_{\bar{a}}R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M).$$

We establish (8.5) below. By our choice, the multiplication map

$$\bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})}/P_i^{e_i} \xrightarrow{x_{\tilde{a}}} \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})}/P_i^{e_i}$$

is injective. So the snake lemma yields the following exact sequence:

$$(8.6) \quad 0 \rightarrow N[x_{\tilde{a}}] \rightarrow M/(x_{\tilde{a}}M + M_{\text{null}}) \rightarrow \bigoplus_{i=1}^m R_{W(\overline{\mathbb{F}})}/(x_{\tilde{a}}R_{W(\overline{\mathbb{F}})} + P_i^{e_i}) \rightarrow N/x_{\tilde{a}}N \rightarrow 0.$$

There is a short exact sequence:

$$(8.7) \quad 0 \rightarrow N[x_{\tilde{a}}] \rightarrow N \xrightarrow{x_{\tilde{a}}} N \rightarrow N/x_{\tilde{a}}N \rightarrow 0$$

of pseudo-null $R_{W(\overline{\mathbb{F}})}$ -modules, where both $N[x_{\tilde{a}}]$ and $N/x_{\tilde{a}}N$ are naturally $R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})}$ -modules. Localizing both (8.6) and (8.7) at all height-two primes P of $R_{W(\overline{\mathbb{F}})}$ containing $x_{\tilde{a}}$, a length computation for the sequence localized at P reveals that

$$(8.8) \quad \text{char}_{R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})}}(M/(x_{\tilde{a}}M + M_{\text{null}})) = \text{char}_{R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})}}\left(\bigoplus_{1 \leq i \leq m} R_{W(\overline{\mathbb{F}})}/(x_{\tilde{a}}R_{W(\overline{\mathbb{F}})} + P_i^{e_i})\right).$$

On the other hand, Lemma 8.4 shows that

$$(8.9) \quad \left(\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)(R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})})\right)^{\text{rc}} = \text{char}_{R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})}}\left(\bigoplus_{1 \leq i \leq m} R_{W(\overline{\mathbb{F}})}/(x_{\tilde{a}}R_{W(\overline{\mathbb{F}})} + P_i^{e_i})\right).$$

Then combining both (8.8) and (8.9), we get

$$(8.10) \quad \left(\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)(R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})})\right)^{\text{rc}} = \text{char}_{R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})}}(M/(x_{\tilde{a}}M + M_{\text{null}}))$$

Finally, since the multiplication on M/M_{null} by $x_{\tilde{a}}$ is injective, we have an exact sequence:

$$(8.11) \quad 0 \rightarrow M_{\text{null}}/x_{\tilde{a}}M_{\text{null}} \rightarrow M/x_{\tilde{a}}M \rightarrow M/(x_{\tilde{a}}M + M_{\text{null}}) \rightarrow 0.$$

Taking characteristic ideals to (8.11), in view of (8.10) and the condition **(B)** in Definition 8.1, it follows that the desired implication (8.5) holds if $M_{\text{null}}/x_{\tilde{a}}M_{\text{null}}$ is a pseudo-null $R_{W(\overline{\mathbb{F}})}/x_{\tilde{a}}R_{W(\overline{\mathbb{F}})}$ -module, which is true if $x_{\tilde{a}}R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{\text{null}})$. This completes the proof of (1).

(2) This part is done in ([14], Lemma 3.5 together with Lemma 3.1) in case $R_{W(\overline{\mathbb{F}})}$ is regular, so we leave the proof with necessary modifications to the reader. \square

The next purpose is to show that $\mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ is an infinite set for a finitely generated torsion $R_{W(\overline{\mathbb{F}})}$ -module M . We need to have sufficiently many specializations of $R_{W(\overline{\mathbb{F}})}$ that are normal in order to prove the control theorem by combining Proposition 5.1, Lemma 8.2, and Lemma 8.5.

Lemma 8.6. *Under the notation and the hypothesis as in Corollary 4.6, assume that M is a finitely generated torsion $R_{W(\overline{\mathbb{F}})}$ -module. Then we have the following assertions:*

- (1) The subset $\mathcal{L}_{W(\overline{\mathbb{F}})}(M) \subseteq \theta_{W(\overline{\mathbb{F}})}(U)$ may be identified with a non-empty open subset of $\mathbb{P}^d(\overline{\mathbb{F}})$ under the mapping $\theta_{W(\overline{\mathbb{F}})} : \mathbb{P}^d(\overline{\mathbb{F}}) \rightarrow \mathbb{P}^d(W(\overline{\mathbb{F}}))$. In particular, it is infinite.
- (2) Let P be a fixed height-one prime ideal appearing in $\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)$. Then one can find an infinite sequence $\{\mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}\}_{i \in \mathbb{N}} \subseteq \mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ such that the union:

$$\bigcup_{i \in \mathbb{N}} \text{Min}_{R_{W(\overline{\mathbb{F}})}}(P + \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})})$$

is an infinite set.

Proof. (1) By Lemma 8.5, we have $\mathbf{x}_{\tilde{a}} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{\text{null}})$ if and only if $\mathbf{x}_{\tilde{a}} \in \mathcal{L}_{W(\overline{\mathbb{F}})}$ is contained in none of height-two primes $Q'_1, \dots, Q'_k \in \text{Ass}_{R_{W(\overline{\mathbb{F}})}}(M_{\text{null}})$. Let Q_1, \dots, Q_ℓ be prime ideals attached to M as in Lemma 8.2. By the assumption that $\dim R \geq 3$, all these primes together with the set of height-one primes $P_1, \dots, P_m \in \text{Supp } M$ are strictly contained in the maximal ideal of $R_{W(\overline{\mathbb{F}})}$. So let us find a non-empty Zariski open subset of $\mathbb{P}^d(\overline{\mathbb{F}})$ with the required properties via Lemma 8.5. In other words, it suffices to choose $\mathbf{x}_{\tilde{a}}$ such that

$$(8.12) \quad \mathbf{x}_{\tilde{a}} \notin \left(\bigcup_{1 \leq i \leq m} P_i \right) \cup \left(\bigcup_{1 \leq i \leq k} Q'_i \right) \cup \left(\bigcup_{1 \leq i \leq \ell} Q_i \right).$$

Then applying Lemma 4.2 to each prime appearing in (8.12), we may find a Zariski open subset $U \subseteq \mathbb{P}^d(\overline{\mathbb{F}})$ such that the condition (8.12) holds for $a = (a_0 : \dots : a_d) \in \text{Sp}_{W(\overline{\mathbb{F}})}^{-1}(U)$. Then combining the conclusion of Proposition 5.1, we complete the proof of (1).

(2) The proof will be completed through inductive steps using (1) as follows. Let P be a fixed height-one prime ideal appearing in $\text{char}_{R_{W(\overline{\mathbb{F}})}}(M)$. Since P is a non-maximal prime ideal of $R_{W(\overline{\mathbb{F}})}$, we can attach a non-empty Zariski open subset of $\mathbb{P}^d(\overline{\mathbb{F}})$ by Lemma 4.2 and we may find $\mathbf{x}_{\tilde{a}_0} R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M)$, which is not contained in P . Then this initial choice satisfies our requirement.

Choose $\mathbf{x}_{\tilde{a}_1} R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ such that $\mathbf{x}_{\tilde{a}_1}$ is contained in no prime of the set:

$$\text{Min}_{R_{W(\overline{\mathbb{F}})}}(P + \mathbf{x}_{\tilde{a}_0} R_{W(\overline{\mathbb{F}})})$$

(which is a finite set of primes strictly contained in the maximal ideal of $R_{W(\overline{\mathbb{F}})}$ due to $\dim R \geq 3$). Next, choose $\mathbf{x}_{\tilde{a}_2} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M)$ such that $\mathbf{x}_{\tilde{a}_2}$ is in no prime ideals contained in the set:

$$\text{Min}_{R_{W(\overline{\mathbb{F}})}}(P + \mathbf{x}_{\tilde{a}_0} R_{W(\overline{\mathbb{F}})}) \cup \text{Min}_{R_{W(\overline{\mathbb{F}})}}(P + \mathbf{x}_{\tilde{a}_1} R_{W(\overline{\mathbb{F}})}).$$

By continuing this process, we will eventually obtain a sequence $\mathbf{x}_{\tilde{a}_0}, \mathbf{x}_{\tilde{a}_1}, \mathbf{x}_{\tilde{a}_2}, \dots$ with the required properties. \square

Let M be a finitely generated module over a complete local normal domain R with $W(\mathbb{F})$ its coefficient ring and let $W(\mathbb{F}) \rightarrow W(\mathbb{F}')$ be an extension induced by an algebraic

extension $\mathbb{F} \rightarrow \mathbb{F}'$. We set $M_{W(\mathbb{F}')} := M \widehat{\otimes}_{W(\mathbb{F})} W(\mathbb{F}')$, which coincides with our previous notation. Let M be a finitely generated torsion R -module. Then

$$\mathrm{char}_{R_{W(\mathbb{F}')}}(M_{W(\mathbb{F}')}) = \mathrm{char}_R(M)R_{W(\mathbb{F}')},$$

which may be verified directly from the definition.

Lemma 8.7. *Let M, N be finitely generated torsion R -modules. Then we have*

$$\mathrm{char}_R(M) \subseteq \mathrm{char}_R(N) \iff \mathrm{char}_{R_{W(\overline{\mathbb{F}})}}(M_{W(\overline{\mathbb{F}})}) \subseteq \mathrm{char}_{R_{W(\overline{\mathbb{F}})}}(N_{W(\overline{\mathbb{F}})}).$$

Proof. The implication \Rightarrow is obvious. So let us prove the other implication. Since $R \rightarrow R_{W(\mathbb{F})\mathrm{ur}}$ is ind-étale, it suffices to show that

$$\mathrm{char}_{R_{W(\mathbb{F})\mathrm{ur}}}(M) \subseteq \mathrm{char}_{R_{W(\mathbb{F})\mathrm{ur}}}(N) \iff \mathrm{char}_{R_{W(\overline{\mathbb{F}})}}(M_{W(\overline{\mathbb{F}})}) \subseteq \mathrm{char}_{R_{W(\overline{\mathbb{F}})}}(N_{W(\overline{\mathbb{F}})}).$$

Note that $R_{W(\mathbb{F})\mathrm{ur}} \rightarrow R_{W(\overline{\mathbb{F}})}$ is faithfully flat, with trivial residue field extension. Then the claim follows from this. \square

Now we prove the following theorem.

Theorem 8.8 (Control Theorem for Characteristic Ideals). *With the notation and the hypothesis as in Corollary 4.6, assume that M and N are finitely generated torsion R -modules. Then the following statements are equivalent:*

- (1) $\mathrm{char}_R(M) \subseteq \mathrm{char}_R(N)$.
- (2) For all but finitely many height-one primes:

$$\mathbf{x}_{\tilde{a}}R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{W(\overline{\mathbb{F}})}) \cap \mathcal{L}_{W(\overline{\mathbb{F}})}(N_{W(\overline{\mathbb{F}})}),$$

we have

$$\mathrm{char}_{R_{W(\mathbb{F}')}/\mathbf{x}_{\tilde{a}}R_{W(\mathbb{F}')}}(M_{W(\mathbb{F}')}/\mathbf{x}_{\tilde{a}}M_{W(\mathbb{F}')}) \subseteq \mathrm{char}_{R_{W(\mathbb{F}')}/\mathbf{x}_{\tilde{a}}R_{W(\mathbb{F}')}}(N_{W(\mathbb{F}')}/\mathbf{x}_{\tilde{a}}N_{W(\mathbb{F}')}),$$

where \mathbb{F}' is any finite field extension of \mathbb{F} depending on \tilde{a} such that $\mathbf{x}_{\tilde{a}} \in R_{W(\mathbb{F}')$.

- (3) For all but finitely many height-one primes:

$$\mathbf{x}_{\tilde{a}}R_{W(\overline{\mathbb{F}})} \in \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{W(\overline{\mathbb{F}})}) \cap \mathcal{L}_{W(\overline{\mathbb{F}})}(N_{W(\overline{\mathbb{F}})}),$$

we have

$$\mathrm{char}_{R_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\tilde{a}}R_{W(\overline{\mathbb{F}})}}(M_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\tilde{a}}M_{W(\overline{\mathbb{F}})}) \subseteq \mathrm{char}_{R_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\tilde{a}}R_{W(\overline{\mathbb{F}})}}(N_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\tilde{a}}N_{W(\overline{\mathbb{F}})}).$$

Note that $\dim R \geq 3$ holds automatically, due to the hypothesis $\mathrm{depth} R \geq 3$.

Proof. The implications (1) \Rightarrow (2) \Rightarrow (3) are obvious in view of Definition 8.1. So it remains to prove (3) \Rightarrow (1). By Lemma 8.7, it suffices to show that

$$\mathrm{char}_{R_{W(\overline{\mathbb{F}})}}(M_{W(\overline{\mathbb{F}})}) \subseteq \mathrm{char}_{R_{W(\overline{\mathbb{F}})}}(N_{W(\overline{\mathbb{F}})}).$$

Take fundamental pseudo-isomorphisms for M and N :

$$M \rightarrow \bigoplus_i R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \quad (\text{resp. } N \rightarrow \bigoplus_j R_{W(\overline{\mathbb{F}})} / Q_j^{f_j})$$

for a (not necessarily redundant) finite set of height-one primes $\{P_i\}$ (resp. $\{Q_j\}$) of $R_{W(\overline{\mathbb{F}})}$. Put $I_M := (\prod_i P_i^{e_i})^{\text{rc}}$ and $I_N := (\prod_j Q_j^{f_j})^{\text{rc}}$ and the condition **(B)** in Definition 8.1 allows one to assume that

$$(8.13) \quad M = \bigoplus_i R_{W(\overline{\mathbb{F}})} / P_i^{e_i} \quad (\text{resp. } N = \bigoplus_j R_{W(\overline{\mathbb{F}})} / Q_j^{f_j}).$$

To simplify the notation, assume that $\{P_i\}$ (resp. $\{Q_j\}$) is a redundant set of prime ideals and all relevant modules are defined over $R_{W(\overline{\mathbb{F}})}$. We require several steps to complete the proof of the theorem. Let

$$\{\mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}\}_{i \in \mathbb{N}} \subseteq \mathcal{L}_{W(\overline{\mathbb{F}})}(M_{W(\overline{\mathbb{F}})}) \cap \mathcal{L}_{W(\overline{\mathbb{F}})}(N_{W(\overline{\mathbb{F}})})$$

be any infinite sequence of distinct primes of $R_{W(\overline{\mathbb{F}})}$ satisfying the condition (3). In particular, we have $\bigcap_{i \in \mathbb{N}} \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})} = 0$.

Step1: In this step, we establish $\text{Supp}_{\text{ht}=1} N \subseteq \text{Supp}_{\text{ht}=1} M$, where $\text{Supp}_{\text{ht}=1}(-)$ is the set of height-one primes contained in the support of a module. By assumption, we have

$$\left(I_M (R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}} \subseteq \left(I_N (R_{W(\overline{\mathbb{F}})} / \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}) \right)^{\text{rc}}$$

for all $i \in \mathbb{N}$. Rewriting this inclusion, we get

$$\left((I_M + \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}) / \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})} \right)^{\text{rc}} \subseteq \left((I_N + \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}) / \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})} \right)^{\text{rc}}.$$

From this description, we deduce the following fact. Fix a height-one prime ideal Q_k from (8.13). We may choose the set $\{\mathbf{x}_{\tilde{a}_i}\}_{i \in \mathbb{N}}$ such that

$$(8.14) \quad \bigcup_{i \in \mathbb{N}} \text{Min}_{R_{W(\overline{\mathbb{F}})}}(Q_k + \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})})$$

is an infinite set in view of Lemma 8.6. For every fixed $i \in \mathbb{N}$, we have

$$I_M \subseteq \mathfrak{p}_i,$$

where $\mathfrak{p}_i \in \text{Min}_{R_{W(\overline{\mathbb{F}})}}(Q_k + \mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})})$ is chosen to be an arbitrary fixed element.

On the other hand, since (8.14) is infinite, we may find an infinite subset $\{\mathfrak{p}_i\}_{i \in \mathbb{N}}$ of (8.14) and we fix it once and for all. Since $R_{W(\overline{\mathbb{F}})} / Q_k$ is an integral domain and $\{\mathfrak{p}_i\}_{i \in \mathbb{N}}$ is an infinite set of height-two primes containing Q_k , it follows from Lemma 5.3 that

$$Q_k = \bigcap_{i \in \mathbb{N}} \mathfrak{p}_i.$$

Since $I_M \subseteq \mathfrak{p}_i$ for all $i \in \mathbb{N}$, we have $I_M \subseteq Q_k$. Since Q_k is arbitrary, $I_M \subseteq (\prod_j Q_j)^{\text{rc}}$, or equivalently, $\text{Supp}_{\text{ht}=1} N \subseteq \text{Supp}_{\text{ht}=1} M$.

Step2: In this step, we deal with multiplicities of divisors in the characteristic ideal and we complete this step by induction on the number of divisors appearing in I_M .

First, assume $I_M = (P^e)^{\text{rc}}$ for $e \geq 1$. Then we have $\text{Supp}_{\text{ht}=1} N = \emptyset$ or $\{P\}$ because of **Step1**. If $\text{Supp}_{\text{ht}=1} N = \emptyset$, there is nothing to prove. So assume $\text{Supp}_{\text{ht}=1} N = \{P\}$. Both M and N are assumed to be fundamental torsion $R_{W(\overline{\mathbb{F}})}$ -modules, thus $M[\mathbf{x}_{\tilde{a}_i}]$ and $N[\mathbf{x}_{\tilde{a}_i}]$ are trivial modules and Proposition 7.5 yields that $\ell_{(R_{W(\overline{\mathbb{F}})})_P}(M_P) \geq \ell_{(R_{W(\overline{\mathbb{F}})})_P}(N_P)$.

In the general case, we prove by contradiction and thus, assume that $I_M \not\subseteq I_N$. Then this implies that we have $e_k < f_k$ for some k , where e_k, f_k are coming from (8.13). Put

$$(8.15) \quad \tilde{I}_M := (P_k^{-e_k} \cdot I_M)^{\text{rc}} \quad (\text{resp. } \tilde{I}_N := (P_k^{-e_k} \cdot I_N)^{\text{rc}}),$$

which are both integral reflexive ideals. There are short exact sequences:

$$0 \rightarrow \tilde{I}_M/I_M \rightarrow R_{W(\overline{\mathbb{F}})}/I_M \rightarrow R_{W(\overline{\mathbb{F}})}/\tilde{I}_M \rightarrow 0$$

and

$$0 \rightarrow \tilde{I}_N/I_N \rightarrow R_{W(\overline{\mathbb{F}})}/I_N \rightarrow R_{W(\overline{\mathbb{F}})}/\tilde{I}_N \rightarrow 0$$

and it is clear that

$$\text{char}_{R_{W(\overline{\mathbb{F}})}}(\tilde{I}_M/I_M) = \text{char}_{R_{W(\overline{\mathbb{F}})}}(\tilde{I}_N/I_N) = (P_k^{e_k})^{\text{rc}},$$

which induces the following short exact sequences by the snake lemma:

$$0 \rightarrow \tilde{I}_M/(I_M, \mathbf{x}_{\tilde{a}_i} \tilde{I}_M) \rightarrow R_{W(\overline{\mathbb{F}})}/(I_M, \mathbf{x}_{\tilde{a}_i}) \rightarrow R_{W(\overline{\mathbb{F}})}/(\tilde{I}_M, \mathbf{x}_{\tilde{a}_i}) \rightarrow 0$$

and

$$0 \rightarrow \tilde{I}_N/(I_N, \mathbf{x}_{\tilde{a}_i} \tilde{I}_N) \rightarrow R_{W(\overline{\mathbb{F}})}/(I_N, \mathbf{x}_{\tilde{a}_i}) \rightarrow R_{W(\overline{\mathbb{F}})}/(\tilde{I}_N, \mathbf{x}_{\tilde{a}_i}) \rightarrow 0.$$

Taking characteristic ideals, we get from the condition (3) that

$$\text{char}_{R_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}}(R_{W(\overline{\mathbb{F}})}/(\tilde{I}_M, \mathbf{x}_{\tilde{a}_i})) \subseteq \text{char}_{R_{W(\overline{\mathbb{F}})}/\mathbf{x}_{\tilde{a}_i} R_{W(\overline{\mathbb{F}})}}(R_{W(\overline{\mathbb{F}})}/(\tilde{I}_N, \mathbf{x}_{\tilde{a}_i})).$$

Since the number of primes ideals in $\text{Ass}_{R_{W(\overline{\mathbb{F}})}}(R_{W(\overline{\mathbb{F}})}/\tilde{I}_M)$ is exactly one less than that of components of prime ideals in I_M , the induction hypothesis on \tilde{I}_M yields that

$$\text{char}_{R_{W(\overline{\mathbb{F}})}}(R_{W(\overline{\mathbb{F}})}/\tilde{I}_M) \subseteq \text{char}_{R_{W(\overline{\mathbb{F}})}}(R_{W(\overline{\mathbb{F}})}/\tilde{I}_N).$$

However, we deduce from these observations and (8.15) that $I_M \subseteq I_N$, which is a contradiction to our assumption $I_M \not\subseteq I_N$. Hence, we obtain $I_M \subseteq I_N$, as desired. \square

Remark 8.9. It is worth pointing out that Theorem 8.8 holds for complete local normal rings of mixed characteristic with arbitrary infinite perfect residue field as well. More precisely, it can be proven that $\text{char}_R(M) \subseteq \text{char}_R(N) \iff \text{char}_{R/xR}(M/xM) \subseteq \text{char}_{R/xR}(N/xN)$ for sufficiently many $x \in R$.

In this article, we presented an application of the local Bertini theorem to characteristic ideals. However, we believe that the main theorem has more interesting applications such as the study of the restriction map on divisor class (Chow) groups.

9. APPENDIX

In this appendix, we study the relationship between Fitting ideals and characteristic ideals. For Fitting ideals, we refer the reader to Northcott's book [13], but we review the basic part of the theory. For reflexive sheaves on normal schemes, we refer te reader to [9]. Throughout, we assume that R is a Noetherian ring and M is a finitely generated R -module.

Definition 9.1 (Fitting ideal). Let the notation be as above and assume that

$$F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$$

is a finite free resolution of the R -module M , where the mapping $F_1 \rightarrow F_0$ is defined via a $m \times n$ -matrix X with $\text{rank}(F_1) = n$ and $\text{rank}(F_0) = m$. Then $\text{Fitt}_R(M)$ is defined as an ideal of R generated by all m -minors of X .

The Fitting ideal does not depend on the choice of a free resolution and it enjoys the following properties.

Proposition 9.2. *Let M be a finitely generated module over a Noetherian ring R . Then we have the following properties.*

- (1) *Let $I \subseteq R$ be an ideal. Then $\text{Fitt}_R(R/I) = I$.*
- (2) *Let S be any Noetherian R -algebra. Then $\text{Fitt}_S(M \otimes_R S) = \text{Fitt}_R(M)S$.*
- (3) *Let $\text{Ann}_R(M)$ be the annihilator of the R -module M . Then*

$$\text{Fitt}_R(M) \subseteq \text{Ann}_R(M).$$

- (4) *If $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is a short exact sequence of R -modules, then*

$$\text{Fitt}_R(L) \cdot \text{Fitt}_R(N) \subseteq \text{Fitt}_R(M).$$

- (5) *Assume that R is a discrete valuation ring with its uniformizing parameter b and M is a torsion R -module. Then $\text{Fitt}_R(M) = (b)^{\ell_R(M)}$.*

Proof. These facts are all well known. For (5), it simply follows from the elementary divisors of modules over a principal ideal domain. \square

For a Noetherian domain R and an R -module M , let $M^* := \text{Hom}_R(M, R)$, the dual of M . We say that $M^{\text{rc}} := (M^*)^*$ is the *reflexive closure* of M . Then we have the following lemma.

Lemma 9.3. *Let R be a Noetherian domain and let I be a fractional ideal of R . Then the reflexive closure I^{rc} is naturally regarded as a fractional ideal of R .*

Proof. By assumption, there exists $\alpha \in R$ such that $I \simeq \alpha \cdot I \subseteq R$. Let $J := \alpha \cdot I$, an ideal of R . The short exact sequence $0 \rightarrow J \rightarrow R \rightarrow R/J \rightarrow 0$ induces a short exact sequence

$$0 = \text{Hom}_R(R/J, R) \rightarrow R \rightarrow \text{Hom}_R(J, R) \rightarrow N \rightarrow 0,$$

with $N \subseteq \text{Ext}_R^1(R/J, R)$ cokernel of $R \rightarrow \text{Hom}_R(J, R)$. Then applying $\text{Hom}_R(-, R)$ twice, we get an exact sequence

$$0 = \text{Hom}_R(N, R) \rightarrow J^{\text{rc}} \rightarrow R,$$

because $J \cdot N = 0$. This implies that J^{rc} is an ideal of R . Then $I^{\text{rc}} = \alpha^{-1} \cdot J^{\text{rc}}$ is a fractional ideal. \square

Note that even when I and J are reflexive, the product $I \cdot J$ need not be reflexive. Any principal ideal is reflexive. Let I be an ideal of a normal domain R . Then we recall the following fact:

$$I^{\text{rc}} = \bigcap_P I_P,$$

where P ranges over all height-one primes of R . The natural inclusion $I \rightarrow I^{\text{rc}}$ is a pseudo-isomorphism, since $I_P \rightarrow (I^{\text{rc}})_P = (I_P)^{\text{rc}}$ for every height-one prime $P \subseteq R$ and every ideal in a discrete valuation ring is principal. The following lemma explains the naturality of reflexive ideals and gives a way to investigate the inclusion relation between characteristic ideals.

Lemma 9.4. *Let R be a Noetherian normal domain and let I and J be reflexive ideals. Then $I \subseteq J$ if and only if $v_P(I) \geq v_P(J)$ for a valuation v attached to every height-one prime P of R . In particular, the only reflexive integral ideal containing a prime ideal of R properly is R itself.*

If R is only assumed to be Cohen-Macaulay, a similar result holds for invertible modules ([3], Lemma 5.3). We defined characteristic ideals as reflexive ideals and this is natural from the viewpoint of Iwasawa's main conjecture, because the most interesting arithmetic information may be captured at height-one primes. For finitely generated torsion R -modules M, N , it follows from the above lemma that $\text{char}_R(M) \subseteq \text{char}_R(N)$ if and only if $\text{char}_R(M)_P \subseteq \text{char}_R(N)_P$ for every height-one prime $P \in \text{Supp}(M) \cup \text{Supp}(N)$.

Example 9.5. Suppose I is reflexive and let $\mathfrak{a} \subseteq R$ be such that R/\mathfrak{a} is a normal domain. Then $I(R/\mathfrak{a})$ need not be reflexive. For a general ideal $I \subseteq R$, it can happen that

$$(I(R/\mathfrak{a}))^{\text{rc}} \neq (I^{\text{rc}}(R/\mathfrak{a}))^{\text{rc}}.$$

Here, I^{rc} is the reflexive closure with respect to R and $(I(R/\mathfrak{a}))^{\text{rc}}$ is the reflexive closure with respect to R/\mathfrak{a} . Let us take a look at the following simple example. Take $R =$

$\mathbb{Z}_p[[x, y]]$, $I = (x, y)$, and $\mathfrak{a} = (x)$. Then $(I^{\text{rc}}(R/\mathfrak{a}))^{\text{rc}} = R/xR$, since there is no height-one prime of R containing I . But $(I(R/\mathfrak{a}))^{\text{rc}} = y(R/xR)$.

Proposition 9.6. *Let R be a Noetherian normal domain. Then the following hold.*

(1) *Let M be a finitely generated torsion R -module. Then we have*

$$\text{char}_R(M) = \left(\prod_{\text{ht } \mathfrak{p}=1} \mathfrak{p}^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} \right)^{\text{rc}} = \text{Fitt}_R(M)^{\text{rc}}.$$

In particular, $\text{Fitt}_R(M) \subseteq \text{char}_R(M)$ and if R is a UFD, then

$$\text{Fitt}_R(M) \subseteq \prod_{\text{ht } \mathfrak{p}=1} \mathfrak{p}^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} = \text{char}_R(M).$$

(2) *Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be a short exact sequence of finitely generated R -modules. Then*

$$\text{char}_R(M) = (\text{char}_R(L) \cdot \text{char}_R(N))^{\text{rc}}.$$

Proof. (1) Since the characteristic ideal is reflexive, the first equality follows by taking localization at all height-one primes of R . The second equality follows from the fact

$$\text{Fitt}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) = (PR_{\mathfrak{p}})^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})}$$

for any height-one prime ideal $P \subseteq R$. The second assertion is due to the fact that a height-one prime in a UFD is principal.

(2) This is clear, since the length is additive with respect to short exact sequences. The equality continues to hold true without torsion property of modules. \square

Example 9.7. The ordinary power of a height-one prime in a normal domain is not necessarily reflexive. Here is an example. Let $R = \mathbb{Z}_p[[x^2, xy, y^2]]$ and let $\mathfrak{p} = (x^2, xy)$. Then R is a normal domain and $\text{ht}(\mathfrak{p}) = 1$. Then $\mathfrak{p}^2 = (x^4, x^3y, x^2y^2)$ and $\mathfrak{p}^2 \neq \mathfrak{p}^{(2)}$. In fact, $x^2 \notin \mathfrak{p}^2$, but $\mathfrak{p}^{(2)} = (x^2)$.

Now let $\mathfrak{q} = (xy, y^2)$ and $M = R/(\mathfrak{p} \cap \mathfrak{q})$, which is torsion over R . Then one verifies that

$$\text{Fitt}_R(M) = \mathfrak{p} \cap \mathfrak{q} \not\subseteq \prod_{\text{ht } \mathfrak{p}=1} \mathfrak{p}^{\ell_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})} = \mathfrak{p}\mathfrak{q},$$

which tells us that Proposition 9.6 (1) is the most optimal.

Acknowledgement. The authors thank the anonymous referees for reading the manuscript carefully. The second-named author thanks Prof. Trivedi for explaining the proof of a lemma in [4].

REFERENCES

- [1] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge University Press. **39** (1993).
- [2] D. Eisenbud and E. G. Evans, *Generating modules efficiently; theorems from algebraic K-theory*, *J. Algebra* **49** (1977) 276–303.
- [3] M. Flach, *The equivariant Tamagawa number conjectures: a survey*, *Contemp. Math.* **358** (2004), 79–125.
- [4] H. Flenner, *Die Sätze von Bertini für lokale Ringe*, *Math. Ann.* **229** (1977), 97–111.
- [5] J. Fresnel and M. van der Put, *Rigid Analytic Geometry and Its Applications*, Birkhäuser, Boston (2003).
- [6] S. Greco, *Two theorems on excellent rings*, *Nagoya Math. J.* **60** (1976), 139–149.
- [7] A. Grothendieck, *Éléments de Géométrie Algébrique IV*, *Publications Math. I.H.E.S.* **24** (1965).
- [8] A. Grothendieck, *Cohomologie locale des faisceaux cohérents et théorèmes des Lefschetz locaux et globaux*, (SGA2) Amsterdam: North-Holland, (1968).
- [9] R. Hartshorne, *Stable reflexive sheaves*, *Math. Ann.* **254** (1980), 121–176.
- [10] E. Kunz, *Kähler Differentials*, *Advanced Lectures in Mathematics*, Vieweg+Teubner Verlag (1986).
- [11] H. Matsumura, *Commutative ring theory*, Cambridge University Press. **8** (1986).
- [12] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*, *Grundlehren der Mathematischen Wissenschaften.* **323**, Berlin, Springer (2000).
- [13] D. G. Northcott, *Finite Free Resolutions*, Cambridge University Press (Cambridge Tracts in Mathematics) **71** (1976).
- [14] T. Ochiai, *Euler system for Galois deformations*, *Annales de l’Institut Fourier*, **55**, fascicule **1** (2005), 113–146.
- [15] T. Ochiai and K. Shimomoto, in preparation.
- [16] G. Scheja and U. Storch, *Differentielle Eigenschaften der Lokalisierungen analytischer Algebren*, *Math. Ann.* **197** (1972), 137–170.
- [17] J.-P. Serre, *Local Fields*, *Graduate Texts in Mathematics* **67**, Springer-Verlag, New York (1973).
- [18] C. Skinner and E. Urban, *The Main Conjecture for $GL(2)$* , *Invent. Math.* **195** (2014), 1–277.
- [19] V. Trivedi, *A local Bertini theorem in mixed characteristic*, *Comm. in Algebra* **22** (1994), 823–827.

GRADUATE SCHOOL OF SCIENCE OSAKA UNIVERSITY 1-1 MACHIKANNEYAMA TOYONAKA OSAKA 560-0043 JAPAN

E-mail address: ochiai@math.sci.osaka-u.ac.jp

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY MEIJI UNIVERSITY 1-1-1 HIGASHIMITA TAMA-KU KAWASAKI 214-8571 JAPAN

E-mail address: shimomotokazuma@gmail.com