

# A bound on the number of points of a curve in projective space over a finite field

Masaaki Homma \*

Department of Mathematics, Kanagawa University  
Yokohama 221-8686, Japan  
homma@kanagawa-u.ac.jp

## Abstract

For a nondegenerate irreducible curve  $C$  of degree  $d$  in  $\mathbb{P}^r$  over  $\mathbb{F}_q$  with  $r \geq 3$ , we prove that the number  $N_q(C)$  of  $\mathbb{F}_q$ -points of  $C$  satisfies the inequality  $N_q(C) \leq (d-1)q+1$ , which is known as Sziklai's bound if  $r = 2$ .

*Key Words:* Curve, Rational Point, Finite field, Sziklai's bound

*MSC:* 14G15, 11G20, 14H25

## 1 Introduction

In the series of papers [3, 4, 5], we proved that for any plane curve  $C$  of degree  $d$  over  $\mathbb{F}_q$  without  $\mathbb{F}_q$ -linear components, the number  $N_q(C)$  of  $\mathbb{F}_q$ -points of  $C$  is bounded by

$$N_q(C) \leq (d-1)q+1 \quad (1)$$

except for the curve over  $\mathbb{F}_4$  defined by

$$K : (X+Y+Z)^4 + (XY+YZ+ZX)^2 + XYZ(X+Y+Z) = 0.$$

Indeed,  $N_4(K) = 14$ .

The bound (1) was originally conjectured by Sziklai [8], and he found that some curves actually achieve this bound.

The question we are interesting in is whether the bound (1) is valid for curves in higher dimensional projective space.

**Theorem 1.1** *Let  $C$  be an absolutely irreducible curve of degree  $d$  defined over  $\mathbb{F}_q$  in  $\mathbb{P}^r$  with  $r \geq 3$ , which is not contained in any planes. Then*

$$N_q(C) \leq (d-1)q+1.$$

---

\*Partially supported by Grant-in-Aid for Scientific Research (21540051), JSPS.

The main ingredient of our proof of this theorem is the order-sequence<sup>1</sup> of a projective curve like Stöhr-Voloch theory [7], however, our bound does not involve the genus of the curve.

As a corollary of this theorem, we have the following fact.

**Corollary 1.2** *Let  $C$  be a curve, which may have several components, of degree  $d$  in  $\mathbb{P}^r$  over  $\mathbb{F}_q$  without  $\mathbb{F}_q$ -linear components. In addition, when  $q = d = 4$ ,  $C$  is not a planar curve which is isomorphic to  $K$  over  $\mathbb{F}_4$ . Then*

$$N_q(C) \leq (d - 1)q + 1.$$

Throughout this paper,  $C(\mathbb{F}_q)$  denotes the set of  $\mathbb{F}_q$ -points of  $C$ , in other words,  $C(\mathbb{F}_q) = C \cap \mathbb{P}^r(\mathbb{F}_q)$ , where  $\mathbb{P}^r(\mathbb{F}_q)$  is the set of  $\mathbb{F}_q$ -points of  $\mathbb{P}^r$ .

## 2 Combinatorial approach

We regard  $\mathbb{P}^r(\mathbb{F}_q)$  as the  $r$ -dimensional finite projective space over  $\mathbb{F}_q$ .

**Definition 2.1** Suppose  $r \geq 2$ . For a subset  $X \subset \mathbb{P}^r(\mathbb{F}_q)$ , the s-degree<sup>2</sup> of  $X$  is the maximum number of points of  $X$  that lie on a hyperplane of  $\mathbb{P}^r(\mathbb{F}_q)$ . The s-degree of  $X$  is denoted by s-deg  $X$ .

The total number of points of  $X$  is denoted by  $N$ . If  $r = 2$  and s-deg  $X = d$ ,  $X$  is called an  $(N, d)$ -arc [1, (12.1)].

In the following lemma,  $\lfloor \alpha \rfloor$  denotes the integer part of a real number  $\alpha$ .

**Proposition 2.2** *For  $X \subset \mathbb{P}^r(\mathbb{F}_q)$  of s-degree  $d$ , the cardinality  $N$  of  $X$  is bounded by*

$$N \leq (d - 1)q + 1 + \left\lfloor \frac{d - 1}{q^{r-2} + q^{r-3} + \dots + q + 1} \right\rfloor. \quad (2)$$

*Proof.* Fix a point  $P_0 \in X$ . Let  $\check{P}_0 = \{H \in \check{\mathbb{P}}^r(\mathbb{F}_q) \mid P_0 \in H\}$ , where  $\check{\mathbb{P}}^r(\mathbb{F}_q)$  denotes the set of hyperplanes of  $\mathbb{P}^r(\mathbb{F}_q)$ . Let

$$\mathcal{P} = \{(P, H) \in (X \setminus \{P_0\}) \times \check{P}_0 \mid P \in H\}.$$

Moreover,  $\pi_1 : \mathcal{P} \rightarrow X \setminus \{P_0\}$  denotes the first projection and  $\pi_2 : \mathcal{P} \rightarrow \check{P}_0$  the second projection.

Let  $P \in X \setminus \{P_0\}$ . Since  $\pi_1^{-1}(P)$  is the set of hyperplanes that contain the line  $P_0P$ ,  $\#\pi_1^{-1}(P) = q^{r-2} + q^{r-3} + \dots + 1$ . Hence

$$\#\mathcal{P} = \sum_{P \in X \setminus \{P_0\}} \#\pi_1^{-1}(P) = (N - 1)(q^{r-2} + q^{r-3} + \dots + 1).$$

---

<sup>1</sup> As for the definition and the basic properties of order-sequence, see [2, 7.6]

<sup>2</sup>This jargon is an abbreviation for ‘set-theoretic degree’. We want to reserve the simple terminology ‘degree’ for the degree of a curve.

On the other hand, since  $\text{s-deg } X = d$ ,  $\#(H \cap (X \setminus \{P_0\})) \leq d - 1$  for any  $H \in \check{P}_0$ . Hence

$$\#\mathcal{P} \leq (d - 1)\#\check{P}_0 = (d - 1)(q^{r-1} + q^{r-2} + \cdots + 1).$$

Therefore

$$\begin{aligned} N &\leq (d - 1)\frac{q^{r-1} + \cdots + 1}{q^{r-2} + \cdots + 1} + 1 \\ &= (d - 1)q + 1 + \frac{d - 1}{q^{r-2} + \cdots + 1}. \end{aligned}$$

This completes the proof.  $\square$

**Remark 2.3** When  $r = 2$ , the bound (2) is rather trivial, that is,  $N \leq (d - 1)q + d$  (see [1, (12.5)]).

### 3 Number of points of a nondegenerate irreducible curve

In this section, we consider an irreducible curve  $C$  in  $\mathbb{P}^r$  with  $r \geq 3$  defined over  $\mathbb{F}_q$ . Moreover we assume  $C$  to be nondegenerate, that is, no hyperplane of  $\mathbb{P}^r$  contains  $C$ . For a point  $P \in C$  and a hyperplane  $H$  of  $\mathbb{P}^r$  with  $H \ni P$ , let  $h$  be a local equation of  $H$  around  $P$ . Under this situation,  $V(h)$  denotes the hyperplane  $H$ . The intersection multiplicity  $i(H.C; P)$  of  $C$  with  $H$  at  $P$  is

$$i(H.C; P) = \dim \mathcal{O}_{P,C}/(\bar{h}),$$

where  $\bar{h}$  is the image of  $h$  in the local ring  $\mathcal{O}_{P,C}$  of  $P \in C$ .

**Lemma 3.1** For a point  $P \in \mathbb{P}^r(\mathbb{F}_q)$ ,

$$\sum_{H \in \check{P}} (i(H.C; P) - 1) \geq \frac{q^{r-1} + q^{r-2} + \cdots + q + 1 - r}{q - 1},$$

where  $\check{P}$  is the set of  $\mathbb{F}_q$ -hyperplanes passing through  $P$ .

*Proof.* First suppose  $P$  is a nonsingular point of  $C$ . Without loss of generality, we may assume that  $P = (1, 0, \dots, 0)$ . Let  $x_1, \dots, x_r$  be a system of affine coordinate functions around  $P$  with  $x_1(P) = \cdots = x_r(P) = 0$ , each of which is defined over  $\mathbb{F}_q$ . Then  $\check{P} = \{V(\alpha_1 x_1 + \cdots + \alpha_r x_r) \mid (\alpha_1, \dots, \alpha_r) \in \mathbb{P}^{r-1}(\mathbb{F}_q)\}$ . We choose a local parameter  $t$  at  $P \in C$  which is defined over  $\mathbb{F}_q$ . Through the identification  $\hat{\mathcal{O}}_{P,C} = \bar{\mathbb{F}}_q[[t]]$ ,  $x_i$  can be written as

$$x_i = a_{i1}t + a_{i2}t^2 + \cdots \quad (i = 1, 2, \dots, r)$$

in  $\bar{\mathbb{F}}_q[[t]]$ , where  $a_{ij} \in \mathbb{F}_q$ . Applying elementary row-operations over  $\mathbb{F}_q$  to  $(a_{ij})_{\substack{i=1,2,\dots,r \\ j=1,2,\dots}}$ , we have the following form:

$$\begin{cases} x'_1 = t^{j_1} + \dots & \dots & \dots \\ \vdots & & \\ x'_i = & t^{j_i} + \dots & \dots \\ \vdots & & \\ x'_r = & & t^{j_r} + \dots \end{cases},$$

where  $0 < j_1 = 1 < j_2 < \dots < j_r$  and the  $\mathbb{F}_q$ -vector space spanned by  $x'_1, \dots, x'_r$  is the original space spanned by  $x_1, \dots, x_r$ . By using this renewed system of affine coordinate functions around  $P$ , we have a filtration  $\check{P} = V_1 \supset V_2 \supset V_3 \supset \dots \supset V_r$ , where

$$V_i = \{V(\alpha_i x'_i + \dots + \alpha_r x'_r) \mid (\alpha_i, \dots, \alpha_r) \in \mathbb{P}^{r-i}(\mathbb{F}_q)\}.$$

If  $H = V(h) \in V_i \setminus V_{i+1}$ , then  $h = \alpha_i t^{j_i} + \dots$  with  $\alpha_i \neq 0$ . Hence  $i(H.C; P) = j_i \geq i$ . Therefore

$$\begin{aligned} \sum_{H \in \check{P}} (i(H.C; P) - 1) &\geq \sum_{i=1}^r (i-1)q^{r-i} \\ &= \frac{q^{r-1} + q^{r-2} + \dots + 1 - r}{q-1}. \end{aligned}$$

Secondly, suppose  $P$  is a singular point of  $C$ . Hence  $i(H.C; P) \geq 2$  for any  $H \in \check{P}$ . Therefore

$$\begin{aligned} \sum_{H \in \check{P}} (i(H.C; P) - 1) &\geq \#\check{P} = q^{r-1} + q^{r-2} + \dots + 1 \\ &> \frac{q^{r-1} + q^{r-2} + \dots + 1 - r}{q-1}. \end{aligned}$$

This completes the proof.  $\square$

**Theorem 3.2** *Let  $C$  be a nondegenerate irreducible curve of degree  $d$  in  $\mathbb{P}^r$  over  $\mathbb{F}_q$ . Then*

$$N_q(C) \leq \frac{(q-1)(q^r + q^{r-1} + \dots + 1)}{q^r + q^{r-1} + \dots + q - r} d = \frac{(q-1)(q^{r+1} - 1)}{q(q^r - 1) - r(q-1)} d.$$

*Proof.* Let us consider the point-hyperplane correspondence with respect to  $C$  over  $\mathbb{F}_q$ :

$$\mathcal{Q} := \{(P, H) \in C(\mathbb{F}_q) \times \check{\mathbb{P}}^r(\mathbb{F}_q) \mid P \in H\}.$$

Let  $\pi_1 : \mathcal{Q} \rightarrow C(\mathbb{F}_q)$  and  $\pi_2 : \mathcal{Q} \rightarrow \check{\mathbb{P}}^r(\mathbb{F}_q)$  be the first and second projections respectively. If  $H \in \pi_2(\mathcal{Q}) \subseteq \check{\mathbb{P}}^r(\mathbb{F}_q)$ , then  $\pi_2^{-1}(H) = (H \cap C(\mathbb{F}_q)) \times \{H\}$ , and  $d - \sum_{P \in H \cap C(\mathbb{F}_q)} i(H.C; P) \geq d - (H.C) = 0$ . Hence

$$\begin{aligned} \#\pi_2^{-1}(H) &\leq \#(H \cap C(\mathbb{F}_q)) + d - \sum_{P \in H \cap C(\mathbb{F}_q)} i(H.C; P) \\ &= d - \sum_{P \in H \cap C(\mathbb{F}_q)} (i(H.C; P) - 1). \end{aligned}$$

Hence

$$\begin{aligned} \#\mathcal{Q} &= \sum_{H \in \pi_2(\mathcal{Q})} \#\pi_2^{-1}(H) \\ &\leq \sum_{H \in \pi_2(\mathcal{Q})} \left( d - \sum_{P \in H \cap C(\mathbb{F}_q)} (i(H.C; P) - 1) \right) \\ &\leq d(q^r + q^{r-1} + \cdots + 1) - \sum_{P \in C(\mathbb{F}_q)} \sum_{H \in \check{P}} (i(H.C; P) - 1) \quad (3) \end{aligned}$$

because  $\#\pi_2(\mathcal{Q}) \leq q^r + q^{r-1} + \cdots + 1$  and

$$\mathcal{Q} = \bigsqcup_{H \in \pi_2(\mathcal{Q})} \left( \bigsqcup_{P \in H \cap C(\mathbb{F}_q)} \{(P, H)\} \right) = \bigsqcup_{P \in C(\mathbb{F}_q)} \left( \bigsqcup_{H \in \check{P}} \{(P, H)\} \right).$$

Applying Lemma 3.1 to (3), we have

$$\#\mathcal{Q} \leq d(q^r + q^{r-1} + \cdots + 1) - \frac{q^{r-1} + q^{r-2} + \cdots + 1 - r}{q - 1} N_q(C). \quad (4)$$

On the other hand,

$$\#\mathcal{Q} = \sum_{P \in C(\mathbb{F}_q)} \#\pi_1^{-1}(P) = (q^{r-1} + q^{r-2} + \cdots + 1) N_q(C). \quad (5)$$

From (4) and (5), we have the desired bound for  $N_q(C)$ .  $\square$

**Corollary 3.3** *Under the same assumption as Theorem 3.2,*

$$N_q(C) \leq (q - 1)d + \frac{r + 1}{q^{r-1} + 2q^{r-2} + \cdots + (r - 1)q + r} d.$$

*Proof.* Let  $N = N_q(C)$ . By Theorem 3.2,

$$\frac{q^r + q^{r-1} + \cdots + q - r}{q - 1} N \leq d(q^r + q^{r-1} + \cdots + 1).$$

Note that

$$\begin{aligned}
q^r + q^{r-1} + \cdots + q - r &= \sum_{i=1}^r (q^i - 1) \\
&= \sum_{i=1}^r (q-1)(q^{i-1} + q^{i-2} + \cdots + 1) \\
&= (q-1) \left( \sum_{j=1}^r jq^{r-j} \right).
\end{aligned}$$

Therefore, if we put  $S = \sum_{j=1}^r jq^{r-j}$ , then  $SN \leq d((q-1)S + r + 1)$ . This completes the proof.  $\square$

## 4 Proof of Theorem 1.1 and Corollary 1.2

Now we give proofs of the main theorem and its corollary.

*Proof of Theorem 1.1.* Let  $L$  be the minimal linear subspace of  $\mathbb{P}^r$  so that  $L \supset C$ . Since  $C$  is defined over  $\mathbb{F}_q$ ,  $L \cap L^{(q)} \supset C$ , where  $L^{(q)}$  is the image of  $L$  by the  $q$ -Frobenius map. By the minimality of  $L$ ,  $L = L^{(q)}$ , that is,  $L$  is an  $\mathbb{F}_q$ -space. Since  $C$  is not contained any plane,  $\dim L \geq 3$ . Therefore we may assume that  $C$  is nondegenerate in  $\mathbb{P}^r$ .

(i) Suppose  $d \leq q^{r-2} + q^{r-3} + \cdots + 1$ . Since  $\deg C = d$ , the  $s$ -degree  $d'$  of  $C(\mathbb{F}_q)$  is at most  $d$ . By the combinatorial bound (2) with our assumption,

$$N_q(C) \leq (d' - 1)q + 1 \leq (d - 1)q + 1.$$

(ii) Suppose  $d \geq q$ . In this case, we have  $N_q(C) \leq (d - 1)q + 1$  by Corollary 3.3. In fact,

$$\begin{aligned}
&(d - 1)q + 1 - \left( (q - 1)d + \frac{r + 1}{q^{r-1} + 2q^{r-2} + \cdots + (r - 1)q + r} d \right) \\
&= \left( 1 - \frac{r + 1}{q^{r-1} + 2q^{r-2} + \cdots + (r - 1)q + r} \right) d - q + 1. \tag{6}
\end{aligned}$$

Since the coefficient of  $d$  is positive and  $d \geq q$ ,

$$\text{the quantity (6)} \geq 1 - \frac{r + 1}{q^{r-1} + 2q^{r-2} + \cdots + (r - 1)q + r} q. \tag{7}$$

Since  $r \geq 3$ ,

$$\begin{aligned}
&(q^{r-1} + 2q^{r-2} + \cdots + (r - 1)q + r) - (r + 1)q = \\
&q^{r-1} + \cdots + (r - 3)q^3 + (r - 3)q^2 + (q - 1)^2 + (r - 1) > 0.
\end{aligned}$$

Hence (7) is positive.

Obviously,  $q < q^{r-2} + q^{r-3} + \dots + 1$  because  $r \geq 3$ . Hence (i) and (ii) imply the desired bound.  $\square$

*Proof of Corollary 1.2.* If  $r = 2$ , this is nothing but the main theorem of [5]. So we assume that  $r \geq 3$ .

(i) First we show that we may assume  $C$  to be irreducible over  $\mathbb{F}_q$ . Let  $C = C_1 \cup \dots \cup C_s$  be the decomposition of  $C$  into  $\mathbb{F}_q$ -irreducible components, and  $\deg C_i = d_i$  ( $i = 1, \dots, s$ ). If  $N_q(C_i) \leq (d_i - 1)q + 1$  holds true for any  $C_i$ , then

$$\begin{aligned} N_q(C) &\leq \sum_{i=1}^s N_q(C_i) \leq \sum_{i=1}^s ((d_i - 1)q + 1) \\ &= (d - s)q + s < (d - 1)q + 1. \end{aligned}$$

When  $q = 4$  and  $s \geq 2$ , suppose each of the first  $s'$  components  $C_1, \dots, C_{s'}$  is contained in a plane and isomorphic to  $K$  over  $\mathbb{F}_4$ , and the remaining  $s - s'$  components are not. Then  $d_1 = \dots = d_{s'} = 4$ ,  $d = 4s' + \sum_{i=s'+1}^s d_i$  and  $N_q(C_1) = \dots = N_q(C_{s'}) = 14$ . Hence

$$\begin{aligned} N_q(C) &\leq 14s' + \sum_{i=s'+1}^s ((d_i - 1)4 + 1) \\ &= (d - 1)4 + 1 + 3 - 3s + s' \\ &\leq (d - 1)4 + 1 + 3 - 2s \quad (\text{because } s' \leq s) \\ &< (d - 1)4 + 1 \quad (\text{because } s \geq 2). \end{aligned}$$

(ii) Suppose  $C$  is not absolutely irreducible. As the preliminary step of the proof of Theorem 1.1, we may assume that  $C$  is nondegenerate in  $\mathbb{P}^r$ . Let  $D$  be an irreducible component of  $C$ . Then  $C = D \cup D^{(q)} \cup \dots \cup D^{(q^{t-1})}$  for some  $t \geq 2$ , because  $C$  is irreducible over  $\mathbb{F}_q$ . Hence  $\deg D = \frac{d}{t} \leq \frac{d}{2}$ , and  $C(\mathbb{F}_q) \subset D \cap D^{(q)} \cap \dots \cap D^{(q^{t-1})}$ . When  $C(\mathbb{F}_q)$  does not span  $\mathbb{P}^r$ , choose a hyperplane  $H$  over  $\mathbb{F}_q$  such that  $H \supset C(\mathbb{F}_q)$ . Since  $C$  is nondegenerate,  $H$  does not contain any components of  $C$  because  $H$  is defined over  $\mathbb{F}_q$ . Hence we have

$$N_q(C) \leq (D.H) = \deg D \leq \frac{d}{2} < (d - 1)q + 1,$$

which is the desired bound. Therefore we may assume that  $C(\mathbb{F}_q)$  spans  $\mathbb{P}^r$ . Hence we can pick up  $r - 1$  points  $Q_1, \dots, Q_{r-1} \in C(\mathbb{F}_q)$  such that the linear space  $L_0$  spanned by these  $r - 1$  points is an  $\mathbb{F}_q$ -linear subspace of codimension 2. Put  $\#(L_0 \cap C(\mathbb{F}_q)) = r'$ . Obviously  $r' \geq r - 1$ . Let  $\{H_0, \dots, H_q\}$  be the set of  $\mathbb{F}_q$ -hyperplanes, each of which contains  $L_0$ . Since  $C(\mathbb{F}_q) \setminus L_0 \subset \cup_{i=0}^q (D \cap H_i \setminus L_0)$ ,

$$N_q(C) \leq \left(\frac{d}{t} - r'\right)(q + 1) + r' \leq \left(\frac{d}{2} - r'\right)(q + 1) + r'. \quad (8)$$

Since

$$(d-1)q+1 - \left( \left( \frac{d}{2} - r' \right) (q+1) + r' \right) = r'q + \left( \frac{d}{2} - 1 \right) (q-1) > 0,$$

(8) is bounded by  $(d-1)q+1$ .

Therefore we may assume that  $C$  is absolutely irreducible, which is the case we already considered in Theorem 1.1.  $\square$

## 5 Asymptotic behavior

In this section, we introduce an analogue of Ihara's constant<sup>3</sup>  $A(q)$ .

**Notation 5.1**  $\tilde{\mathcal{C}}_d^i(\mathbb{F}_q)$  denotes the set of irreducible curve over  $\mathbb{F}_q$  of degree  $d$  in a projective space of some dimensions.

**Remark 5.2** The set  $\tilde{\mathcal{C}}_d^i(\mathbb{F}_q)$  consists of finitely many elements. In fact, any member  $C \in \tilde{\mathcal{C}}_d^i(\mathbb{F}_q)$  can be embedded into  $\mathbb{P}^s$  with  $s \leq d$  over  $\mathbb{F}_q$  as a degree  $d$  curve.

**Definition 5.3** Let  $\tilde{M}_q^i(d) := \max\{N_q(C) \mid C \in \tilde{\mathcal{C}}_d^i(\mathbb{F}_q)\}$ , which makes sense because of the finiteness of  $\tilde{\mathcal{C}}_d^i(\mathbb{F}_q)$ . The quantity

$$D(q) := \limsup_{d \rightarrow \infty} \tilde{M}_q^i(d)/d$$

measures the asymptotic behavior of  $\tilde{M}_q^i(d)$ .

We don't know yet the exact value of  $D(q)$  for any  $q$ . Here we state just an observation.

**Proposition 5.4**

$$\frac{1}{2}A(q) \leq D(q) \leq q.$$

*Proof.* Since  $\tilde{M}_q^i(d) \leq (d-1)q+1$  by Theorem 1.1, we have  $D(q) \leq q$ . Let  $N_q(g)$  be the maximum number of  $\mathbb{F}_q$ -points on a nonsingular curve of genus  $g$ . By definition,  $A(q) = \limsup_{g \rightarrow \infty} N_q(g)/g$ . It is known that  $A(q) > 0$  by Serre (for more and precise information on the Ihara's constant, see [9, Chap. 3]). Hence, for most  $g$ 's,  $N_q(g)$  is achieved by a nonhyperelliptic curve, which can be embedded into  $\mathbb{P}^{g-1}$  over  $\mathbb{F}_q$  as a degree  $2g-2$  curve. Therefore

$$\limsup_{g \rightarrow \infty} N_q(g)/(2g-2) = \frac{1}{2}A(q)$$

is a lower bound for  $D(q)$ .  $\square$

---

<sup>3</sup>We use this terminology after [6, 7.1.1].



## References

- [1] J. W. P. Hirschfeld, *Projective geometries over finite fields* (second edition), Oxford University Press, Oxford, 1998.
- [2] J. W. P. Hirschfeld, G. Korchmáros and F. Torres, *Algebraic curves over a finite field*, Princeton Univ. Press, Princeton and Oxford, 2008.
- [3] M. Homma and S. J. Kim, *Around Sziklai's conjecture on the number of points of a plane curve over a finite field*, *Finite Fields Appl.* 15 (2009), 468–474.
- [4] M. Homma and S. J. Kim, *Sziklai's conjecture on the number of points of a plane curve over a finite field II*, in: G. McGuire, G.L. Mullen, D. Panario, I.E. Shparlinski (Eds.), *Finite Fields: Theory and Applications*, in: *Contemp. Math.*, vol. 518, AMS, Providence, 2010, 225–234.
- [5] M. Homma and S. J. Kim, *Sziklai's conjecture on the number of points of a plane curve over a finite field III*, *Finite Fields Appl.* 16 (2010) 315–319.
- [6] H. Stichtenoth, *Algebraic function fields and codes* (second edition), GTM 254, Springer-Verlag, Berlin and Heidelberg, 2008.
- [7] K.-O. Stöhr and J. F. Voloch, *Weierstrass points and curves over finite fields*, *Proc. London Math. Soc.* (3) 52 (1986) 1–19.
- [8] P. Sziklai, *A bound on the number of points of a plane curve*, *Finite Fields Appl.* 14 (2008) 41–43.
- [9] M. Tsfasman, S. Vlăduț and D. Nogin, *Algebraic geometric codes: Basic notions*, SURV 139, AMS, Providence, 2007.