

On a class of degenerate parabolic equations with dynamic boundary conditions

Ciprian G. Gal
 Department of Mathematics
 Florida International University,
 Miami, FL 33199, USA
 cgal@fiu.edu

Abstract

We consider a quasi-linear parabolic (possibly, degenerate) equation with nonlinear dynamic boundary conditions. The corresponding class of initial and boundary value problems has already been studied previously, proving well-posedness of weak solutions and the existence of the global attractor, assuming that the nonlinearities are *subcritical* to a given exponent. The goal of this article is to show that the previous analysis can be redone for *supercritical* nonlinearities by proving an additional L^∞ -estimate on the solutions. In particular, we derive new conditions which reflect an exact balance between the internal and the boundary mechanisms involved, even when both the nonlinear sources contribute in opposite directions. Then, we show how to construct a trajectory attractor for the weak solutions of the associated parabolic system, and prove that any solution belonging to the attractor is bounded, which implies uniqueness. Finally, we also prove for the (semilinear) reaction-diffusion equation with nonlinear dynamic boundary conditions, that the fractal dimension of the global attractor is of the order $\nu^{-(N-1)}$, as diffusion $\nu \rightarrow 0^+$, in any space dimension $N \geq 2$, improving some recent results in [23].

1. Introduction

Let us consider the following partial differential equation

$$\partial_t u - \operatorname{div} (a (|\nabla u|^2) \nabla u) + f(u) = h_1(x), \quad (1.1)$$

in $\Omega \times (0, +\infty)$, where Ω is a bounded domain in \mathbb{R}^N , $N \geq 1$, with smooth boundary $\Gamma := \partial\Omega$, a is a given nonnegative function, and f and h_1 are suitable functions. The mathematical literature regarding equation (1.1) subject to all kinds of homogeneous boundary conditions is fairly vast. We recall that global well-posedness results for (1.1) with Dirichlet or Neumann type of boundary conditions can be found in [1, 6, 7, 8] (see also [3, 5, 9, 37, 46, 52]). In addition, the analysis of dissipative dynamical systems generated by equations like (1.1) was carried out in a number of papers mainly devoted to the asymptotic behavior of strong solutions [2, 8, 31, 41], and to establish the existence of global and/or exponential attractors (see, for instance, [1, 9, 15, 37, 46, 52]). For other classical results concerning the long term dynamics of (1.1) we also refer the reader to [6, 15].

All the mentioned results are mainly concerned with standard boundary conditions (that is, Dirichlet's and/or Neumann's). Let us now consider *dynamic* boundary conditions of the form

$$\partial_t u + b(x) a(|\nabla u|^2) \partial_{\mathbf{n}} u + g(u) = h_2(x), \quad (1.2)$$

on $\Gamma \times (0, +\infty)$, where g and h_2 are suitable functions defined on Γ , and $b \in L^\infty(\Gamma)$, $b \geq b_0 > 0$. This type of boundary conditions arises for many known equations of mathematical physics. They are motivated by problems in diffusion phenomena [12, 13, 18, 19, 22, 30, 49, 51], reaction-diffusion systems in phase-transition phenomena [10, 25, 26, 47], special flows in hydrodynamics [32, 39], models in climatology [40], and many others. For possible physical interpretations of (1.2) for problem (1.1), we refer the reader to [24] (cf. [30] also).

Problems such as (1.1)-(1.2) have already been investigated in a number of papers [11, 16, 17, 27, 36, 49]. Constantin and Escher deal with non-degenerate boundary value problems with smooth nonlinearities (in particular, $\langle a(|\xi|^2) \xi, \xi \rangle \geq c|\xi|^2$, with $c > 0$) and show that unique (classical) maximal solutions exist in some Bessel potential spaces [16, 17]. Such results enable the authors to investigate other qualitative properties concerning global existence and blow-up phenomena (see, also [11]). These results are also improved by Meyries [36], still in the non-degenerate case, by assuming more general boundary conditions and by requiring that $f(s)/s$ and $g(s)/s$ are dissipative as $|s| \rightarrow \infty$. A first analysis, which aimed at deducing only a minimal number of assumptions on the data and nonlinearities, was done in [27] by assuming that f and g are subcritical polynomial nonlinearities and by allowing $a(s)$ to have a polynomial degeneracy at zero. For instance, one can take

$$a(s) = |s|^{(p-2)/2}, \quad \text{for } p \neq 2. \quad (1.3)$$

In particular, we proved that problem (1.1)-(1.2) with $a(s)$ as in (1.3), subject to square-integrable initial data $u|_{t=0} = u_0$ is well-posed, and then we established the existence of a global attractor bounded in $W^{1,p}(\Omega)$. Well-posedness for problem (1.1), (1.2) for $a(s) = |s|^{(p-2)/2}$, assuming monotone functions f, g was considered in [49]. The non-degenerate case $a(s) \equiv \nu > 0$ when $g = 0$, is discussed in detail in [23]. The stationary case associated with (1.1)-(1.2) is treated in [29].

It is well-known that when at least one of the source terms, the bulk nonlinear term f or the boundary term g is present in (1.1)-(1.2), conditions can be derived on their growth rates which imply either the global existence of solutions or blow-up in finite time [21]. Namely in the non-degenerate case, for $\lambda, \mu \in \{0, \pm 1\}$ with $\max\{\lambda, \mu\} = 1$, $f(s) := -\lambda |s|^{r_1-1} s$ and $g(s) := -\mu |s|^{r_2-1} s$, solutions of

$$\partial_t u - \nu \Delta u + f(u) = h_1(x), \quad \text{in } \Omega \times (0, +\infty), \quad (1.4)$$

subject to the dynamic condition

$$\partial_t u + \nu b \partial_{\mathbf{n}} u + g(u) = h_2(x), \quad \text{on } \Gamma \times (0, \infty), \quad (1.5)$$

are globally well-defined, for every given (sufficiently smooth) initial data $u|_{t=0} = u_0$, if $r_1 r_2 > 1$ and $\lambda r_1 + \mu r_2 > 0$. Furthermore, [21] shows that if we further restrict the growths

of r_1, r_2 so that $r_1 < (N + 2) / (N - 2)$ and $r_2 \leq N / (N - 2)$, then the global solutions are also bounded. On the other hand, if $\lambda = 0$, $\mu = 1$, then some solutions blowup in finite time with blowup occurring in the L^∞ -norm at a rate $(t - T_*)^{-(r_2-1)}$, for some additional conditions on u_0 and r_2 . In the same way, when $\mu = 0$ and $\lambda = 1$, then some solutions blowup in finite time with a blowup rate which depends on r_1 and u_0 (see [4]). The occurrence of blow up phenomena is closely related to the blowup problem for the ordinary differential equation

$$u_t + h(u) = 0, \tag{1.6}$$

where either $h = f$ or $h = g$. More precisely, it is easy to see that solutions of the ODE (1.6) are spatially homogeneous solutions of either equation (1.4) or equation (1.5), and so if these solutions blowup in finite time so do the solutions of (1.4), (1.5) (see [43] for further details, and additional references). Similar results showing the same behavior are also derived for the parabolic system (1.1)-(1.2) in [48], stating sufficient conditions for the functions a , f and g so that blowup in finite time occurs in the L^∞ -norm. In particular, it was shown, for odd functions f , g and initial data $u_0 \in L^\infty(\Omega) \cap W^{1,p}(\Omega)$ and some additional conditions on u_0 , that there are solutions that blow up in finite time with an upper bound on the blowup time which can be determined precisely.

The main goal of this paper is to deduce more general conditions (when compared to conditions deduced in [11, 16, 17, 22, 27, 36, 49, 53, 54]) on the reactive and radiation terms f and g , respectively, which imply that problem (1.1)-(1.2) is dissipative in a suitable sense, and that it possesses a (possibly, finite dimensional) global attractor which characterizes the long-term behavior of the parabolic system under consideration. Recently in [43] (see also [55] for some extensions), the authors have considered the semilinear parabolic equation (1.4) subject to nonlinear Robin boundary conditions

$$\nu \partial_{\mathbf{n}} u + g(u) = 0 \text{ on } \Omega \times (0, +\infty), \tag{1.7}$$

and they derived sufficient conditions on f and g , which imply dissipativity for such problems. In particular, they have obtained a general balance between f and g , allowing for a real competition between both the two nonlinear mechanisms which may work in opposite directions, one fighting for blow-up in finite time, the other for dissipativity. Then, they also proved the existence of a compact attractor in $H^1(\Omega)$, assuming that the growth of f and g is subcritical. Their method relies essentially on the fact that problem (1.4), (1.7) possesses a Lyapunov functional, which can then be used to show either dissipativity of (1.4), (1.7), by exploiting some Poincare type inequality (see (2.9) below), or blow up of some solutions.

Our goal is to extend these results in several directions, by working instead with a class of degenerate parabolic equations, such as (1.1), and then by subjecting (1.1) to dynamic boundary conditions of the form (1.2). Moreover, we also wish to consider nonlinearities with *arbitrary* polynomial growth at infinity. We aim to construct weak (energy) solutions with the help from a different (than in [27]) approximation scheme, which is based on the existence of classical (smooth) solutions for a (strictly) non-degenerate system associated with (1.1)-(1.2). Let F and G be the primitives of f and g , respectively, such that $F(0) = 0$, $G(0) = 0$. Even though a natural energy functional exists for suitable approximates of the

problem (1.1)-(1.2), at the moment it is not clear how to prove that this energy (see [27, (1.5)]), namely,

$$\mathcal{E}_{\Omega,\Gamma}(u) := \int_{\Omega} [a(|\nabla u|^2) |\nabla u|^2 + F(u) - h_1(x)u] dx + \int_{\Gamma} [G(u) - h_2(x)u] \frac{dS}{b},$$

is in fact a Lyapunov function for (1.1)-(1.2) when $a(s) = |s|^{(p-2)/2}$, $p \neq 2$, due to a lack of regularity of the weak solutions (see, however [23], and its references, when $a(s) = \nu > 0$). Therefore, the method in [43] which relies on the use of a Lyapunov function does not seem applicable to our situation here. Indeed, when one is dealing with gradient systems with a set of equilibria which is bounded in the phase space where $\mathcal{E}_{\Omega,\Gamma}(u_0) < \infty$, one could avoid to prove the existence of a bounded absorbing set and directly show the existence of the global attractor for *subcritical* nonlinearities. However, since we wish to construct global attractors for (1.1)-(1.2), under no essential growth assumptions on the nonlinearities, we prefer to prove the existence of a uniform dissipative estimate which can be also easily adapted to nonautonomous generalizations. Another difficult step that we need to overcome is the uniqueness problem for the weak (energy) solutions considered here. Indeed, we wish to deduce sufficiently general conditions on f and g without excluding the scenario based on which these functions are competing at infinity. When f and g are both dissipative, i.e., when f and g are both polynomial potentials of odd degree with a positive leading coefficient, uniqueness holds, and thus, the dynamical system associated with the parabolic system (1.1)-(1.2) can be defined in the classical sense. However, if at least one of the polynomial nonlinearities possesses a negative leading coefficient (for instance, suppose that

$$g(s) \sim c_g |s|^{r_2-1} s, \text{ as } |s| \rightarrow \infty, \quad (1.8)$$

for $c_g < 0$), uniqueness is not known, and in this case the classical semigroup can be defined as a semigroup of multi-valued maps only. In order to investigate the long-term behavior of the degenerate parabolic system (1.1)-(1.2), we will employ the trajectory dynamical approach, which allows us to avoid the use of unfriendly multivalued maps, and to apply the usual theory of global attractors (see, e.g., [14] for the general theory). We strongly emphasize that non-uniqueness of the weak solutions constructed here is only a feature of the nonlinear interplay between the two nonlinear mechanisms, and is not related to the smoothness of the functions involved, as this is usually the case for other PDE's. See, e.g., [14, 38, 45] and references therein, for applications for which the uniqueness is not yet solved, such as, hyperbolic equations with supercritical nonlinearities, reaction-diffusion systems, and so on. In fact, even when our functions $f(s)$ and $g(s)$ are assumed to be (locally) Lipschitz, uniqueness of the weak energy solutions is not known unless $f(s)$ and $g(s)$ are monotone increasing for all $|s| \geq s_0$ (for some $s_0 > 0$), which is quite restrictive. Finally, to make matters worse, for boundary nonlinearities that satisfy (1.8), there may be solutions of (1.1)-(1.2) which blowup in finite time at some points in $\overline{\Omega}$, unless the internal mechanism governed by nonlinear flux and reaction is sufficiently strong to overcome the boundary reaction. Therefore, it is also essential to deduce some kind of *optimal* general

conditions between the bulk and boundary nonlinearities that will only produce bounded (non-singular) solutions for all positive times and arbitrary initial data. The main difficulty here is, of course, to establish the asymptotic compactness for the system (1.1)-(1.2) under some general conditions involving f, g , even when at least one has a bad sign at infinity, and to verify that any solution belonging to the attractor, is bounded. Consequently, we obtain the uniqueness on the trajectory attractor even for competing nonlinear mechanisms.

To better understand the larger scope of our results, we shall illustrate their application to the reaction-diffusion equation (1.4), (1.5) for *supercritical* nonlinearities, that is, for functions f and g satisfying the growth assumptions

$$\lim_{|y| \rightarrow \infty} \frac{f'(y)}{|y|^{r_1-2}} = (r_1 - 1) c_f, \quad \lim_{|y| \rightarrow \infty} \frac{g'(y)}{|y|^{r_2-2}} = (r_2 - 1) c_g, \quad (1.9)$$

for some arbitrary $r_2, r_1 \geq 1$, with $\max(r_1, r_2) \geq 2$, and some $c_f, c_g \in \mathbb{R} \setminus \{0\}$. Of course, our results below hold under more general assumptions on f, g , see Section 3. In (1.9), we say that f is dissipative if $c_f > 0$ and non-dissipative if $c_f < 0$ (the same applies to g). Let us assume bounded $h_1(x)$ and $h_2(x)$. When both nonlinear terms cooperate, i.e., both f and g are dissipative and

$$f'(y) \geq -\tilde{c}_f, \quad g'(y) \geq -\tilde{c}_g, \quad \text{for all } y \in \mathbb{R} \quad (1.10)$$

(for some $\tilde{c}_f, \tilde{c}_g > 0$), then problem (1.4), (1.5) is well-posed and possesses a global attractor \mathcal{A}_{gl} in the classical sense, bounded in $\mathcal{Z} := W^{2,2}(\Omega) \cap L^\infty(\overline{\Omega})$, regardless of the size of r_1 and r_2 (see Section 3.1; cf. also [23], when $g = 0$). When the conditions (1.10) do *not* hold, we recall that uniqueness of weak solutions is not known in general. However, if f and g are still dissipative, we can prove that the reaction-diffusion system (1.4), (1.5) possesses a (strong) trajectory attractor \mathcal{A}_{tr} , which is bounded in $L^\infty(\mathbb{R}_+; \mathcal{Z})$. Moreover, uniqueness holds on the attractor \mathcal{A}_{tr} so that the long-term behavior of (1.4), (1.5) can be also characterized by a regular global attractor

$$\mathcal{A}_{gl} := \mathcal{A}_{tr}(0), \quad (1.11)$$

which can be defined in the usual sense of dynamical systems (cf. Section 3.2).

For the case of competing nonlinearities, the following scenarios are possible:

Case (i): For the case of bulk dissipation (i.e., $c_f > 0$) and anti-dissipative behavior at the boundary Γ (i.e., $c_g < 0$), problem (1.4), (1.5) has at least one globally-defined weak solution, which is bounded, if

$$\max(r_2, 2(r_2 - 1)) < r_1.$$

Equality can be also allowed if the boundary condition is homogeneous, i.e., if $h_2 = 0$. Moreover, there exists a (strong) trajectory attractor \mathcal{A}_{tr} , bounded in $L^\infty(\mathbb{R}_+; \mathcal{Z})$, such that solutions are unique on the attractor. Thus, (1.4)-(1.5) also possesses the (smooth) global attractor \mathcal{A}_{gl} , defined as in (1.11).

Case (ii): On the other hand, in the case of boundary dissipation ($c_g > 0$) and internal non-dissipation ($c_f < 0$), for every L^2 -data we obtain that, if $r_2 = r_1 = 2$ (which imply that f and g are sublinear) and $h_2 = 0$, and

$$(c_f + (|\Omega| \alpha)^{-1} c_g) \nu > 2(\tilde{C}_\Omega c_g r_2)^2, \quad (1.12)$$

where

$$\alpha^{-1} := \int_{\Gamma} (b(x))^{-1} dS,$$

and \tilde{C}_Ω is a proper Sobolev-Poincaré constant (see Section 2, assumption (H4)), then (1.4)-(1.5) is well-posed in the classical sense and has a global attractor \mathcal{A}_{gl} , bounded in \mathcal{Z} (in that case, $|f'(y)|$ and $|g'(y)|$ are bounded for all $y \in \mathbb{R}$ by (1.9), see Proposition 2.8). We note that the nonlinear balance condition established in Section 3 can only be used to show the existence of a (strong) trajectory attractor, whenever the interior non-dissipative term f is sublinear as long as the boundary mechanism stays dissipative (i.e., $c_g > 0$), such that g suitably dominates f (see, (1.12)). It would be interesting to see whether one can still construct attractors for the case of a superlinear non-dissipative function f , and some dissipative boundary function g of arbitrary growth. However, as we show at the end of Section 3, we will see that, in this case, the superlinear growth of the nondissipative function f produces blowup in L^∞ -norm of some solutions even for arbitrary nonlinearities g (see Section 3.4). Hence, the nonlinear boundary conditions (1.5) for equation (1.4) *cannot* prevent blowup of some solutions of (1.4)-(1.5) as long as the internal nonlinearity is strongly non-dissipative (for instance, when f satisfies (1.9) with $r_1 > 2$, such that $c_f < 0$). Our main result in Section 3.4 (see, Theorem 3.28) extends some results in [4] for the parabolic equation (1.4) subject to linear dynamic boundary conditions (i.e., when $g \equiv 0$ in (1.5)), and the special cases treated in [21, 35]. In this sense, the nonlinear balance conditions derived in Section 3, which imply that the system (1.4)-(1.5) is dissipative, are optimal.

Finally, exploiting known parabolic regularity theory for PDE's of the form (1.4)-(1.5), the regularity of the solution for (1.4)-(1.5) increases as the functions f , g and the domain Ω become more regular (see Remark 3.22; cf. also [23] and references therein). In particular, the global attractor \mathcal{A}_{gl} consists of (smooth) classical solutions which are defined for all times. Thus, using this additional regularity that solutions of (1.4)-(1.5) enjoy on \mathcal{A}_{gl} (for all the above cases), we obtain an explicit upper bound on the fractal dimension of \mathcal{A}_{gl} for this reaction-diffusion system by imposing weaker assumptions on the nonlinearities than in [23]. In particular, for any $N \geq 2$ and for as long as (1.9) with $c_f > 0$ and $c_g > 0$ holds, we have

$$\mathcal{C}_0 \nu^{-(N-1)} \leq \dim_F \mathcal{A}_{gl} \leq \mathcal{C}_1 (1 + \nu^{-(N-1)}), \quad (1.13)$$

for some positive constants $\mathcal{C}_0, \mathcal{C}_1$ which can be computed explicitly (see Section 3.3). The lower bound in (1.13) was established in [23], assuming dissipative f and (homogeneous) linear boundary equations (i.e., $g = 0$). We note that, for each fixed $\nu > 0$, there is a discrepancy between the upper and lower bounds in (1.13) as \mathcal{C}_0 depends only on Ω, Γ, f, g , whereas \mathcal{C}_1 is also a function of the L^∞ -norms of the sources h_1, h_2 . However, we observe that both the upper and lower bounds are of the order $\nu^{-(N-1)}$ as $\nu \rightarrow 0^+$, cf. (1.13). When $N = 1$, the dimension of \mathcal{A}_{gl} is of the order $\nu^{-1/2}$, as $\nu \rightarrow 0^+$. We recall that, for the reaction-diffusion equation (1.4) with the usual Dirichlet or Neumann-Robin boundary condition (1.7), we have upper and lower bounds of the order $\nu^{-N/2}$, for any $N \geq 1$ (see, e.g., [42, 45]). Thus, we have a much larger estimate (as a function of diffusion, as $\nu \rightarrow 0^+$) for the global attractor \mathcal{A}_{gl} in dimension $N \geq 3$ (see also [23]).

We outline the plan of the paper, as follows. In Section 2, we introduce some notations and preliminary facts, then we recall how to prove the existence and stability of L^2 -energy solutions to our system (1.1)-(1.2). Section 3 is devoted to the existence of a bounded absorbing set and, then, of the global attractor \mathcal{A}_{gl} for unique L^2 -energy solutions. Then, we show that weak solutions possess the L^2 - L^∞ smoothing property exploiting some kind of iterative argument, and consequently, deduce the existence of an absorbing set in L^∞ . In the final part of Section 3, we establish the existence of the trajectory attractor \mathcal{A}_{tr} for our problem and deduce some additional properties for \mathcal{A}_{tr} , especially in the non-degenerate case when $a(s) \equiv \nu > 0$. A blowup result for (1.4)-(1.5) is also established in the case when $g \neq 0$. Finally, in the Appendix we give some auxiliary results which are essential in the proofs.

2. Well-posedness in L^2 -space

We use the standard notation and facts from the dynamic theory of parabolic equations (see, for instance, [27]). The natural space for our problem is

$$\mathbb{X}^{s_1, s_2} := L^{s_1}(\Omega) \oplus L^{s_2}(\Gamma) = \left\{ U = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} : u_1 \in L^{s_1}(\Omega), u_2 \in L^{s_2}(\Gamma) \right\},$$

$s_1, s_2 \in [1, +\infty]$, endowed with norm

$$\|U\|_{\mathbb{X}^{s_1, s_2}} = \left(\int_{\Omega} |u_1(x)|^{s_1} dx \right)^{1/s_1} + \left(\int_{\Gamma} |u_2(x)|^{s_2} \frac{dS_x}{b(x)} \right)^{1/s_2}, \quad (2.1)$$

if $s_1, s_2 \in [1, \infty)$, and

$$\begin{aligned} \|U\|_{\mathbb{X}^\infty} &:= \max\{\|u_1\|_{L^\infty(\Omega)}, \|u_2\|_{L^\infty(\Gamma)}\} \\ &\simeq \|u_1\|_{L^\infty(\Omega)} + \|u_2\|_{L^\infty(\Gamma)}. \end{aligned}$$

We agree to denote by \mathbb{X}^s the space $\mathbb{X}^{s,s}$. Identifying each function $u \in W^{1,p}(\Omega)$, with the vector $U := \begin{pmatrix} u \\ u|_{\Gamma} \end{pmatrix}$, it is easy to see that $W^{1,p}(\Omega)$ is a dense subspace of \mathbb{X}^s for $s \in [1, \infty)$. Moreover, we have

$$\mathbb{X}^s = L^s(\overline{\Omega}, d\mu), \quad s \in [1, +\infty],$$

where the measure $d\mu = dx|_{\Omega} \oplus dS_x/b(x)|_{\Gamma}$ on $\overline{\Omega}$ is defined for any measurable set $A \subset \overline{\Omega}$ by $\mu(A) = |A \cap \Omega| + S(A \cap \Gamma)$. Identifying each function $\theta \in C(\overline{\Omega})$ with the vector $\Theta = \begin{pmatrix} \theta|_{\Omega} \\ \theta|_{\Gamma} \end{pmatrix}$, we have that $C(\overline{\Omega})$ is a dense subspace of \mathbb{X}^s for every $s \in [1, \infty)$ and a closed subspace of \mathbb{X}^∞ . In general, any vector $\theta \in \mathbb{X}^s$ will be of the form $\begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$ with $\theta_1 \in L^s(\Omega, dx)$ and $\theta_2 \in L^s(\Gamma, dS/b(x))$, and there need not be any connection between θ_1 and θ_2 . For domains Ω with Lipschitz boundary Γ , recall that we have $W^{k,p}(\Omega) \subset L^{p_s}(\Omega)$, with $p_s = \frac{pN}{N-pk}$ if $pk < N$, and $1 \leq p_s < \infty$, if $N = pk$. Moreover the trace operator $Tr_D(u) := u|_{\Gamma}$, initially defined for $u \in C^k(\overline{\Omega})$, has an extension to a bounded linear operator from $W^{k,p}(\Omega)$ into $L^{q_s}(\partial\Omega)$, where $q_s := \frac{p(N-1)}{N-pk}$ if $pk < N$, and $1 \leq q_s < \infty$ if $N = pk$. For $p > Nk$, we have

$W^{k,p}(\Omega) \subset C^{\zeta, \tilde{l}}(\overline{\Omega})$, for some ζ, \tilde{l} . We also recall, on account of well-known generalized Poincaré-type inequalities (see, e.g., [33]), that

$$\|u\|_{W^{1,p}(\Omega)} := \|\nabla u\|_{(L^p(\Omega))^N} + l(u) \quad (2.2)$$

is a norm on $W^{1,p}(\Omega)$, which is equivalent to the usual one, for any of the following expressions for l :

$$l(u) := \left(\int_{\Gamma} |u|^s \frac{dS_x}{b} \right)^{1/s}, \quad l(u) := \left(\int_{\Omega} |u|^s dx \right)^{1/s},$$

for any $1 \leq s \leq p$. Next, for each $p > 1$, we let

$$\mathbb{V}^{k,p} = \left\{ U := \begin{pmatrix} u \\ u|_{\Gamma} \end{pmatrix} : u \in W^{k,p}(\Omega) \right\}$$

and endow it with the norm $\|\cdot\|_{\mathbb{V}^{k,p}}$ given by

$$\|U\|_{\mathbb{V}^{k,p}} = \|u\|_{W^{k,p}(\Omega)} + \|u|_{\Gamma}\|_{W^{k-1/p,p}(\Gamma)}.$$

It is easy to see that we can identify $\mathbb{V}^{k,p}$ with $W^{k,p}(\Omega) \oplus W^{k-1/p,p}(\Gamma)$ under this norm. Moreover, we emphasize that $\mathbb{V}^{k,p}$ is not a product space and since $W^{k,p}(\Omega) \hookrightarrow W^{k-1/p,p}(\Gamma)$ by trace theory, $\mathbb{V}^{k,p}$ is topologically isomorphic to $W^{k,p}(\Omega)$ in the obvious way. It is also immediate that $\mathbb{V}^{k,p}$ is compactly embedded into \mathbb{X}^2 , for any $p > p_0 := 2N/(N+2)$ and $k \geq 1$. From now on, we denote by $\|\cdot\|_{W^{k,p}(\Omega)}$ and $\|\cdot\|_{W^{k,q}(\Gamma)}$ the norms on $W^{k,p}(\Omega)$ and $W^{k,q}(\Gamma)$, respectively. Also, $\langle \cdot, \cdot \rangle_s$ and $\langle \cdot, \cdot \rangle_{s,\Gamma}$ stand for the usual scalar product in $L^s(\Omega)$ and $L^s(\Gamma)$, respectively. We also agree to denote by \mathbb{V}^p the space $\mathbb{V}^{1,p}$, and $\langle \cdot, \cdot \rangle$ the duality between X and X^* , for some generic Banach space X .

Our first goal in this paper is to give a nonlinear balance between f and g which implies dissipativity of (1.1)-(1.2), even when both the nonlinear terms contribute in opposite directions. More precisely, we wish to prove the existence of (globally well-defined) weak solutions, provided that the nonlinearities satisfy (possibly part of) the assumptions listed below:

(H1) Let $b(y) := a(|y|^2)y$, $y \in \mathbb{R}^N$ and assume that $a \in C(\mathbb{R}, \mathbb{R})$, $b \in C^1(\mathbb{R}^N, \mathbb{R}^N)$ satisfy the following conditions:

$$\begin{cases} |a(|y|^2)| \leq c_1(1 + |y|^{p-2}), & \forall y \in \mathbb{R}^N, \\ \langle b(y), y \rangle_{\mathbb{R}^N} = a(|y|^2)|y|^2 \geq \nu|y|^p, & \forall y \in \mathbb{R}^N, \end{cases} \quad (2.3)$$

for some constants $c_1, \nu > 0$. Moreover, assume that b is monotone nondecreasing, i.e.,

$$\langle a(|y_1|^2)y_1 - a(|y_2|^2)y_2, y_1 - y_2 \rangle_{\mathbb{R}^N} \geq 0, \quad \text{for all } y_1, y_2 \in \mathbb{R}^N. \quad (2.4)$$

(H2) $f, g \in C^1(\mathbb{R}, \mathbb{R})$ satisfy

$$\liminf_{|y| \rightarrow +\infty} f'(y) > 0, \quad \liminf_{|y| \rightarrow +\infty} g'(y) > 0. \quad (2.5)$$

(H3) (a) $f, g \in C(\mathbb{R}, \mathbb{R})$ satisfy the growth assumptions

$$|f(y)| \leq c_f (1 + |y|^{r_1-1}), \quad |g(y)| \leq c_g (1 + |y|^{r_2-1}), \quad \forall y \in \mathbb{R}, \quad (2.6)$$

for some positive constants c_f, c_g , and some $r_1, r_2 \geq 1$.

(b) $f, g \in C(\mathbb{R}, \mathbb{R})$ satisfy

$$\begin{aligned} |f(y)| &\leq c_f (1 + |y|^{r_1-1}), \\ c_g |y|^{r_2} - c &\leq g(y) y \leq \tilde{c}_g |y|^{r_2} + c, \quad \forall y \in \mathbb{R}, \end{aligned}$$

for some appropriate positive constants and some $r_1, r_2 \geq 1$.

(H4) Let $\lambda = (\int_{\Gamma} b^{-1} dS)^{-1}$, and suppose that $g \in C^1(\mathbb{R}, \mathbb{R})$. There exists $\varepsilon \in (0, \nu/q)$, with ν as in (2.3), such that,

$$\liminf_{|y| \rightarrow +\infty} \frac{f(y) y + (|\Omega| \lambda)^{-1} g(y) y - \frac{\tilde{C}_{\Omega}^q}{(\varepsilon p)^{q/pq}} |g'(y) y + g(y)|^q}{|y|^{r_1}} > 0, \quad (2.7)$$

for some $r_1 \geq p$. Here

$$\tilde{C}_{\Omega} = \begin{cases} C_{\Omega, b} (\lambda |\Omega|)^{-1}, & \text{if (H3a) holds} \\ \frac{C_{\Omega, b}}{2} (\lambda |\Omega|)^{-1}, & \text{if (H3b) holds,} \end{cases} \quad (2.8)$$

and $C_{\Omega, b}$ is the best Sobolev constant in the following Poincaré's inequality:

$$\|\phi - \lambda \langle \phi/b, 1 \rangle_{1, \Gamma}\|_{L^s(\Omega)} \leq C_{\Omega, b} \|\nabla \phi\|_{L^s(\Omega)}, \quad s \geq 1 \quad (2.9)$$

(see, e.g., [43, Lemma 3.1]).

We observe that condition (H4) provides an exact balance between the two nonlinear mechanisms. As we shall see, this balance will depend both upon the sign and growth rate of f and g at infinity (cf. also, [43]).

We have the following rigorous notion of weak solution to (1.1)-(1.2), with initial condition $u(0) = u_0$, similar to [27].

Definition 2.1. Let $p \in (\frac{2N}{N+2}, +\infty) \cap (1, +\infty)$, and let $h_1(x) \in L^{r'_1}(\Omega)$, $h_2(x) \in L^{r'_2}(\Gamma)$, where r'_i is the dual conjugate of r_i . The pair $U(t) = \begin{pmatrix} u(t) \\ v(t) \end{pmatrix}$ is said to be a weak solution if $v(t) = u(t)|_{\Gamma}$, in the trace sense, for a.e. $t \in (0, T)$, for any $T > 0$, and U fulfills

$$\begin{cases} U(t) \in L^{\infty}([0, +\infty); \mathbb{X}^2) \cap W_{loc}^{1, s}([0, +\infty); (\mathbb{V}^{k, p})^*) \\ u(t) \in L_{loc}^p([0, +\infty); W^{1, p}(\Omega)), \\ v(t) \in L_{loc}^p([0, +\infty); W^{1-1/p, p}(\Gamma)), \end{cases}$$

for $s = \min(q, r'_1, r'_2)$, $q := p/(p-1)$, and

$$k = \max\left(1, \frac{N}{p} - \frac{N}{r_1}, \frac{N}{p} - \frac{N-1}{r_2}\right). \quad (2.10)$$

Moreover, the following identity

$$\begin{aligned} & \langle \partial_t U, \Xi \rangle_{\mathbb{X}^2} + \langle a (|\nabla u|^2) \nabla u, \nabla \sigma \rangle_2 + \langle f(u), \sigma \rangle_2 + \langle g(v)/b, \sigma|_{\Gamma} \rangle_{2,\Gamma} \\ & = \langle h_1, \sigma \rangle_2 + \langle h_2/b, \sigma|_{\Gamma} \rangle_{2,\Gamma}, \end{aligned} \quad (2.11)$$

holds for all $\Xi = \begin{pmatrix} \sigma \\ \sigma|_{\Gamma} \end{pmatrix} \in \mathbb{V}^{k,p}$, a.e. $t \in (0, T)$. Finally, we have, in the space \mathbb{X}^2 ,

$$U(0) = \begin{pmatrix} u_0 \\ v_0 \end{pmatrix} =: U_0, \quad (2.12)$$

where $u(0) = u_0$ almost everywhere in Ω , and $v(0) = v_0$ almost everywhere in Γ . Note that in this setting, v_0 need not be the trace of u_0 at the boundary.

We can cast the weak formulation (2.11) into a proper functional equation by defining suitable operators. To this end, let $\langle \cdot, \cdot \rangle$ denote the duality between \mathbb{V}^p and $(\mathbb{V}^p)^*$. Define the form

$$\tilde{\mathcal{B}}_p(U_1, U_2) := \int_{\Omega} a (|\nabla u_1|^2) \nabla u_1 \cdot \nabla u_2 dx + \int_{\Omega} |u_1|^{p-2} u_1 u_2 dx,$$

for all $U_i = \begin{pmatrix} u_i \\ u_i|_{\Gamma} \end{pmatrix} \in \mathbb{V}^p$, $i = 1, 2$. Note that

$$\begin{aligned} \tilde{\mathcal{B}}_p(U_1, U_2) & = - \int_{\Omega} \operatorname{div} (a (|\nabla u_1|^2) \nabla u_1) u_2 dx \\ & \quad + \int_{\Gamma} b(x) a (|\nabla u_1|^2) \partial_{\mathbf{n}} u_1 u_2 \frac{dS}{b(x)} + \int_{\Omega} |u_1|^{p-2} u_1 u_2 dx. \end{aligned} \quad (2.13)$$

It follows from Lemma 4.1 (see Appendix), that for each $U = \begin{pmatrix} u \\ v \end{pmatrix} \in \mathbb{V}^p$, there exists $\mathcal{B}_p(U) \in (\mathbb{V}^p)^*$ such that

$$\tilde{\mathcal{B}}_p(U, W) = \langle \mathcal{B}_p(U), W \rangle, \quad (2.14)$$

for every $W \in \mathbb{V}^p$. Hence, this relation defines an operator $\mathcal{B}_p : \mathbb{V}^p \rightarrow (\mathbb{V}^p)^*$, which is bounded. Exploiting Lemma 4.1 once again, it is easy to see that \mathcal{B}_p is monotone and coercive. It also follows that $\mathcal{B}_p(\mathbb{V}^p) = (\mathbb{V}^p)^*$ (see, e.g., [5]). Thus, we end up with the following functional form

$$\partial_t U + \mathcal{B}_p U + \mathcal{F}(U) = \mathcal{G}(x), \quad (2.15)$$

where $\mathcal{G}(x) = \begin{pmatrix} h_1(x) \\ h_2(x) \end{pmatrix}$, and the operators $\mathcal{B}_p : D(\mathcal{B}_p) \rightarrow \mathbb{X}^2$, $\mathcal{F} : D(\mathcal{F}) \subset \mathbb{X}^2 \rightarrow \mathbb{X}^2$ are given, formally, by

$$\begin{aligned} \mathcal{B}_p U & = \begin{pmatrix} -\operatorname{div} (a (|\nabla u|^2) \nabla u) + |u|^{p-2} u \\ b(x) a (|\nabla u|^2) \partial_{\mathbf{n}} u \end{pmatrix}, \\ \mathcal{F}(U) & = \begin{pmatrix} f(u) - |u|^{p-2} u \\ g(v) \end{pmatrix}. \end{aligned} \quad (2.16)$$

We aim to prove some regularity results for the weak solutions constructed in Definition 2.1. In [27], solutions were constructed with aid from a Galerkin approximation scheme

by imposing additional growth restrictions on the nonlinearities f, g so that they are essentially dominated by a monotone operator associated with the p -Laplacian. However, the application of this scheme seems to be problematic in our context since the solutions constructed with Definition 2.1 are much weaker than those constructed in [27, Definition 2.3]. Therefore, we need to rely on another scheme which is based on the construction of classical (smooth) solutions to the non-degenerate analogue of (1.1)-(1.2). One of the advantages of this construction is that now every weak solution can be approximated by regular ones and the justification of our estimates for such solutions is immediate. Thus, for each $\epsilon > 0$, let us consider the following non-degenerate parabolic problem:

$$\begin{aligned} \partial_t u - \operatorname{div} (a_\epsilon (|\nabla u|^2) \nabla u) + f(u) &= h_1(x), \text{ in } \Omega \times (0, \infty), \\ \partial_t u + b(x) a_\epsilon (|\nabla u|^2) \partial_{\mathbf{n}} u + g(u) &= h_2(x), \text{ on } \Gamma \times (0, \infty), \end{aligned} \quad (2.17)$$

where $a_\epsilon(s) = a(s + \epsilon) > 0$, for all $s \in \mathbb{R}$, subject to the initial conditions

$$u_\epsilon(0) = u_{0\epsilon}, \quad v_\epsilon(0) = u_{0\epsilon|_\Gamma}. \quad (2.18)$$

Let $u_{0\epsilon} \in C^\infty(\overline{\Omega})$ such that

$$U_\epsilon(0) \rightarrow U(0) = U_0 \text{ in } \mathbb{X}^2.$$

Then, the approximate problem (2.17)-(2.18) admits a unique (smooth) classical solution with

$$u_\epsilon \in C^1([0, t_*]; C^\infty(\overline{\Omega})) \quad (2.19)$$

for some $t_* > 0$ and each $\epsilon > 0$ (see [16, 17, 11]). Being pedants, we cannot apply the main results of [16] (cf. also [17]) directly to equations (2.17)-(2.18) since the functions a_ϵ, f, g and the external forces h_1, h_2 are not smooth enough. Moreover, the solutions constructed this way may only exist locally in time for some interval $[0, t_*)$. However, taking sequences $h_{1\epsilon} \in C^\infty(\overline{\Omega}), h_{2\epsilon} \in C^\infty(\Gamma)$ such that $h_{1\epsilon} \rightarrow h_1$ in $L^\infty(\Omega)$, and $h_{2\epsilon} \rightarrow h_2$ in $L^\infty(\Gamma)$, respectively, and by approximating the functions a_ϵ, f, g by smooth ones, say, in $C^\infty(\mathbb{R}, \mathbb{R})$, we may apply Remark 3.3 below for the solutions of the approximate equations, and deduce the existence of a globally defined in \mathbb{X}^∞ -norm solution to (2.17)-(2.18). Indeed, taking advantage of the fact that $u_{0\epsilon} \in C^\infty(\overline{\Omega})$, the global \mathbb{X}^∞ -a priori bound for u_ϵ guarantees its global existence in at least $\mathbb{V}^p \cap \mathbb{X}^\infty$ -norm, which turns out to be sufficient for our purpose. As we shall see in the next section, this bound can be naturally obtained under the above assumptions on the nonlinearities by performing a modified Alikakos-Moser iteration argument (see Theorem 3.2 and Remark 3.3 below).

We shall now deduce the first result concerning the solvability of problem (1.1)-(1.2).

Theorem 2.2. *Let a, f and g satisfy either the assumptions (H1), (H3a), (H4) with*

$$\max(r_2, q(r_2 - 1)) < r_1, \quad (2.20)$$

or (H1), (H3b) and (H4). Then, for any initial data $U_0 \in \mathbb{X}^2$, there exists at least one (globally defined) weak solution $U(t)$ in the sense of Definition 2.1.

Proof. We divide the proof into several steps according to the different hypotheses being used.

Step 1. (i) We shall now derive some basic apriori estimates for $U = U_\epsilon(t)$ when $p \leq N$, assuming that (H1), (H3a) and (H4) are satisfied. The case $p > N$ can be treated analogously. The following estimates will be deduced by a formal argument, which can be justified by means of the approximation procedure devised above. Also, for practical purposes c will denote a positive constant that is independent of time, $\epsilon > 0$ and initial data, but which only depends on the other structural parameters. Such a constant may vary even from line to line. Note that the smooth solutions constructed in (2.17)-(2.18) also satisfy the weak formulation (2.11). Thus, the key choices $\sigma = u_\epsilon(t)$, $\sigma_\Gamma = v_\epsilon(t)/b$ in (2.11) are justified. After standard transformations, in view of assumption (H1), we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + \nu \|\nabla u(t)\|_{L^p(\Omega)}^p + \langle f(u(t)), u(t) \rangle_2 + \langle g(v(t)), v(t)/b \rangle_{2,\Gamma} \\ & \leq \langle h_1, u(t) \rangle_2 + \langle h_2/b, v(t) \rangle_{2,\Gamma}. \end{aligned} \quad (2.21)$$

Following [43], we can now write

$$\begin{aligned} & \langle f(u), u \rangle_2 + \langle g(v), v/b \rangle_{2,\Gamma} \\ & = \langle f(u)u + c_B g(u)u, 1 \rangle_2 - c_B \left\langle g(u)u - \lambda \langle g(v)v/b, 1 \rangle_{2,\Gamma} \right\rangle_2, \end{aligned} \quad (2.22)$$

where

$$c_B := \frac{\lambda^{-1}}{|\Omega|}, \quad \lambda = \left(\int_\Gamma b^{-1} dS \right)^{-1}.$$

Applying inequality (2.9) to the last term on the right-hand side of (2.22) yields

$$\begin{aligned} & c_B \left| \left\langle g(u)u - \lambda \langle g(v)v/b, 1 \rangle_{2,\Gamma} \right\rangle_2 \right| \\ & \leq \tilde{C}_\Omega \|\nabla(g(u)u)\|_{L^1(\Omega)} = \tilde{C}_\Omega \left\| \left(g'(u)u + g(u) \right) \nabla u \right\|_{L^1(\Omega)} \\ & \leq \varepsilon \|\nabla u\|_{L^p(\Omega)}^p + \frac{\tilde{C}_\Omega^q}{(\varepsilon p)^{q/p} q} \left\| g'(u)u + g(u) \right\|_{L^q(\Omega)}^q \end{aligned} \quad (2.23)$$

with $\tilde{C}_\Omega = C_{\Omega,b}(\lambda|\Omega|)^{-1}$, and we recall that q is conjugate to p . Since assumption (H4) holds for some $\varepsilon \in (0, \nu/q)$, then from (2.22)-(2.23), we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + (\nu - \varepsilon) \|\nabla u(t)\|_{L^p(\Omega)}^p + c \|u(t)\|_{L^{r_1}(\Omega)}^{r_1} \\ & \leq \langle h_1, u(t) \rangle_2 + \langle h_2/b, v(t) \rangle_{2,\Gamma} + c, \end{aligned} \quad (2.24)$$

for some positive constant $c > 0$, independent of U , t and ϵ . Exploiting the estimate in Lemma 4.2 (see Appendix), and then using Hölder and Young inequalities, we can bound

the term on the right-hand side of (2.24) by

$$\begin{aligned} & \left(c \|h_1\|_{L^{r_1}'}^{r_1'} + \frac{c}{2} \|u\|_{L^{r_1}(\Omega)}^{r_1} \right) \\ & + \left[\delta \|\nabla u\|_{L^p(\Omega)}^p + C_\delta \left(\|u\|_{L^\gamma(\Omega)}^\gamma + \|h_2/b\|_{L^{r_2}(\Gamma)}^{r_2'} + 1 \right) \right], \end{aligned}$$

with $\gamma = \max(r_2, p(r_2 - 1)/(p - 1))$, for a sufficiently small $\delta > 0$, and sufficiently large $C_\delta > 0$. Since by assumption $\gamma < r_1$, we can control the L^γ -norm of u in terms of the L^{r_1} -norm of the solution (i.e., $\|u\|_{L^\gamma}^\gamma \leq \kappa \|u\|_{L^{r_1}}^{r_1} + C_\kappa$, $\kappa \ll 1$). Thus, we get for suitable choices of $\varepsilon \in (0, \nu/q)$ and $\delta \in (0, \nu/q)$, the following inequality

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + \frac{c}{p} \|\nabla u(t)\|_{L^p(\Omega)}^p + \frac{c}{2} \|u(t)\|_{L^{r_1}(\Omega)}^{r_1} \\ & \leq c \left(1 + \|h_1\|_{L^{r_1}'}^{r_1'} + \|h_2/b\|_{L^{r_2}(\Gamma)}^{r_2'} \right), \end{aligned} \quad (2.25)$$

for almost all $t \geq 0$. Recalling that $L^{r_1}(\Omega) \subset L^p(\Omega)$ (we have, $r_1 \geq p$), we can now integrate this inequality over $[0, T]$ to deduce

$$\begin{aligned} U_\epsilon & \in L^\infty([0, T]; \mathbb{X}^2) \cap L^p([0, T]; \mathbb{V}^p), \\ u_\epsilon & \in L^{r_1}([0, T] \times \Omega), \end{aligned} \quad (2.26)$$

uniformly with respect to $\epsilon > 0$. On account of these bounds, we get

$$\mathcal{B}_{p,\epsilon}(U_\epsilon) \in L^q([0, T]; (\mathbb{V}^p)^*) \subseteq L^q([0, T]; (\mathbb{V}^{k,p})^*),$$

uniformly in $\epsilon > 0$, for any $k \geq 1$ (cf. Lemma 4.1; see also [27]). Here $\mathcal{B}_{p,\epsilon}$ is the monotone operator associated with the function a_ϵ (see (2.14), (2.16)). Moreover, by Lemma 4.2 in the Appendix, we get at once

$$v_\epsilon = u_{\epsilon|\Gamma} \in L^{r_2}([0, T] \times \Gamma) \cap L^p([0, T] \times \Gamma), \quad (2.27)$$

uniformly in ϵ . Due to assumption (H3a), from (2.26)-(2.27), we deduce

$$\begin{pmatrix} f(u_\epsilon) \\ g(v_\epsilon) \end{pmatrix} \in L^{r_1'}([0, T] \times \Omega) \times L^{r_2'}([0, T] \times \Gamma). \quad (2.28)$$

Thus, $\mathcal{F}(U_\epsilon)$ is uniformly (in ϵ) bounded in $L^s([0, T]; \mathbb{X}^s)$, which implies

$$\mathcal{G}(x) - \mathcal{B}_{p,\epsilon}U_\epsilon - \mathcal{F}(U_\epsilon) \in L^s([0, T]; (\mathbb{V}^{k,p})^*),$$

with $s = \min(q, r_1', r_2') > 1$. Therefore, $\partial_t U_\epsilon$ is bounded in $L^s([0, T]; (\mathbb{V}^{k,p})^*)$, uniformly with respect to $\epsilon > 0$, for some $k \geq 1$. Indeed, having chosen k so that $W^{k,p} \subset L^{r_1}(\Omega) \subset L^p(\Omega)$ and $W^{k-1/p,p} \subset L^{r_2}(\Gamma)$ (in particular, it holds $\mathbb{V}^{k,p} \subset \mathbb{X}^{r_1, r_2}$ with continuous inclusion),

so if $V \in (\mathbb{X}^{r_1, r_2})^*$, then $V \in (\mathbb{V}^{k, p})^*$. Thus, equation (2.15) holds as an equality in $L^s([0, T]; (\mathbb{V}^{k, p})^*)$ and it can be considered distributionally in the space $\mathcal{D}'([0, T]; (\mathbb{V}^{k, p})^*)$. The existence of a weak solution is then based on monotone operator arguments, followed by a passage to limit as $\epsilon \rightarrow 0$, and can be carried over exactly as in the proof of [27, Theorem 2.6]. We shall briefly describe the details below in **Step 2**.

Step 1. (ii) We will now deduce the apriori bounds (2.26)-(2.27), if one assumes (H3b) instead of (H3a). According to (2.21), in light of inequality (H3b) for g , we have the following:

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + \nu \|\nabla u(t)\|_{L^p(\Omega)}^p + \langle f(u(t)), u(t) \rangle_2 \\ & + \frac{1}{2} \langle g(v(t)), v(t)/b \rangle_{2, \Gamma} + \frac{c_g}{2} \|v(t)\|_{L^{r_2}(\Gamma, b^{-1}dS)}^{r_2} \\ & \leq \langle h_1, u(t) \rangle_2 + \langle h_2/b, v(t) \rangle_{2, \Gamma} + c, \end{aligned} \quad (2.29)$$

for some positive constant c . We can write, as in (2.22),

$$\begin{aligned} & \langle f(u), u \rangle_2 + \frac{1}{2} \langle g(v), v/b \rangle_{2, \Gamma} \\ & = \left\langle f(u)u + \frac{c_B}{2} g(u)u, 1 \right\rangle_2 - \frac{c_B}{2} \left\langle g(u)u - \lambda \langle g(v)v/b, 1 \rangle_{2, \Gamma} \right\rangle_2, \end{aligned}$$

and argue exactly as above to get the following estimate:

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + (\nu - \varepsilon) \|\nabla u(t)\|_{L^p(\Omega)}^p + c \left(\|u(t)\|_{L^{r_1}(\Omega)}^{r_1} + \|v(t)\|_{L^{r_2}(\Gamma, b^{-1}dS)}^{r_2} \right) \\ & \leq \langle h_1, u(t) \rangle_2 + \langle h_2/b, v(t) \rangle_{2, \Gamma} + c. \end{aligned} \quad (2.30)$$

The desired control of $U = U_\epsilon(t)$ in (2.26)-(2.27) can be obtained immediately from a simple application of Hölder and Young inequalities on the terms on the right-hand side of (2.30). Thus the proof is the same as in Step 1-(i).

Step 2. It is obvious that

$$\partial_t U_\epsilon + \mathcal{B}_{p, \epsilon} U_\epsilon + \mathcal{F}(U_\epsilon) = \mathcal{G} \quad (2.31)$$

holds as an equality in $L^s([0, T]; (\mathbb{V}^{k, p})^*)$ (this is the same as equation (2.15) with U and $a(\cdot)$ replaced by U_ϵ and $a_\epsilon(\cdot)$, respectively). From the estimates that we deduced in Step 1, (i)-(ii), we see that there exists a subsequence $\{U_\epsilon\} = \left\{ \begin{pmatrix} u_\epsilon \\ v_\epsilon \end{pmatrix} \right\}$ (still denoted by $\{U_\epsilon\}$), such that as $\epsilon \rightarrow 0$,

$$\begin{aligned} & U_\epsilon \rightarrow U \text{ weakly star in } L^\infty([0, T]; \mathbb{X}^2), \\ & U_\epsilon \rightarrow U \text{ weakly in } L^p([0, T]; \mathbb{V}^p), \\ & \partial_t U_\epsilon \rightarrow \partial_t U \text{ weakly in } L^s([0, T]; (\mathbb{V}^{k, p})^*). \end{aligned} \quad (2.32)$$

On the basis on standard interpolation and compact embedding results for vector valued functions (see [27]), we also have

$$U_\epsilon \rightarrow U \text{ strongly in } L^p([0, T]; \mathbb{X}^2). \quad (2.33)$$

Clearly, $U \in C([0, T]; (\mathbb{V}^{k,p})^*)$. By refining in (2.33), u_ϵ converges to u a.e. in $\Omega \times (0, T)$ and v_ϵ converges to v a.e. in $\Gamma \times (0, T)$, respectively. Then, by means of known results in measure theory (see, e.g., [45]), the continuity of f , g , and the convergence of (2.33) imply that $f(u_\epsilon)$ converges weakly to $f(u)$ in $L^{r_1}(\Omega \times (0, T))$. Moreover, $g(v_\epsilon)$ converges weakly to $g(v)$ in $L^{r_2}(\Gamma \times (0, T))$, and thus, $\mathcal{F}(U_\epsilon)$ converges weakly star to $\mathcal{F}(U)$ in $L^s([0, T]; (\mathbb{V}^{k,p})^*)$. Since $\mathcal{B}_{p,\epsilon}U_\epsilon$ is bounded in $L^q([0, T]; (\mathbb{V}^p)^*)$, we further see that

$$\mathcal{B}_{p,\epsilon}U_\epsilon \rightarrow \Xi \text{ weakly star in } L^q([0, T]; (\mathbb{V}^p)^*), \quad (2.34)$$

and thus weakly star in $L^s([0, T]; (\mathbb{V}^{k,p})^*)$, since $s \leq q$ and $k \geq 1$. We are now ready to pass to the limit as $\epsilon \rightarrow 0$ in equation (2.31). We have

$$\partial_t U + \Xi + \mathcal{F}(U) = \mathcal{G} \quad (2.35)$$

as an equality in $L^s([0, T]; (\mathbb{V}^{k,p})^*)$. It remains to show $\Xi = \mathcal{B}_p U$, which can be proved by a standard monotonicity argument exactly as in [27, Theorem 2.6]. We leave the details to the interested reader. ■

The following proposition is also immediate.

Proposition 2.3. *Let $a(\cdot)$ satisfy (H1). In addition, assume that $f, g \in C(\mathbb{R}, \mathbb{R})$ satisfy*

$$\begin{aligned} c_1 |y|^{r_1} - c &\leq f(y) y \leq c_2 |y|^{r_1} + c, & \forall y \in \mathbb{R}, \\ c_3 |y|^{r_2} - c &\leq g(y) y \leq c_4 |y|^{r_2} + c, & \forall y \in \mathbb{R}, \end{aligned} \quad (2.36)$$

for some appropriate positive constants and some $r_1, r_2 \geq 1$ such that $\max(r_1, r_2) \geq p$. Then, for any initial data $U_0 \in \mathbb{X}^2$, there exists at least one (globally defined) weak solution $U(t)$ in the sense of Definition 2.1.

Proof. In this case, both f and g are dissipative so that we do *not* need to exploit the validity of assumption (H4). Indeed, it follows from (2.21) that

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + \nu \|\nabla u(t)\|_{L^p(\Omega)}^p + c_1 \|u(t)\|_{L^{r_1}(\Omega)}^{r_1} + c_3 \|v(t)\|_{L^{r_2}(\Gamma, b^{-1}dS)}^{r_2} \\ &\leq \langle h_1, u(t) \rangle_2 + \langle h_2/b, v(t) \rangle_{2,\Gamma} + c, \end{aligned} \quad (2.37)$$

which yields the desired control of $U(t)$ in the corresponding spaces (2.26)-(2.27) with relative ease (see [27], for further details). Thus, the proof is the same as in Theorem 2.2. ■

Remark 2.4. (i) *If $U(t)$ is a weak solution of problem (1.1)-(1.2), in the sense of Definition 2.1, then clearly $U(t) \in C([0, T]; (\mathbb{V}^{k,p})^*)$. Since by duality,*

$$\mathbb{V}^{k,p} \subset \mathbb{X}^2 = (\mathbb{X}^2)^* \subset (\mathbb{V}^{k,p})^*,$$

for any $p \in (p_0, \infty) \cap (1, \infty)$, $k \geq 1$, and recalling that $U(t) \in L^\infty([0, T]; \mathbb{X}^2)$, it follows

$$U(t) \in C_w([0, T]; \mathbb{X}^2)$$

(see, e.g., [14, Theorem II.1.7]). Therefore the initial value $U|_{t=0} = U_0$ is meaningful when $U_0 \in \mathbb{X}^2$. Finally, we note that in general, the assumptions (H3)-(H4) alone do not ensure the uniqueness of the Cauchy problem (1.1)-(1.2).

(ii) We are also allowed to have equality in (2.20), i.e., $\gamma = \max(r_2, q(r_2 - 1)) \leq r_1$ if $h_2 \equiv 0$. Indeed, this follows once again from (2.24) and Lemma 4.2, which allows us to control surface integrals in terms of volume integrals.

Proposition 2.5. *Let the assumptions of Theorem 2.2 be satisfied. Then any weak solution $U(t) = \begin{pmatrix} u(t) \\ v(t) \end{pmatrix}$ of (1.1)-(1.2) belongs to $C([0, T]; \mathbb{X}^2)$, such that $\|U(t)\|_{\mathbb{X}^2}^2$ is absolutely continuous on $[0, T]$, and*

$$\frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 = -\langle \mathcal{B}_p(U(t)), U(t) \rangle - \langle \mathcal{F}(U(t)) - \mathcal{G}, U(t) \rangle, \quad (2.38)$$

for almost all $t \in [0, T]$. We will refer to (2.38) as the energy identity for the parabolic system (1.1)-(1.2).

Proof. This statement follows from a generalization of a known interpolation result (see, e.g., [14, 34, 45]). Indeed, identifying the space $H^* = (\mathbb{X}^2)^*$ with $H = \mathbb{X}^2$, we have

$$V := \mathbb{V}^{1,p} \subset \mathbb{X}^2 \subset (\mathbb{V}^{1,p})^* \subseteq (\mathbb{V}^{k,p})^* =: W,$$

for any $k \geq 1$. Moreover, the following inclusion $E := \mathbb{X}^{r_1, r_2} \subseteq \mathbb{X}^2 \subseteq E^* \subseteq W$ also holds (indeed, the dual of E is the space $E^* = \mathbb{X}^{r_1', r_2'}$, and by (2.10), $\mathbb{V}^{k,p} \subset E$). By virtue of equation (2.15), any distributional derivative $\partial_t U(t)$ from $\mathcal{D}'([0, T]; W)$ can be represented as $\partial_t U(t) = \mathcal{Z}_1(t) + \mathcal{Z}_2(t)$, where

$$\mathcal{Z}_1(t) := -\mathcal{B}_p(U(t)), \quad \mathcal{Z}_2(t) := -\mathcal{F}(U(t)) + \mathcal{G}.$$

According to (2.26)-(2.28), $\mathcal{Z}_1(t) \in L^q([0, T]; V^*) = (L^p([0, T]; V))^*$, $q = p/(p-1)$, for

$$U(t) \in L^p([0, T]; V) + (L^{r_1}([0, T] \times \Omega) \times L^{r_2}([0, T] \times \Gamma)), \quad (2.39)$$

while from (2.28), $\mathcal{Z}_2(t) \in L^{r_1'}([0, T] \times \Omega) \times L^{r_2'}([0, T] \times \Gamma)$, which is precisely the dual of the product space in (2.39). Thus, the claim follows, for instance, from [14, Theorem II.1.8] (see also [57, Proposition 23.23]). ■

We will now state some results which reflect the applicability of assumption (H4) to a wide range of situations. In particular, it applies to the case of competing nonlinearities f and g , that is, nonlinearities with arbitrary *polynomial* growth which satisfy (H3), but when either one exhibits a non-dissipative behavior at infinity. Recall that $p \in (\frac{2N}{N+2}, \infty) \cap (1, \infty)$. In the interesting case of an internal dissipation mechanism, and non-dissipative boundary conditions, we have the following.

Corollary 2.6. *Assume that $f, g \in C^1(\mathbb{R}, \mathbb{R})$ satisfy*

$$\lim_{|y| \rightarrow \infty} \frac{f'(s)}{|s|^{r_1-2}} = (r_1 - 1) c_f > 0 \quad \text{and} \quad \lim_{|y| \rightarrow \infty} \frac{g'(s)}{|s|^{r_2-2}} = (r_2 - 1) c_g < 0 \quad (2.40)$$

with $r_1 \geq p$, $r_2 > 1$. Suppose that one of the following conditions holds:

- (i) (2.20) holds, i.e., $\max(r_2, q(r_2 - 1)) < r_1$.
- (ii) $h_2 = 0$, $p < r_2 < q(r_2 - 1) = r_1$ and

$$c_f \nu p^{-q} q > \tilde{C}_\Omega^q c_g^q (r_2)^q. \quad (2.41)$$

Then, in each case the conclusion of Theorem 2.2 applies.

Proof. We begin by noting that (H3a) is immediately satisfied. For sufficiently large y , we have

$$f(y) \sim c_f |y|^{r_1-2} y, \quad g(y) \sim c_g |y|^{r_2-2} y,$$

and $f(y)y \sim c_f |y|^{r_1}$, $g(y)y \sim c_g |y|^{r_2}$. Thus, the leading terms in (2.7) are

$$c_f |y|^{r_1} + (|\Omega| \lambda)^{-1} c_g |y|^{r_2} - \frac{\tilde{C}_\Omega^q}{(\varepsilon p)^{q/p} q} c_g^q (r_2)^q |y|^{q(r_2-1)}, \quad (2.42)$$

for some $\varepsilon \in (0, \nu/q)$. By assumption (i), it holds $\gamma = \max(r_2, q(r_2 - 1)) < r_1$, so the coefficient of the highest order term in (2.42) is c_f , which is positive. If (ii) holds, it is obvious that $r_2 < \max(r_2, q(r_2 - 1)) = r_1$, so the coefficient of the highest order term in (2.42) is

$$c_f - \frac{\tilde{C}_\Omega^q}{(\varepsilon p)^{q/p} q} c_g^q (r_2)^q,$$

which is positive for some $\varepsilon \in (0, \nu/q)$, if (2.41) is satisfied. Therefore, the conditions of Theorem 2.2 are met, and the proof is finished. ■

In the case of non-dissipative *polynomial* behavior for f , we have the following.

Corollary 2.7. Assume $h_2 = 0$ and $f, g \in C^1(\mathbb{R}, \mathbb{R})$ satisfy

$$\lim_{|y| \rightarrow \infty} \frac{f'(s)}{|s|^{p-2}} = (r_1 - 1) c_f < 0 \quad \text{and} \quad \lim_{|y| \rightarrow \infty} \frac{g'(s)}{|s|^{p-2}} = (r_2 - 1) c_g > 0 \quad (2.43)$$

for some $p \in (\frac{2N}{N+2}, \infty) \cap (1, \infty)$, and let

$$(c_f + (|\Omega| \lambda)^{-1} c_g) \nu p^{-q} q > \tilde{C}_\Omega^q c_g^q (r_2)^q. \quad (2.44)$$

Then, the conclusion of Theorem 2.2 applies.

Proof. First, it is immediate that (H3b) holds with $r_1 = r_2 = p$. Obviously, in this case $r_2 = r_1 = q(r_2 - 1)$. The coefficient of the highest order term in (2.42) is then

$$(c_f + (|\Omega| \lambda)^{-1} c_g) - \frac{\tilde{C}_\Omega^q}{(\varepsilon p)^{q/p} q} c_g^q (r_2)^q,$$

which is positive for some $\varepsilon \in (0, \nu/q)$, if (2.44) is satisfied. Therefore, condition (H4) holds true and, thus, the assumptions of Theorem 2.2 are verified. ■

In the case of simultaneous internal and boundary dissipation, we can prove the following stability result. Note that in this proposition, the uniqueness holds in the class of all solutions which are constructed by means of Definition 2.1, and not only for solutions which can be obtained as the limit, as $\varepsilon \rightarrow 0$, of the (strictly) non-degenerate parabolic system introduced earlier.

Proposition 2.8. *Let the assumptions of Theorem 2.2 be satisfied, and in addition, assume that (H2) holds. Then, there exists a unique weak solution to problem (1.1)-(1.2), which depends continuously on the initial data in a Lipschitz way.*

Proof. Let us consider two weak solutions $U_1 = \begin{pmatrix} u_1 \\ u_{1\Gamma} \end{pmatrix}$ and $U_2 = \begin{pmatrix} u_2 \\ u_{2\Gamma} \end{pmatrix}$, which according to Proposition 2.5 belong to the spaces in (2.39), and set $U(t) = U_1(t) - U_2(t)$. Thus, $U(t)$ satisfies the equation

$$\partial_t U + (\mathcal{B}_p(U_1(t)) - \mathcal{B}_p(U_2(t))) + \mathcal{F}(U_1(t)) - \mathcal{F}(U_2(t)) = 0,$$

and

$$U(0) = U_1(0) - U_2(0).$$

Since

$$U(t) \in L^p([0, T]; \mathbb{V}^p) \cap (L^{r_1}([0, T] \times \Omega) \times L^{r_2}([0, T] \times \Gamma)),$$

and

$$\partial_t U(t) \in L^q([0, T]; (\mathbb{V}^p)^*) + (L^{r_1'}([0, T] \times \Omega) \times L^{r_2'}([0, T] \times \Gamma)),$$

Proposition 2.5 is indeed applicable, and we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 + \langle \mathcal{B}_p(U_1(t)) - \mathcal{B}_p(U_2(t)), U(t) \rangle \\ &= - \langle \mathcal{F}(U_1(t)) - \mathcal{F}(U_2(t)), U(t) \rangle, \end{aligned}$$

for almost all $t \in [0, T]$. Recalling that $\mathcal{B}_p(\cdot) \in \mathcal{L}(\mathbb{V}^p, (\mathbb{V}^p)^*)$ is monotone and coercive (see (2.14); cf. also Appendix), we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 \\ & \leq - \langle f(u_1(t)) - f(u_2(t)), u(t) \rangle - \langle g(u_1(t)) - g(u_2(t)), u(t) \rangle. \end{aligned} \tag{2.45}$$

Exploiting assumption (H2) (which implies, $f'(y) \geq -c_f$ and $g'(y) \geq -c_g$, $\forall y \in \mathbb{R}$, for some $c_f, c_g > 0$), we obtain

$$\frac{d}{dt} \|U(t)\|_{\mathbb{X}^2}^2 \leq 2(c_f + c_g) \|U(t)\|_{\mathbb{X}^2}^2,$$

for almost all $t \in [0, T]$. Integrating this inequality over $[0, T]$ and applying Gronwall's inequality, we deduce

$$\|U(t)\|_{\mathbb{X}^2}^2 \leq e^{ct} \|U(0)\|_{\mathbb{X}^2}^2, \tag{2.46}$$

which yields the desired result. ■

Remark 2.9. (i) Note that assumption (H2) is only required to prove uniqueness of the weak solution, and is usually not required for the theory of attractors. Moreover, this assumption is actually too restrictive so that nonlinearities that satisfy it are not allowed to carry a bad sign at infinity, and thus this would automatically eliminate the scenario proposed by the statements of Corollaries 2.6 and 2.7. Indeed, a simple observation that will be made in Section 3 is that actually uniqueness is necessary on the attractor only, and this can be obtained by deducing additional regularity estimates for the solutions. This observation is in particular very useful if one needs to consider entropy-related nonlinearities of the form $f(y) = y^l \log(y)$, for $y > 0$, and $f(y) = 0$, for $y \leq 0$, for some $l \geq 1$.

(ii) Obviously, estimate (2.46) also holds if we assume that $|f'(y)|$ and $|g'(y)|$ are bounded for all $y \in \mathbb{R}$.

As an immediate consequence of the stability result just proven above, problem (1.1)-(1.2), (2.12) defines a dynamical system in the classical sense.

Corollary 2.10. *Let the assumptions of Proposition 2.8 be satisfied. The problem (1.1)-(1.2), (2.12) defines a (nonlinear) continuous semigroup $\mathcal{S}_2(t)$ on the phase space \mathbb{X}^2 ,*

$$\mathcal{S}_2(t) : \mathbb{X}^2 \rightarrow \mathbb{X}^2,$$

given by

$$\mathcal{S}_2(t) U_0 = U(t), \tag{2.47}$$

where $U(t)$ is the (unique) weak solution which satisfies the energy identity (2.38).

3. Global Attractors

3.1. Attractors for $(\mathcal{S}_2(t), \mathbb{X}^2)$ revisited

In order to study the asymptotic behavior of (1.1)-(1.2), (2.12), we need to derive some additional apriori estimates for the solutions. We shall focus our study on the case $p \geq 2$ only, since for $p \in (\frac{2N}{N+2}, 2)$ we need to impose slightly different assumptions on the nonlinearities, and so we will pursue this question elsewhere. We first aim to improve some results from [27] for the weak solutions constructed in Definition 2.1, which are unique by Proposition 2.8, and to show the existence of the (classical) global attractor, bounded in $\mathbb{X}^\infty \cap \mathbb{V}^p$. We emphasize again that all the results below hold for any $p \geq 2$.

The next result is a direct consequence of estimate (2.25) of Theorem 2.2 (see [27, Section 2, Proposition 3.3], for details).

Proposition 3.1. *Let the assumptions of either Theorem 2.2 or Proposition 2.3 be satisfied. The solution semigroup $\{\mathcal{S}_2(t)\}_{t \geq 0}$ has a $(\mathbb{X}^2, \mathbb{X}^2)$ -bounded absorbing set. More precisely, there is a positive constant C_0 , depending only on the physical parameters of the problem, such that for any bounded subset $B \subset \mathbb{X}^2$, there exists a positive constant $t^\# = t^\#(\|B\|_{\mathbb{X}^2})$ such that*

$$\sup_{t \geq t^\#} \left[\|U(t)\|_{\mathbb{X}^2} + \int_t^{t+1} \left(\int_{\Omega} a(|\nabla u(s)|^2) |\nabla u(s)|^2 + |u(s)|^{r_1} \right) dx ds \right] \leq C_0. \tag{3.1}$$

Our next goal is to establish the existence of a bounded absorbing set in \mathbb{X}^∞ , which has an interest on its own. The following result extends [27, Theorem 3.7] by removing the additional growth conditions that were imposed on f, g in [27, Theorem 3.7, (3.17)].

Theorem 3.2. *Let the assumptions of Proposition 3.1 be satisfied. Let $h_1 \in L^\infty(\Omega)$, $h_2 \in L^\infty(\Gamma)$, and suppose*

$$\liminf_{|y| \rightarrow \infty} \frac{f(y)}{y} > 0, \quad \liminf_{|y| \rightarrow \infty} \frac{g(y)}{y} > 0. \quad (3.2)$$

Then, given any initial data U_0 in \mathbb{X}^2 , the corresponding solution $U(t)$ of (1.1)-(1.2), (2.12) belongs to \mathbb{X}^∞ , for each $t > 0$. Moreover, there exists a positive constant C_1 , independent of t and the initial data, and a positive constant t_+ depending on $t^\#$, such that

$$\sup_{t \geq t_+} \|U(t)\|_{\mathbb{X}^\infty} \leq C_1. \quad (3.3)$$

Proof. All the calculations below are formal. However, they can be rigorously justified by means of the approximation procedure devised in Section 2 (see (2.17)-(2.18)). From now on, c will denote a positive constant that is independent of t, ϵ, m and initial data, which only depends on the other structural parameters of the problem. Such a constant may vary even from line to line. Moreover, we shall denote by $Q_\tau(m)$ a monotone nondecreasing function in m of order τ , for some nonnegative constant τ , independent of m . More precisely, $Q_\tau(m) \sim cm^\tau$ as $m \rightarrow +\infty$.

We begin by showing that the \mathbb{X}^m -norm of U satisfies a local recursive relation which can be used to perform an iterative argument. We divide the proof of (3.3) into several steps.

Step 1 (The basic energy estimate in \mathbb{X}^{m+1}). We multiply (1.1) by $|u|^{m-1}u$, $m \geq 1$, and integrate over Ω . We obtain

$$\begin{aligned} & \frac{1}{(m+1)} \frac{d}{dt} \|u\|_{m+1}^{m+1} + \langle f(u), |u|^{m-1}u \rangle_2 + m \int_\Omega a(|\nabla u|^2) |\nabla u|^2 |u|^{m-1} dx \\ &= \int_\Gamma a(|\nabla u|^2) \partial_n u |v|^{m-1} v dS + \langle h_1(x), |u|^{m-1}u \rangle_2. \end{aligned} \quad (3.4)$$

Similarly, we multiply (1.2) by $|v|^{m-1}v/b$ and integrate over Γ . We have

$$\begin{aligned} & \frac{1}{(m+1)} \frac{d}{dt} \left\| \frac{v}{b} \right\|_{m+1, \Gamma}^{m+1} + \int_\Gamma b(x) a(|\nabla u|^2) \partial_n u |v|^{m-1} v \frac{dS}{b(x)} + \left\langle g(v), \frac{|v|^{m-1}v}{b} \right\rangle_{2, \Gamma} \\ &= \left\langle h_2(x), \frac{|v|^{m-1}v}{b} \right\rangle_{2, \Gamma}. \end{aligned} \quad (3.5)$$

Let us first observe that, in light of assumption (2.3), it is easy to check

$$m \int_\Omega a(|\nabla u|^2) |\nabla u|^2 |u|^{m-1} dx \geq \nu m \left(\frac{p}{p+m-1} \right)^p \int_\Omega \left| \nabla |u|^{\frac{p+m-1}{p}} \right|^p dx. \quad (3.6)$$

Adding relations (3.4)-(3.5), we deduce on account of the assumptions (3.2) (indeed, it holds

$$f(y)y \geq -c_1 s^2 - c_2, g(y)y \geq -c_3 s^2 - c_4, \quad (3.7)$$

for all $y \in \mathbb{R}$, and some $c_i > 0$) and an application of basic Hölder and Young inequalities, the following inequality

$$\frac{d}{dt} \|U\|_{\mathbb{X}^{m+1}}^{m+1} + \gamma \nu \int_{\Omega} \left| \nabla |u|^{\frac{p+m-1}{p}} \right|^p dx \leq Q_1(m) (\|U\|_{\mathbb{X}^{m+1}}^{m+1} + 1). \quad (3.8)$$

Here the positive constant $\gamma \sim m^{-(p-2)}$, $p \geq 2$, and the function $Q_1(m) \sim m$ depends also on the L^∞ -norms of h_1, h_2 , and of b_0 .

Step 2 (The local relation). Set $m_k = p^k$, and define

$$\mathcal{Y}_k(t) := \int_{\Omega} |u(t, \cdot)|^{1+m_k} dx + \int_{\Gamma} |v(t, \cdot)|^{1+m_k} \frac{dS}{b} = \|U(t)\|_{\mathbb{X}^{m_k+1}}^{m_k+1}, \quad (3.9)$$

for all $k \geq 0$. Let t, μ be two positive constants such that $t - \mu/m_k > 0$. Their precise values will be chosen later. We claim that

$$\mathcal{Y}_k(t) \leq M_k(t, \mu) := c (m^k)^\sigma \left(\sup_{s \geq t - \mu/m_k} \mathcal{Y}_{k-1}(s) + 1 \right)^{n_k}, \quad \forall k \geq 1, \quad (3.10)$$

where c, σ are positive constants independent of k , and $n_k := \max\{z_k, l_k\} \geq 1$ is a bounded sequence for all k .

We will now prove (3.10) when $p < N$. The case $p \geq N$ shall require only minor adjustments (in fact, in this case we can choose any arbitrary, but fixed, $p_s, q_s > p$ in the embedding $\mathbb{V}^{1,p}(\Omega) \subset \mathbb{X}^{p_s, q_s}$). For each $k \geq 0$, we define

$$r_k := \frac{N(p + m_k - 1) - (N - p)(1 + m_k)}{N(p + m_k - 1) - (N - p)(1 + m_{k-1})}, \quad s_k := 1 - r_k.$$

We aim to estimate the term on the right-hand side of (3.8) in terms of the $\mathbb{X}^{1+m_{k-1}}$ -norm of U . First, Hölder and Sobolev inequalities (with the equivalent norm of Sobolev spaces in $W^{1,p}(\Omega) \subset L^{p_s}(\Omega)$, $p_s = pN/(N - p)$) yield

$$\begin{aligned} \int_{\Omega} |u|^{1+m_k} dx &\leq \left(\int_{\Omega} |u|^{\frac{(p+m_k-1)N}{N-p}} dx \right)^{s_k} \left(\int_{\Omega} |u|^{1+m_{k-1}} dx \right)^{r_k} \\ &\leq c \left(\int_{\Omega} \left| \nabla |u|^{\frac{(p+m_k-1)}{p}} \right|^p dx + \left(\int_{\Omega} |u|^{1+m_{k-1}} dx \right)^{\alpha_k} \right)^{\bar{s}_k} \left(\int_{\Omega} |u|^{1+m_{k-1}} dx \right)^{r_k}, \end{aligned} \quad (3.11)$$

with

$$\begin{aligned} \bar{s}_k &:= s_k \frac{N}{N-p} = \frac{(p-1)Nm_{k-1}}{(p-1)Nm_{k-1} + m_k + p_1} \in (0, 1), \\ p_1 &:= (p-1)N - (N-p) > 0, \quad \alpha_k := \frac{p+m_k-1}{1+m_{k-1}} \in [1, p]. \end{aligned}$$

Applying Young's inequality on the right-hand side of (3.11), we get

$$Q_{\tau_1}(m_k) \int_{\Omega} |u|^{1+m_k} dx \leq \frac{\gamma_k}{4} \int_{\Omega} \left| \nabla |u|^{\frac{p+m_k-1}{p}} \right|^p dx + Q_{\tau_2}(m_k) \left(\int_{\Omega} |u|^{1+m_{k-1}} dx \right)^{\max\{z_k, \alpha_k\}}, \quad (3.12)$$

for some positive constants τ_i independent of m , and where

$$z_k := r_k / (1 - \bar{s}_k) = (p_1 + m_{k+1}) / (p_1 + m_k) \geq 1$$

is bounded for all k . Note that we can choose τ_2 to be some fixed positive number since Q_{τ_2} also depends on $\gamma_k^{-1} \sim m_k^{p-2}$. To treat the boundary terms in (3.8), we define for $k \geq 1$,

$$y_k := \frac{(N-1)(p+m_k-1) - (N-p)(1+m_k)}{(N-1)(p+m_k-1) - (N-p)(1+m_{k-1})}, \quad x_k := 1 - y_k.$$

On account of Hölder and Sobolev inequalities (see Section 2), we obtain

$$\begin{aligned} \int_{\Gamma} |v|^{1+m_k} \frac{dS}{b} &\leq c \left(\int_{\Gamma} |v|^{\frac{(N-1)(p+m_k-1)}{N-p}} dS \right)^{x_k} \left(\int_{\Gamma} |v|^{1+m_{k-1}} \frac{dS}{b} \right)^{y_k} \\ &\leq c \left(\int_{\Omega} \left| \nabla |u|^{\frac{p+m_k-1}{p}} \right|^p dx + \left(\int_{\Omega} |u|^{1+m_{k-1}} dx \right)^{\alpha_k} \right)^{\bar{x}_k} \left(\int_{\Gamma} |v|^{1+m_{k-1}} \frac{dS}{b} \right)^{y_k}, \end{aligned} \quad (3.13)$$

with

$$\begin{aligned} \bar{x}_k &:= \frac{N-1}{N-p} x_k = \frac{(N-1)(p-1)m_{k-1}}{(N-1)(p-1)m_{k-1} + (p-1)m_{k-1} + p_2}, \\ p_2 &:= (N-1)(p-1) - (N-p) > 0. \end{aligned}$$

Since $\bar{x}_k \in (0, 1)$, we can apply Young's inequality on the right-hand side of (3.13), use the estimate for the $L^{1+m_k}(\Omega)$ -norm of u from (3.12) in order to deduce the following estimate:

$$\begin{aligned} Q_{\tau_3}(m_k) \int_{\Gamma} |v|^{1+m_k} \frac{dS}{b} \\ \leq \frac{\gamma_k}{4} \int_{\Omega} \left| \nabla |u|^{\frac{p+m_k-1}{p}} \right|^p dx + Q_{\tau_4}(m_k) \left(\int_{\Omega} |u|^{1+m_{k-1}} dx \right)^{\max\{l_k, \alpha_k\}}, \end{aligned} \quad (3.14)$$

for some positive constants τ_3, τ_4 depending on τ_1, τ_2 , but which are independent of m . The sequence

$$l_k := \frac{y_k}{(1 - \bar{x}_k)} = \frac{(p-1)m_{k+1} + pp_2}{(p-1)m_k + pp_2} \geq 1$$

is bounded for all $k \geq 1$. Inserting estimates (3.12)-(3.14) on the right-hand side of (3.8), we obtain the following inequality:

$$\partial_t \mathcal{Y}_k + \gamma_k \int_{\Omega} \left| \nabla |u|^{\frac{p+m_k-1}{p}} \right|^p dx \leq c (m_k)^{\sigma_1} (\mathcal{Y}_{k-1} + 1)^{n_k}, \quad (3.15)$$

for some positive constant σ_1 that depends on τ_i ; we recall that $n_k = \max\{z_k, l_k, \alpha_k\} \geq 1$, and $\gamma_k \sim m_k^{2-p}$.

Let now $\zeta(s)$ be a positive function $\zeta : \mathbb{R}_+ \rightarrow [0, 1]$ such that $\zeta(s) = 0$ for $s \in [0, t - \mu/r_k]$, $\zeta(s) = 1$ if $s \in [t, +\infty)$ and $|d\zeta/ds| \leq m_k/\mu$, if $s \in (t - \mu/r_k, t)$. We define $Z_k(s) = \zeta(s) \mathcal{Y}_k(s)$ and notice that

$$\begin{aligned} \frac{d}{ds} Z_k(s) &\leq \zeta(s) \frac{d}{ds} \mathcal{Y}_k(s) + \frac{m_k}{\mu} \mathcal{Y}_k(s) \\ &= \zeta(s) \frac{d}{ds} \mathcal{Y}_k(s) + Q_1(m_k) \left(\int_{\Omega} |u|^{1+m_k} dx + \int_{\Gamma} |v|^{1+m_k} \frac{dS}{b} \right). \end{aligned} \quad (3.16)$$

The last two integrals in (3.16) can be estimated as in (3.12) and (3.14). Combining the above estimates and the fact that $Z_k \leq \mathcal{Y}_k$, we deduce the following inequality:

$$\frac{d}{ds} Z_k(s) + c m_k Z_k(s) \leq M_k(t, \mu), \text{ for all } s \in [t - \mu/r_k, +\infty). \quad (3.17)$$

Note that $c = c(\mu) \sim \mu^{-1}$ as $\mu \rightarrow 0$, and $c(\mu)$ is bounded if μ is bounded away from zero. Integrating (3.17) with respect to s from $t - \mu/r_k$ to t , and taking into account the fact that $Z_k(t - \mu/r_k) = 0$, we obtain that $\mathcal{Y}_k(t) = Z_k(t) \leq M_k(t, \mu) (1 - e^{-c\mu})$, which proves the claim (3.10).

Step 3 (The iterative argument). Let now $\tau' > \tau > 0$ be given with $\tau = t^\#$ as in (3.1), and define $\mu = p(\tau' - \tau) \geq 1$, $t_0 = \tau' = t^\# + 1$ and $t_k = t_{k-1} - \mu/m_k$, $k \geq 1$. Using (3.10), we have

$$\sup_{t \geq t_{k-1}} \mathcal{Y}_k(t) \leq c(\mu) (m_k)^\sigma (\sup_{s \geq t_k} \mathcal{Y}_{k-1}(s) + 1)^{n_k}, \quad k \geq 1. \quad (3.18)$$

Note that from (3.1), we have

$$\sup_{s \geq t_1 = \tau} (\mathcal{Y}_0(s) + 1) \leq C_0 + 1 =: \overline{C}, \quad (3.19)$$

and $c = c(\mu)$ is bounded away from zero. Thus, we can iterate in (3.18) with respect to $k \geq 1$ and obtain that

$$\begin{aligned} \sup_{t \geq t_{k-1}} \mathcal{Y}_k(t) &\leq (c m_k^\sigma) (c m_{k-1}^\sigma)^{n_k} (c m_{k-2}^\sigma)^{n_k n_{k-1}} \cdot \dots \cdot (c m_0^\sigma)^{n_k n_{k-1} \dots n_0} (\overline{C})^{\xi_k} \\ &\leq c^{A_k} p^{\sigma B_k} (\overline{C})^{\xi_k}, \end{aligned} \quad (3.20)$$

where $\xi_k := n_k n_{k-1} \dots n_0$, and

$$A_k := 1 + n_k + n_k n_{k-1} + \dots + n_k n_{k-1} \dots n_0, \quad (3.21)$$

$$B_k := k + n_k(k-1) + n_k n_{k-1}(k-2) + \dots + n_k n_{k-1} \dots n_0. \quad (3.22)$$

Without loss of generality, let us assume that $z_k \geq l_k \geq 1$, for each k . Then, $n_k = z_k$, and $\xi_k = (p_1 + m_{k+1}) / (p_1 + m_0)$. The argument below also applies to the case when $n_k = l_k$. Thus, we have

$$A_k \leq (p_1 + m_k) \sum_{i=1}^{\infty} \frac{1}{p_1 + m_i} \text{ and } B_k \leq (p_1 + m_k) \sum_{i=1}^{\infty} \frac{i}{p_1 + m_i}. \quad (3.23)$$

Therefore, since

$$\sup_{t \geq t_0} \mathcal{Y}_k(t) \leq \sup_{t \geq t_{k-1}} \mathcal{Y}_k(t) \leq c^{A_k} p^{\sigma B_k} (\overline{C})^{\xi_k} \quad (3.24)$$

and the series in (3.23) are convergent, we can take the $1 + m_k$ -root on both sides of (3.24) and let $k \rightarrow +\infty$. We deduce

$$\sup_{t \geq t_0 = \tau'} \|U(t)\|_{\mathbb{X}^\infty} \leq \lim_{k \rightarrow +\infty} \sup_{t \geq t_0} (\mathcal{Y}_k(t))^{1/(1+m_k)} \leq C_1, \quad (3.25)$$

for some positive constant C_1 independent of t, k, U, ϵ and initial data. The proof of Theorem 3.2 is now complete. ■

Remark 3.3. (i) We can easily modify our argument in the proof of (3.3) in order to show that the \mathbb{X}^∞ -norm of the solution $U(t)$ stays bounded for all time $t \geq 0$, if U_0 is bounded in the \mathbb{X}^∞ -norm. It suffices to note that in place of the inequality (3.10), we may use instead the inequality

$$\mathcal{Y}_k(t) \leq Q(\|U_0\|_{\mathbb{X}^\infty}, \sup_{t > 0} M_k(t, \mu)),$$

which is an immediate consequence of (3.15). Then arguing as in the proof of Theorem 3.2, we also have the estimate:

$$\sup_{t \geq 0} \|U(t)\|_{\mathbb{X}^\infty} \leq Q(\|U_0\|_{\mathbb{X}^\infty}, \sup_{t \geq 0} \|U(t)\|_{\mathbb{X}^2}), \quad (3.26)$$

for some positive monotone nondecreasing (in each of its variables) function $Q : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ independent of ϵ .

(ii) By slightly refining the arguments in the proof of Theorem 3.2 (in Step 3), it is also easy to show that, for each $\tau > 0$,

$$\sup_{t \geq 2\tau} \|U(t)\|_{\mathbb{X}^\infty} \leq Q(\tau^{-1}, \sup_{t \geq \tau} \|U(t)\|_{\mathbb{X}^2}).$$

Arguing exactly as in the proof of Theorem 2.2, we can also obtain the following *general* balance condition between the functions f, g , implying boundedness of the solution.

Proposition 3.4. Let the assumptions of Theorem 2.2 be satisfied, and let $h_1 \in L^\infty(\Omega)$, $h_2 \in L^\infty(\Gamma)$. Suppose that there exist $\tau \geq 0$ and $y_0 > 0$, such that for any $m \geq 1$ and $|y| \geq y_0$, it holds

$$\begin{aligned} & f(y) |y|^{m-1} y + (|\Omega| \lambda)^{-1} g(y) |y|^{m-1} y - \frac{\tilde{C}_\Omega^q m^{-q/p}}{(\varepsilon p)^{q/p} q} |y|^{m-1} \left| g'(y) y + m g(y) \right|^q \\ & \geq -Q_\tau(m) (|y|^{m+1} + 1), \end{aligned} \quad (3.27)$$

for some $\varepsilon \in (0, \frac{\nu}{q})$, and some positive function $Q_\tau : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, $Q_\tau(m) \sim m^\tau$, as $m \rightarrow \infty$. Then, the same conclusion of Theorem 3.2 applies to any weak solution of problem (1.1)-(1.2), (2.12).

Proof. Let us return to equations (3.4)-(3.6). We can write

$$\begin{aligned}
& \langle f(u), |u|^{m-1} u \rangle_2 + \left\langle g(v), \frac{|v|^{m-1} v}{b} \right\rangle_{2,\Gamma} \\
&= \langle f(u) + c_B g(u), |u|^{m-1} u \rangle_2 \\
& - c_B \left\langle g(u) u |u|^{m-1} - \lambda \langle g(u) u |u|^{m-1}, 1/b \rangle_{1,\Gamma} \right\rangle_2,
\end{aligned} \tag{3.28}$$

where c_B and λ are as in Theorem 2.2. Applying the Poincaré's inequality (2.9), we have

$$\begin{aligned}
& c_B \left| \left\langle g(u) u |u|^{m-1} - \lambda \langle g(u) u |u|^{m-1}, 1/b \rangle_{1,\Gamma} \right\rangle_2 \right| \\
& \leq \tilde{C} \left\| \nabla (g(u) u |u|^{m-1}) \right\|_{L^1(\Omega)} \\
& = \tilde{C} \left\| |u|^{m-1} \nabla u (g'(u) u + mg(u)) \right\|_{L^1(\Omega)} \\
& = \tilde{C} \int_{\Omega} \left| (|u|^{\frac{m-1}{p}} \nabla u) |u|^{\frac{m-1}{q}} (g'(u) u + mg(u)) \right| dx.
\end{aligned}$$

On account of standard Hölder and Young inequalities, we can estimate the term on the right-hand side in terms of

$$\begin{aligned}
& \tilde{C} \left(\int_{\Omega} |u|^{m-1} |\nabla u|^p dx \right)^{1/p} \left(\int_{\Omega} |u|^{m-1} |g'(u) u + mg(u)|^q dx \right)^{1/q} \\
& = \tilde{C} \left(\frac{p}{p+m-1} \right) \left(m \int_{\Omega} |\nabla |u|^{\frac{p+m-1}{p}}|^p dx \right)^{1/p} \\
& \times \left(\int_{\Omega} |u|^{m-1} |g'(u) u + mg(u)|^q dx \right)^{1/q} m^{-1/p} \\
& \leq \varepsilon m \left(\frac{p}{p+m-1} \right)^p \int_{\Omega} |\nabla |u|^{\frac{p+m-1}{p}}|^p dx \\
& + \frac{\tilde{C}^q m^{-q/p}}{(\varepsilon p)^{q/p} q} \int_{\Omega} |u|^{m-1} |g'(u) u + mg(u)|^q dx.
\end{aligned} \tag{3.29}$$

Recalling (3.28), on the basis of (3.29), we can estimate

$$\begin{aligned}
& \langle f(u), |u|^{m-1} u \rangle_2 + \left\langle g(v), \frac{|v|^{m-1} v}{b} \right\rangle_{2,\Gamma} \\
& \geq \langle f(u) + c_B g(u), |u|^{m-1} u \rangle_2 - \frac{\tilde{C}^q m^{-q/p}}{(\varepsilon p)^{q/p} q} \left\langle |g'(u) u + mg(u)|^q, |u|^{m-1} \right\rangle_2 \\
& - \varepsilon m \left(\frac{p}{p+m-1} \right)^p \int_{\Omega} |\nabla |u|^{\frac{p+m-1}{p}}|^p dx.
\end{aligned} \tag{3.30}$$

Therefore, combining (3.4) with (3.5), then using (3.6) and (3.27), we arrive at the following inequality

$$\frac{d}{dt} \|U\|_{\mathbb{X}^{m+1}}^{m+1} + \gamma(\nu - \varepsilon) \int_{\Omega} \left| \nabla |u|^{\frac{p+m-1}{p}} \right|^p dx \leq Q_{\tau}(m) (\|U\|_{\mathbb{X}^{m+1}}^{m+1} + 1), \quad (3.31)$$

for some $\varepsilon \in (0, \nu/q)$. From this point on, the proof goes on exactly as in the proof of Theorem 3.2 (cf. **Steps 2** and **3**). We omit the details. ■

We will now verify the hypothesis in Proposition 3.4 for functions that satisfy the assumptions of Corollaries 2.6 and 2.7.

Corollary 3.5. *Assume that (H1) holds, and the functions $f, g \in C^1(\mathbb{R}, \mathbb{R})$ satisfy all the assumptions of Corollary 2.6-(i). Then, for any initial data U_0 in \mathbb{X}^2 , the corresponding solution $U(t)$ of (1.1)-(1.2), (2.12) belongs to \mathbb{X}^{∞} , for each $t > 0$, and estimate (3.3) holds.*

Proof. As in the proof of Corollary 2.6, the leading terms on the left-hand side of (3.27) are, for sufficiently large $|y| \gg 1$ and any $m \geq 1$,

$$c_f |y|^{r_1+m-1} + (|\Omega| \lambda)^{-1} c_g |y|^{r_2+m-1} - \frac{\tilde{C}_{\Omega}^q m^{-q/p}}{(\varepsilon p)^{q/p} q} c_g^q (r_2 + m - 1)^q |y|^{q(r_2-1)+m-1}, \quad (3.32)$$

for some $\varepsilon \in (0, \nu/q)$. From Corollary 2.6-(i), it holds $\gamma = \max(r_2, q(r_2 - 1)) < r_1$, so the coefficient of the highest order term in (3.32), for any $m \geq 1$, is $c_f > 0$. Therefore, the desired claim follows immediately from (3.27). ■

Corollary 3.6. *Let $h_2 = 0$ and assume $f, g \in C^1(\mathbb{R}, \mathbb{R})$ satisfy*

$$\lim_{|y| \rightarrow \infty} f'(s) = (r_1 - 1) c_f < 0 \text{ and } \lim_{|y| \rightarrow \infty} g'(s) = (r_2 - 1) c_g > 0.$$

Then, the conclusion of Theorem 3.2 applies.

Proof. The proof follows, for instance, from Theorem 3.2 since the functions f, g satisfy (3.2). ■

Having established that the weak solution is bounded for any positive times, we also have the following.

Proposition 3.7. *Let the assumptions of either Theorem 3.2 or Proposition 3.4 be satisfied. Then, any solution $U(t)$ of (1.1)-(1.2), (2.12) belongs to \mathbb{V}^p , for each $t > 0$, and the following estimate holds:*

$$\sup_{t \geq t_1} \left(\|U(t)\|_{\mathbb{V}^p}^p + \int_t^{t+1} \|\partial_t U(s)\|_{\mathbb{X}^2}^2 ds \right) \leq C_2, \quad (3.33)$$

for some positive constant C_2 , independent of t, ε and initial data.

Proof. It suffices to show (3.33). We first recall that, using assumption (2.3) and the fact that $Tr_D : W^{1,p}(\Omega) \rightarrow W^{1-1/p,p}(\Gamma)$ is a bounded map, from (3.1) we get

$$\sup_{t \geq t^\#} \int_t^{t+1} \left(\|U(s)\|_{\mathbb{V}^p}^p + \|u(s)\|_{L^{r_1}(\Omega)}^{r_1} \right) ds \leq C'_0, \quad (3.34)$$

for some positive constant C'_0 independent of time and initial data.

Let us now multiply equation (1.1) by $\partial_t u(t)$, then integrate over Ω , and multiply equation (1.2) by $\partial_t v(t)/b(x)$ and integrate over Γ . Adding the relations that we obtain, we deduce after standard transformations,

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left[\langle A(|\nabla u(t)|^2), 1 \rangle_2 + 2 \langle F(u(t)), 1 \rangle_2 + 2 \left\langle G(v(t)), \frac{1}{b} \right\rangle_{2,\Gamma} \right. \\ & \quad \left. - 2 \langle h_1, u(t) \rangle_2 - 2 \left\langle \frac{h_2}{b}, v(t) \right\rangle_{2,\Gamma} \right] \\ & = - \|\partial_t u\|_{L^2(\Omega)}^2 - \|\partial_t v\|_{L^2(\Gamma, dS/b)}^2, \end{aligned} \quad (3.35)$$

for all $t \geq t_+$ (with t_+ as in (3.3)), where we have set

$$A(|y|^2) = \int_0^{|y|^2} a(s) ds, \quad F(y) = \int_0^y f(s) ds, \quad G(y) = \int_0^y g(s) ds.$$

Next, let us define

$$\begin{aligned} \mathcal{E}(t) & := \langle A(|\nabla u(t)|^2), 1 \rangle_2 + 2 \langle F(u(t)), 1 \rangle_2 + 2 \langle G(v(t)), 1/b \rangle_{2,\Gamma} \\ & \quad - 2 \langle h_1, u(t) \rangle_2 - 2 \langle h_2/b, v(t) \rangle_{2,\Gamma} + C_{F,G}. \end{aligned} \quad (3.36)$$

Here the constant $C_{F,G} > 0$ is taken large enough in order to ensure that $\mathcal{E}(t)$ is nonnegative (recall that $F(u)$ and $G(v)$ are both bounded by (3.3)). On the other hand, on account of (3.3), one can easily check, using the fact

$$\langle A(|\nabla u|^2), 1 \rangle_2 \geq c_p \|\nabla u\|_{L^p(\Omega)}^p, \quad c_p > 0,$$

that there exists a positive constant c , independent of t and the initial data, such that

$$\|\nabla u(t)\|_{L^p}^p - c \leq \mathcal{E}(t), \quad (3.37)$$

for $t \geq \max\{t_+, t^\#\}$. From (3.35), we have

$$\frac{d\mathcal{E}(t)}{dt} + 2 \|\partial_t u\|_{L^2(\Omega)}^2 + 2 \|\partial_t v\|_{L^2(\Gamma, dS/b)}^2 = 0, \quad \forall t \geq \max\{t_+, t^\#\}. \quad (3.38)$$

Then, exploiting estimates (3.3), (3.34) and (3.37), we can apply to (3.38) the uniform Gronwall's lemma (see, e.g., [45]) and find a time $t_1 \geq 1$, depending on t_+ , $t^\#$, such that

$$\|\nabla u(t)\|_{L^p(\Omega)}^p \leq c, \quad \forall t \geq t_1, \quad (3.39)$$

for some positive constant c . Summing up, we conclude by observing that (3.33) follows from (3.39) and the boundedness of the trace map $Tr_D : W^{1,p}(\Omega) \rightarrow W^{1-1/p,p}(\Gamma)$. The proof is finished. ■

Finally, the above dissipative estimates and standard compactness results (see, e.g., [27, Section 2]), allow us to conclude the following.

Theorem 3.8. *Let the assumptions of either Theorem 3.2 or Proposition 3.4 be satisfied and, in addition, let (H2) hold. Then, the dynamical system $(\mathcal{S}_2(t), \mathbb{X}^2)$ generated by the initial value problem (1.1)-(1.2), (2.12) possesses the global attractor $\mathcal{A}_{gl} \subset \mathbb{X}^2$, which is a bounded subset of $\mathbb{V}^p \cap \mathbb{X}^\infty$. Moreover,*

$$\lim_{t \rightarrow +\infty} \text{dist}_{\mathbb{X}^{s_1, s_2}}(\mathcal{S}_2(t)B, \mathcal{A}_{gl}) = 0, \quad (3.40)$$

for any finite $s_1, s_2 \geq 2$, for all bounded subsets B of \mathbb{X}^2 .

3.2. Trajectory dynamical systems

In the final part of this section, we shall devote our attention to constructing the “usual” weak trajectory attractor and verify (using the maximum principle established in Theorem 3.2) that any solution, belonging to the attractor, is bounded so that uniqueness holds on the attractor. We will employ a slightly different construction (compared to e.g., [14] and references therein) of the trajectory attractor, which also looks more natural from the physical point of view. Namely any weak solution $U(t)$ of (1.1)-(1.2), (2.12) is included in the trajectory phase-space of the problem if and only if it can be obtained in the limit, as $\epsilon \rightarrow 0$, of the corresponding solutions $U_\epsilon(t)$ of the approximate system (2.17)-(2.18).

In order to define the trajectory dynamical system for weak solutions without uniqueness we need to introduce first the appropriate functional framework. First, let us recall estimates (2.25) and (2.30) which hold for any smooth solution $U = U_\epsilon$ of the approximate problem (2.17)-(2.18). By a standard application of Gronwall’s inequality (see, e.g., [45]), we get the well-known estimate

$$\begin{aligned} & \|U(t)\|_{\mathbb{X}^2}^2 + \int_s^t \left(\|U(r)\|_{\mathbb{V}^p}^p + \|u(r)\|_{L^{r_1}(\Omega)}^{r_1} + \tilde{c} \|v(r)\|_{L^{r_2}(\Gamma)}^{r_2} \right) dr \\ & \leq \|U(s)\|_{\mathbb{X}^2}^2 e^{-c(t-s)} + c \left(1 + \|h_1\|_{L^{r_1}'}^{r_1'} + \|h_2\|_{L^{r_2}(\Gamma)}^{r_2'} \right) (1 - e^{-c(t-s)}), \end{aligned} \quad (3.41)$$

for all $t \geq s \geq 0$, and some appropriate positive constant c . Here and below $\tilde{c} = 1$, if (H3b) is assumed and $\tilde{c} = 0$, when (H3a) holds. Let $\Theta_+^{w,loc}$ denote the local weak topology in the space

$$L^\infty(\mathbb{R}_+; \mathbb{X}^2) \cap L^p(\mathbb{R}_+; \mathbb{V}^p) \cap (L^{r_1}(\mathbb{R}_+; L^{r_1}(\Omega)) \times L^{r_2}(\mathbb{R}_+; L^{r_2}(\Gamma))).$$

By definition, a sequence $U_n(t) \rightarrow U(t)$, as $n \rightarrow \infty$, in the topology of $\Theta_+^{w,loc}$ if, for every $T > 0$,

$$\begin{aligned} U_n(t) & \rightarrow U(t) \quad \text{* -weakly in } L^\infty([0, T]; \mathbb{X}^2), \\ U_n(t) & \rightarrow U(t) \quad \text{weakly in } L^p([0, T]; \mathbb{V}^p), \\ U_n(t) & \rightarrow U(t) \quad \text{weakly in } L^{r_1}([0, T]; L^{r_1}(\Omega)) \times L^{r_2}([0, T]; L^{r_2}(\Gamma)). \end{aligned}$$

We recall that $\Theta_+^{w,loc}$ is a Hausdorff and Frechet-Urysohn space with a countable topology space (see, e.g., [44]). Next, let Θ_+^b be the Banach space defined as

$$\Theta_+^b := L^\infty(\mathbb{R}_+; \mathbb{X}^2) \cap L^p(\mathbb{R}_+; \mathbb{V}^p) \cap (L^{r_1}(\mathbb{R}_+ \times \Omega) \times L^{r_2}(\mathbb{R}_+ \times \Gamma)).$$

Note that the unit ball of Θ_+^b is compact in the local weak topology of $\Theta_+^{w,loc}$ (see [44]).

Definition 3.9. *A function $U \in \Theta_+^b$ is a solution of (1.1)-(1.2), (2.12) with $U_0 \in \mathbb{X}^2$ if it solves (1.1)-(1.2) in the sense of Definition 2.1, and there exists a sequence $\epsilon_n \rightarrow 0$, a sequence $u_{0\epsilon_n} \in C^\infty(\bar{\Omega})$, $U_{0\epsilon_n} = \begin{pmatrix} u_{0\epsilon_n} \\ u_{0\epsilon_n}|_\Gamma \end{pmatrix}$ and a sequence U_{ϵ_n} of classical solutions of the approximate problem (2.17)-(2.18) with $\epsilon = \epsilon_n$ such that*

$$U_0 = \mathbb{X}_w^2 - \lim_{n \rightarrow \infty} U_{0\epsilon_n} \text{ and } U = \Theta_+^{w,loc} - \lim_{n \rightarrow \infty} U_{\epsilon_n}. \quad (3.42)$$

Note that (3.42) implies in a standard way the weak convergence of

$$\mathcal{B}_{p,\epsilon_n} U_{\epsilon_n} \rightarrow \mathcal{B}_p U \text{ in } L_{loc}^q(\mathbb{R}_+; (\mathbb{V}^p)^*), \quad (3.43)$$

and consequently, the weak-star convergence of $\partial_t U_{\epsilon_n} \rightarrow \partial_t U$ in

$$L_{loc}^q(\mathbb{R}_+; (\mathbb{V}^p)^*) + (L_{loc}^{r_1'}(\mathbb{R}_+; L^{r_1}(\Omega)) \times L_{loc}^{r_2'}(\mathbb{R}_+; L^{r_2}(\Gamma))). \quad (3.44)$$

This gives the strong convergence $U_{\epsilon_n}(t) \rightarrow U(t)$ in $C_{loc}(\mathbb{R}_+; (\mathbb{V}^{k,p})^*)$ (see Section 2). Thus, any solution U of (1.1)-(1.2), (2.12) is weakly continuous with values in \mathbb{X}^2 (see Remark 2.4), and for any $t \geq 0$, we have the weak-convergence

$$U_{\epsilon_n}(t) \rightharpoonup U(t) \quad (3.45)$$

in the space \mathbb{X}^2 . It is important that we do not require the strong convergence in (3.45) even for $t = 0$.

We can now summarize the results in Section 2 by stating the following.

Proposition 3.10. *Let the assumptions of Theorem 2.2 hold. Then, for every $U_0 \in \mathbb{X}^2$, there exists at least one globally defined weak solution $U(t)$, $t \in \mathbb{R}_+$, of the degenerate problem (1.1)-(1.2), with $U(0) = U_0$, which can be obtained as a weak limit (3.42) of the corresponding solutions $U_{\epsilon_n}(t)$ of the approximate non-degenerate parabolic system (2.17)-(2.18).*

In order to construct the global attractor for the dynamical system associated with the degenerate parabolic system (1.1)-(1.2), we need the following definition.

Definition 3.11. *Let*

$$M_U(t) := \inf \left\{ \liminf_{n \rightarrow \infty} \|U_{\epsilon_n}(t)\|_{\mathbb{X}^2} : U = \Theta_+^{w,loc} - \lim_{n \rightarrow \infty} U_{\epsilon_n}, U_0 = \mathbb{X}_w^2 - \lim_{n \rightarrow \infty} U_{\epsilon_n}(0) \right\},$$

where the external infimum is taken over all possible sequences of solutions of the approximate problem (2.17)-(2.18), which converges as $\epsilon_n \rightarrow 0$ to the given solution U of the limit problem (1.1)-(1.2), (2.12).

Some simple properties of this M -functional are stated below.

Proposition 3.12. *Let U be a solution of problem (1.1)-(1.2), (2.12) and let $M_U(t)$ be the associated functional. Then,*

- (a) $\|U(t)\|_{\mathbb{X}^2} \leq M_U(t)$, for all $t \in \mathbb{R}_+$.
- (b) *The following estimate holds:*

$$\begin{aligned} & (M_U(t))^2 + \int_s^t \left(\|U(r)\|_{\mathbb{V}^p}^p + \|u(r)\|_{L^{r_1}(\Omega)}^{r_1} + \tilde{c} \|v(r)\|_{L^{r_2}(\Gamma)}^{r_2} \right) dr \\ & \leq (M_U(s))^2 e^{-c(t-s)} + c \left(1 + \|h_1\|_{L^{r_1}'}^{r_1'} + \|h_2\|_{L^{r_2}(\Gamma)'}^{r_2'} \right) (1 - e^{-c(t-s)}), \end{aligned} \quad (3.46)$$

where $t \geq s \geq 0$.

- (c) $M_{T(h)U}(t) \leq M_U(t+h)$, for all $h \geq 0$, where $(T(h)U)(t) := U(t+h)$.

Proof. (a) is immediate since the norm $\|\cdot\|_{\mathbb{X}^2}$ is weakly lower semicontinuous and the convergence of U_{ϵ_n} to U in $\Theta_+^{w,loc}$ implies the weak convergence $U_{\epsilon_n}(t) \rightarrow U(t)$ for every t . To prove (b), we note that due to the energy estimates for the approximate parabolic system (2.17)-(2.18) (cf. Section 2), we have

$$\begin{aligned} & \|U_{\epsilon_n}(t)\|_{\mathbb{X}^2}^2 + \int_s^t \left(\|U_{\epsilon_n}(r)\|_{\mathbb{V}^p}^p + \|u_{\epsilon_n}(r)\|_{L^{r_1}(\Omega)}^{r_1} + \tilde{c} \|v_{\epsilon_n}(r)\|_{L^{r_2}(\Gamma)}^{r_2} \right) dr \\ & \leq \|U_{\epsilon_n}(s)\|_{\mathbb{X}^2}^2 e^{-c(t-s)} + c \left(1 + \|h_1\|_{L^{r_1}'}^{r_1'} + \|h_2\|_{L^{r_2}(\Gamma)'}^{r_2'} \right) (1 - e^{-c(t-s)}), \end{aligned} \quad (3.47)$$

for every U_{ϵ_n} . By definition of M_U , for every $\delta > 0$, we can find an approximating sequence U_{ϵ_n} such that

$$\liminf_{n \rightarrow \infty} \|U_{\epsilon_n}(t)\|_{\mathbb{X}^2} \leq M_U(t) + \delta.$$

Passing to the limit, as $n \rightarrow \infty$, in (3.47), we have

$$\begin{aligned} & (M_U(t))^2 + \int_s^t \left(\|U(r)\|_{\mathbb{V}^p}^p + \|u(r)\|_{L^{r_1}(\Omega)}^{r_1} + \tilde{c} \|v(r)\|_{L^{r_2}(\Gamma)}^{r_2} \right) dr \\ & \leq \liminf_{n \rightarrow \infty} \left(\|U_{\epsilon_n}(t)\|_{\mathbb{X}^2}^2 + \int_s^t \left(\|U_{\epsilon_n}(r)\|_{\mathbb{V}^p}^p + \|u_{\epsilon_n}(r)\|_{L^{r_1}(\Omega)}^{r_1} + \tilde{c} \|v_{\epsilon_n}(r)\|_{L^{r_2}(\Gamma)}^{r_2} \right) dr \right) \\ & \leq (M_U(s) + \delta)^2 e^{-c(t-s)} + c \left(1 + \|h_1\|_{L^{r_1}'}^{r_1'} + \|h_2\|_{L^{r_2}(\Gamma)'}^{r_2'} \right) (1 - e^{-c(t-s)}), \end{aligned}$$

and since $\delta \rightarrow 0$ is arbitrary, we get the desired inequality. The third assertion is also immediate since the infimum in the definition of $M_{T(h)U}(t)$ is taken over the larger set of admissible approximating sequences than the infimum in the definition of $M_{U(t+h)}$. ■

We are now ready to construct the trajectory-phase space, the trajectory semigroup and the kernel associated with the degenerate problem (1.1)-(1.2). To this end, let $\mathcal{K}_+ \subset \Theta_+^b$ be

the set of all solutions of (1.1)-(1.2), in the sense of Definition 2.1, which corresponds to all $U_0 \in \mathbb{X}^2$, and let

$$T(h) : \mathcal{K}_+ \rightarrow \mathcal{K}_+, \quad h \geq 0, \quad (3.48)$$

be

$$(T(h)U)(t) := U(t+h). \quad (3.49)$$

We shall refer to \mathcal{K}_+ and $T(h) : \mathcal{K}_+ \rightarrow \mathcal{K}_+$ as the trajectory phase space and the trajectory dynamical system, respectively, associated with the degenerate parabolic system (1.1)-(1.2). In addition, we endow the set \mathcal{K}_+ with the topology induced by the embedding $\mathcal{K}_+ \subset \Theta_+^{w,loc}$ and we will say that a set $B \subset \mathcal{K}_+$ is M -bounded if

$$M_B(0) := \sup_{U \in B} M_U(0) < \infty.$$

Note that any M -bounded set $B \subset \mathcal{K}_+$ is bounded in the norm of Θ_+^b . Finally, a kernel $\mathcal{K} \subset \Theta^b$,

$$\Theta^b := L^\infty(\mathbb{R}; \mathbb{X}^2) \cap L^p(\mathbb{R}; \mathbb{V}^p) \cap (L^{r_1}(\mathbb{R} \times \Omega) \times L^{r_2}(\mathbb{R} \times \Gamma))$$

consists of all complete (defined for all $t \in \mathbb{R}$) bounded solutions of (1.1)-(1.2), (2.12) which can be obtained as the weak limit, as $\epsilon_n \rightarrow 0$, of the appropriate solutions of the approximate non-degenerate parabolic system (2.17)-(2.18). Namely, $U \in \mathcal{K}$ if and only if there exists a sequence $\epsilon_n \rightarrow 0$, a sequence of times $t_n \rightarrow -\infty$, and a bounded sequence of initial data $u_{0\epsilon_n} \in C^\infty(\overline{\Omega})$, $\|U_{0\epsilon_n}\|_{\mathbb{X}^2} \leq C$, such that the corresponding solutions U_{ϵ_n} of (2.17) on the interval $[t_n, +\infty)$ with initial data $U_{\epsilon_n}(t_n) = U_{0\epsilon_n}$ converges weakly in Θ^b to the complete solution U considered.

We now recall the definition of the global attractor for the trajectory dynamical system $(T(h), \mathcal{K}_+)$ (see [14] for more details; cf. also [38, 56]).

Definition 3.13. *A set $\mathcal{A}_{tr} \subset \mathcal{K}_+$ is a (weak) trajectory attractor associated with the degenerate parabolic system (1.1)-(1.2) (= global attractor for the trajectory dynamical system $(T(h), \mathcal{K}_+)$) if the following conditions are satisfied:*

- (i) \mathcal{A}_{tr} is compact in \mathcal{K}_+ and is M -bounded;
- (ii) It is strictly invariant, i.e., $T(h)\mathcal{A}_{tr} = \mathcal{A}_{tr}$, $h > 0$;
- (iii) It attracts the images of bounded (M -bounded) sets as $h \rightarrow \infty$, i.e., for every B bounded in \mathcal{K}_+ and every neighborhood $O(\mathcal{A}_{tr})$ of \mathcal{A}_{tr} (in the topology of $\Theta_+^{w,loc}$), there exists $h_0 = h_0(B, O)$ such that $T(h)B \subset O(\mathcal{A}_{tr})$, $\forall h \geq h_0$.

The next theorem can be considered as the second main result of this section.

Theorem 3.14. *Let $p \in (\frac{2N}{N+2}, \infty) \cap (1, \infty)$. Let all the assumptions of Theorem 2.2 be satisfied. Then, the degenerate parabolic problem (1.1)-(1.2), (2.12) possesses a trajectory attractor $\mathcal{A}_{tr} \subset \Theta_+^b$ and the following description holds:*

$$\mathcal{A}_{tr} = \Pi_{t \geq 0}(\mathcal{K}). \quad (3.50)$$

Here and below, $\Pi_{t \in I}(f)$ denotes the restriction on I of a function f defined on \mathbb{R} .

Proof. According to general theory [14, 38, 56], we are only required to check that the trajectory dynamical system is continuous and that it possesses a compact and M -bounded absorbing set. The continuity is immediate since $T(h)$ are continuous on $\Theta_+^{w,loc}$. The estimate (b) in Proposition 3.12 guarantees that the set

$$\mathcal{B} := \left\{ U \in \mathcal{K}_+ : (M_U(0))^2 \leq 2c \left(1 + \|h_1\|_{L^{r'_1}}^{r'_1} + \|h_2\|_{L^{r'_2}(\Gamma)}^{r'_2} \right) \right\} \quad (3.51)$$

will be absorbing for the semigroup $T(h) : \mathcal{K}_+ \rightarrow \mathcal{K}_+$. Moreover, this set is semi-invariant. This follows from Proposition 3.12, (c) since

$$\begin{aligned} (M_{T(h)U}(0))^2 &\leq (M_U(h))^2 \\ &\leq (M_U(0))^2 e^{-c(t-s)} + c \left(1 + \|h_1\|_{L^{r'_1}}^{r'_1} + \|h_2\|_{L^{r'_2}(\Gamma)}^{r'_2} \right) (1 - e^{-c(t-s)}) \\ &\leq 2c \left(1 + \|h_1\|_{L^{r'_1}}^{r'_1} + \|h_2\|_{L^{r'_2}(\Gamma)}^{r'_2} \right), \end{aligned}$$

for all $U \in \mathcal{B}$. Therefore, $T(h)\mathcal{B} \subset \mathcal{B}$. It remains to show that \mathcal{B} is compact. Due to the inequality (b) in Proposition 3.12, the set \mathcal{B} is bounded in Θ_+^b , and therefore precompact in $\Theta_+^{w,loc}$. Thus, we only need to show that \mathcal{B} is sequentially closed, i.e., if $U_n \in \mathcal{B}$ and $U = \Theta_+^{w,loc} - \lim_{n \rightarrow \infty} U_n$, then $U \in \mathcal{B}$ as well.

For $U_n \in \mathcal{B}$, $M_{U_n}(0)$ is bounded. By estimate (b) in Proposition 3.12 the sequence $M_{U_n}(t)$ is bounded for all $t \geq 0$. Moreover, since every U_n is a solution of the degenerate problem (1.1)-(1.2), there exists a sequence U_{n,ϵ_l} of solutions to the approximate non-degenerate problem (2.17)-(2.18) such that $\epsilon_l = \epsilon_{n,l} \rightarrow 0$ as $k \rightarrow \infty$, and

$$U_n = \Theta_+^{w,loc} - \lim_{k \rightarrow \infty} U_{n,\epsilon_l}.$$

Without loss of generality, we may assume that

$$|M_{U_n}(0) - \|U_{n,\epsilon_l}(0)\|_{\mathbb{X}^2}| \leq \frac{1}{n}, \quad \epsilon_l \leq \frac{1}{n}, \quad (3.52)$$

for all $l \in \mathbb{N}$, and we may also suppose that

$$\liminf_{n \rightarrow \infty} M_{U_n}(0) = \lim_{n \rightarrow \infty} M_{U_n}(0)$$

(we may pass to a subsequence in n if necessary). It remains to show that we can extract from $\{U_{n,\epsilon_l}\}_{n,l \in \mathbb{N}}$ a one parametric sequence which will converge to the limit function U . To this end, recall that the topology of $\Theta_+^{w,loc}$ is metrizable on every bounded set of Θ_+^b (see [44]). Let $\delta > 0$ be such that all U_n belong to the closed ball B_δ of Θ_+^b . Evidently, $U \in B_\delta \subset B_{2\delta}$, and we may also assume that $U_{n,\epsilon_l} \in B_{2\delta}$, for all $n, l \in \mathbb{N}$. Indeed, the sequence $U_{n,\epsilon_l}(0)$ is uniformly bounded in n, l , due to (3.52), and since $M_{U_n}(0)$ is bounded, then recalling

estimate (3.47), we also get that $U_{n,\epsilon_l}(t)$ is uniformly bounded with respect to n, l and t . Let $d(\cdot, \cdot)$ be a metric on $B_{2\delta}$. Thus,

$$\lim_{n \rightarrow \infty} d(U_n, U) = 0, \quad \lim_{k \rightarrow \infty} d(U_n, U_{n,\epsilon_l}) = 0,$$

for every n . Therefore, for any n , there exists $l_0 = l_0(n)$ such that $d(U, U_{n,\epsilon_l}) \leq 1/n$, for all $l \geq l_0$. Thus, we have $d(U, U_{n,\epsilon_{l_0}}) \rightarrow 0$ as $n \rightarrow \infty$, and therefore

$$U = \Theta_+^{w,loc} - \lim_{n \rightarrow \infty} U_{n,\epsilon_{l_0}}.$$

Moreover, thanks to (3.52), $\epsilon_{l_0} \rightarrow 0$ as $n \rightarrow \infty$ and so U is a solution of the degenerate problem, and

$$M_U(0) \leq \liminf_{n \rightarrow \infty} \|U_{n,\epsilon_l}(0)\|_{\mathbb{X}^2} = \liminf_{n \rightarrow \infty} M_{U_n}(0).$$

Thus, \mathcal{B} is indeed a compact semi-invariant absorbing set for $(T(h), \mathcal{K}_+)$, and the desired attractor can now be found in a standard way, as the ω -limit set of \mathcal{B} :

$$\mathcal{A}_{tr} = \omega(\mathcal{B}) = \bigcap_{h \geq 0} T(h)\mathcal{B}.$$

The description (3.50) is a standard corollary of this explicit formula and the diagonalization procedure described above. This completes the proof. ■

What is the connection between the dynamical system $(\mathcal{S}_2(t), \mathbb{X}^2)$ introduced at the end of Section 2, and the trajectory dynamical system $(T(h), \mathcal{K}_+)$ constructed here? It turns out that, under the assumptions of Proposition 2.8, the solution $U(t)$ of the system (1.1)-(1.2) is unique and, consequently, this parabolic system generates a semigroup in the classical phase space \mathbb{X}^2 , in a standard way by the formula (2.47). If we define the map

$$\Pi_0 : \mathcal{K}_+ \rightarrow \mathbb{X}^2, \quad \Pi_0(U(t)) := U(0),$$

we see that the map Π_0 is one-to-one and, in fact, Π_0 defines a Lipschitz homeomorphism between \mathcal{K}_+ and \mathbb{X}_w^2 (i.e., \mathbb{X}^2 endowed with the weak topology). Therefore, when uniqueness holds (for instance, if we require that the functions f, g satisfy (H2)),

$$\mathcal{S}_2(t) = \Pi_0 T(t) \Pi_0^{-1},$$

the trajectory dynamical system $(T(t), \mathcal{K}_+)$ is conjugated to the classical dynamical system $\mathcal{S}_2(t)$ defined on the phase space \mathbb{X}^2 endowed with the weak topology. However, we note that assumption (H2) is quite restrictive as it does not allow for a competing scenario between the nonlinearities f, g , as proposed in e.g., Corollaries 2.6 and 2.7. So without assumption (H2) the uniqueness problem for (1.1)-(1.2) is not known, and the classical semigroup $\mathcal{S}_2(t)$ can be defined as a semigroup of multi-valued maps only (see, e.g., [38] for further details). The trajectory dynamical approach allows us to avoid the use of multivalued maps and to apply the usual theory of global attractors to investigate the long term behavior of the degenerate parabolic system (1.1)-(1.2), (2.12).

As a consequence of Theorem 3.14, we can also state the following.

Corollary 3.15. *Let the assumptions of Theorem 3.14 hold and let $B \subset \mathcal{K}_+$ be an arbitrary M -bounded set. Then, for every $T \in \mathbb{R}_+$ and every $s \in (0, k)$, the following convergence holds:*

$$\lim_{h \rightarrow \infty} \text{dist}_{C((h, T+h); (\mathbb{V}^{s,p})^*)} (B|_{(h, T+h)}, \mathcal{A}_{tr}|_{(h, T+h)}) = 0. \quad (3.53)$$

Proof. Indeed, from the fact that every $U \in \mathcal{K}_+$ is a weak solution in the sense of Definition 2.1 (cf. (2.11)), we can express and estimate (as in Section 2) the time derivative of U , i.e., $\partial_t U \in L_{loc}^s(\mathbb{R}_+; (\mathbb{V}^{k,p})^*)$. Next, since the embedding

$$\left\{ U : U \in L_{loc}^\infty(\mathbb{R}_+; \mathbb{X}^2), \partial_t U \in L_{loc}^s(\mathbb{R}_+; (\mathbb{V}^{k,p})^*) \right\} \subset C_{loc}(\mathbb{R}_+; (\mathbb{V}^{s,p})^*), \quad (3.54)$$

is compact for every $0 < s < k$, then $\mathcal{K}_+ \subset C_{loc}(\mathbb{R}_+; (\mathbb{V}^{s,p})^*)$ is also compact in the sense that every M -bounded subset of \mathcal{K}_+ is a precompact set in the set $C_{loc}(\mathbb{R}_+; (\mathbb{V}^{s,p})^*)$. Thus, the above convergence (3.53) is an immediate corollary of Definition 3.13, (iii). ■

In the sequel, we shall also verify, under additional assumptions on the nonlinearities (which still allow for a competing behavior between f and g), that every solution U of the degenerate parabolic system is uniformly bounded on the (weak) trajectory attractor \mathcal{A}_{tr} . Then, using this fact we can establish that the solution of (1.1)-(1.2), (2.12) is unique on the attractor. From now, we will always assume that $p \geq 2$.

Theorem 3.16. *Let the assumptions of either Theorem 3.2 or Proposition 3.4 be satisfied. Then, for every complete solution $U \in \mathcal{K}$, we have $U \in L^\infty(\mathbb{R}; \mathbb{X}^\infty \cap \mathbb{V}^p)$ and the following estimate holds:*

$$\|U(t)\|_{\mathbb{X}^\infty \cap \mathbb{V}^p} \leq Q(1 + \|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Gamma)}), \quad (3.55)$$

for all $t \in \mathbb{R}$, for some monotone non-decreasing function Q independent of U , t and initial data. Moreover, $U \in C_{loc}(\mathbb{R}; \mathbb{X}^2)$, for every $U \in \mathcal{K}$.

Proof. The proof is essentially based on the maximum principle deduced in Section 2, and the description of \mathcal{K} from Theorem 3.14. Let $U \in \mathcal{K}$ be an arbitrary complete solution, i.e., let $\epsilon_n \rightarrow 0$, $t_n \rightarrow -\infty$ and $U_{\epsilon_n}(t)$, $t \geq t_n$ be the sequence of solutions of the approximate parabolic system (2.17) with $U_{\epsilon_n}(t_n) = U_{0\epsilon_n}$, where $\|U_{0\epsilon_n}\|_{\mathbb{X}^2} \leq C$, uniformly with respect to n . Let us now fix an arbitrary $T \in \mathbb{R}$. Then from the convergence $U_{\epsilon_n} \rightarrow U$, we also know that

$$U_{\epsilon_n}(T) \rightarrow U(T) \quad (3.56)$$

strongly in \mathbb{X}^2 (passing to a subsequence in n , if necessary), since the embedding $\mathcal{K}_+ \subset L_{loc}^p(\mathbb{R}_+; \mathbb{X}^2)$ is compact (i.e., any M -bounded subset of \mathcal{K}_+ is precompact in $L_{loc}^p(\mathbb{R}_+; \mathbb{X}^2)$). It follows from Theorem 3.14 and estimate (3.47) that

$$\begin{aligned} & \|U_{\epsilon_n}(T)\|_{\mathbb{X}^2}^2 + \int_T^{T+1} \left(\|U_{\epsilon_n}(r)\|_{\mathbb{V}^p}^p + \|u_{\epsilon_n}(r)\|_{L^{r_1}(\Omega)}^{r_1} + \tilde{c} \|v_{\epsilon_n}(r)\|_{L^{r_2}(\Gamma)}^{r_2} \right) dr \\ & \leq 2c \left(1 + \|h_1\|_{L^{r_1}'}^{r_1'} + \|h_2\|_{L^{r_2}(\Gamma)}^{r_2'} \right), \end{aligned} \quad (3.57)$$

where the constant $c > 0$ is independent of ϵ_n and T . By modifying the proof of Theorem 3.2 in a suitable way, we arrive at the following inequality for the approximate solutions U_{ϵ_n} ,

$$\begin{aligned} \sup_{t \geq t_n} \|U_{\epsilon_n}(t)\|_{\mathbb{X}^\infty} &\leq Q(c + \|U\|_{L^\infty(\mathbb{R}; \mathbb{X}^2)}), \\ &\leq Q(1 + \|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Gamma)}), \end{aligned} \quad (3.58)$$

for some monotone function Q independent of ϵ_n , T , t . Passing to the limit, in a standard way in (3.58), we may think that $U_{\epsilon_n}(T) \rightharpoonup U_0(T)$ weakly-star in \mathbb{X}^∞ , for some $U_0 \in \mathbb{X}^\infty$ such that U_0 satisfies

$$\|U_0(T)\|_{\mathbb{X}^\infty} \leq Q(1 + \|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Gamma)}). \quad (3.59)$$

This together with (3.56) gives $U(T) \in \mathbb{X}^\infty$ such that $U(T)$ satisfies the analogue of (3.59).

Finally, it remains to prove that $U(T)$ is also bounded in \mathbb{V}^p by essentially arguing as in the proof of Proposition 3.7. Note once again that in contrast to the limit case $\epsilon_n = 0$, the parabolic system (2.17) is non-degenerate if $\epsilon_n > 0$, and we have enough regularity of U_{ϵ_n} to justify the multiplication by the test functions $\partial_t U_{\epsilon_n}$ in the weak formulation (2.11) (cf. Definition 2.1). Analogous to (3.35)-(3.38), we get

$$\partial_t \mathcal{E}_{\epsilon_n}(t) + 2 \|\partial_t u_{\epsilon_n}(t)\|_{L^2(\Omega)}^2 + 2 \|\partial_t v_{\epsilon_n}(t)\|_{L^2(\Gamma, dS/b)}^2 = 0, \quad (3.60)$$

for the energy \mathcal{E}_{ϵ_n} defined in (3.36). Therefore, Gronwall's inequality applied to (3.60) yields as in (3.38)-(3.39),

$$\begin{aligned} \mathcal{E}_{\epsilon_n}(T) + \int_T^{T+1} \left(\|\partial_t u_{\epsilon_n}(r)\|_{L^2(\Omega)}^2 + \|\partial_t v_{\epsilon_n}(r)\|_{L^2(\Gamma, dS/b)}^2 \right) dr \\ \leq Q(1 + \|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Gamma)}), \end{aligned} \quad (3.61)$$

which in light of (3.37) gives that $U_{\epsilon_n}(T)$ is uniformly (in T, ϵ_n) bounded in \mathbb{V}^p . Passing now to the limit as $\epsilon_n \rightarrow 0$, we obtain in a standard way that $U(T)$ satisfies (3.61) as well. The desired inequality (3.55) follows immediately from (3.59) and (3.61). The last assertion in the theorem is a standard corollary of the energy identity (2.38). So, the proof is complete. ■

The (forward) uniqueness theorem holds for bounded solutions of (1.1)-(1.2), (2.12).

Theorem 3.17. *Let the assumptions of Theorem 3.16 be satisfied, and let $f, g \in C^1(\mathbb{R}, \mathbb{R})$. Consider two functions $U_1(t), U_2(t) \in C([0, T]; \mathbb{X}^2)$ which solve (1.1)-(1.2), (2.12) in the sense of Definition 2.1. In addition, let $U_1, U_2 \in L^\infty([0, T]; \mathbb{X}^\infty)$. Then, $U_1(0) = U_2(0)$ implies that $U_1(t) = U_2(t)$, for all $t \in [0, T]$.*

Proof. *The proof is an immediate consequence of inequality (2.45) (cf. Proposition 2.8). Indeed, for $E(t) := \|U_1(t) - U_2(t)\|_{\mathbb{X}^2}^2$, we have $E \in C^1(0, T)$, and*

$$\frac{1}{2} \partial_t E(t) \leq Q(\|U_1(t)\|_{\mathbb{X}^\infty}, \|U_2(t)\|_{\mathbb{X}^\infty}) E(t) \leq cE(t), \quad (3.62)$$

which yields the desired claim on account of the application of Gronwall's inequality. ■

Remark 3.18. We note that in contrast to the non-degenerate case of $a(s) \equiv \nu$, we do not know whether the backwards uniqueness theorem holds for bounded solutions of the parabolic system (1.1)-(1.2) for $p \neq 2$. Namely, if the equality $U_1(T) = U_2(T)$ holds for some $T \geq 0$, then we have $U_1(t) = U_2(t)$, for $t \leq T$, as well. Indeed, in the former case the parabolic system (1.1)-(1.2) is just a reaction-diffusion equation with dynamic boundary conditions, so we can establish additional regularity of the weak solutions in $L^\infty(\mathbb{R}_+; \mathbb{V}^{2,2}) \cap W^{1,\infty}(\mathbb{R}_+; \mathbb{V}^{1,2})$, following [23, Theorem 2.3] (see also below). Thus, exploiting a well-known theorem (see, e.g., [42, Theorem 11.10]; cf. also [45, Chapter III]), we can easily establish the backwards uniqueness result in this case.

Finally, we observe that since a bounded weak solution $U(t) \in L^\infty(\mathbb{R}_+; \mathbb{X}^\infty \cap \mathbb{V}^p)$ is unique, we may define a global attractor \mathcal{A}_{gl} for the parabolic system (1.1)-(1.2), (2.12) by the standard expression

$$\mathcal{A}_{gl} := \Pi_0 \mathcal{A}_{tr}, \quad (3.63)$$

and define a classical semigroup on this attractor via

$$\mathcal{S}_t : \mathcal{A}_{gl} \rightarrow \mathcal{A}_{gl}, \quad \mathcal{S}_t U(0) = U(t), \quad (3.64)$$

Here, $U(t)$ is the unique (bounded) weak solution of (1.1)-(1.2), (2.12), such that $U(t)$ satisfies the energy identity (2.38). We also note that estimate (3.55) gives a uniform estimate of the L^∞ -norm of the trajectories belonging to the attractor \mathcal{A}_{gl} . Therefore, the growth rate of the nonlinearities f, g with respect to U becomes nonessential for further investigations of global attractors and we can study them exactly as in, e.g., [42, 45].

3.3. Strong trajectory attractors: the semilinear case

Let $(T_2(t), \mathcal{K}_+)$ be the trajectory dynamical system associated with the reaction-diffusion equation (1.4), subject to the dynamic boundary condition (1.5) (see Section 1). In this section, we shall establish additional regularity estimates for the weak solutions of (1.4)-(1.5), and obtain as a by-product, that the weak trajectory attractor \mathcal{A}_{tr} , constructed in Theorem 3.14, is in fact a *strong* trajectory attractor. In order to do so we will verify, for every $U \in \mathcal{A}_{tr}$, that the attraction property and the compactness holds not only in the weak topology of $\Theta_+^{w,loc}$, but also in the strong topology of $\Theta_+^{s,loc}$. The definition of a strong trajectory attractor is obtained by replacing the weak attraction condition (iii) in Definition 3.13 by the condition of strong attraction in the topology of $\Theta_+^{s,loc}$.

We have the following proposition, whose proof goes essentially as in [23, Theorem 2.3].

Proposition 3.19. *Let the assumptions of either Theorem 3.2 or Proposition 3.4 be satisfied, and $f, g \in C^1(\mathbb{R}, \mathbb{R})$. Every bounded complete weak solution $U \in \mathcal{K}$ of problem (1.4)-(1.5) belongs to $L^\infty(\mathbb{R}; \mathbb{V}^{2,2}) \cap W^{1,\infty}(\mathbb{R}; \mathbb{V}^{1,2})$, and the following inequality holds for $t \in \mathbb{R}$,*

$$\begin{aligned} & \|U(t)\|_{\mathbb{V}^{2,2}}^2 + \|\partial_t U(t)\|_{\mathbb{V}^{1,2}}^2 + \int_t^{t+1} \|\partial_t^2 U(r)\|_{\mathbb{X}^2}^2 dr \\ & \leq Q(1 + \|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Gamma)}), \end{aligned} \quad (3.65)$$

where Q is a monotone nondecreasing function, independent of t, U and the data.

Proof. Indeed, having established the L^∞ -estimate (3.55), (3.65) can be easily derived using a standard technique for parabolic equations with dynamic boundary conditions (see [23, Theorem 2.3] for further details; cf. also [10, 25, 26]). ■

Consequently, we have shown the following.

Theorem 3.20. *Under the assumptions of Proposition 3.19, the weak attractor \mathcal{A}_{tr} , constructed in Theorem 3.14, is a strong trajectory attractor for the trajectory dynamical system $(T_2(t), \mathcal{K}_+)$. Moreover, \mathcal{A}_{tr} is compact in $C_{loc}(\mathbb{R}_+; \mathbb{V}^{2-s,2})$, for any $s \in (0, 1]$, and the set*

$$\partial_t \mathcal{A}_{tr} := \{\partial_t U : U \in \mathcal{A}_{tr}\}$$

is compact in $C_{loc}(\mathbb{R}_+; \mathbb{V}^{1-l,2})$, for any $l \in (0, 1/2)$.

Proof. It is sufficient to show that the set $T(t_\#)\mathcal{B}$, where $t_\# \geq 1$ is sufficiently large and \mathcal{B} is the absorbing set (3.51) for the semigroup $\{T(t)\}$ is compact in the strong topology of the space $\Theta_+^{w,loc}$. We observe that for every $t \geq t_\#$ and any $U \in \mathcal{K}_+$, we have

$$T(t)U \in L^\infty(\mathbb{R}_+; \mathbb{V}^{2,2} \cap \mathbb{X}^\infty) \cap W^{1,\infty}(\mathbb{R}_+; \mathbb{V}^{1,2}) \cap W^{2,2}(\mathbb{R}_+; \mathbb{X}^2). \quad (3.66)$$

Thus, it is enough to prove that the kernel \mathcal{K}_+ , for the attractor \mathcal{A}_{tr} defined by (3.50), is compact in the strong local topology of the space Θ_+^b . We first note that, due to estimates (3.55), (3.65) and (3.66), we have

$$\mathcal{K}_+ \text{ is bounded in } C_b(\mathbb{R}_+; \mathbb{V}^{2-s,2}), \text{ for any } s \in (0, 1],$$

and moreover, it is compact in the local topology

$$\mathcal{K}_+ \subset C_{loc}(\mathbb{R}_+; \mathbb{V}^{2-s,2}). \quad (3.67)$$

This follows from the following embedding

$$\{U : U \in L_{loc}^\infty(\mathbb{R}_+; \mathbb{V}^{2,2}), \partial_t U \in L_{loc}^\infty(\mathbb{R}_+; \mathbb{V}^{1,2})\} \subset C_{loc}(\mathbb{R}_+; \mathbb{V}^{2-s,2}).$$

which is compact. Thus, in view of (3.67) and the boundedness of \mathcal{K}_+ in $L^\infty(\mathbb{R}_+; \mathbb{X}^\infty)$, we immediately see that \mathcal{K}_+ is also compact in the (strong) local topology of Θ_+^b , i.e., we have

$$\mathcal{K}_+ \stackrel{c}{\subset} \Theta_{+,loc}^b := L_{loc}^\infty(\mathbb{R}_+; \mathbb{X}^2) \cap L_{loc}^2(\mathbb{R}_+; \mathbb{V}^{1,2}) \cap (L_{loc}^{r_1}(\mathbb{R}_+; L^{r_1}(\Omega)) \times L_{loc}^{r_2}(\mathbb{R}_+; L^{r_2}(\Gamma))).$$

The second statement follows analogously using the compactness of the following embedding

$$\{\partial_t U : U \in L_{loc}^\infty(\mathbb{R}_+; \mathbb{V}^{1,2}), \partial_t^2 U \in L_{loc}^2(\mathbb{R}_+; \mathbb{X}^2)\} \subset C_{loc}(\mathbb{R}_+; \mathbb{V}^{1-l,2}).$$

The proof is finished. ■

Corollary 3.21. *Under the validity of assumptions of Theorem 3.20, the reaction-diffusion equation (1.4), with the dynamic boundary condition (1.5) possesses a global attractor $\mathcal{A}_{gl}(= \mathcal{K}(0))$, defined by (3.63)-(3.64), with \mathcal{A}_{gl} bounded in $\mathbb{V}^{2,2} \cap \mathbb{X}^\infty$.*

Remark 3.22. *It is worth mentioning that one can also establish more regularity of the weak solution $u \in C^s(\overline{\Omega})$ as much as it is allowed by the regularity of Ω , f , g and the external sources h_1, h_2 . Taking advantage of the regularity result in Corollary 3.21, we can prove that the global attractor \mathcal{A}_{gl} is finite dimensional, by establishing the existence of a more refined object called exponential attractor \mathcal{E}_{gl} . However, since the associated solution semigroup \mathcal{S}_t happens to be (uniformly quasi-) differentiable with respect to the initial data, on the attractor \mathcal{A}_{gl} , we can instead employ a volume contraction argument (see, e.g., [23]).*

We can now extend the results in [23] for the case of nonlinear boundary conditions, without requiring that the restrictive condition (H2) holds. More precisely, we can establish the following upper-bound on the dimension of the global attractor \mathcal{A}_{gl} .

Theorem 3.23. *Provided that $f, g \in C^2(\mathbb{R}, \mathbb{R})$ satisfy the assumptions of Corollary 3.21 for as long as (3.2) holds, the fractal dimension of $\mathcal{A}_{gl} = \mathcal{K}(0)$ admits the estimate*

$$\dim_F \mathcal{A}_{gl} \leq \mathcal{C}_1 (1 + \nu^{-(N-1)}), \text{ for } N \geq 2 \quad (3.68)$$

and

$$\dim_F \mathcal{A}_{gl} \leq \mathcal{C}_1 (1 + \nu^{-1/2}), \text{ for } N = 1. \quad (3.69)$$

where \mathcal{C}_1 depends only on Ω , Γ and the sources h_1, h_2 .

Proof. First, it is easy to establish that the flow $\mathcal{S}_t : \mathcal{A}_{gl} \rightarrow \mathcal{A}_{gl}$ generated by the reaction-diffusion equation (1.4) and dynamic boundary condition (1.5) is uniformly differentiable on \mathcal{A}_{gl} , with differential

$$\mathbf{L}(t, U(t)) : \Phi = \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} \in \mathbb{X}^2 \mapsto V = \begin{pmatrix} v \\ \varphi \end{pmatrix} \in \mathbb{X}^2, \quad (3.70)$$

where V is the unique solution to

$$\partial_t v = \nu \Delta v - f'(u(t))v, \quad \left(\partial_t \varphi + \nu b \partial_{\mathbf{n}} v + g'(u(t)|_{\Gamma})v \right)_{|\Gamma} = 0, \quad (3.71)$$

$$V(0) = \Phi.$$

Indeed, the uniform differentiability result follows from the assumptions on f, g and is a consequence of the boundedness of \mathcal{A}_{gl} into $\mathbb{V}^{2,2} \cap \mathbb{X}^\infty$ (see [23]). In order to deduce (3.68)-(3.69), it is sufficient (see, e.g., [14, Chapter III, Definition 4.1]) to estimate the j -trace of the operator

$$\mathbf{L}(t, U(t)) = \begin{pmatrix} \nu \Delta - f'(u(t)) & 0 \\ -b\nu \partial_{\mathbf{n}} & -g'(v(t)) \end{pmatrix}.$$

We have

$$\begin{aligned}
Trace(\mathbf{L}(t, U(t)) Q_m) &= \sum_{j=1}^m \langle \mathbf{L}(t, U(t)) \varphi_j, \varphi_j \rangle_{\mathbb{X}^2} \\
&= \sum_{j=1}^m \langle \nu \Delta \varphi_j, \varphi_j \rangle_2 - \sum_{j=1}^m \langle \nu \partial_{\mathbf{n}} \varphi_j, \varphi_j \rangle_{2, \Gamma} \\
&\quad - \sum_{j=1}^m \langle f'(u(t)) \varphi_j, \varphi_j \rangle_2 - \sum_{j=1}^m \langle g'(v(t)) \varphi_j, \varphi_j \rangle_{2, \Gamma},
\end{aligned}$$

where the set of real-valued functions $\varphi_j \in \mathbb{X}^2 \cap \mathbb{V}^{1,2}$ is an orthonormal basis in $Q_m \mathbb{X}^2$. By Theorem 3.16, it follows that every bounded complete trajectory $U(t)$, $t \in \mathbb{R}$, for the dynamical system $(\mathcal{S}_t, \mathcal{K}(0))$ is uniformly bounded in $\mathbb{V}^{2,2} \cap \mathbb{X}^\infty$, namely, it holds:

$$\sup_{t \in \mathbb{R}} \|U(t)\|_{\mathbb{X}^\infty} \leq \mathcal{C}, \quad (3.72)$$

where the positive constant \mathcal{C} is independent of $U(t)$, ν , Ω , Γ , but depends on the L^∞ -norms of the external forces h_1 , h_2 and the constants in (3.7) (this follows from the usual description of the global attractor, see, e.g., [6], and the main estimates). Thus, exploiting (2.36) once more there holds

$$\max_{t \in \mathbb{R}} \sup \left\{ \left\| f'(u(t)) \right\|_{L^\infty(\Omega)}, \left\| g'(v(t)) \right\|_{L^\infty(\Gamma)} \right\} \leq C_\star, \quad (3.73)$$

for some $C_\star > 0$ which depends on \mathcal{C} and the growth rates $r_1, r_2 \geq 1$; we find

$$Trace(\mathbf{L}(t, U) Q_m) \leq -\nu \sum_{j=1}^m \|\nabla \varphi_j\|_2^2 + C_\star m.$$

From [23, Proposition 5.5], we obtain

$$\begin{aligned}
Trace(\mathbf{L}(t, U) Q_m) &\leq -\nu c_1 C_W(\Omega, \Gamma) m^{\frac{1}{N-1}+1} + (c_1 \nu C_W(\Omega, \Gamma) + C_\star) m \\
&=: \rho(m),
\end{aligned}$$

for some absolute positive constant c_1 independent of the parameters of the problem (C_W is given explicitly in [23, Theorem 5.4]). The function $\rho(y)$ is concave. The root of the equation $\rho(y) = 0$ is

$$y^* = \left(1 + \frac{C_\star}{\nu c_1 C_W(\Omega, \Gamma)} \right)^{N-1}. \quad (3.74)$$

Thus, we can apply a well-known result [14, Chapter VIII, Theorem 3.1] to deduce that $\dim_F \mathcal{A}_{gl} \leq y^*$, from which (3.68) follows. The case $N = 1$ is similar. ■

Remark 3.24. In the competing scenario (2.7), we can also explicitly estimate from above the dimension of the global attractor $\mathcal{A}_{gl} = \mathcal{K}(0)$ for the dynamical system $(\mathcal{S}_t, \mathcal{A}_{gl})$ defined by (3.63)-(3.64). However, in this case the upper bound does not seem to be as sharp as in (3.68)-(3.69). Indeed, by Theorem 3.16 (see also Theorem 3.2), it follows that every bounded complete trajectory $U(t)$, $t \in \mathbb{R}$, for the dynamical system $(\mathcal{S}_t, \mathcal{K}(0))$ satisfies (3.72) with constant $\mathcal{C} = \mathcal{C}_\nu \sim \nu^{\sigma_1}$ as $\nu \rightarrow 0^+$, for some $\sigma_1 < 0$; by assumption (H3), this implies that the constant C_\star in (3.73) behaves as $C_\star = C_\star(\nu) \sim \nu^{\sigma_2}$, for some $\sigma_2 < -1$ depending on σ_1 and r_1, r_2 . Hence, in this case there seems to be a discrepancy between the upper bound (3.74) and the lower bound in (1.13).

3.4. A blowup result

As pointed out at the beginning of this article, nonlinear dissipative boundary conditions cannot prevent blowup of some solutions of (1.4)-(1.5) when the non-dissipative interior term f is superlinear, i.e., when f satisfies (1.9) for some $r_1 > 2$. We will follow some arguments similar to ones presented in [43] for nonlinear Robin boundary conditions, by constructing some subsolutions to (1.4)-(1.5) which become unbounded in finite time at some points of the boundary Γ . We begin with the following notion.

Definition 3.25. A function $v : \overline{\Omega} \times (0, T) \rightarrow \mathbb{R}$ is a subsolution of (1.4)-(1.5) if it satisfies

$$\begin{cases} \partial_t v - \nu \Delta v + f(v) - h_1(x) \leq 0, & \text{in } \Omega \times (0, T), \\ \partial_t v + \nu b \partial_{\mathbf{n}} v + g(v) - h_2(x) \leq 0, & \text{on } \Gamma \times (0, T) \end{cases} \quad (3.75)$$

and

$$v(0) \leq u_0 \text{ in } \overline{\Omega}. \quad (3.76)$$

Analogously, the function v is called a supersolution if the inequalities in (3.75)-(3.76) are reversed.

From [4, Section 7], we have the following

Proposition 3.26. Let u be a solution of (1.4)-(1.5), and let v and \tilde{v} be a subsolution and supersolution, respectively, of (1.4)-(1.5), in the sense of Definition 3.25. Then,

$$v(x, t) \leq u(x, t) \leq \tilde{v}(x, t),$$

for all $x \in \overline{\Omega}$ and for as long as they exist. In particular, if $f(0) \leq 0$ and $g(0) \leq 0$, and if $u_0 \geq 0$, then the solution of (1.4)-(1.5) satisfies $u(x, t) \geq 0$, for all $x \in \overline{\Omega}$, for as long as it exists.

We aim to construct subsolutions by comparing solutions of (1.4)-(1.5) with classical solutions fulfilling certain Dirichlet conditions on the time lateral boundary $\Gamma \times (0, T)$. For that purpose, the following result is very useful (see, [20, 50]; cf. also [4, 43]). In what follows, it suffices to consider the case $h_1 \equiv 0$, $h_2 \equiv 0$.

Proposition 3.27. *Assume that there are a C^1 -concave decreasing function $h(s)$ and a number $s_0 \geq 0$ such that*

$$\limsup_{s \rightarrow \infty} h'(s) < -\lambda_{1,D},$$

where $\lambda_{1,D} > 0$ is the first eigenvalue of $-\nu\Delta$ with Dirichlet boundary conditions. Moreover, suppose that $h(s) < 0$ for all $s > s_0$ such that

$$\int_{s_0}^{\infty} \frac{ds}{|h(s)|} < \infty. \quad (3.77)$$

Then, there are positive (smooth and locally well-defined) solutions v of the reaction-diffusion equation

$$\partial_t v - \nu\Delta v + f(v) = 0, \quad \text{in } \Omega \times (0, T), \quad (3.78)$$

subject to the boundary and initial conditions

$$\begin{cases} v = 0, & \text{on } \Gamma \times (0, T), \\ v(0) := v_0, & \text{in } \Omega, \end{cases} \quad (3.79)$$

that blowup in finite time.

Theorem 3.28. *Let the assumptions of Proposition 3.27 be satisfied, and assume in addition that $f(s) \leq h(s) < 0$ for all $s > s_0$ such that (3.77) holds. Then, for any nonlinear function g , there exist solutions of (1.4)-(1.5) that blowup in finite time. Moreover, there exists a positive function $w_0(x)$ such that all solutions with initial data u_0 greater or equal than $w_0 + v_0$, blowup in finite time.*

Proof. Let φ_1 be the principal eigenfunction associated with $\lambda_{1,D} > 0$ such that $\|\varphi_1\|_{L^1(\Omega)} = 1$. It is well-known that $\varphi_1 > 0$ in Ω by the maximum principle. Thus, by choosing $A := \max\{s_0, s'_0, 0\}$ such that $h'(s) < -\lambda_{1,D}$, for all $s > s'_0$, we can define $w(x) := \delta\varphi_1(x) > 0$, for some $\delta > 0$, such that

$$\nu b \partial_{\mathbf{n}} w + g(A) \leq 0 \text{ on } \Gamma. \quad (3.80)$$

Note that it is always possible to fix $\delta > 0$ since, by the maximum principle once again, we have $\partial_{\mathbf{n}}\varphi_1 < 0$ on Γ . Let us now define

$$\underline{u}(x, t) = w_0(x) + v(x, t),$$

where $w_0(x) := w(x) + A$ and v is a solution of (3.78)-(3.79). Arguing as in the proof of [43, Theorem 4.3], we can easily establish that

$$\partial_t \underline{u} - \nu\Delta \underline{u} + f(\underline{u}) \leq 0, \text{ in } \Omega \times (0, T).$$

On the other hand, on $\Gamma \times (0, T)$, we have

$$\partial_t \underline{u} + \nu b \partial_{\mathbf{n}} \underline{u} + g(\underline{u}) = \nu b \partial_{\mathbf{n}} w + \nu b \partial_{\mathbf{n}} v + g(A),$$

but since $\partial_{\mathbf{n}}v \leq 0$ on Γ , by the choice of A and w (cf. (3.80)), it follows

$$\partial_t \underline{u} + \nu b \partial_{\mathbf{n}} \underline{u} + g(\underline{u}) \leq 0, \text{ on } \Gamma \times (0, T).$$

Consequently, we deduce that \underline{u} is a subsolution of (1.4)-(1.5). Therefore, on account of Propositions 3.26 and 3.27, all solutions $u = u(x, t)$ of (1.4)-(1.5) that satisfy $u_0 \geq w_0 + v_0$ in $\overline{\Omega}$, blowup in finite time. The proof is complete. ■

Corollary 3.29. *Assume that f satisfies*

$$\limsup_{s \rightarrow \infty} \frac{f(s)}{s(\ln s)^l} < 0,$$

for some $l > 1$. Then the conclusion of Theorem 3.28 applies.

4. Appendix

In this section, we state two basic results which are essential to the analysis of problem (1.1)-(1.2). The first lemma is just a variation of a result in [28, Section 5] using well-known facts about nonlinear forms and maximal monotone operators in Sobolev spaces. For the convenience of the reader, we give below a simple proof of that result.

Lemma 4.1. *Let $a(s)$ satisfy all the assumptions in Theorem 2.2, and let $U \in \mathbb{V}^p$ be fixed. Then the functional $V \mapsto \mathcal{B}_p(U, V)$, defined by (2.13), belongs to $(\mathbb{V}^p)^*$. Moreover, \mathcal{B}_p is strictly monotone, hemicontinuous and coercive.*

Proof. We will make use of [28, Definitions 2.1 and 2.2]. In our case, for $p > p_0$,

$$V := \mathbb{V}^p \subset \mathbb{X}^2 = (\mathbb{X}^2)^* \subset V^* = (\mathbb{V}^p)^*.$$

Let $U = \begin{pmatrix} u \\ u|_{\Gamma} \end{pmatrix} \in \mathbb{V}^p$ be fixed. It is clear that $\mathcal{B}_p(U, \cdot)$ is linear. Let $V = \begin{pmatrix} v \\ v|_{\Gamma} \end{pmatrix} \in \mathbb{V}^p$. Exploiting (2.3), we obtain

$$|\mathcal{B}_p(U, V)| \leq \|u\|_{W^{1,p}(\Omega)}^{p-1} \|V\|_{\mathbb{V}^p}. \quad (4.1)$$

This implies that $\mathcal{B}_p(U, \cdot) \in (\mathbb{V}^p)^*$, for every $U \in \mathbb{V}^p$.

Next, let $U, V \in \mathbb{V}^p$. Then, recalling (2.4), we have

$$\begin{aligned} & \mathcal{B}_p(U, U - V) - \mathcal{B}_p(V, U - V) \\ &= \int_{\Omega} (a(|\nabla u|^2) \nabla u - a(|\nabla v|^2) \nabla v) \cdot \nabla(u - v) dx + \int_{\Omega} (|u|^{p-2} u - |v|^{p-2} v) (u - v) dx \\ &\geq \int_{\Omega} (|u| + |v|)^{p-2} |u - v|^2 dx \geq 0, \end{aligned} \quad (4.2)$$

which shows that \mathcal{B}_p is monotone. This estimate also shows that

$$\mathcal{B}_p(U, U - V) - \mathcal{B}_p(V, U - V) > 0,$$

for all $U, V \in \mathbb{V}^p$ with $U \neq V$. Thus, \mathcal{B}_p is strictly monotone. The continuity of the norm function implies that \mathcal{B}_p is hemicontinuous. Finally, it is easy to deduce that

$$\lim_{\|U\|_{\mathbb{V}^p} \rightarrow +\infty} \frac{\mathcal{B}_p(U, U)}{\|U\|_{\mathbb{V}^p}} = +\infty, \quad (4.3)$$

which shows that \mathcal{B}_p is coercive. The proof is finished. ■

The following lemma is useful in controlling surface integrals by means of volume integrals. The proof follows from an application of the divergence theorem to the function $w = (|u|^s \xi) \cdot \mathbf{n}$, for a smooth vector field $\xi \in C^1(\overline{\Omega}, \mathbb{R}^N)$ such that $\xi \cdot \mathbf{n} = 1$.

Lemma 4.2. *Let $p > 1$, $s > 1$ and $u \in W^{1,p}(\Omega)$. Then for every $\varepsilon > 0$, there exists a positive constant $C_\varepsilon = C(\varepsilon, s, p)$, independent of u , such that*

$$\|u\|_{L^s(\Gamma)}^s \leq \varepsilon \|\nabla u\|_{L^p(\Omega)}^p + C_\varepsilon \left(\|u\|_{L^\gamma(\Omega)}^\gamma + 1 \right),$$

where $\gamma = \max(s, p(s-1)/(p-1))$.

Acknowledgement 4.3. *The author wishes to express his thanks to Martin Meyries for bringing his attention to paper [43], and for all the interesting and fruitful discussions on the issues presented in this article.*

References

- [1] M. Aida, M. Efendiev, A. Yagi, *Quasi-linear abstract parabolic evolution equations and exponential attractors*, Osaka J. Math., 42 (2005), 101–132.
- [2] G. Akagi, M. Ôtani, *Evolution inclusions governed by the difference of two subdifferentials in reflexive Banach spaces*, J. Differential Equations, 209 (2005), 392–415.
- [3] E. DiBenedetto, “Degenerate Parabolic Equations,” Universitext, Springer-Verlag, New York, 1993.
- [4] J. von Below, M.G. Maily, *Blow up for reaction diffusion equations under dynamical boundary conditions*, Comm. Partial Differential Equations 28 (2003), no. 1-2, 223–247.
- [5] H. Brezis, “Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert,” North-Holland Math. Stud., vol. 5, North-Holland, Amsterdam, 1973.
- [6] A. V. Babin, M. I. Vishik, “Attractors of Evolutions Equations,” North-Holland, Amsterdam, 1992.
- [7] J. W. Cholewa, T. Dlotko, “Global Attractors in Abstract Parabolic Problems,” Cambridge University Press, 2000.

- [8] R. Chill, A. Fiorenza, *Convergence and decay rate to equilibrium of bounded solutions of quasi-linear parabolic equations*, J. Differential Equations, 228 (2006), 611–632.
- [9] A. N. Carvalho, C. B. Gentile, *Asymptotic behavior of nonlinear parabolic equations with monotone principal part*, J. Math. Anal. Appl., 280 (2003), 252–272.
- [10] C. Cavaterra, C.G. Gal, M. Grasselli, A. Miranville, *Phase-field systems with nonlinear coupling and dynamic boundary conditions*, Nonlinear Anal. 72 (2010), no. 5, 2375–2399.
- [11] A. Constantin, J. Escher, *Global existence for fully parabolic boundary value problems*, NoDEA Nonlinear Differential Equations Appl. 13 (2006), no. 1, 91–118.
- [12] J. Crank, “The mathematics of Diffusion,” Second edition, Clarendon Press, Oxford, 1975.
- [13] I. Chueshov, B. Schmalfuss, *Qualitative behavior of a class of stochastic parabolic PDEs with dynamical boundary conditions*, Discrete Contin. Dyn. Syst., 18 (2007), 315–338.
- [14] V.V. Chepyzhov, M.I. Vishik, *Attractors for equations of mathematical physics*, Amer. Math. Soc., Providence, RI, 2002.
- [15] M. A. Efendiev, M. Otani, *Infinite-dimensional attractors for evolution equations with p -Laplacian and their Kolmogorov entropy*, Diff. Int. Eq., 20 (2007), 1201–1209.
- [16] J. Escher, *Quasilinear parabolic systems with dynamical boundary conditions*, Comm. Partial Differential Equations, 18 (1993), 1309–1364.
- [17] J. Escher, *On quasilinear fully parabolic boundary value problems*, Differential Integral Equations, 7 (1994), 1325–1343.
- [18] A. Favini, G. Ruiz Goldstein, J. A. Goldstein, S. Romanelli, *The heat equation with general Wentzell boundary conditions*, J. Evol. Equations, 2 (2002), 1–19.
- [19] A. Favini, G. Ruiz Goldstein, J. A. Goldstein, S. Romanelli, *The heat equation with nonlinear general Wentzell boundary condition*, Adv. Differential Equations, 11 (2006), 481–510.
- [20] A. Friedman, B. McLeod, *Blow-up of positive solutions of semilinear heat equations*, Indiana Univ. Math. J. 34 (1985), 425–447.
- [21] M. Fila, P. Quittner, *Large time behavior of solutions of a semilinear parabolic equation with a nonlinear dynamical boundary condition*, Topics in nonlinear analysis, 251–272, Progr. Nonlinear Differential Equations Appl., 35, Birkhäuser, Basel, 1999.
- [22] Z. H. Fan, C. K. Zhong, *Attractors for parabolic equations with dynamic boundary conditions*, Nonlinear Anal., 68 (2008), 1723–1732.

- [23] C. G. Gal, *Sharp estimates for the global attractor of scalar reaction-diffusion equations with a Wentzell boundary condition*, J. Nonlinear Science, 22 (2012), 85-106.
- [24] C. G. Gal, *Sup-norm estimates for parabolic systems with dynamic boundary conditions*, submitted.
- [25] C. G. Gal, M. Grasselli, *The non-isothermal Allen-Cahn equation with dynamic boundary conditions*, Discrete and Contin. Dyn. Syst., 12 (2008), 1009–1040.
- [26] C. G. Gal, M. Grasselli, *On the asymptotic behavior for the Caginalp system with dynamic boundary conditions*, Communications in Pure and Applied Analysis, 8 (2009), 689–710.
- [27] C. G. Gal, M. Warma, *Well-posedness and long term behavior of quasilinear parabolic equations with nonlinear dynamic boundary conditions*, Differential and Integral Equations 23 (2010), 327-358.
- [28] C. Gal and M. Warma, *Nonlinear elliptic boundary value problems at resonance with nonlinear Wentzell-Robin type boundary conditions*, preprint, arXiv:1105.3870v1.
- [29] C. G. Gal, M. Warma, *Existence of bounded solutions for a class of quasilinear elliptic systems on manifolds with boundary*, submitted.
- [30] G. R. Goldstein, *Derivation of dynamical boundary conditions*, Adv. Differential Equations, 11 (2006), 457–480.
- [31] H. Ishii, *Asymptotic stability and blowing up of solutions of some nonlinear equations*, J. Differential Equations, 26 (1977), 291–319.
- [32] R. E. Langer, *A problem in diffusion or in the flow of heat for a solid in contact with a fluid*, Tohoku Math. J., 35 (1932), 260-275.
- [33] G. Leoni, *A first course in Sobolev spaces*, Graduate Studies in Mathematics 105, American Mathematical Society, 2009.
- [34] J.L. Lions, E. Magenes, *Problemes aux limites non homogenes at applications*, vol. 1, Paris, Dunod, 1968.
- [35] T.A. Kwembe, Z. Zhang, *A semilinear equation with generalized Wentzell boundary condition*, Nonlinear Anal. 73 (2010), 3162–3170.
- [36] M. Meyries, *Maximal regularity in weighted spaces, nonlinear boundary conditions, and global attractors*, PhD thesis, 2010.
- [37] K. Matsuura, M. Ôtani, *Exponential attractors for a quasi-linear parabolic equation*, Discrete and Continuous Dynamical Systems, supplement, (2007), 713–720.

- [38] A. Miranville, S. Zelik, Attractors for dissipative partial differential equations in bounded and unbounded domains, in: Handbook of Differential Equations: Evolutionary Equations, vol. IV, Elsevier/North-Holland, Amsterdam, 2008.
- [39] H. W. March, W. Weaver, *The diffusion problem for a solid in contact with a stirred fluid*, Physical Review, 31 (1928), 1072-1082.
- [40] M. Morantine, R.G. Watts, *Rapid climate change and the deep ocean*, Clim. Change 16 (1990), 83-97.
- [41] M. Ôtani, *Nonmonotone perturbations for nonlinear parabolic equations associated with subdifferential operators, Cauchy problems*, J. Differential Equations, 46 (1982), 268–299.
- [42] J.C. Robinson, Infinite-dimensional dynamical systems. An introduction to dissipative parabolic PDEs and the theory of global attractors, Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2001.
- [43] A. R. Bernal, A. Tajdine, *Nonlinear Balance for Reaction-Diffusion Equations under Nonlinear Boundary Conditions: Dissipativity and Blow-up*, Journal of Differential Equations 169, (2001), 332-372.
- [44] A. Robertson, W. Robertson, Topological Vector Spaces, Reprint of the second edition, Cambridge Tracts in Mathematics, vol. 53, Cambridge University Press, Cambridge, New York, 1980.
- [45] R. Temam, “Infinite-Dimensional Dynamical Systems in Mechanics and Physics,” Springer-Verlag, New York, 1997.
- [46] S. Takeuchi, T. Yokota, *Global attractors for a class of degenerate diffusion equations*, Electron. J. Differential Equations, 76 (2003), 13 pp. (electronic).
- [47] G. Schimperna, *Weak solution to a phase-field transmission problem in a concentrated capacity*, Math. Methods Appl. Sci., 22 (1999), 1235–1254.
- [48] L.G. Vulkov, *Blow up for some quasilinear equations with dynamical boundary conditions of parabolic type*, Appl. Math. Comput. 191 (2007), no. 1, 89–99.
- [49] M. Warma, *Quasilinear parabolic equations with nonlinear Wentzell-Robin type boundary conditions*, J. Math. Anal. Appl. 336 (2007), no. 2, 1132—1148.
- [50] F. B. Weissler, *Existence and non-existence of global solutions for semilinear heat equation*, Israel J. Math. 6 (1981), 29-40.
- [51] H. Wu, *Convergence to equilibrium for the semilinear parabolic equation with dynamic boundary condition*, Adv. Math. Sci. Appl., 17 (2007), 67-88.

- [52] M. Yang, C. Sun, C. Zhong, *Global attractors for p -Laplacian equation*, J. Math. Anal. Appl., 327 (2007), 1130–1142.
- [53] L. Yang, *Uniform attractors for the closed process and applications to the reaction-diffusion equation with dynamical boundary condition*, Nonlinear Anal. 71 (2009), no. 9, 4012–4025.
- [54] L. Yang, M. Yang, Meihua, *Long-time behavior of reaction-diffusion equations with dynamical boundary condition*, Nonlinear Anal. 74 (2011), no. 12, 3876–3883.
- [55] L. Yang, M. Yang, *Attractors of the non-autonomous reaction-diffusion equation with nonlinear boundary condition*, Nonlinear Anal. Real World Appl. 11 (2010), no. 5, 3946–3954.
- [56] S. Zelik, *Asymptotic regularity of solutions of singularly perturbed damped wave equations with supercritical nonlinearities*, Discr. Cont. Dyn. Syst. 11 (2004) 351–392.
- [57] E. Zeidler, *Nonlinear functional analysis and its applications. II/A*. Springer-Verlag, New York, 1990. Linear monotone operators, Translated from the German by the author and Leo F. Boron.