Competing superconducting, magnetic and charge orderings in the AF Heisenberg-Kondo lattice with Dirac electrons

E. C. MARINO¹ and LIZARDO H. C. M. NUNES²

¹ Instituto de Física, Universidade Federal do Rio de Janeiro, Caixa Postal 68528, Rio de Janeiro, RJ, 21941-972, Brazil

² Departamento de Ciências Naturais, Universidade Federal de São João del Rei, 36301-000 São João del Rei, MG, Brazil

PACS 71.10.Fd – Lattice fermion models

PACS 71.10.Li - Excited states and pairing interactions in model systems

PACS 75.10.-b – General theory and models of magnetic ordering

PACS 71.27.+a - Strongly correlated electron systems

Abstract. - Many recently discovered advanced materials, such as high-Tc cuprates, iron pnictides and several heavy-fermions, exhibit a rich phase diagram suggesting the presence of different competing interactions that would lead to various types of ordering. Nevertheless, there is not yet a clear unifying picture allowing the understanding of the detailed mechanisms that generate such competing interactions. Having such a picture, however, could quite well be at the very roots of the requirements for understanding high-Tc superconductivity in cuprates and pnictides, for instance. In this work we consider the antiferromagnetic (AF) Heisenberg-Kondo lattice, consisting of localized spins with AF exchange interactions between nearest neighbors on a square lattice and itinerant electrons, which undergo a magnetic Kondo interaction with the localized spins, but are otherwise non-interacting. Using the Schwinger-boson (CP¹) formalism and assuming the electrons are Dirac-like, we integrate on the localized degrees of freedom thereby obtaining the effective interaction among the itinerant electrons. This contains a BCS-like superconducting term, a Nambu-Jona-Lasinio-like, charge gap term and a Ising and Heisenberg-like magnetic terms. All these four competing interactions, therefore are generated by the original Kondo magnetic interaction.

Introduction. – Many recently discovered systems in condensed matter physics present a deep interplay among different types of orderings such as superconducting, magnetic or charge ordering. Also in some of them the electronic excitations behave as Dirac fermions These include iron pnictides, superconducting cuprates, transition metal dichalcogenides (TMD) and possibly graphene, if the material is doped, strained or has atoms adsorbed on its surface.

Iron based pnictide materials, for instance, undergo a transition from a magnetically ordered state to a superconducting one upon doping [1–3]. For the particular case of the 122 materials, magnetic order and superconductivity coexist in a small region of the phase diagram [4] and the new quasiparticles in that region exhibit a Dirac-like linear energy dispersion [5–7] with Cooper pairing possibly promoted by short wavelength antiferromagnetic spin fluctuations.

On the same token, the parent compounds of the cuprate superconductors are insulators presenting AF order. As charge carriers are added to the CuO₂ planes, there is the onset of superconductivity, with the characteristic dome-shaped superconducting phase diagram [16,17]. Dirac points appear in the intersection of the nodes of the *d*-wave superconducting gap and the two-dimensional (2D) Fermi surface [18–20]. Strongly interacting 2D Dirac fermion systems also exhibit a dome structure in their superconducting phase diagram [21, 22], so we may specculate whether Dirac fermions may play any role in the description of the cuprate superconductors.

Moreover, the quasi 2D TMD are layered compounds where *s*-wave superconductivity coexists with a charge density wave (CDW) at low temperatures and applied pressure [8-10]. A theory has been proposed in which Dirac electrons appear close to the nodes of the CDW gap and form Cooper pairs [11, 12]. The theory is consistent with the linear decay of the temperature dependent critical field [14], which is observed experimentally in the copper doped dichalcogenide $Cu_x TiSe_2$ [15].

Graphene, on the other hand, is a semi-metal with gapless electronic quasiparticles, which due to the peculiar lattice structure behave as Dirac fermions. Pure graphene does not exhibit superconductivity or magnetism, however, upon doping and/or straining may display competing orders, such as local magnetic moments, superconductivity or an excitonic gap [23].

Heavy fermions is another vast class of materials presenting a rich competition of different types of order in their phase diagrams [24–26]

The above mentioned systems have been extensively studied both theoretically and experimentally but, nonetheless, we still do not have a clear unified picture affording a detailed understanding of the microscopic mechanisms that lead to each kind of ordering. It would be very instructive and inspiring, therefore, to find fully controllable models for systems displaying phases with the aforementioned different types of ordering, where one could precisely trace back the original interaction and the mechanisms leading to such phases.

In this work, we investigate the AF Heisenberg-Kondo lattice model [27], a system containing both localized spins and itinerant electrons with a mutual Kondo-like magnetic interaction. Our main aim is to determine what is the effective net interaction among the conduction electrons, which results from their magnetic interaction with the AF substrate of localized spins. For this purpose, we use the well-known Nonlinear Sigma Model (NLSM) description of the latter [28]. The itinerant electrons, conversely, are supposed to have a tight-binding band structure showing the presence of Dirac cones whose vertices touch at the interface between the valence and conduction bands. We assume the system to be close to half-filling and therefore describe the kinematics of the itinerant electrons by the Dirac hamiltonian. In order to describe the magnetic Kondo interaction it is convenient to re-phrase the NLSM in the CP^1 language [29], whereby that interaction becomes a quartic term involving two Dirac fermion fields and two bosonic (Schwinger Boson) fields.

Our strategy will be to functionally integrate over the bosonic fields in order to derive the resulting effective interaction existing among the fermion fields. In this process, we show that the Kondo interaction among itinerant and localized electrons can be completely expressed as a gauge coupling between the Dirac fermions and the Schwinger bosons, mediated by the CP¹ vector gauge field. Our final result for the effective electron interaction is then obtained upon integration over this gauge field.

The resulting interaction possesses four pieces: a superconducting BCS-type term, an Ising and a Heisenberg-like magnetic interactions and a Nambu-Jona-Lasinio-type interaction. These terms will favor respectively superconducting, magnetic and insulating charge-gapped ordering. We conclude that the original system of localized and itinerant spins yields ultimately an interacting electronic system with those types of competing orders.

The Model and Its Continuum Limit. – We consider a single layered system containing both localized and itinerant spins in which the former are located at the sites of a square lattice and have an antiferromagnetic exchange integral while the latter are conduction electrons with a tight-binding dispersion relation, which is assumed to be Dirac-like. The localized spins mutual interaction will be described by an AF Heisenberg hamiltonian on a 2D square lattice, whereas their interaction with the itinerant ones, by a Kondo-like term. The complete hamiltonian, therefore, contains three parts, namely

$$H = J \sum_{\langle ij \rangle} \mathbf{S}_{i} \cdot \mathbf{S}_{j} - t \sum_{\langle ij \rangle} \left(c_{i\alpha}^{\dagger} c_{j\alpha} + hc \right) + J_{K} \sum_{i} \mathbf{S}_{i} \cdot \left(c_{i\alpha}^{\dagger} \vec{\sigma}_{\alpha\beta} c_{i\beta} \right), \qquad (1)$$

where \mathbf{S}_i is the localized spin operator and $c_{i\alpha}^{\dagger}$ is the creation operator of an itinerant electron of spin $\alpha = \uparrow, \downarrow$, both at site *i*. Frequently we have materials for which there are electrons coming from different bands or even from inequivalent regions of the Brillouin zone. In these cases we would add an extra label a = 1, ..., N to the electron operators. Here, for the sake of simplicity, we shall omit such a label, this fact having no effect in our conclusions.

In order to obtain the partition function, we employ the continuum path integral approach. By using a basis of spin coherent states we have the localized spins \mathbf{S}_i replaced by their correspondidng eigenvalues: $S\mathbf{N}(\mathbf{x})$, where S is the spin quantum number and $|\mathbf{N}(\mathbf{x})|^2 = 1$. N is then decomposed into two perpendicular components associated respectively with ferromagnetic and antiferromagnetic fluctuations [30],

$$\mathbf{N}(\mathbf{x}) = a\mathbf{L}(\mathbf{x}) + (-1)^{|\mathbf{x}|} \sqrt{1 - a^2 |\mathbf{L}|^2} \,\mathbf{n}(\mathbf{x}) \,, \qquad (2)$$

where a is the lattice parameter. In the continuum limit $(a \rightarrow 0)$, this becomes

$$\mathbf{N}(\mathbf{x}) = a\mathbf{L}(\mathbf{x}) + (-1)^{|\mathbf{x}|} \mathbf{n}(\mathbf{x}) + O(a^2).$$
 (3)

Notice that we always have $|\mathbf{n}(\mathbf{x})|^2 = 1$.

In terms of these and of the continuum fermion field $\psi_{\alpha}(\mathbf{x})$ corresponding to $c_{i\alpha}$ we can express the partition function as the functional integral

$$\mathcal{Z} = \int \mathcal{D}\psi \,\mathcal{D}\psi^{\dagger} \,\mathcal{D}\mathbf{L} \,\mathcal{D}\mathbf{n} \,\,\delta\left[|\mathbf{n}|^{2}-1\right] \\ \times \exp\left[-\int_{0}^{\beta} d\tau \int d^{2}x \,\left(\mathcal{H}-\psi^{\dagger}i\partial_{\tau}\psi\right)\right], \,\,(4)$$

where the continuum hamiltonian density reads

$$\mathcal{H} = \psi^{\dagger} \left(i \vec{\sigma} \cdot \vec{\nabla} - \mu \right) \psi + \rho_s |\nabla \mathbf{n}|^2 + \chi_{\perp} S^2 |\mathbf{L}|^2 + S \mathbf{L} \cdot \left[J_{\mathrm{K}} \mathbf{s} + i \left(\mathbf{n} \times \partial_{\tau} \mathbf{n} \right) \right] + (-1)^{|\mathbf{x}|} S J_{\mathrm{K}} \mathbf{n} \cdot \mathbf{s}, \qquad (5)$$

with $\rho_s = JS^2$ as the spin stiffness, $\chi_{\perp} = 4J$ the transverse susceptibility and **s** the itinerant spin operator, given by

$$\mathbf{s} = \psi_{\delta}^{\dagger} \left(\vec{\sigma} \right)_{\delta \gamma} \psi_{\gamma} \,. \tag{6}$$

In expression (5), the first term is the continuum electron kinetic hamiltonian density, derived from the tightbinding energy assuming the system has a Dirac-like dispersion relation near the Fermi points. As the system is doped, charge carriers are added or removed from the conduction band. Their total number is controlled by a chemical potential μ , which has, therefore, been included in the previous equation.

Integrating over \mathbf{L} in (4) we obtain the resulting effective lagrangian density

$$\mathcal{L}_{\text{eff}} = \psi^{\dagger} \left[i \gamma^{0} \gamma^{\mu} \partial_{\mu} - \mu \right] \psi + \frac{\rho_{s}}{2} \left(|\nabla \mathbf{n}|^{2} - \frac{1}{c^{2}} |\partial_{t} \mathbf{n}|^{2} \right) + J_{\text{K}} \left\{ (-1)^{|\mathbf{x}|} S \mathbf{n} + \frac{i}{\chi_{\perp}} \left(\mathbf{n} \times \partial_{\tau} \mathbf{n} \right) + \frac{J_{\text{K}}}{2\chi_{\perp}} \mathbf{s} \right\} \cdot \mathbf{s} , (7)$$

where $(\gamma^0)^2 = 1$, $\gamma^0 \gamma^i = \sigma^i$ and $c = \sqrt{\rho_s \chi_{\perp}}$ is the spinwave velocity.

It will be convenient to use the CP^1 (Schwinger Boson) formulation of the O(3) NLSM, in which the AF fluctuation field is written as

$$n_i = z_{\alpha}^* \left(\sigma_i \right)_{\alpha\beta} z_{\beta} \,, \quad i = x, y, z, \tag{8}$$

in terms of the two complex fields z_{α} , $\alpha = 1, 2$, satisfying the constraint $|z_1|^2 + |z_2|^2 = 1$. In the CP¹ language the effective lagrangian density (7) is rewritten as

$$\mathcal{L}_{\text{eff}} = \psi^{\dagger} \left[i \gamma^{0} \gamma^{\mu} \partial_{\mu} - \mu \right] \psi + 2\rho_{s} |D_{\mu} z_{i}|^{2} + J_{\text{K}} \psi^{\dagger} \left[(-1)^{|\mathbf{x}|} S \ \vec{\sigma} \cdot \mathbf{n} + \frac{i}{\chi_{\perp}} \vec{\sigma} \cdot (\mathbf{n} \times \partial_{\tau} \mathbf{n}) \right] \psi + \frac{J_{\text{K}}^{2}}{2\chi_{\perp}} \mathbf{s} \cdot \mathbf{s} , \qquad (9)$$

where the components of **n** are given by (8) and $D_{\mu} = \partial_{\mu} - iA_{\mu}$.

A Gauge Coupling Replaces the Magnetic Interaction . – We now perform a canonical transformation [31] on the electron field, namely

$$\psi_{\alpha} \to U_{\alpha\beta} \,\psi_{\beta} \,, \tag{10}$$

where the unitary matrix U is written in terms of the z_{α} -fields as

$$U = \begin{pmatrix} z_1 & -z_2^* \\ z_2 & z_1^* \end{pmatrix} .$$
 (11)

This matrix has the following property

$$U^{\dagger}\vec{\sigma}\cdot\mathbf{n}U=\sigma^{z}\tag{12}$$

and therefore the first term in the second line in (9) can be expressed, up to a sign, as the density difference of electrons with opposite spins:

$$(-1)^{|\mathbf{x}|} J_{\mathrm{K}} S \left(\psi_{\uparrow}^{\dagger} \psi_{\uparrow} - \psi_{\downarrow}^{\dagger} \psi_{\downarrow} \right).$$
 (13)

Assuming a uniform density of electrons, we conclude that this term will vanish upon integration in \mathbf{x} because of the rapidly oscillating pre-factor.

Now, since U represents a local operation, it follows that, under the transformation (10), the electron kinetic term generates the additional interaction

$$i \psi^{\dagger} \gamma^{0} \gamma^{\mu} \left(U^{\dagger} \partial_{\mu} U \right) \psi \,. \tag{14}$$

From (11), we obtain

$$U^{\dagger} \partial_{\mu} U = i \sigma^{z} A_{\mu} + \begin{pmatrix} 0 & z_{2}^{*} \partial_{\mu} z_{1}^{*} - z_{1}^{*} \partial_{\mu} z_{2}^{*} \\ -z_{2} \partial_{\mu} z_{1} + z_{1} \partial_{\mu} z_{2} & 0 \end{pmatrix}, (15)$$

where we used the fact that $A_{\mu} = -iz_i^* \partial_{\mu} z_i$, which follows from (9).

Now, consider the polar representations of the fields z_{α} ,

$$z_{\alpha} = \frac{\rho_{\alpha}}{\sqrt{2}} e^{i \theta_{\alpha}}, \quad \alpha = 1, 2.$$
 (16)

Integration over the θ_i fields eliminates rapidly oscillating phase dependent terms, such as the second term in (15). We show in the Appendix that the same happens to the second term in the second line in (9). The effective lagrangian density, therefore, becomes

$$\mathcal{L}_{\text{eff}} = \psi^{\dagger} \left[i \gamma^{0} \gamma^{\mu} \partial_{\mu} - \mu \right] \psi + 2\rho_{s} |D_{\mu} z_{i}|^{2} + \psi^{\dagger}_{\alpha} \gamma^{0} \gamma^{\mu} \sigma^{z}_{\alpha\beta} \psi_{\beta} A_{\mu} + \frac{J_{\text{K}}^{2}}{2\chi_{\perp}} \mathbf{s} \cdot \mathbf{s} \,.$$
(17)

Observe that the magnetic interaction between the itinerant electrons and the localized spins manifests ultimately as a gauge coupling between the electrons and the Schwinger boson fields, mediated by the CP¹ vector field A_{μ} , which becomes a gauge field. Indeed, (17) is invariant under the gauge transformation

$$\begin{aligned}
\psi &\to e^{i\Lambda}\psi, \\
\theta_i &\to \theta_i + \Lambda, \\
A_\mu &\to A_\mu - \partial_\mu\Lambda.
\end{aligned}$$
(18)

Our aim is to obtain the net effective interaction among the conduction electrons, associated to the fermion fields ψ . For this purpose we are going to functionally integrate over the CP¹ fields, in order to derive the resulting effective interaction. Before doing that, however we shall express the effective lagrangian in an explicitly gauge invariant way.

We first introduce the gauge invariant phase-fields [33]

$$\chi_i = \theta_i + \frac{\partial_\mu A^\mu}{\Box},\tag{19}$$

which are clearly invariant under (18). In (19), $1/\Box$ is the Green function of the $\Box = \partial_{\mu}\partial^{\mu}$ operator.

We can now re-write the effective lagrangian density (17) in an explicitly gauge invariant form given by [33]

$$\mathcal{L}_{\text{eff}} = \psi^{\dagger} \left[i \gamma^{0} \gamma^{\mu} \partial_{\mu} - \mu \right] \psi + \frac{1}{2} \sum_{i=1}^{2} \rho_{i}^{2} \partial_{\mu} \chi_{i} \partial^{\mu} \chi_{i} + \frac{1}{4} F_{\mu\nu} \left[\frac{4\rho_{s}}{-\Box} \right] F^{\mu\nu} + \psi^{\dagger}_{\alpha} \gamma^{0} \gamma^{\mu} \sigma^{z}_{\alpha\beta} \psi_{\beta} A_{\mu} + \frac{J_{\text{K}}^{2}}{2\chi_{\perp}} \mathbf{s} \cdot \mathbf{s} \,.$$
(20)

where we have used the constant ρ_i approximation for i = 1, 2 and $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$.

The Effective Electron Interaction. – Let us now perform the functional integration over the bosonic fields ρ , χ and A_{μ} . This will ultimately generate the final effective interaction among the conduction electrons.

We shall adopt the constant ρ_i (i = 1, 2) approximation for performing the functional integration over the Schwinger boson fields, namely, ρ_i 's and χ_i 's. This approximation usually reproduces the physical situation found in many materials. It implies that integration over the z_i fields would just yield a trivial multiplicative constant in the partition function. The nontrivial interaction effect comes from integration over the gauge field A_{μ} . This can be easily performed given the quadratic dependence of (20) in this field.

The resulting effective lagrangian density for the conduction electrons is

$$\mathcal{L}_{\text{eff},\psi} = \psi^{\dagger} \left[i\gamma^{0}\gamma^{\mu}\partial_{\mu} - \mu \right] \psi + \frac{1}{8\rho_{s}} \left(\psi^{\dagger}_{\alpha}\gamma^{0}\gamma^{\mu}\sigma^{z}_{\alpha\beta}\psi_{\beta} \right) \left(\psi^{\dagger}_{\alpha}\gamma^{0}\gamma_{\mu}\sigma^{z}_{\alpha\beta}\psi_{\beta} \right) + \frac{J_{K}^{2}}{2\chi_{\perp}} \mathbf{s} \cdot \mathbf{s} .$$
(21)

Explicitly writing the components of the Dirac field we may, after some algebra, express the effective interaction term above as

$$\mathcal{L}_{\mathrm{I},\psi} = \frac{1}{4\rho_s} \left(\psi_{1\uparrow}^{\dagger} \ \psi_{2\downarrow}^{\dagger} + \psi_{2\uparrow}^{\dagger} \ \psi_{1\downarrow}^{\dagger} \right) \left(\psi_{2\downarrow} \ \psi_{1\uparrow} + \psi_{1\downarrow} \ \psi_{2\uparrow} \right) + \frac{1}{8\rho_s} s_z^2 + \frac{1}{8\rho_s} \left[\left(\bar{\psi}\psi \right)^2 - \left(\bar{\psi}\gamma^0\psi \right)^2 \right] + \frac{J_{\mathrm{K}}^2}{2\chi_{\perp}} \mathbf{s} \cdot \mathbf{s} \,.$$
(22)

The first term above is a superconducting, *s*-wave BCStype interaction. It would lead to a superconducting phase with an order parameter $\Delta = \langle \psi_{1\uparrow}^{\dagger} \psi_{2\downarrow}^{\dagger} + \psi_{2\uparrow}^{\dagger} \psi_{1\downarrow}^{\dagger} \rangle$. The second term is a Nambu-Jona-Lasinio-type interaction, that may produce an insulating charge-gapped phase showing an excitonic condensate. The gap parameter would be $M = \langle \bar{\psi} \psi \rangle$. Finally, the second and fourth terms in (22) are, respectively, Ising and Heisenberg-like, magnetic interactions. These may potentially lead to a magnetically ordered phase with the nonzero order parameter being the average magnetization $\vec{m} = \langle \vec{s} \rangle$.

A model containing the interactions of the first and third terms in (22) has been investigated in a mean-field approximation [22]. We have shown that, even if the excitonic interaction strength is larger than the superconducting interaction, as the chemical potential increases, superconductivity eventually suppresses the excitonic order parameter, which means that the system goes from an insulating state to a superconducting one as charge carrriers are added to the system.

A model with Dirac fermions subject to the interaction described by the first term in (22) was studied in [13,14].

Conclusions. – We have provided a concrete example of a model containing both localized spins and itinerant electrons, where different competing interactions are generated out of the original purely magnetic interactions. The localized spins present an AF Heisenberg interaction on a square lattice, the itinerant electrons are Dirac-like and the mutual localized-itinerant interaction is magnetic, Kondo-like.

Our results yield an unified, controllable picture for the the common magnetic origin of three competing types of order in a strongly correlated system: superconducting, magnetic and charge ordering.

Appendix. – Let us show here that the second term in the second line in (9) only contains rapidly oscillating phase dependent terms, which are, therefore, eliminated through functional integration over the phase fields.

We may write this term as

$$\begin{pmatrix} \psi^{\dagger} \vec{\sigma} \psi \end{pmatrix} \cdot (\mathbf{n} \times \partial_{\tau} \mathbf{n}) = \psi^{\dagger}_{\alpha} \psi_{\beta} z_{\mu}^{*} z_{\nu} \partial_{\tau} (z_{\lambda}^{*} z_{\rho}) \\ \times \epsilon^{ijk} \sigma^{i}_{\alpha\beta} \sigma^{j}_{\mu\nu} \sigma^{k}_{\lambda\rho},$$
(23)

where we have used (8).

Now, we have the following identity for Pauli matrices

$$\sigma^{i}_{\alpha\beta}\sigma^{j}_{\mu\nu} = \frac{\delta^{ij}}{3} \left[2\delta_{\alpha\nu}\delta_{\beta\mu} - \delta_{\alpha\beta}\delta_{\mu\nu} \right] + i\epsilon^{ijk} \left[\delta_{\beta\mu}\sigma^{k}_{\alpha\nu} - \delta_{\alpha\mu}\sigma^{k}_{\beta\nu} \right].$$
(24)

Inserting (24) in (23), we immediately obtain

$$\mathbf{s} \cdot (\mathbf{n} \times \partial_{\tau} \mathbf{n}) = \psi_{\alpha}^{\dagger} \psi_{\alpha} z_{\mu}^{*} z_{\mu} \partial_{\tau} \left(z_{\lambda}^{*} z_{\lambda} \right) + p dt, \qquad (25)$$

where pdt stands for "phase dependent terms".

The first term on the r.h.s. above vanishes because $z_{\lambda}^* z_{\lambda} = 1$ and, therefore, only phase dependent terms are left, as we have asserted.

* * *

[33] MARINO E. C., Int. J. Mod. Phys., 1019954311.

This work has been supported by CNPq, FAPERJ and FAPEMIG. We would like to thank H. Caldas, A. L. Mota, R. L. S. Farias and M. B. Silva Neto for discussions.

References

- [1] DE LA CRUZ C. et al., Nature, 453 (2008) 899.
- [2] KAMIHARA Y. et al., J. Am. Chem. Soc., **130** (2008) 3296.
- [3] ROTTER M., TEGEL M. and JOHRENDT D., Phys. Rev. Lett., 101 (2008) 107006.
- [4] GOKO T. et al., Phys. Rev. B, 80 (2009) 024508.
- [5] RICHARD P. et al., Phys. Rev. Lett., **104** (2010) 137001.
- [6] HUYNH K. K., TANABE Y. and TANIGAKI K., Phys. Rev. Lett., 106 (2011) 217004.
- [7] DA CONCEIÇÃO C. M. S., SILVA NETO M. B. and MARINO E. C., *Phys. Rev. Lett.*, **106** (2011) 117002.
- [8] WITHERS R. L. and WILSON J. A., J. Phys. C: Solid State Phys., 19 (1986) 4809.
- [9] WILSON J. A. and YOFFE A. D., Adv. Phys., 18 (1969) 193.
- [10] WILSON J. A., DISALVO F. J. and MAHAJAN S., Adv. Phys., 24 (1975) 117.
- [11] CASTRO NETO A. H., Phys. Rev. Lett., 86 (2001) 4382.
- [12] UCHOA B., CABRERA G. G. and CASTRO NETO A. H., *Phys. Rev. B*, **71** (2005) 184509.
- [13] MARINO E. C. and NUNES L. H. C. M., Nucl. Phys. B, 741 (2006) 404.
- [14] MARINO E. C. and NUNES L. H. C. M., Nucl. Phys. B, 769 (2007) 275.
- [15] MOROSAN E. et al., Nature Physics, 2 (2006) 544.
- [16] DAGOTTO E., Rev. Mod. Phys., 66 (1994) 763.
- [17] LEE P. A., NAGAOSA N. and WEN X-G., Rev. Mod. Phys., 78 (2006) 17.
- [18] AFFLECK I. and MARSTON J. B., Phys. Rev. B, 37 (1988) 3774.
- [19] AFFLECK I. and MARSTON J. B., Phys. Rev. B, **39** (1989) 11538
- [20] WEN X-G. and LEE P. A., Phys. Rev. Lett., 76 (1996) 503.
- [21] LIM L.-K. et al., Eur. Phys. Lett., 88 (2009) 36001.
- [22] NUNES L. H. C. M., FARIAS R. L. S. and MARINO E. C., cond-mat/1010.4279 preprint, 2010.
- [23] KOTOV V. N. et al., cond-mat/1012.3484.preprint, 2010.
- [24] STEGLICH F. et al., J. Phys.: Condens. Matter, 22 (2010) 164202.
- [25] STEWART G. R., Rev. Mod. Phys., 73 (2001) 797.
- [26] STEWART G. R., Rev. Mod. Phys., 78 (2006) 743.
- [27] KAMPF A.P., Phys. Rep., 249 (1994) 219.
- [28] CHAKRAVARTY S., HALPERIN B. and NELSON D., Phys. Rev. B, 39 (1989) 2344.
- [29] AUERBACH A., Interacting Electrons and Quantum Magnetism, edited by SPRINGER-VERLAG, BERLIN 1994.
- [30] TSVELIK A., Quantum Field Theory in Condensed Matter Physics, edited by CAMBRIDGE UNIVERSITY PRESS, CAM-BRIDGE, UK 1995.
- [31] MARINO E. C. and SILVA NETO M. B., Phys. Rev. B, 66 (2002) 224512.
- [32] KÜBERT C. and MURAMATSU A., Phys. Rev. B, 47 (1993) 787.