

Eco-Evolutionary Feedback in Host–Pathogen Spatial Dynamics

Blake C. Stacey,^{1,2} Andreas Gros,² and Yaneer Bar-Yam²

¹*Martin A. Fisher School of Physics, Brandeis University*

²*New England Complex Systems Institute*

(Dated: October 21, 2018)

Abstract

Spatial extent is a complicating factor in mathematical biology. The possibility that an action at point A cannot immediately affect what happens at point B creates the opportunity for spatial nonuniformity. This nonuniformity must change our understanding of evolutionary dynamics, as the same organism in different places can have different expected evolutionary outcomes. Since organism origins and fates are both determined locally, we must consider heterogeneity explicitly to determine its effects. We use simulations of spatially extended host–pathogen and predator–prey ecosystems to reveal the limitations of standard mathematical treatments of spatial heterogeneity. Our model ecosystem generates heterogeneity dynamically; an adaptive network of hosts on which pathogens are transmitted arises as an emergent phenomenon. The structure and dynamics of this network differ in significant ways from those of related models studied in the adaptive-network field. We use a new technique, organism swapping, to test the efficacy of both simple approximations and more elaborate moment-closure methods, and a new measure to reveal the timescale dependence of invasive-strain behavior. Our results demonstrate the failure not only of the most straightforward (“mean field”) approximation, which smooths over heterogeneity entirely, but also of the standard correction (“pair approximation”) to the mean field treatment. In spatial contexts, invasive pathogen varieties can prosper initially but perish in the medium term, implying that the concepts of reproductive fitness and the Evolutionary Stable Strategy have to be modified for such systems.

I. INTRODUCTION

Mathematical modeling of biological systems involves a tradeoff between detail and tractability. Here, we consider evolutionary ecological systems with spatial extent—a complicating factor. Analytical treatments of spatial systems typically treat as equivalent all configurations with the same overall population density, the same allele frequencies, the same pairwise contact probabilities or the like. For ease of analysis, one seeks a simplified analytical model, which *coarse-grains* “microstates” (the complete specification of each organism) to “macrostates” (characterized by quantities like average densities), allowing one to make useful predictions about the model’s behavior [1, 2]. Corrections to simple coarse-grainings can quickly generate an overbearing quantity of algebra. It is fairly well appreciated that the simplest approximations break down in the spatial context. What is less acknowledged and not yet systematically understood is that the extensions of the simpler approximations also fail. Before exhausting ourselves with ever-more-elaborate refinements, it would be useful to have some understanding of when a particular series of approximations is doomed to inadequacy.

In this article, we study the context in which commonly-used coarse-grainings can be expected to fail at capturing the evolutionary dynamics of an ecosystem, and in addition we provide a novel, direct demonstration of that failure. The fundamental issue is *spatial heterogeneity*, a long-recognized concern for mathematical biology [3, 4]. When does spatial heterogeneity significantly impact the choice of appropriate mathematical treatment, and when does a chosen mathematical formalism not capture the full implications of spatial variability? We show that one can test a treatment of heterogeneity by transplanting organisms within a simulated ecosystem in such a way that, were the treatment valid, the modeled behavior of the ecosystem over time would remain essentially unchanged. We demonstrate situations where the system’s behavior changes dramatically and cannot be captured by a conventional treatment. The complications we explore imply that *short-term descriptions* of what is happening in an evolutionary ecological model can be insufficient and, in fact, misleading, with regard not just to quantitative details but also to qualitative characteristics of ecological dynamics.

Many modeling approaches in mathematical biology which appear distinct at first glance turn out to be describing the same phenomenon with different equations [5–7]. What matters

for our purposes is not so much which technique is chosen, but whether the underlying assumptions do, in fact, apply.

“Mean-field theory” is a term from statistical physics [8, 9] which has been adopted in ecology [10–12], referring to an approximation in which each component of a system is modeled as experiencing the same environment as any other. This implies that the probability distribution over all possible states of the system factors into a product of probability distributions for individual components. An example in population genetics is the assumption that a population is panmictic. That is, if a new individual in one generation has an equal chance of receiving an allele from any individual in the previous generation, then we can approximate the ecosystem dynamics using only the proportion of that allele, rather than some more complicated representation of the population’s genetic makeup. Modeling evolution of that population as “change in allele frequencies over time” (per, *e.g.*, [5, 13]) is, implicitly, a mean-field approximation [14]. The mean-field approximation is also in force if one postulates that an individual organism interacts with some subset, chosen at random, of the total population, even if the form and effect of interactions within that subset are complicated (as in, *e.g.*, [15, 16]).

It is well known that real species are not necessarily panmictic. However, many treatments which acknowledge this are still mean-field models. The textbook way of incorporating geographical distance into a population-genetic model is to divide the system into N local subpopulations, “islands,” connected via migration [17–19]. Within each subpopulation, distance is treated as negligible, and organisms are well mixed [4, 20]. This approach makes a simplifying assumption that there is a single distance scale below which panmixia prevails [21], and it relies on well-defined boundaries between panmictic subpopulations which persist over time [20]. Furthermore, the connections among subpopulations are frequently taken to have the topology of a complete graph, *i.e.*, an organism in one subpopulation can migrate to any other with equal ease [4, 18–20]. In this case, each of the N subpopulations do experience the same environment, to within one part in N . Thus, the mean-field approximation is in force at the island level, and the island model incorporates spatial extent without incorporating a full treatment of spatial heterogeneity. For real ecosystems [21–24], one or more of these simplifying assumptions can fail. Long-distance migration is often thought to return a spatial ecosystem to a well-mixed form, but if organisms’ migration habits are themselves adaptive, this is not necessarily so [25]. More complicated population

structures require more sophisticated mathematical treatments of evolution, a fact which has mathematical consequences, but more importantly has real-world implications for practical issues like the evolution of drug-resistant diseases [26].

Where mean field approximations fail, “higher order” approximations may be employed. Rather than individual organisms or islands, a pair approximation considers pairs of organisms or pairs of spatial regions in average contexts. However, this approximation can also fail when local contexts of groups do not reflect the overall system behavior due to heterogeneity across larger domains. Patches of distinct genetic composition in different parts of a spatial system that are well separated cannot be treated correctly by such approximations. Quantitative analyses confirm this inadequacy. We introduce a new approach to analyzing such approximations by swapping pairs of organisms in a way that preserves the pair description. For spatial systems, such swapping events violate the spatial separation between patches and changes the evolutionary behavior of the system. The swapping method therefore serves as a direct test of the (in)adequacy of the pair approximation. For evolution on random networks of sites that do not embody large spatial distances, the pair approximation can work and the swapping test does not change measures of evolutionary dynamics. However, such networks do not capture important properties of spatial heterogeneity.

As one of the key properties of spatial extent is the propagation of organisms from one part of the space to the other over long distances, we show that important insights can be gained by considering models of percolation. Percolation describes the physical propagation of, *e.g.*, fluids through a random medium. In certain limits the evolutionary behavior of spatial systems can be mapped onto percolation behavior, demonstrating that investigations of such systems which go beyond mean-field or scaling studies are relevant to evolutionary dynamics. This and other advances that go beyond the mean field are necessary to fully describe spatial evolutionary dynamics as they are necessary for the description of many physical systems of spatial extent. The complexities of spatially extended evolutionary dynamical systems beyond the prototypical problem of percolation create new demands and opportunities for advancing our insight into the dynamics of heterogenous systems and their implications for evolution.

II. MODEL AND METHODS

We make the issue of spatial heterogeneity concrete by focusing on a specific model of ecological and evolutionary interest. We take a model of hosts and consumers interacting on a 2D spatial lattice. Each lattice site can be empty (0), occupied by a host (H) or occupied by a consumer (C). We use the term *consumer* as a general label to encompass parasites, pathogens and predators. Where convenient for examples, we will specialize to one or another of these terminologies. Hosts reproduce into adjacent empty sites with some probability g per site, taken as a constant for all hosts. Consumers reproduce into adjacent sites occupied by hosts, with probability τ per host; sometimes τ is fixed for all consumers, but we also consider cases in which it is a mutable parameter passed from parent to offspring. We will refer to τ as the *transmissibility*. Hosts do not die of natural causes, while consumers perish with probability v per unit time (leaving empty sites behind). Because consumers can only reproduce into sites where hosts live, the effective graph topology of reproductively available sites experienced by the consumers is constantly changing due to their very presence. This makes the ecosystem an *adaptive network*, a system in which the dynamics *of* a network and the dynamics *on* that network can occur at comparable timescales and reciprocally affect one another [27–30]. In this model, dynamics can be highly complex, including spatial cascades of host and consumer reproduction. Even when a quasi-steady-state behavior emerges, as we shall see, it is a consequence of fluctuations over extended space and time intervals.

Several different types of biological interactions can be treated by this modeling framework. Hosts could represent regions inhabited by autotrophs alone, while consumers represent regions containing a mixture of autotrophs and the heterotrophs which predate upon them [31]. Alternatively, host agents could represent healthy organisms, while consumers represent organisms infected with a parasite or pathogen. Thus, host–consumer models are closely related to Susceptible–Infected–Recovered (SIR) models, which are epidemiological models used to understand the spread of a disease through a population. SIR models describe scenarios in which each individual in a network is either susceptible (S) to a pathogen, infected (I) with it, or recovered (R) from it; susceptible nodes can catch the disease from infected neighbors, becoming infected themselves, while nodes which have become infected can recover from the disease and are then resistant against further infection. Susceptible, infected and recovered individuals roughly correspond to hosts, consumers, and empty cells,

respectively. An important difference between host–consumer models and epidemiological models concerns the issue of *reinfection*. In the host–consumer model, an empty site left behind by a dead consumer can be reoccupied by another consumer, but only if a host reproduces into it first. Other research has considered models where R[ecovered] individuals can also become I[nfected], with a different (typically lower) probability than S[usceptible] ones, thereby incorporating imperfect immunity into the model [32, 33]. The degree of immunity is independent of geography and the environment of the R[ecovered] individual, unlike reoccupation in the host–consumer model. Another application is illustrated by the Amazon molly, *Poecilia formosa*, which is a parthenogenetic species: *P. formosa*, all of which are female, reproduce asexually but require the presence of sperm to carry out egg development. (This kind of sperm-dependent parthenogenesis is also known as gynogenesis.) *P. formosa* are thus dependent on males of other species in the same genus—usually *P. mexicana* or *P. latipinna*—for reproduction. Because *P. formosa* do not incur the cost of sex, they can outcompete the species on which they rely, thereby possibly depleting the resource they require for survival, *i.e.*, male fish [18, 34]. Thus, hosts could be regions containing sexual organisms, with consumers standing for areas containing both sexual and asexual individuals [18].




FIG. 1: Snapshots of a simulated host–consumer ecosystem on a 250×250 lattice, taken at intervals of 100 generations. Consumers are dark gray (red online), hosts are light gray (green online) and empty space is left white. The simulation began with a single consumer at the center of the lattice, which gave rise to an expanding front of consumers. The first image in this sequence shows the state of the ecosystem 100 generations into the simulation. Hosts which survive the consumer wave recolonize the empty sites, leading to pattern formation. Here, the host growth rate is $g = 0.1$, the consumer death rate is $v = 0.2$ and the consumer transmissibility is fixed at $\tau = 0.33$.

This host–consumer model displays waves of colonization, consumption and repopulation. Hosts reproduce into empty sites, and waves of consumers follow, creating new empty regions

open for host colonization. Therefore, clusters of hosts arise dynamically [35–38], a type of pattern formation which can separate regions of the resources available to pathogens into patches without the need for such separation to be inserted manually. Figure 1 illustrates a typical example of this effect. This is a specific example of the general phenomenon of pattern formation in nonequilibrium systems [14]. Consumers are *ecosystem engineers* [39–43] which shape their local environment: an excessively voracious lineage of consumers can deplete the available resources in its vicinity, causing that lineage to suffer a Malthusian catastrophe [18, 24, 35, 44–48]. Because the ecology is spatially extended, this catastrophe is a local niche annihilation, rather than a global collapse [49]. A mutant strain with a high transmissibility can successfully invade in the short term but suffer resource depletion in the medium term, meaning that in a population where consumer transmissibilities evolve, averages taken over long numbers of generations yield a moderate value [10, 50]. This implies that an empirical payoff matrix or reproduction ratio will exhibit nontrivial timescale dependence [10, 51, 52].

This model is distinct from another approach to studying evolutionary dynamics in spatial contexts, that of evolutionary game theory. Game-theoretic models of spatially structured populations have been explored at great length. These investigations have found that breakdowns of mean-field approximations are commonplace. However, evolutionary game theory has its own simplifying assumptions. The vast majority of studies consider only two-player games. Population size is usually taken to be constant, and population structure is typically fixed in place. In game-theoretic models, the benefits and costs of different organism behavioral traits are parameters whose values are chosen by the modeler. By contrast, “benefits” and “costs” in host–consumer models are emergent properties which depend on interactions over many generations. Population size is not fixed, and population structure is dynamical: the environment in which different consumer varieties compete changes stochastically, in ways affected by their presence.

III. RESULTS

A. Evolution of Transmissibility

We investigate evolution in the spatial hostconsumer ecosystem through simulation and analytic discussion. If the transmissibility τ is made a heritable trait, passed from a consumer to its offspring with some chance of mutation, what effect will natural selection have on the consumer population? Figure 2(A) shows the average, minimum and maximum values of the transmissibility τ observed in a population over time. The average τ tends to a quasi-steady-state value dependent on the host growth rate g and the consumer death rate v ; if the simulation is started with τ set to below this value, the average τ will increase, and likewise, the average τ will decrease if the consumer population is initialized with τ over the quasi-steady-state value. Even when the average τ has achieved its quasi-steady-state value, the population displays a wide spread of transmissibilities whose extremes fluctuate over time [31].




FIG. 2: **(A)** Minimum (blue), average (green) and maximum (red) transmissibility τ for a consumer population over time, with $g = 0.1$ and $v = 0.2$. (The mutation rate is $\mu = 0.255$ and the step size is $\Delta\tau = 0.005$, as was used in reference [31].) The average τ tends to a quasi-steady-state value dependent on g and v ; if the simulation is started with τ set to below this value, the average τ will increase, and likewise, the average τ will decrease if the consumer population is initialized with τ over the quasi-steady-state value [31]. The horizontal dotted line indicates the threshold value of τ which, in a mean-field model, is the smallest value at which a consumer population can sustain its numbers. The dashed line indicates the value to which τ would trend in a well-mixed ecosystem. **(B)** Minimum, average and maximum τ as a function of v , with $g = 0.1$. The dotted line shows the minimum sustainable τ as predicted by mean-field approximation. Each point is found by averaging over 15,000 timesteps. Error bars indicate one standard deviation.

In a well-mixed ecosystem, the average τ of the population will tend to 1, maximizing the reproductive rate of the individual consumer. This occurs because each consumer on average experiences the same environment as any other, and thus has the same number of hosts available to reproduce into. A consumer with a higher τ has a higher reproduction rate and therefore evolutionary dominance up to the highest possible value, 1. The observation of a quasi-steady-state value below 1 is an important result. This is the first breakdown of the mean-field approximation, and it indicates the inapplicability of traditional assumptions about fitness optimization, with implications for the origins of reproductive restraint, communication-based altruism and social behaviors in general [10, 31, 49, 51–53].

One can avoid τ tending to 1 in a panmictic system by imposing some extra constraint, such as a tradeoff between transmissibility and lethality, where higher transmissibility becomes impossible due to lethality that prevents transmission. This tradeoff between infectiousness and lethality can be considered as a within-host version of resource overexploitation that here occurs at the population level. Such within-host tradeoffs are difficult to establish empirically in living populations [54, 55]. Often, one lacks pertinent information, such as the functional relationship between pathogen load and disease transmission probability, or the extent to which empirical proxies for pathogen load predict actual host mortality [56]. An empirical observation of low virulence should not by itself be taken as evidence that a tradeoff exists: it may well be that another condition, such as panmixia, fails to obtain. The behavior of spatial models makes clear that the relevant scale of the limiting factor is not necessarily within the individual host.

Another difference between spatial and nonspatial host–consumer systems is the rate at which consumers must reproduce in order to sustain their population. One can calculate the minimum sustainable value of τ in the mean-field approximation [57] by balancing the birth and death rates. If the host population is small compared to the total ecosystem size, then the minimum sustainable τ is the value which satisfies $k\tau = v$, where k is the number of neighbors adjacent to a site. For the parameters used in Figure 2(A), this value would be 0.05, which is substantially smaller—by a factor of 4—than the lowest τ seen in the evolving spatial population. Consumer populations with τ at the mean-field threshold are not sustainable in the spatial case. This is easily verified by numerical simulations or by using the mean-field equations for the host–consumer dynamics [53, 58, 59]. Stochastic fluctuations suppress the active phase, *i.e.*, the range of parameter values which permit a

living consumer population is reduced [57].

B. Timescale Dependence of Invasion Success




FIG. 3: **(A)** Survival probability as a function of time for five scenarios: injecting mutants with the same transmissibility as the native consumers, three examples of injecting mutants with transmissibility higher than the τ of the native consumers, and an example of injecting mutants with lower τ than the native population. **(B)** Time intervals during which the survival-probability curves for the native and invasive strains overlap. $\Delta\tau$ indicates the difference between the invasive and native transmissibilities. The closer the mutant trait value is to the resident, the greater the duration of time over which the survival-probability curves for the native and mutant strains overlap. Here, overlap is defined by probabilities being coincident at the 95% confidence level; using other overlap criteria gives qualitatively the same results. Inset: magnified view of the $\Delta\tau \geq 0.04$ region.

A key question about an ecological system is whether a new variety of organism, having a different genetic character and phenotypic trait values, can successfully invade a native population. If a mutant consumer strain with fixed transmissibility τ_m can successfully invade a population of transmissibility $\tau_0 < \tau_m$, then we expect the time-averaged value of τ seen in the evolving system to be larger than τ_0 . To investigate this, we simulate scenarios where the native population has τ close to the average value seen in the evolutionary case described above. We then inject a mutant consumer strain with significantly larger τ and study the results. For a typical example, we see from Figure 2(A) that when $g = 0.1$ and $v = 0.2$, the average τ is approximately 0.33. So, we simulate $\tau_m = 0.45$ mutants entering an ecosystem whose native population has $\tau_0 = 0.33$. Initially, the mutants prosper,

but they ultimately fail to invade. As shown in Figure 3, the probability of a $\tau_m = 0.45$ strain surviving for tens of generations after injection is larger than that of a $\tau_0 = 0.33$ strain. That is, mutants with the higher τ can out-compete the neutral case. However, after ≈ 74 generations, the survival-probability curves cross. Observed over longer timescales, the mutant strain is less successful than the native variety. This pattern is consistent for $\tau_m > \tau_0$: the average transmissibility seen in the evolutionary case stands up to invasive varieties. This key result manifests the distinctive properties of the spatial structure of the model. The underlying reason for this result is that the mutants encounter the resource limitations imposed by the patchy native population. Over short timescales, the mutant strain enjoys the resources available within the local patch, consuming those resources more rapidly than can be sustained once it encounters the limitations of the local patch size. In this way, the initial generations of the mutant strain “shade” their descendants. Thanks to descendant-shading, short-term prosperity is not a guarantee of medium- or long-term success.

This is to be contrasted with what happens in a well-mixed ecosystem. In the well-mixed scenario, consumer strains with higher τ successfully invade and displace the native population with a high probability. The invasion success is consistent with the dynamics of a continuously evolving ecosystem. If τ is made an evolvable trait in simulated panmictic systems, the average τ of the population will tend to 1, as predicted by the mean-field analytic proof. There is no difference in a well-mixed scenario between short-term and long-term success. Descendant-shading does not occur in the well-mixed case. This follows from the lack of distinction between local patches and large-scale structure.

One common measure of evolutionary success is the expected relative growth rate of the number of offspring of a mutant individual within a native population, *i.e.*, the relative growth rate of a mutant strain. This rate, known as the *invasion fitness*, is often used to investigate the stability of an evolutionary ecosystem [11, 60, 61]. If the invasion fitness is found to be positive, the native variety is judged to be vulnerable to invasion by the mutant. Conversely, if the invasion fitness is found to be negative, the native variety is deemed to be stable. For the spatial host–consumer ecosystem, this method gives qualitatively incorrect predictions for evolutionary dynamics.

Our investigation builds on earlier work which studied the timescale dependence of fitness indicators in spatial host–consumer ecosystems [51, 52]. In this paper we have augmented

the prior work by considering the survival probability to show the effects of varying τ . We have also more systematically shown the number of generations until dominance of the evolutionary stable strain. In addition, we reported the case of a mutant strain invading a background population, clarifying the conceptual and quantitative results of those earlier works, which considered instead scenarios complicated by multiple ongoing mutations.

C. Pair Approximations

The inadequacy of mean-field treatments of spatial systems motivates the development of more elaborate mathematical methods. In this section, we review one such methodology, based on augmenting mean-field approximations with successively higher-order correlations, and we test its applicability to our host–consumer spatial model. The numerical variables used in this methodology are probabilities which encode the state of the ecosystem and can change over time. One such variable is, for example, the probability p_a that a lattice site chosen at random contains an organism of type a . Another is p_{ab} , the probability that a randomly-chosen *pair* of neighboring sites will have one member of type a and the other of type b . The change of these quantities over time is usually described by differential equations, for which analysis tools from nonlinear dynamics are available [11, 35, 53, 61–64].

The importance of the joint probabilities p_{ab} is that they reflect correlations which mean-field approximations neglect. To understand the relevance of the joint probabilities p_{ab} , consider a scenario where an invasive mutant variety forms a spatial cluster near its point of entry. Let p_M be the probability that a lattice site chosen at random contains a mutant-type organism, and let p_{MM} denote the probability that a pair of neighboring sites chosen at random will both be occupied by mutant-type organisms. Then the average density of invasive mutants in the ecosystem, p_M , will be low, while the conditional probability that a neighbor of an invasive individual will also be of the invasive type, $q_{M|M} = p_{MM}/p_M$, will be significantly higher. (It is typical in theoretical spatial ecology to denote conditional probabilities with q , rather than p [65].) A discrepancy between the conditional probability $q_{a|b}$ and the overall probability p_a can persist when the ecosystem has settled into a quasi-steady-state behavior, and is then an indicator of spatial pattern formation.

Applying this idea to the spatial host–consumer model, let p_C be the probability that a lattice site chosen at random contains a consumer, and let $q_{C|C}$ denote the conditional

probability that lattice site adjacent to a consumer will also be occupied by a consumer. Figure 4(A) shows p_C and $q_{C|C}$ measured during the course of numerical simulations. In a well-mixed scenario (where we expect the mean-field approximation to be applicable), the average consumer density p_C and the consumer–consumer pairwise correlation $q_{C|C}$ are essentially equal over time. In the spatial lattice scenario, p_C and $q_{C|C}$ are noticeably different.




FIG. 4: (A) Pairwise conditional probability $q_{C|C}$ plotted against the average density of consumers, p_C , for three variations on the host–consumer model: a well-mixed case in which mean-field theory is applicable, a random regular graph (in which each site has exactly four neighbors) and a 2D square lattice. The dotted line, $p_C = q_{C|C}$, indicates the mean-field approximation. 10^4 timesteps were computed for each case. The well-mixed case is simulated by dynamically rewiring sites at each time step, precluding the generation of spatial heterogeneity; consequently, the pairwise correlation $q_{C|C}$ is within statistical variation equal to p_C ($R^2 = 0.953$). A random regular graph (RRG) with random but static connections does develop spatial heterogeneity so that $q_{C|C}$ is not the same as p_C ($R^2 = 0.581$). The discrepancy is even stronger in the lattice case ($R^2 = 0.304$). (B) Success rate of invasive mutant strains as a function of swapping probability. Voracious mutant strains with $\tau = 0.45$ are introduced into a lattice ecosystem defined by a host growth rate of $g = 0.1$, a consumer death rate $v = 0.2$ (the same for both consumer varieties), and a native consumer transmissibility of $\tau = 0.33$. Average success rates are found by simulating 2000 invasions per value of the swapping probability parameter; error bars indicate 95%-confidence intervals. Increasing the fraction of possible swaps which are actually performed makes the voracious invasive strain more likely to take over the ecosystem.

Treating the correlations $q_{a|b}$ as not wholly determined by the probabilities p_a is a way of allowing spatial heterogeneity to enter an analytical model. Whether it is a *sufficient* extension in any particular circumstance is not, *a priori*, obvious. Typically, the differential equations for the pair probabilities p_{ab} depend on triplet probabilities p_{abc} , which depend upon quadruplet probabilities and so forth. The standard procedure is to truncate this hierarchy at some level, a technique known as *moment closure* [30, 60, 62, 66, 67]. Moment

closures constitute a series of approximations of increasing intricacy [68, 69]. The simplest moment closure is the mean field approximation; going beyond the mean field to include second-order correlations but neglecting correlations of third and higher order constitutes a *pair approximation*. These approximations do not incorporate all of the information about spatial structure which may be necessary to account for real-world ecological effects [60].

D. Organism Swapping

Several factors have been identified which undermine pair approximations [30, 31, 53, 58–60, 63, 70–72]. In our model, we can directly test the efficacy of pair approximations in a completely general way. The key idea is to transplant individuals in such a way that the variables used in the moment-closure analytical treatment remain unchanged. At each timestep, we look through the ecosystem for isolated consumers, that is, for individual consumers surrounded only by a specified number of hosts and empty sites. We can exchange these individuals without affecting the pairwise correlations. For example, if we find a native-type consumer adjacent to three hosts and one empty site, we can swap it with an invasive-type consumer also adjacent to three hosts and one empty site. We can also exchange isolated pairs of consumers in the same way. The variables used in the moment-closure treatment remain the same. Were the moment-closure treatment valid, we would expect the dynamics to remain unchanged when we perform such exchanges.

When we perform the simulation, however, swapping strongly affects the dynamics. With this type of swapping in effect, mutants with higher τ can invade a native population with lower τ . In one typical simulation run with a native τ of 0.33 and an invasive τ of 0.45, the invasive strain succeeded in 1,425 of 10,000 injections. Without swapping, the number of successful invasions is *zero*.

Swapping can be considered as creating a new ecosystem model with the same moment-closure treatment as that of the original. The behavior of invasive strains is different, because transplanting organisms allows invasive varieties to evade localized Malthusian catastrophes. Swapping opens the ecosystem up to invasive strains, since, in essence, it removes individuals from the “scene of the crimes” committed by their ancestors.

This type of swapping is, to our knowledge, a new test of moment-closure validity. Randomized exchanges have been incorporated into computational ecology simulations for differ-

ent purposes. For example, research on dispersal rates in an island model shuffled individuals in such a way that the population size of each island was held constant [73].

If, instead of performing every permissible swap, we transplant organisms with some probability between 0 and 1, we can interpolate between the limit of no swapping, where invasions always fail, and the case where pair approximation is most applicable and invasions succeed significantly often. The results are shown in Figure 4(B) and indicate that the impact of swapping becomes detectable at a probability of ≈ 0.25 and effectively saturates at a probability of ≈ 0.9 .

Our swapping method allows us to test the significance of complications which can undermine pair approximation techniques or make them impractical to apply, several of which have been identified. First, introducing *mutation* into a game-theoretic dynamical system can make pair approximation treatments of that system give inaccurate predictions [63, 71]. Second, when the evolving population has a network structure, the presence of *short loops* in the network often makes pair approximations fail [70]. For example, in a triangular lattice, one can take a walk of three steps and return to one's starting point, whereas on a hexagonal lattice, the shortest closed circuit is six steps long. A pair approximation can work well for a dynamical system defined on the hexagonal lattice but fail when the same dynamics are played out on a triangular one. This happens because the short loops provide opportunities for contact which the coarse-graining necessary for a pair approximation will miss. This effect is amplified in adaptive network models, where the underlying network changes dynamically in response to the population living upon it. In such cases, even extending the moment closure to the triplet level brings little improvement [30]. Third, *fluctuating population sizes* make pair approximations significantly more cumbersome to construct, leading to systems of differential equations which are too intricate to be significantly illuminating. In a game-theoretic model where a lattice is completely filled at all times with cooperators and defectors, there is one independent population density variable and three types of pairs. By contrast, in an ecological model where two consumer varieties are competing within an adaptive network of hosts, a pair approximation requires nine independent variables [53, 58, 59]. Modeling phenomena of biological interest can easily increase the complexity still more. For example, if organism behavior changes in response to social signals [31], the number of possible states per site, and thus the number of dynamical variables in a pair approximation treatment, increases further. Fourth, the pair-approximation philosophy of averaging over

all pairs in the system impedes the incorporation of *environmental heterogeneities*, including biologically crucial factors like variable organism mobility, background toxicity or other localized “costs of living,” and resource availability [72]. Finally, *dynamical pattern formation* creates spatial arrangements which the pair approximation does not describe [60].

E. Effect of Substrate Topology

It is instructive to compare the spatial lattice ecosystem with the host–consumer model defined on a random regular graph (RRG). In an RRG, each node has the same number of neighbors, as they do in a lattice network, but the connections are otherwise random. RRGs have been used as approximations to incorporate the effects of spatial extent into population models, as they make for more tractable mathematical treatments, although they are typically less realistic than spatial lattices [74]. The network structure is set at the beginning of a simulation and does not change over time. The important aspect of this network as compared to the spatial case is that there exist short paths of links that couple all nodes of the network. This is quite different from the spatial case, where strains in one part of the network cannot reach another in only a few generations due to the need to traverse large numbers of spatially local links.

When we simulate our host–consumer ecosystem on an RRG, we find that an invasive consumer strain with higher transmissibility τ can out-compete and overwhelm a native consumer population with lower τ . In one typical simulation run, using the native and invasive τ values of 0.33 and 0.45 respectively, 2,233 out of 10,000 invasions were successful, whereas on the lattice *no* invasion succeeded using the same parameters. Thus, the RRG does not capture the essential features of the spatial scenario. In particular, our results show that the RRG case is more like the well-mixed case than the spatial lattice, as far as stability against invasion is concerned.

Our swapping test provides insight into the utility of the pair approximation, which can be effective for the RRG even though it is not for the spatial case. Consider the pairwise correlation value $q_{C|C}$, which would be a variable for a pair approximation treatment. On an RRG, the underlying network topology provides enough locality that p_C and $q_{C|C}$ are unequal, distinct from the well mixed case as shown in Figure 4(A). This means that the pair approximation is nontrivial for the RRG as it incorporates the difference between $q_{C|C}$

and p_C , which would not be contained in a mean-field treatment. We can also implement swapping on the RRG, where invasions can succeed without it; as expected, swapping does not affect the success rate on the RRG. With 10,000 simulated invasions for each case, the 95%-confidence interval for the difference in success rates between full swapping and none is 0.004 ± 0.01 . Thus, the pair approximation may be successful in this network topology. However, this does not mean that the RRG or the pair approximation capture the full significance of a spatial system, because the RRG network does not embody essential properties of spatial extent—separation by potentially large distances.

F. Percolation

In order to obtain quantitatively or even qualitatively correct predictions for spatial host–consumer evolutionary dynamics, different approaches are needed. Having encountered the limitations of moment closures, we now demonstrate a change of perspective which yields quantitatively useful results. In certain situations, the process of pathogen propagation through the host population distributed in space can be mapped onto a *percolation* problem. A topic widely investigated in mathematics, percolation theory deals with movement through a matrix of randomly placed obstacles. A prototypical percolation problem is a fluid flowing downhill through a regular lattice of channels, with some of the lattice junction points blocked at random. The key parameter is the fraction of blocked junction points. If this fraction is larger than a certain threshold value, the fluid will be contained in a limited part of the system. However, if the blocking fraction is below the threshold, the fluid can percolate arbitrarily far from its starting point. This is a *phase transition*, a shift from one regime of behavior to another, in this case between a phase in which fluid flow can continue indefinitely and one in which flow always halts. Similar issues arise when a pathogen propagates by cross-infection through a set of spatially arranged hosts. Sufficiently many hosts in mutual contact are required for the pathogen to propagate successfully. Pathogen strains therefore survive or die out over time depending on whether percolation is or is not possible [12, 57, 75–78].

One important goal of studying host–pathogen models is knowing the pathogen properties that enable its survival in a population, or equivalently what prevents it from persisting in a population. The growth rate of a pathogen in a population can be an important public health concern. We therefore focus on analyzing the minimum value of the transmissibility

that enables a pathogen population to persist, and the growth dynamics of population sizes near that transition.

Of essential importance to the quantitative theoretical and empirical analysis is the recognition that infected population growth can be described by power laws $n \sim t^z$, with an exponent that differs from that of the mean field. Identifying the value of the power z is important to practical projections of the number of infected individuals. The initial growth curve of infected populations can be correctly extrapolated if the exponent is known, guiding public health responses. Knowing what impediments are needed to prevent further propagation can even better guide public health intervention strategies.




FIG. 5: **(A)** Population size as a function of time, averaged over 10^3 simulation runs, for τ values near the transition points at $g = 0$ and $g = 0.05$ on the spatial lattice, with $v = 1$. Dashed and solid lines indicate the population growth for systems at dynamic percolation and directed percolation transitions respectively, showing that these transitions have the characteristic properties of those universality classes. **(B)** Critical τ for the host–consumer ecosystem with $v = 1$. The transition line crosses over from the dynamic percolation universality class at $g = 0$ to directed percolation between $g = 0.015$ and $g = 0.02$. Red Xs indicate the transition curve for the host–consumer dynamics on a random regular graph (RRG) of uniform degree 4; the dashed line connecting them is to guide the eye. The RRG transition is neither directed percolation nor dynamic percolation.

We show in Figure 5 the results of numerical simulations which indicates that the consumer extinction transition, when the transmissibility τ becomes just large enough that the consumer population sustains itself, lies in the *directed percolation universality class* [33, 79–82]. A similar result has been reported for related models [57, 75], consistent with those models being in the same universality class. The directed percolation universality class is a large set of models, all of which exhibit a phase transition between two regimes of behavior,

and all of which behave in essentially the same way near their respective transition points. The scenario of fluid flow through a random medium considered above is a classic example of a directed percolation-class model, but many others exist as well [33, 79]. The *critical exponents* describe how properties of the modeled system vary over time or as a function of how far the control parameter is from the critical point. They are the same for all systems in the universality class. Other universality classes exist as well, with different classes having different quantitative values for the critical exponents. Identifying the universality class a system belongs to enables us to study a complicated phenomenon by examining a simpler representative of its class instead. This is convenient, because regions of parameter space near phase transitions are precisely where mean-field and moment-closure approximations are least reliable, even for short-timescale modeling. Near the phase transition, stochastic fluctuations create dynamical patterns with a wide range of sizes. In Figure 5(A), we see that percolation theory gives quantitatively correct predictions for the growth of consumer population sizes in the spatial host–consumer model.

We can understand the $g = 0$ and $g = 1$ extremes by mapping the host–consumer model onto other stochastic models for which exact or approximate results are available. When $g = 0$, the host–consumer model maps onto the SIR epidemic process [33]. In turn the SIR model on the square lattice can be understood in terms of bond percolation on the square lattice [83], for which the transition point is known exactly [84, 85]. We can therefore predict analytically that the critical τ on the square lattice is 0.5. Percolation theory also gives a prediction for the critical τ on an RRG: it should be approximately $1/3$ [86]. These both match the simulation results seen in Figure 5(B).

In contrast, when $g = 1$, empty sites are filled as quickly as possible, so the behavior of the host–consumer model should resemble that of an epidemic model with only Susceptible and Infected sites. In this case, the transition point of the epidemic model on the square lattice is only known numerically [83]. The numerical value, ≈ 0.29 , does agree with the critical τ found by simulating the host–consumer model at $g = 1$.

Thus, in the limiting cases of $g = 0$ and $g = 1$, the host–consumer model is roughly equivalent to the SIR and SIS epidemic models. However, a host–consumer model with $0 < g < 1$ has dynamical behavior distinct from an epidemic model which allows reinfection of Recovered sites. The key difference is that reoccupying an empty site with a consumer requires prior recolonization by a host, whereas the vulnerability of a R[ecovered] individual

to becoming I[nfected] is defined as an intrinsic property of the R[ecovered] type. This changes the role of ecology: both models incorporate space, but the effect of spatial extent is different. This manifests as a change in the shape of the critical-threshold curve, as well as a change in universality class [33, 83].

Furthermore, when we use an RRG topology instead, comparing host–consumer dynamics at $g = 1$ with an SIS epidemic model reveals their transitions to take place at different thresholds. For the host-consumer model, the critical τ on an RRG is approximately 0.2615, while the SIS threshold is approximately 0.34 [87].

The analysis of percolation behavior near the transition point from physics, maps directly onto the critical public health problem of the growth of infected populations, and more generally onto the dynamics of evolutionary systems. For these systems the mean field treatment fails and the standard transmission of infectious diseases in a population need not apply. Applications to real world systems must accommodate the actual network of connectivity. This network can also be modified by intervention strategies.

IV. DISCUSSION

Understanding the effects of spatial extent is a vital part of evolutionary ecology. Spatial extent changes the quantitative and qualitative characteristics of a model’s evolutionary behavior, compared to well-mixed models. The short-term success rate of novel genetic varieties is not indicative of their long-term chance of success relative to the prevalent type. Standard stability criteria fail to reflect the actual stability achieved over time. We must instead consider extended timescales because they are determined by spatial patterns, whose ongoing formation is an intrinsic part of nonequilibrium evolutionary dynamics. Our analysis provides a clear understanding of why there are dramatic differences between spatial models and mean-field models, which simplify away heterogeneity through mixing populations, averaging over variations or mandating a globally connected patch structure. We have further shown that transplanting organisms dramatically changes the dynamics of spatial systems, even when we preserve local correlations as would be considered in a pair approximation treatment. Our results prove that any model striving to capture the effects of heterogeneity that does not change its behavior with organism transplanting cannot fully capture the dynamics of spatial evolution. The following subsections summarize the general conclusions

we draw from these results.

A. Defining Fitness

In our host–consumer models, each individual either survives or it does not, and any individual has a specific number of offspring and survives over a certain amount of time; that is to say, an “individual fitness” (in the terminology of [88]) is a well-defined concept. To find *expected* individual fitness, or *average* individual fitness, we must define a set of individual organisms over which to take an average, which is the very concept we have established to be problematic. Consequently, derived notions of fitness, which depend on comparisons between such averages [88], become elusive, context-dependent quantities. The problem is both temporal and spatial: Average relative fitness in one generation is not necessarily a good measure of the long-term success of a strain in one, or a combination of, the broad variety of dynamically-generated niches. This problem is not, however, the same as the traditional concept of variation of fitness across a static set of niches, because the niche dynamics of our spatially explicit model ensures that evolutionary outcomes are not reflected in any standard definition of the average.

One might be tempted to call the behavior of voracious invasive consumers “frequency-dependent fitness” [4, 5], as the invasive strain is successful initially when rare but fails when it becomes more common. The term “frequency-dependent fitness” is, however, a misnomer in this context, because the organism type is rare and successful when it is newly introduced, but as it declines to extinction it becomes rare and unsuccessful. Nor can we attribute the decline to the frequency of hosts: the average population density of hosts remains essentially unchanged, because the boom and the following bust are localized. *Frequency*, being defined by an average over the whole ecosystem, is only a proper variable to use for describing the ecosystem in the panmictic case. One might attempt to refine the concept of global frequency by including local frequencies. However, the breakdown of moment-closure techniques implies that defining fitness as a function of organism type together with average local environment [89] will, in many circumstances, not be an adequate solution.

Consequently, we find that trying to assign a meaningful invasion fitness value to an invasive variety of organism is too drastic a simplification. In turn, this implies that we cannot assign a fitness value to a phenotypic or genetic characteristic such as infectiousness

or transmissibility. To understand this point, we rephrase the spatial host–consumer model in terms of alleles. In an invasion scenario, an individual consumer can have one of two possible alleles of the “transmissibility gene”, one coding for the native value of τ (*e.g.*, $\tau = 0.33$) and the other for the invasive value (*e.g.*, $\tau = 0.45$). A mean-field treatment would then involve specifying the fraction of the population which carries the native allele versus the fraction which carries the invasive variant. We have seen, however, that the predictions based on such a heavily coarse-grained caricature of the original model deviate from its actual behavior. In short, the evolutionary dynamics cannot be characterized using the allele frequencies at a particular time.

If we can no longer summarize the genetic character of a population by an allele frequency—or a set of allele frequencies for well-defined local subpopulations—then computing the fitness of a genotype from its generation-to-generation change in frequency is a fruitless task. In a world which exhibits nonequilibrium spatial pattern formation, allele frequencies are the wrong attribute for understanding the dynamics of natural selection. Formally, the conventional assumption that the allele frequencies are a sufficient set of variables to describe evolutionary dynamics is incorrect. The spatial structure itself is a necessary part of the system description at a particular time in order to determine the subsequent generation outcomes, even in an average sense.

The timescale-dependence issues which arise in spatial host–consumer ecosystems exist in a wider context. Multiple examples indicate that initial success and eventual fixation are only two extremes of a continuum which must be understood in its entirety to grasp the stability of a system. In the study of genetic drift, it has been found that neutral mutations can fixate and beneficial mutations fail to fixate due to stochasticity [4]. Likewise, in the study of clonal interference [90], one beneficial mutation can out-compete another and prevent its fixation.

Furthermore, classical genetics makes much use of the Price Equation for studying the change in a population’s genetic composition over time [6, 7], and it is well known that analytic models built using the Price Equation lack “dynamic sufficiency”. That is, the equation requires more information about the current generation than it produces about the next [5, 7, 91–93], and so predictions for many-generation phenomena must be made carefully, if they can be made at all. Modeling approaches which are fundamentally grounded in the Price Equation, such as “neighbor-modulated” fitness calculations [6, 19, 47, 94, 95]

and their “multilevel” counterparts [6, 15, 94, 96–98], are not likely to work well here, as the analyses in question draw conclusions only from the short-timescale regime. In addition, those particular analyses which address host–consumer-like dynamics either rely on moment closures [47] or they assume a fixed, complete connection topology of local populations which are internally well-mixed [19, 96]. These simplified population structures are quite unlike the dynamical patch formation seen in the host–consumer lattice model. (Wild and Taylor [99] demonstrate an equivalence between stability criteria defined via immediate gains, or “reproductive fitness”, and criteria defined using fixation probability; however, their proofs are explicitly formulated for the case of a well-mixed population of constant size, neither assumption being applicable here. Whether fixation probability is equivalent to any other criterion of evolutionary success generally depends on mutation rates, even in panmixia [93].)

In the adaptive dynamics literature, models have been studied in which “the resident strikes back” [100–102]. That is, an initially rare mutant variety M can invade a resident population of type R , but M does not supplant R and become the new resident variety, even though a population full of type M is robust against incursions by type R . This is often considered a rare occurrence, requiring special conditions to obtain [101, 102], though the theorems proved to that effect apply to nonspatial models, and in adaptive dynamics, it is standard to consider small differences between mutant and resident trait values. The spatial host–consumer ecosystem has the important property that, if mutation is an ongoing process, the spatial extent allows genetic diversity to grow. We initialize the system with all the consumers having the same trait value, but soon enough, different local subpopulations have different trait values. If the effects of single mutations are small, then the different varieties arising have roughly comparable survival probabilities, and so the distribution of extant trait values can spread out. However, the cumulative effect of many mutations which happen to act in the same direction on a trait such as transmissibility creates a variety which may engender its own local Malthusian catastrophe. So, the results of rare, big mutations tell us about the spread of trait values we see in the case of frequent, small mutations.

In our model, transmissibility and consumer death rate are independently adjustable parameters. One can also build a model in which one of these quantities is tied to the other, for example by imposing a tradeoff between transmissibility and virulence of a disease. Different functional forms of such a relationship are appropriate for modeling different ecosystems:

host/pathogen, prey/predator, sexual/parthenogenetic and so forth. As long as spatial pattern formation occurs and organism type impacts on the environment of descendants via ecosystem engineering, the shortcomings of mean-field theory are relevant, as are limitations of pair approximations [48].

B. Pair Approximations and Stability

Pair approximations have been used to test for the existence of an *Evolutionary Stable Strategy* (ESS) in a system—that is, a strategy which, when established, cannot be successfully replaced by another [70]. In addition to the limitations of pair approximation for representing patch structure [61], as we saw in the previous section, the question of whether a mutant strain can initially grow is distinct from the question of whether that strain achieves fixation or goes extinct [40, 50, 72, 90, 103–107]. The former is a question about short-term behavior, and the latter concerns effects apparent at longer timescales. This distinction is often lost or obscured in analytical treatments. The reason is that one typically tests whether a new type can invade by linearizing the corresponding differential equations at a point where its density is negligible. However, this only reveals the initial growth rate (see the fixed-point eigenvalue analysis in [11, 53, 60, 61]).

Our analysis implies that pair approximations are inadequate for analysis of systems with spatial inhomogeneity. Even including including triple and other higher-order corrections does not suffice, as this series approximation is poorly behaved at phase transitions [68]. Such higher order terms continue to reflect only the local structure of the system and not the existence of well separated areas that diverge in their genetic composition. Nonequilibrium pattern formation will necessarily also be poorly described, at least until the order of expansion reaches the characteristic number of elements in a patch, or an area that encompasses any relevant heterogeneity. Given the algebraic intricacy of higher-order corrections to pair approximations [53, 74, 108], it is useful to know in advance whether such elaborations have a chance at success. As approximation techniques based on successively refining mean-field treatments are blind to important phenomena, then we need to build our analytical work on a different conceptual foundation.

C. Percolation

The mathematical connection between pathogen–host and percolation problems can provide insight into the difficulties in analytical treatment of the biological problem. Spatial heterogeneity gives rise to failure of traditional analytic treatments of percolation and a need for new methodologies. Since the pathogen problem maps onto the percolation problem under some circumstances, the same analytic problems must arise in the biological context. While the presence of a nonequilibrium transition point indicates that traditional analysis techniques fail, it raises the possibility that new tools from the theory of phase transitions [33, 68, 79, 109] will become applicable. For example, in Section III F, we saw that percolation theory enables us to make quantitatively accurate predictions of population growth and of the critical parameter values which divide one ecological regime from another. Indeed, specific important problems in public health, such as the growth in number of individuals infected in a pandemic can be considered directly within the context of percolation. Simulations of propagation on approximate of real world networks may help provide accurate predictions, but the general properties of disease propagation can be understood analytically.

D. Adaptive Networks

Our results also have significance in the context of adaptive-network research. This field studies systems in which a network’s wiring pattern and the states of its nodes change in interrelated ways. Prior modeling efforts have considered epidemics on adaptive networks, where the spread of the disease *through* the network changes the connections *of* the network [67, 110–116]. In such models, if a susceptible node has an infected neighbor, it can break that connection by rewiring to another susceptible node. A key point in the analysis is that the new neighbor is chosen at random from the eligible population. This choice of rewiring scheme is exactly what makes a pair approximation work for that epidemic model, because it eliminates higher-order correlations in the system [67]. In our system, by contrast, hosts can form new connections by reproducing into empty sites, but these contacts can only connect geographically proximate individuals.

The difference we have seen between lattice behavior on one hand and RRG or swapping-

enabled behavior on the other emphasizes the need to study the effect of spatial proximity on link rewiring. While the structure-erasing nature of unconstrained rewiring among susceptible hosts has been acknowledged [113, 114], new rewiring rules which reflect spatial and community structure have yet to be systematically investigated. The reason that they have not is naturally related to the need for different analytic approaches. “Myopic” rewiring rules, such as restricting the set of eligible new partners to the neighbors of a node’s current partners, have on occasion been considered, but in contexts other than epidemiology, like evolutionary game theory [117, 118], making the endeavour of exploring such rules in epidemic models all the more worth pursuing.

E. Conclusions

Fisher [119] introduced modern genetic theory in large part motivated by the need to describe the existence of biodiversity. However, the expressions he described which apply in panmictic populations and mean-field treatments lead to a population genetics that rapidly converges to homogeneous populations. Spatial extents and their violation of the mean-field approximations are a key to biodiversity in nature. Their proper theoretical treatment will be a large step forward for evolutionary biology.

Most laboratory experiments, guided by traditional evolutionary thinking, have used well-mixed populations. The results obtained are consistent with theoretical analysis precisely because the conditions are consistent with those assumptions. Such experiments do not provide insight into the role of spatial extent and the implications for real-world biological populations. A growing number of experiments today are going beyond such conditions and, as is to be expected, are obtaining quite different results [24, 43, 45, 50, 76, 77, 107, 120–122].

Mean-field models are often helpful as a first step towards understanding the behavior of systems, but we cannot trust them to provide a complete story, and we should not let mean-field thinking furnish all the concepts we use to reason about evolutionary dynamics. Our analysis of transplanting organisms can be considered parallel to real world concerns and manifest effects of invasive species introduced by human activity and the impact of shipping and air transportation on pathogen evolution [40, 49]. These are among the well-established examples of situations in which spatial extent influences evolutionary dynamics [65, 107, 121, 123–127]. Identifying specific implications of the issues explored in this paper for particular

biological systems [24, 43, 45, 50, 76, 77, 107, 120–122] requires field and laboratory work, as well as theoretical insight to guide the questions that are being asked.

V. ACKNOWLEDGEMENTS

We thank M. de Aguiar, B. Allen, K. Z. Bertrand, E. Downes, A. S. Gard-Murray and D. Harmon for providing helpful comments on the manuscript.

-
- [1] S. Levin, “The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture,” *Ecology* **73** (1992) no. 6, 1943–67.
 - [2] U. Dieckmann and R. Law, “Relaxation projections and the method of moments,” in Dieckmann *et al.* [65], pp. 412–55.
 - [3] S. Wright, “*Tempo and Mode in Evolution: A Critical Review*,” *Ecology* **26** (1945) no. 4, 415–19, [JSTOR:1931666](#).
 - [4] D. L. Hartl and A. G. Clark, *Principles of Population Genetics*. Sinauer Associates, 2007.
 - [5] K. M. Page and M. Nowak, “Unifying Evolutionary Dynamics,” *Journal of Theoretical Biology* **219** (2002) 93–8, [PMID:12392978](#).
 - [6] P. Bijma and M. Wade, “The joint effects of kin, multilevel selection and indirect genetic effects on response to genetic selection,” *Journal of Evolutionary Biology* **21** (2008) 1175–88, [PMID:18547354](#).
 - [7] J. A. Damore and J. Gore, “Understanding microbial cooperation,” *Journal of Theoretical Biology* **299** (2012) 31–41, [PMID:21419783](#).
<http://gorelab.homestead.com/Papers/UnderstandingMicrobialCooperation.pdf>.
 - [8] Y. Bar-Yam, “Formalizing the gene-centered view of evolution,” *Advances in Complex Systems* **2** (1999) 277–81, [arXiv:physics/0002016](#).
<http://necsi.edu/research/evoeco/>.
 - [9] M. Kardar, *Statistical Physics of Particles*. Cambridge University Press, 2007.
<http://www.mit.edu/~kardar/teaching/index.html>.
 - [10] C. Goodnight *et al.*, “Evolution in spatial predator–prey models and the “prudent predator”: the inadequacy of steady-state organism fitness and the concept of individual

- and group selection,” *Complexity* **13** (2008) 23–44. <http://necsi.edu/research/evoeco/>.
- [11] S. Lion and M. van Baalen, “Self-structuring in spatial evolutionary ecology,” *Ecology Letters* **11** (2008) 277–95, PMID:18070102.
- [12] O. Givan *et al.*, “Predicting epidemic thresholds on complex networks: Limitations of mean-field approaches,” *Journal of Theoretical Biology* **288** (2011) 21–28, PMID:21840323.
- [13] G. C. Williams, *Natural Selection: Domains, Levels and Challenges*. Oxford University Press, 1992.
- [14] H. Sayama *et al.*, “Symmetry breaking and coarsening in spatially distributed evolutionary processes including sexual reproduction and disruptive selection,” *Physical Review E* **62** (2000) 7065–69.
- [15] J. D. Van Dyken, T. A. Linksvayer, and M. J. Wade, “Kin selection–mutation balance: A model for the origin, maintenance and consequences of social cheating,” *American Naturalist* **177** (2011) no. 3, 288–300, PMID:21460538.
- [16] M. Archetti and I. Scheuring, “Coexistence of cooperation and defection in public goods games,” *Evolution* **65** (2011) no. 4, 1140–48, PMID:21062277.
- [17] P. D. Taylor and S. A. Frank, “How to Make a Kin Selection Model,” *Journal of Theoretical Biology* **180** (1996) 27–37, PMID:8763356.
- [18] H. Kokko, K. U. Heubel, and D. J. Rankin, “How populations persist when asexuality requires sex: the spatial dynamics of coping with sperm parasites,” *Proceedings of the Royal Society B* **275** (2008) 817–25, PMID:18182369.
- [19] G. Wild, A. Gardner, and S. A. West, “Adaptation and the evolution of parasite virulence in a connected world,” *Nature* **459** (2009) 983–6, PMID:19474791.
- [20] R. Levins, “Some demographic and genetic consequences of environmental heterogeneity for biological control,” *Bulletin of the Entomological Society of America* **15** (1969) 237.
- [21] A. Platt *et al.*, “The scale of population structure in *Arabidopsis thaliana*,” *PLOS Genetics* **6** (2010) no. 2, e1000843.
- [22] J. M. Halley *et al.*, “Uses and abuses of fractal methodology in ecology,” *Ecology Letters* **7** (2004) 254–71.
- [23] T. M. Scanlon, K. K. Caylor, S. A. Levin, and I. Rodriguez-Iturbe, “Positive feedbacks promote power-law clustering of Kalahari vegetation,” *Nature* **449** (2007) no. 7159, 209–12, PMID:17851523.

- [24] C. Reigada and M. A. M. de Aguiar, “Host-parasitoid persistence over variable spatio-temporally susceptible habitats: bottom-up effects of ephemeral resources,” *Oikos* **121** (2012) 1665–79. <http://www.ifi.unicamp.br/~aguiar/Abstracts/2011h.html>.
- [25] G. Ichinose, M. Saito, H. Sayama, and D. S. Wilson, “Adaptive long-range migration promotes cooperation under tempting conditions,” *Scientific Reports* **3** (2013) 2509, [arXiv:1306.0072](https://arxiv.org/abs/1306.0072) [q-bio.PE].
- [26] A. Escalante *et al.*, “The dynamics of mutations associated with anti-malarial drug resistance in *Plasmodium falciparum*,” *Trends in Parasitology* **25** (2009) no. 12, 557–63, [PMC:2881657](https://pubmed.ncbi.nlm.nih.gov/2881657/).
- [27] T. Gross and B. Blasius, “Adaptive coevolutionary networks: a review,” *Journal of the Royal Society Interface* **5** (2008) no. 20, 259–71, [PMC:2405905](https://pubmed.ncbi.nlm.nih.gov/2405905/).
- [28] T. Gross and H. Sayama, eds., *Adaptive Networks: Theory, Models, Applications*. Springer, 2009.
- [29] O. Gräser, C. Xu, and P. M. Hui, “Analytic approach to co-evolving dynamics in complex networks: dissatisfied adaptive snowdrift game,” *New Journal of Physics* **13** (2011) no. 8, 083015.
- [30] G. Demirel, F. Vazquez, G. A. Böhme, and T. Gross, “Moment-closure approximations for discrete adaptive networks,” [arXiv:1211.0449](https://arxiv.org/abs/1211.0449) [nlin.AO].
- [31] J. Werfel and Y. Bar-Yam, “The evolution of reproductive restraint through social communication,” *PNAS* **101** (2004) no. 30, 11019–24, [PMID:15256603](https://pubmed.ncbi.nlm.nih.gov/15256603/).
<http://www.necsi.edu/projects/evolecol/altruismpnaspr.html>.
- [32] A. Jiménez-Dalmaroni and H. Hinrichsen, “Epidemic processes with immunization,” *Physical Review E* **68** (2003) 036103, [arXiv:cond-mat/0304113](https://arxiv.org/abs/cond-mat/0304113).
- [33] M. Henkel, H. Hinrichsen, and S. Lübeck, *Non-Equilibrium Phase Transitions, Volume 1: Absorbing Phase Transitions*. Springer, 2008.
- [34] H. Kokko and K. U. Heubel, “Prudent males, group adaptation, and the tragedy of the commons,” *Oikos* **120** (2011) 641–56.
- [35] Y. Haraguchi and A. Sasaki, “The Evolution of Parasite Virulence and Transmission Rate in a Spatially Structured Population,” *Journal of Theoretical Biology* **203** (2000) 85–96, [PMID:10704294](https://pubmed.ncbi.nlm.nih.gov/10704294/).
- [36] H. Sayama, M. A. M. de Aguiar, Y. Bar-Yam, and M. Baranger, “Spontaneous pattern

- formation and genetic invasion in locally mating and competing populations,” *Physical Review E* **65** (2002) 051919.
- [37] M. A. M. de Aguiar, M. Baranger, Y. Bar-Yam, and H. Sayama, “Robustness of spontaneous pattern formation in spatially distributed genetic populations,” *Brazilian Journal of Physics* **33** (2003) no. 3, 514–20.
- [38] H. Sayama, L. Kaufman, and Y. Bar-Yam, “Spontaneous pattern formation and genetic diversity in habitats with irregular geographical features,” *Conservation Biology* **17** (2003) 893–900.
- [39] C. G. Jones, J. H. Lawton, and M. Shachak, “Organisms as ecosystem engineers,” *Oikos* **69** (1994) no. 3, 373–86, [JSTOR:3545850](#).
- [40] D. L. Strayer, V. T. Eviner, J. M. Jeschke, and M. L. Pace, “Understanding the long-term effects of species invasions,” *Trends in Ecology and Evolution* **21** (2006) no. 11, 645–51, [PMID:16859805](#).
- [41] D. M. Post and E. P. Palkovacs, “Eco-evolutionary feedbacks in community and ecosystem ecology: interactions between the ecological theatre and the evolutionary play,” *Philosophical Transactions of the Royal Society B* **364** (2009) 1629–40.
- [42] R. M. Pringle *et al.*, “Spatial pattern enhances ecosystem functioning in an african savanna,” *PLOS Biology* **8** (2010) no. 5, e1000377.
- [43] B. Allen and M. A. Nowak, “Cooperation and the fate of microbial societies,” *PLOS Biology* **11** (2013) no. 4, e1001549.
- [44] J. T. Cronin and J. D. Reeve, “Host-parasitoid spatial ecology: a plea for a landscape-level synthesis,” *Proceedings of the Royal Society B* **272** (2005) no. 1578, 2225–35, [PMID:16191634](#).
- [45] B. Kerr, C. Neuhauser, B. J. M. Bohannan, and A. M. Dean, “Local migration promotes competitive restraint in a host–pathogen ‘tragedy of the commons’,” *Nature* **442** (2006) 75–8, [PMID:16823452](#).
- [46] S. M. Messinger and A. Ostling, “The consequences of spatial structure for the evolution of pathogen transmission rate and virulence,” *American Naturalist* **174** (2009) no. 4, 441–54.
- [47] S. Lion and M. Boots, “Are parasites “prudent” in space?,” *Ecology Letters* **13** (2010) 1245–55.
- [48] S. M. Messinger and A. Ostling, “The influence of host demography, pathogen virulence,

- and relationships with pathogen virulence on the evolution of pathogen transmission in a spatial context,” *Evolutionary Ecology* **27** (2012) no. 2, 353–80.
- [49] E. Rauch and Y. Bar-Yam, “Long-range interactions and evolutionary stability in a predator–prey system,” *Physical Review E* **73** (2006) 020903, PMID:16605322. <http://necsi.edu/research/evoeco/>.
- [50] S. Heilmann, K. Sneppen, and S. Krishna, “Sustainability of virulence in a phage-bacteria ecosystem,” *Journal of Virology* **84** (2010) no. 6, 3016–22.
- [51] E. Rauch, H. Sayama, and Y. Bar-Yam, “Relationship between Measures of Fitness and Time Scale in Evolution,” *Physical Review Letters* **88** (2002) 228101, PMID:12059453.
- [52] E. Rauch, H. Sayama, and Y. Bar-Yam, “Dynamics and Genealogy of Strains in Spatially Extended Host–Pathogen Models,” *Journal of Theoretical Biology* **221** (2003) 655–64, PMID:12713947.
- [53] M. A. M. de Aguiar, E. Rauch, and Y. Bar-Yam, “Invasion and Extinction in the Mean Field Approximation for a Spatial Host–Pathogen Model,” *Journal of Statistical Physics* **114** (2004) no. 5/6, 1417–51.
- [54] R. Froissart, J. Doumayrou, F. Vuillaume, S. Alizon, and Y. Michalakis, “The virulence–transmission trade-off in vector-borne plant viruses: a review of (non-)existing studies,” *Philosophical Transactions of the Royal Society B* **365** (2010) no. 1548, 1907–18, PMC:2880117.
- [55] M. K. Asplen *et al.*, “Do trade-offs have explanatory power for the evolution of organismal interactions?,” *Evolution* **66** (2012) no. 5, 1297–1307.
- [56] D. M. Hawley *et al.*, “Parallel patterns of increased virulence in a recently emerged wildlife pathogen,” *PLOS Biology* **11** (2013) no. 5, e1001570.
- [57] M. Mobilia, I. T. Georgiev, and U. C. Täuber, “Phase transitions and spatio-temporal fluctuations in stochastic lattice Lotka–Volterra models,” *Journal of Statistical Physics* **128** (2006) no. 1–2, 447–83, arXiv:q-bio/0512039.
- [58] M. A. M. de Aguiar, E. M. Rauch, and Y. Bar-Yam, “Mean-field approximation to a spatial host-pathogen model,” *Physical Review E* **67** (2003) no. 4, 047102, arXiv:1307.5335 [q-bio.PE]. <http://necsi.edu/research/evoeco/>.
- [59] M. A. M. de Aguiar, E. Rauch, B. C. Stacey, and Y. Bar-Yam, “Erratum: Mean-field approximation to a spatial host-pathogen model,” *Physical Review E* **88** (2013) no. 3,

- 039901(E), [arXiv:1307.5335 \[q-bio.PE\]](#).
- [60] M. van Baalen and D. A. Rand, “The unit of selection in viscous populations and the evolution of altruism,” *Journal of Theoretical Biology* **193** (1998) 631–48, [PMID:9750181](#).
- [61] M. van Baalen, “Pair Approximations for Different Spatial Geometries,” in Dieckmann *et al.* [65], pp. 359–85.
- [62] G. Rozhnova and A. Nunes, “Population dynamics on random networks: simulations and analytical models,” *European Physical Journal B* **74** (2010) no. 2, 235–42, [arXiv:0907.0335 \[q-bio\]](#).
- [63] B. Allen, *Studies in the Mathematics of Evolution and Biodiversity*. PhD thesis, Boston University, 2010.
<http://proquest.umi.com/pqdweb?did=2071736811&Fmt=2&clientId=5482&RQT=309>.
- [64] S. B. L. Araujo, G. M. Viswanathan, and M. A. M. de Aguiar, “Home range evolution and its implication in population outbreaks,” *Philosophical Transactions of the Royal Society A* **368** (2010) 5661–77.
- [65] U. Dieckmann, R. Law, and J. A. J. Metz, eds., *The Geometry of Ecological Interactions: Simplifying Spatial Complexity*. Cambridge University Press, 2000.
- [66] H. Matsuda, N. Ogita, A. Sasaki, and K. Sato, “Statistical mechanics of population,” *Progress of Theoretical Physics* **88** (1992) no. 6, 1035–49.
- [67] A.-L. Do and T. Gross, “Contact processes and moment closure on adaptive networks,” in Gross and Sayama [28], pp. 191–208.
- [68] M. A. Buice and J. D. Cowan, “Statistical mechanics of the neocortex,” *Progress in Biophysics and Molecular Biology* **99** (2009) 53–86, [PMID:19695282](#).
- [69] P. J. Dodd and N. M. Ferguson, “A many-body field theory approach to stochastic models in population biology,” *PLOS ONE* **4** (2009) no. 9, e6855, [PMID:19730742](#).
- [70] G. Szabó and G. Fáth, “Evolutionary games on graphs,” *Physics Reports* **446** (2007) 97–216, [arXiv:cond-mat/0607344](#).
- [71] B. Allen, A. Traulsen, C. E. Tarnita, and M. A. Nowak, “How mutation affects evolutionary games on graphs,” *Journal of Theoretical Biology* **299** (2012) 97–105, [PMID:21473871](#). <http://www.people.fas.harvard.edu/~ballen/GraphMutInPress.pdf>.
- [72] P. E. Smaldino, J. C. Schank, and R. McElreath, “Increased costs of cooperation help cooperators in the long run,” *American Naturalist* **181** (2013) no. 4, 451–63.

- <http://xcelab.net/rmpubs/Smaldino%20Am%20Nat%202013.pdf>.
- [73] H. J. Poethke, B. Pfenning, and T. Hovestadt, “The relative contribution of individual and kin selection to the evolution of density-dependent dispersal rates,” *Evolutionary Ecology Research* **9** (2007) 41–50, [OPUS:2010/4822](#).
- [74] S. Lion, “Relatedness in spatially structured populations with empty sites: An approach based on spatial moment equations,” *Journal of Theoretical Biology* **260** (2009) 121–31.
- [75] E. Arashiro and T. Tomé, “Threshold of coexistence and critical behavior of a predator-prey cellular automaton,” *Journal of Physics A* **40** (2007) 887–900, [arXiv:cond-mat/0607360](#).
- [76] S. Davis, P. Trapman, H. Leirs, M. Begon, and J. A. P. Heesterbeek, “The abundance threshold for plague as a critical percolation phenomenon,” *Nature* **454** (2008) 634–37.
- [77] D. J. Salkeld, M. Salathé, P. Stapp, and J. H. Jones, “Plague outbreaks in prairie dog populations explained by percolation thresholds of alternate host abundance,” *Proceedings of the National Academy of Sciences* **107** (2010) no. 32, 14247–50.
- [78] F. M. Neri *et al.*, “The effect of heterogeneity on invasion in spatial epidemics: from theory to evidence in a model system,” *PLOS Computational Biology* **7** (2011) no. 9, e1002174.
- [79] H. Hinrichsen, “Nonequilibrium critical phenomena and phase transitions into absorbing states,” *Advances in Physics* **49** (2000) no. 7, 815–958, [arXiv:cond-mat/0001070](#).
- [80] M. A. Saif and P. M. Gade, “Dynamic phase transition in the prisoner’s dilemma on a lattice with stochastic modifications,” *Journal of Statistical Mechanics* (2010) P03016, [arXiv:0910.0955 \[cond-mat.stat-mech\]](#).
- [81] J. Wendykier, A. Lipowski, and A. L. Ferreira, “Coexistence and critical behaviour in a lattice model of competing species,” *Physical Review E* **83** (2011) no. 3, 031904, [arXiv:1010.2538 \[q-bio.PE\]](#).
- [82] A. Szolnoki, G. Szabó, and M. Perc, “Phase diagrams for the spatial public good game with pool-punishment,” *Physical Review E* **83** (2011) no. 3, 036101, [arXiv:1102.0624 \[physics.soc-ph\]](#).
- [83] P. Grassberger, H. Chaté, and G. Rousseau, “Spreading in media with long-time memory,” *Physical Review E* **55** (1997) no. 3, 2488–95.
- [84] H. Kesten, “The critical probability of bond percolation on the square lattice equals $1/2$,” *Communications in Mathematical Physics* **74** (1980) no. 1, 41–59.
- [85] G. Grimmett, *Percolation*. Springer, 1999.

- [86] M. E. J. Newman, “Spread of epidemic disease on networks,” *Physical Review E* **66** (2002) 016128, [arXiv:cond-mat/0205009](#).
- [87] J. P. Gleeson, “High-accuracy approximation of binary-state dynamics on networks,” *Physical Review Letters* **107** (2011) 068701, [arXiv:1104.1537](#) [physics.soc-ph].
- [88] H. A. Orr, “Fitness and its role in evolutionary genetics,” *Nature Reviews Genetics* **10** (2009) 531–39.
- [89] J. Smith, J. D. Van Dyken, and P. C. Zee, “A generalization of Hamilton’s rule for the evolution of microbial cooperation,” *Science* **328** (2010) 1700–03.
- [90] C. A. Fogle, J. L. Nagle, and M. M. Desai, “Clonal interference, multiple mutations and adaptation in large asexual populations,” *Genetics* **180** (2008) no. 4, 2163–73, [PMID:18832359](#).
- [91] A. Grafen, “A geometric view of relatedness,” *Oxford Surveys in Evolutionary Biology* **2** (1985) 28–89.
- [92] B. Simon, J. A. Fletcher, and M. Doebeli, “Hamilton’s rule in multi-level selection models,” *Journal of Theoretical Biology* **299** (2012) 55–63, [PMID:21820447](#).
- [93] B. Allen and C. E. Tarnita, “Measures of success in a class of evolutionary models with fixed population size and structure,” *Journal of Mathematical Biology* **66** (2012) online before print. <http://scholar.princeton.edu/ctarnita/files/AxiomSuccessPublished.pdf>.
- [94] H. K. Reeve and B. Hölldobler, “The emergence of a superorganism through intergroup competition,” *PNAS* **104** (2007) no. 23, 9736–40, [PMID:1887545](#).
- [95] B. Allen, M. A. Nowak, and E. O. Wilson, “Limitations of inclusive fitness,” *Proceedings of the National Academy of Sciences* **110** (2013) online before print. <http://plektix.fieldofscience.com/2013/12/whats-deal-with-inclusive-fitness-theory.html>.
- [96] M. Wade *et al.*, “Multilevel and kin selection in a connected world,” *Nature* **463** (2010) E8. <http://necsi.edu/research/evoeco/>.
- [97] B. Simon, J. A. Fletcher, and M. Doebeli, “Towards a general theory of group selection,” *Evolution* **67** (2013) no. 6, 1561–72. <http://math.ucdenver.edu/~bsimon/papers.html>.
- [98] A. Nielsen and B. Simon, “Numerical solutions and animations of group selection dynamics,” *Evolutionary Ecology Research* **44** (2012) 757–68. <http://math.ucdenver.edu/~bsimon/papers.html>.

- [99] G. Wild and P. D. Taylor, “Fitness and evolutionary stability in game theoretic models of finite populations,” *Proceedings of the Royal Society of London B* **271** (2004) 2345–9, PMID:15590589.
- [100] S. D. Mylius and O. Diekmann, “The resident strikes back: Invader-induced switching of resident attractor,” *Journal of Theoretical Biology* **211** (2001) 297–311. <http://sido.mylius.nl/downloads/rstrbftp.pdf>.
- [101] S. A. H. Geritz, M. Gyllenberg, F. J. A. Jacobs, and K. Parvinen, “Invasion dynamics and attractor inheritance,” *Journal of Mathematical Biology* **44** (2002) 548–60.
- [102] S. A. H. Geritz, “Resident-invader dynamics and the coexistence of similar strategies,” *Journal of Mathematical Biology* **50** (2005) 67–82.
- [103] M. Doebeli, “Invasion of rare mutants does not imply their evolutionary success: a counterexample from metapopulation theory,” *Journal of Evolutionary Biology* **11** (1997) no. 3, 389–401. <http://www.math.ubc.ca/~doebeli/reprints/Doe26.pdf>.
- [104] S. Lessard, “Long-term stability from fixation probabilities in finite populations: New perspectives for ESS theory,” *Theoretical Population Biology* **68** (2005) 19–27, PMID:16023912.
- [105] T. Antal and I. Scheuring, “Fixation of strategies in an evolutionary game in finite populations,” *Bulletin of Mathematical Biology* **68** (2006) no. 8, 1923–44, [arXiv:q-bio/0509008](http://arxiv.org/abs/q-bio/0509008).
- [106] C. J. Paley, S. N. Taraskin, and S. R. Elliott, “Temporal and dimensional effects in evolutionary graph theory,” *Physical Review Letters* **98** (2007) 098103, [arXiv:q-bio/0604009](http://arxiv.org/abs/q-bio/0604009).
- [107] J. M. Ponciano, H. J. La, P. Joyce, and L. J. Forney, “Evolution of diversity in spatially structured *Escherichia coli* populations,” *Applied and Environmental Microbiology* **75** (2009) no. 19, 6047–54, PMID:19648364.
- [108] M. Raghil, N. A. Hill, and U. Dieckmann, “Multiscale maximum entropy closure for locally regulated space-time point process models of plant population dynamics,” *Journal of Mathematical Biology* **62** (2010) no. 5, 605–53, PMID:20446087.
- [109] H.-K. Janssen and U. C. Täuber, “The field theory approach to percolation processes,” *Annals of Physics* **315** (2005) 147–92, [arXiv:cond-mat/0409670](http://arxiv.org/abs/cond-mat/0409670).
- [110] T. Gross, C. J. D. D’Lima, and B. Blasius, “Epidemic dynamics on an adaptive network,”

- Physical Review Letters* **96** (2006) no. 20, 208701, [arXiv:q-bio/0512037](#).
- [111] L. B. Shaw and I. B. Schwartz, “Noise induced dynamics in adaptive networks with applications to epidemiology,” in Gross and Sayama [28], pp. 209–27.
- [112] L. B. Shaw and I. B. Schwartz, “Enhanced vaccine control of epidemics in adaptive networks,” *Physical Review E* **81** (2010) no. 4, 046120, [PMC:2931598](#).
- [113] C. Kamp, “Untangling the interplay between epidemic spread and transmission network dynamics,” *PLOS Computational Biology* **6** (2010) no. 11, e1000984, [PMC:2987842](#).
- [114] S. Van Segbroeck, F. C. Santos, and J. M. Pacheco, “Adaptive contact networks change effective disease infectiousness and dynamics,” *PLOS Computational Biology* **6** (2010) no. 8, e1000895, [PMC:2924249](#).
- [115] B. Wu *et al.*, “Evolution of cooperation on stochastic dynamical networks,” *PLOS ONE* **5** (2010) no. 6, e11187, [PMC:2894855](#).
- [116] K. Fehl, D. J. van der Post, and D. Semmann, “Co-evolution of behaviour and social network structure promotes human cooperation,” *Ecology Letters* **14** (2011) 546–51, [PMID:21463459](#).
- [117] A. Traulsen, F. C. Santos, and J. M. Pacheco, “Evolutionary games in self-organizing populations,” in Gross and Sayama [28], pp. 253–68.
- [118] P. Holme and G. Ghoshal, “The diplomat’s dilemma: Maximal power for minimal effort in social networks,” in Gross and Sayama [28], pp. 269–88.
- [119] R. A. Fisher, *The Genetical Theory of Natural Selection*. Clarendon, 1930.
- [120] J. B. Xavier and K. R. Foster, “Cooperation and conflict in microbial biofilms,” *PNAS* **104** (2007) no. 3, 876–81, [PMID:17210916](#).
- [121] B. Göncü *et al.*, “Viral epidemics in a cell culture: Novel high resolution data and their interpretation by a percolation theory based model,” *PLOS ONE* **5** (2010) no. 12, e15571, [PMC:3004943](#).
- [122] D. Wodarz *et al.*, “Complex spatial dynamics of oncolytic viruses in vitro: Mathematical and experimental approaches,” *PLOS Computational Biology* **8** (2012) no. 6, e1002547.
- [123] M. A. M. de Aguiar, M. Baranger, E. M. Baptestini, L. Kaufman, and Y. Bar-Yam, “Global patterns of speciation and diversity,” *Nature* **460** (2009) 384–87.
<http://www.necsi.edu/research/evoeco/globalpatternsofspeci/index.html>.
- [124] M. A. M. de Aguiar and Y. Bar-Yam, “The Moran model as a dynamical process on

- networks and its implications for neutral speciation,” *Physical Review E* **84** (2011) 031901, [arXiv:1012.3913](https://arxiv.org/abs/1012.3913) [q-bio.PE].
- [125] A. B. Martins, M. A. M. de Aguiar, and Y. Bar-Yam, “Evolution and stability of ring species,” *PNAS* **110** (2013) no. 13, 5080–84.
<http://www.necsi.edu/research/evoeco/ringspecies/>.
- [126] J. D. Van Dyken, M. J. I. Müller, K. M. L. Mack, and M. M. Desai, “Spatial population expansion promotes the evolution of cooperation in an experimental prisoner’s dilemma,” *Current Biology* **23** (2013) no. 10, 919–23, [arXiv:1311.2646](https://arxiv.org/abs/1311.2646) [q-bio.PE].
- [127] E. M. Baptestini, M. A. M. de Aguiar, and Y. Bar-Yam, “Conditions for neutral speciation via isolation by distance,” *Journal of Theoretical Biology* **335** (2013) 51–6.
<http://necsi.edu/research/evoeco/>.