

# Beyond the Heisenberg time: Semiclassical treatment of spectral correlations in chaotic systems with spin 1/2

Petr Braun

*Fachbereich Physik, Universität Duisburg–Essen, 47048 Duisburg, GERMANY  
Institute of Physics, Saint-Petersburg University, 198504 Saint-Petersburg, RUSSIA*

The two-point correlation function of chaotic systems with spin 1/2 is evaluated using periodic orbits. The spectral form factor for all times thus becomes accessible. Equivalence with the predictions of random matrix theory for the Gaussian symplectic ensemble is demonstrated. A duality between the underlying generating functions of the orthogonal and symplectic symmetry classes is semiclassically established.

PACS numbers: 05.45.-a, 05.45.Mt

## I. INTRODUCTION

Energy levels of classically chaotic systems exhibit correlation only slowly subsiding with the energy offset [1]. Around 1980 it became clear, after extensive numerical experiments, that with as few as two degrees of freedom, spectral correlations in the highly excited energy domain have universal properties and obey the same laws as the eigenvalues of the Gaussian random matrix ensembles of the appropriate symmetry class [2–5]. This assertion, known as the Bohigas-Giannoni-Schmitt (BGS) conjecture took a surprisingly long time to be proven. In most cases the tool used was the Gutzwiller formula giving the spectral density  $\rho(E)$  in a chaotic system as sum over the classical periodic orbits, each orbit creating a contribution  $\sim e^{iS_\gamma/\hbar}$  where  $S_\gamma$  is the action of the orbit  $\gamma$ . Its substitution into the spectral correlation function and its Fourier transform  $K(\tau)$ , the spectral form factor, leads to double sums over orbit pairs with summands proportional to  $e^{i(S_\gamma - S_{\gamma'})/\hbar}$ . Significant contributions can be expected only from pairs with the action difference not large compared with  $\hbar$ .

The first success in the proof of BGS was connected with the diagonal approximation [6] which takes into account only pairs with  $S_\gamma = S_{\gamma'}$ ; it explained the fact that at small times  $K(\tau) \approx 2\tau$  (time reversal allowed, orthogonal universality class) or  $\tau$  (time reversal forbidden, unitary class). Fifteen years later came the realization that a long periodic orbit with a small-angle self-crossing dividing it into two pieces, has a “partner” orbit with the crossing avoided, but otherwise almost unchanged, up to the sense of traversal of one of the pieces [7]. Contributions of such “Sieber-Richter pairs” sum up the next-to-leading term  $-2\tau^2$  in the form factor. Summation over pairs in which the partner consists of pieces of the original orbit reconnected in all thinkable ways, restores the small-time form factor to all orders [8, 9].

The form factor experiences a break-up of analyticity at  $\tau = 1$ , i.e., at the Heisenberg time  $T_H = 2\pi\hbar\bar{\rho}$ , which reflects the existence of an oscillatory component of the correlation function with the period of the mean level spacing  $\Delta = 1/\bar{\rho}$ . That component is overlooked in the straightforward semiclassical approach providing correlation functions as asymptotic power series in  $1/\varepsilon$ ; the reason is that semiclassical sums need for their convergence a non-vanishing positive imaginary part of the energy parameters, however  $e^{i\varepsilon} \sim 0 \cdot \varepsilon^{-1} + 0 \cdot \varepsilon^{-2} + \dots$  if  $\text{Im } \varepsilon > \delta > 0$ . Early estimates of the oscillatory components are contained in [10, 11]. Systematic approach is based on the formalism of generating functions, i.e., averaged ratios of the spectral determinants. In this approach partnership of more than two classical orbits is taken into account, and use is made of the semiclassical approximation of the spectral determinant known as the Riemann-Siegel look-alike [12–15]. As a result complete agreement of the semiclassical correlation functions with RMT for spinless systems was demonstrated [16, 17].

Systems with half-integer spin belong to the symplectic universality class whose RMT counterpart is the Gaussian symplectic ensemble. The spin coupling to chaotic translational motion leads to randomization of the spin evolution [18]. The ergodicity of that evolution is instrumental for the evaluation of the relevant periodic orbit expansions such as the diagonal sum for the form factor [19] and the contribution of the Sieber-Richter pairs which changes its sign in the presence of a half-integer spin [20]. The full expansion of the form factor of systems with symplectic symmetry for times smaller than  $T_H$  was obtained for the quantum graphs in [21] and for general dynamical systems in [9].

Here we close the gap in the proof of BGS for the systems with half-integer spin by showing the equivalence of their semiclassical correlation function with the RMT predictions including the oscillatory terms; the corresponding form factors coincide for all times. Analytical properties of the correlation function are used to recover the oscillatory term with smaller frequency responsible for the well-known logarithmic singularity of the form factor. The derivation employs the simple duality discovered between the semiclassical 4-determinant generating functions of chaotic systems

with symplectic and orthogonal symmetry. We talk mostly about the spin 1/2 case although the results remain true for other half-integer spins.

## II. COMPLEX CORRELATOR, FORM FACTOR AND GENERATING FUNCTION

Our main object of study will be the complex two-point spectral correlation function (complex correlator, for short). This is an analytic function of the complex dimensionless variable  $\varepsilon$  which is the double spectral sum,

$$C(\varepsilon) = \frac{\Delta^2}{2\pi^2} \left\langle \sum_{i \neq k} \frac{1}{(E_k - E - \frac{\varepsilon \Delta}{2\pi})} \frac{1}{(E_i - E + \frac{\varepsilon \Delta}{2\pi})} \right\rangle - \frac{1}{2} \quad (1)$$

where  $\Delta$  stands for the mean level spacing;  $\langle \dots \rangle$  denotes averaging over an interval of the reference energy  $E$ , classically small but large compared with  $\Delta$ . The complex correlator is defined in the half plane  $\text{Im } \varepsilon > 0$  where it is analytic and tends to zero when  $|\varepsilon| \rightarrow \infty$ , and can be continued to the lower half plane where (1) would no longer be true. The real part of  $C(\varepsilon)$  at the positive real axis coincides with the real level-level correlation function [1] while its Fourier transform is the spectral form factor  $K(\tau)$ ; the connection between the two functions is given by

$$K(\tau) = \mathcal{F}(C) = \frac{1}{2\pi\tau_H} \int_{-\infty+i0}^{\infty+i0} C(\varepsilon) e^{-i2\varepsilon\tau/\tau_H} d\varepsilon, \quad \tau > 0; \quad (2)$$

$$C(\varepsilon) = \mathcal{F}^{-1}(K) = 2 \int_0^{\infty} e^{i2\varepsilon\tau/\tau_H} K(\tau) d\tau, \quad \text{Im } \varepsilon > 0. \quad (3)$$

Here  $\tau_H$  stands for 1 for the orthogonal universality class and 2 for the symplectic class; such choice is equivalent to the replacement  $2\varepsilon \rightarrow \varepsilon$  in the symplectic case motivated by the Kramers degeneracy [11].

The semiclassical evaluation of the complex correlator is based on the generating function defined as the averaged ratio of four spectral determinants,

$$Z(\hat{\varepsilon}) = \left\langle \frac{\det(H - E - \frac{\varepsilon_C \Delta}{2\pi}) \det(H - E - \frac{\varepsilon_D \Delta}{2\pi})}{\det(H - E - \frac{\varepsilon_A \Delta}{2\pi}) \det(H - E - \frac{\varepsilon_B \Delta}{2\pi})} \right\rangle. \quad (4)$$

where  $(\hat{\varepsilon}) \equiv (\varepsilon_A, \varepsilon_B, \varepsilon_C, \varepsilon_D)$ . The complex correlator can be obtained from  $Z$  as

$$C(\varepsilon) = \lim_{\substack{\varepsilon_A, \varepsilon_C \rightarrow \varepsilon \\ \varepsilon_B, \varepsilon_D \rightarrow -\varepsilon}} 2 \frac{\partial^2}{\partial \varepsilon_A \partial \varepsilon_B} Z \Big|_{\parallel}. \quad (5)$$

## III. SEMICLASSICAL GENERATING FUNCTION FOR ORTHOGONAL SYMMETRY CLASS

Here briefly we recapitulate the results for systems with orthogonal symmetry. The semiclassical representation of the generating function follows from the chain of relations [16, 17],

$$\det(H - E) \sim \exp \left[ - \int^E dE' \text{Tr} (H - E')^{-1} \right] \sim \exp \left[ - \sum_{\gamma} f_{\gamma} e^{iS_{\gamma}(E)/\hbar} \right];$$

the last step is the Gutzwiller expansion of the exponent into a sum over periodic orbits  $\gamma$  with the actions  $S_{\gamma}$  and stability coefficients  $f_{\gamma}$ . Expanding all four exponentials we get a sum over quadruplets of “pseudo-orbits”  $A, B, C, D$ ,

$$Z^{(1)}(\hat{\varepsilon}) = \left\langle \sum_{A, B, C, D} F_A F_C F_B^* F_D^* (-1)^{\nu_C + \nu_D} \right. \\ \left. \times e^{i\Delta S/\hbar} e^{i(T_A \varepsilon_A + T_C \varepsilon_C - T_B \varepsilon_B - T_D \varepsilon_D)/T_H} \right\rangle; \quad (6)$$

a pseudo-orbit, say  $A$ , is a set of  $\nu_A$  periodic orbits whose actions and periods sum up to  $S_A, T_A$  and whose product of stability coefficients is  $F_A$ . The difference of actions  $\Delta S = S_A + S_C - S_B - S_D$  must be small compared with  $\hbar$  for the quadruplets making meaningful contributions. Note the sign factor  $(-1)^{\nu_C + \nu_D}$  depending on the number of orbits in the pseudo-orbits associated with the numerator of the generating function.

The leading contribution is created by the diagonal quadruplets in which the pseudo-orbit pair  $(B, D)$  contains the same periodic orbits as  $(A, C)$  and consequently  $\Delta S = 0$ . The diagonal contributions sum up to

$$Z_{\text{diag}} = e^{\frac{i}{2}(\varepsilon_A - \varepsilon_B - \varepsilon_C + \varepsilon_D)} \frac{(\varepsilon_A - \varepsilon_D)^2 (\varepsilon_C - \varepsilon_B)^2}{(\varepsilon_A - \varepsilon_B)^2 (\varepsilon_C - \varepsilon_D)^2} \quad (7)$$

and can be factored out like  $Z^{(1)} = Z_{\text{diag}} (1 + Z_{\text{off}})$ ; the off-diagonal part  $Z_{\text{off}}$  stands for a sum similar to (6) but with all orbits of  $(A, C)$  different from those of  $(B, D)$ . Contributions with a small action mismatch now come from the quadruplets in which the periodic orbits of  $(B, D)$  are “partners” of those of  $(A, C)$ , i.e., consist of practically the same but differently connected pieces. Reconnections occur in “encounters” which are places of close approach of  $l \geq 2$  almost parallel stretches of the same or different orbits; the possibility of such a reconnection is a fundamental property of chaotic motion [22]. Using ergodicity summation over all pseudo-orbit quadruplets could be reduced to summation over their topological families (“structures”). In the end the expansion  $Z_{\text{off}} = \sum_{n=1}^{\infty} Z_n$ , was obtained [16, 17]. Here  $Z_n \sim \varepsilon^{-n}$  accumulates contributions of quadruplets with  $L - V = n$ ;  $V$  is the number of encounters containing  $L = \sum_{i=1}^V l_i$  stretches. The explicit expression of  $Z_n$  for the orthogonal class is (“O”=orthogonal),

$$Z_{n,O}(\hat{\varepsilon}) = \frac{(\varepsilon_A - \varepsilon_C)(\varepsilon_B - \varepsilon_D)}{(\varepsilon_A - \varepsilon_D)(\varepsilon_B - \varepsilon_C)} \quad (8)$$

$$\times \frac{(-i)^n (n-1)! 2^n}{(\varepsilon_A - \varepsilon_B)^{n-1}} \left( \frac{1}{\varepsilon_C - \varepsilon_D} + \frac{n}{\varepsilon_A - \varepsilon_B} \right).$$

The true high-energy asymptotics of the generating function is not exhausted by  $Z^{(1)}$ . Recovery of the missing component was achieved on the basis of the so called “Riemann-Siegel look-alike” representation [12, 14, 15] of the spectral determinant, due to the basic quantum mechanical symmetry properties. The additional summand is obtained from  $Z^{(1)}(\hat{\varepsilon})$  by the “Weyl transposition”  $w(\hat{\varepsilon}) \equiv (\varepsilon_A, \varepsilon_B, \varepsilon_D, \varepsilon_C)$  of its arguments, ,

$$Z_O(\hat{\varepsilon}) \sim Z^{(12)} \equiv Z^{(1)}(\hat{\varepsilon}) + Z^{(2)}(\hat{\varepsilon}), \quad (9)$$

$$Z^{(2)}(\hat{\varepsilon}) = Z^{(1)}(w(\hat{\varepsilon})). \quad (10)$$

As shown in [17],  $Z^{(12)}$  can be summed up and then yields the RMT generating function of the orthogonal ensemble  $Z_{\text{GOE}}$ .

#### IV. SYSTEMS WITH SPIN 1/2. DUALITY WITH SPINLESS CASE

The spin-1/2 evolution must be treated quantum mechanically while the orbital motion still allows semiclassical description. We assume that the spin is driven by the interaction with the translational motion while neglecting the back reaction of the spin [18]. The van Vleck propagator of the two-component wave function then falls into a product of the semiclassical propagator of a spinless particle, and a  $2 \times 2$  matrix of the spin evolution. The Gutzwiller formula follows from the van Vleck propagator after going to the energy representation and taking a trace; consequently contribution of a periodic orbit  $\gamma$  in the Gutzwiller expansions is now to be multiplied by  $\text{Tr } U_\gamma$  where  $U_\gamma$  is an  $SU_2$  matrix describing the change of the spin state after a single traversal of  $\gamma$  [19].

The contribution of a quadruplet  $(AC)(BD)$  in (6) is a product of Gutzwiller amplitudes of all the orbits constituting the quadruplet. Therefore, in the presence of spin it has to be multiplied by the product of traces of the spin evolution matrices for all its orbits,  $\Xi_{AC,BD} \equiv \prod_{\gamma \in (AC)} \text{Tr } U_\gamma \prod_{\gamma' \in (BD)} \text{Tr } U_{\gamma'}$ . We recall that the orbits in  $(AC)$  and  $(BD)$  are constructed of the same  $L$  pieces. Therefore due to the group property of the propagator,  $U_\gamma, U_{\gamma'}$  can be replaced by products of the matrices  $\mathcal{D}_i$  describing the spin evolution after traversal of the  $i$ -th piece. Each  $\mathcal{D}_i$ ,  $i = 1, \dots, L$ , occurs once in  $(AC)$ , once in  $(BD)$ ; if the sense of traversal of the  $i$ -th piece is reversed in the partner, the second entry would be  $\mathcal{D}_i^{-1}$ .

The factor  $\Xi_{AC,BD}$  has to be averaged over an interval of the reference energy  $E$ . Interaction with the chaotic translational motion makes the spin evolution ergodic [18]. Assuming that  $\mathcal{D}_i$  associated with different non-overlapping pieces are independent quasi-random  $SU_2$  matrices, we can thus replace averaging by the integration over all  $\mathcal{D}_i$  over the group  $SU_2$ . The result for the partnership of just two orbits (the only one of interest at the sub-Heisenberg times) is well-known [20, 21],

$$\langle \text{Tr } U_\gamma \text{Tr } U_{\gamma'} \rangle = \frac{(-1)^{L-V}}{2^{L-V}}$$




FIG. 1: An orbit separates into two orbits after reconnection in a 2-encounter

where  $V, L$  are the number of encounters and encounter stretches in the orbit pair. We generalize it to the pseudo-orbit quadruplets containing an arbitrary number of orbits. The most important new element is however that the orbit number can change after reconnection in the encounters; see the elementary example Fig. 1 where two pieces of the figure-8 orbit become two separate orbits after reconnection. Calculations in Appendix VIII B show that

$$\langle \Xi_{AC, BD} \rangle = \frac{(-1)^{L-V+\nu_B+\nu_D-\nu_A-\nu_C}}{2^{L-V}} \quad (11)$$

where  $V, L$  are now the number of encounters and the encounter stretches in the quadruplet  $(A, C), (B, D)$ . The additional sign factor  $(-1)^{\nu_B+\nu_D-\nu_A-\nu_C}$  reflects the difference in the number of orbits after the reconnection  $(A, C) \rightarrow (B, D)$ .

The factor (11) leads to important consequences. Namely, inserting it into the semiclassical expansion (6) of the generating function and combining with  $(-1)^{\nu_C+\nu_D}$  we obtain  $(-1)^{L-V+\nu_B+\nu_D}$ . The replacement  $A, C \rightarrow B, D$  in the exponent shows that the roles of the numerator and denominator in the generating function are reversed. We recall that a quadruplet with  $L - V = n$  contributes to  $Z^{(1)}$  in the order  $\varepsilon^{-n}$ ; our rescaling  $2\varepsilon \rightarrow \varepsilon$  in the symplectic case absorbs  $2^{L-V}$ . Consequently all expansion terms of the symplectic off-diagonal sum  $Z_{\text{off}}$  are obtained from their orthogonal counterparts (8) by the interchange  $A \rightleftharpoons C, B \rightleftharpoons D$  and the sign change of all arguments,

$$Z_{n,S}^{(1)}(\hat{\varepsilon}) = Z_{n,O}^{(1)}(-\varepsilon_C, -\varepsilon_D, -\varepsilon_A, -\varepsilon_B). \quad (12)$$

The same substitution connects the full periodic orbit expansions  $Z^{(1)} = Z_{\text{diag}} \left( 1 + \sum_{n=1}^{\infty} Z_n^{(1)} \right)$ ,

$$Z_S^{(1)}(\hat{\varepsilon}) = Z_O^{(1)}(-\varepsilon_C, -\varepsilon_D, -\varepsilon_A, -\varepsilon_B). \quad (13)$$

We get an important and remarkably simple relation (13) between the semiclassical generating functions of systems with or without half-integer spin. (Turning from semiclassics to RMT we note that numerous identities of that kind between the GOE- and GSE-associated functions are well-known under the name of duality relations.) In Appendix VIII C we check that a duality relation analogous to (13) does exist between the 4-determinant generating functions of GOE and GSE. A seeming contradiction arises: whereas the high-energy asymptotic expansion of  $Z_{\text{GOE}}$  is identical with its semiclassical counterpart  $Z_O^{(12)}$ , the analogous expansion of  $Z_{\text{GSE}}$  is not: It differs from  $Z_S^{(12)}(\hat{\varepsilon}) = Z_S^{(1)}(\hat{\varepsilon}) + Z_S^{(1)}(w(\hat{\varepsilon}))$  by an additional elementary summand proportional to  $e^{i(\varepsilon_A - \varepsilon_B)/2}$ , see (32). The reason is purely mathematical, and the missing component of the semiclassical generating function can be recovered by Borel summation. We prefer to demonstrate the method on the less cumbersome example of the symplectic complex correlator, see the next Section.

The averaged spin factor for spins different from  $1/2$  is given below in (30). For half-integer spins the result differs by the replacement of 2 in the denominator of (11) by  $2S + 1$ , however this is compensated by the changed mean level spacing and Heisenberg time [9]; our equation (13) remains in force. For integer spins the sign of the averaged spin factor is always positive while  $2S + 1$  in the denominator is compensated in the way just described, therefore the generating function is the same as without spin.

## V. BOREL SUMMATION; MISSING OSCILLATORY COMPONENT AS “STOKES’S SATELLITE”

Applying  $\partial^2 \varepsilon_A \varepsilon_B$  to  $Z^{(1)}(\hat{\varepsilon}) + Z^{(2)}(\hat{\varepsilon})$  and going to the limit in (5), we obtain the asymptotic expansion of the complex correlator for both symmetry classes,

$$C(\varepsilon) \sim \sum_{n \geq 2} \frac{a_n}{\varepsilon^n} + e^{i2\varepsilon} \sum_{n \geq 4} \frac{b_n}{\varepsilon^n} \quad (14)$$

where the non-oscillatory and oscillatory parts are generated by  $Z^{(1)}(\hat{\varepsilon})$  and  $Z^{(2)}(\hat{\varepsilon})$  respectively. The coefficients are easily calculated from (8) and the duality relation; for both symmetries  $a_2 = -1$ , while for  $n > 2$ ,

$$a_{n,O} = \frac{(n-3)!(n-1)}{2i^n}, \quad b_{n,O} = \frac{(n-3)!(n-3)}{2i^n},$$

$$a_{n,S} = a_{n,O}(-1)^n, \quad b_{n,S} = b_{n,O}.$$

The factorial growth of the coefficients signals that the asymptotic series diverge for all  $\varepsilon$ .

Suppose we want to restore the analytic functions behind these series by means of the Borel method [23]. The first stage would be the term-by-term Fourier transform (2) employing

$$\mathcal{F}(\varepsilon^{-n}) = \frac{\tau^{n-1}}{2i^n (n-1)!} \theta(\tau), \quad \mathcal{F}(e^{i2\varepsilon} \varepsilon^{-n}) = \frac{(\tau-2)^{n-1}}{2i^n (n-1)!} \theta(\tau-2);$$

the resulting series in  $\tau$  and  $(\tau-2)$  converge to analytic functions. On the second stage the inverse Fourier transform produces a closed expression for the complex correlator. This is easily done in the orthogonal case and leads to the form factor  $K_{GOE}(\tau)$  and then to the exact  $C_{GOE}$ , see Appendix VIII A, Eq. (22).

The symplectic case is more interesting. On the first stage we obtain the form factor as

$$K = \theta(\tau) [\tau/2 - (\tau/4) \ln(1-\tau)]$$

$$+ \theta(\tau-2) [1 - \tau/2 + (\tau/4) \ln(\tau-1)];$$

The first summand has a branch cut  $(1, +\infty)$ ; to proceed with Borel we need to continue  $\ln(1-\tau)$  to all  $\tau > 1$ . There are three obvious choices: use the logarithm values  $\ln|1-\tau| \pm i\pi$  at the lower or upper lip of the cut, or their average  $\ln|1-\tau|$ . Only the last option is admissible since the form factor must be real, in view of reality of the energy eigenvalues [1]; the factor at  $\theta(\tau)$  will then be  $K^< \equiv \tau/2 - (\tau/4) \ln|1-\tau|$ . The Fourier transform of  $K$  with this choice produces the *exact* GSE correlator (23) whose asymptotic expansion contains an additional oscillatory summand,

$$C_{GSE} \sim \sum_{n=2}^{\infty} \frac{a_{n,S}}{\varepsilon^n} + e^{2i\varepsilon} \sum_{n=2}^{\infty} \frac{b_{n,S}}{\varepsilon^n} + \frac{\pi}{2\varepsilon^2} (\varepsilon + i) e^{i\varepsilon};$$

incidentally, the latter would be generated by the term  $\propto e^{i(\varepsilon_A - \varepsilon_B)/2}$  of  $Z_{GSE}$  (32) after application of (5).

It may seem strange that the back-and-forth Fourier transform recovered, free of charge, the missing oscillatory contribution to the correlator. In fact, restoration of an oscillatory term, given an asymptotic power series, is a legitimate mathematical tool described in detail in the book [24]; see Berry and coauthors [25, 26] for further important developments. The key idea is that the manner in which coefficients of an asymptotic series tend to infinity contains information about the exponentially small terms disregarded in the classical Poincaré approach. Such terms become oscillatory and of crucial importance when the asymptotics is continued to the anti-Stokes lines.

Here is the barest minimum of detail on the method. Consider a diverging asymptotic expansion of an analytic function  $g(z)$ ,  $z = x + iy$ ,

$$g(z) \sim \sum_{n=1}^{\infty} \frac{c_n}{z^n}, \quad |z| \rightarrow \infty, \quad (15)$$

and assume that in the limit of large  $n$  its coefficients tend to

$$c_n \rightarrow (n - \beta)! \quad (16)$$




FIG. 2: The exponential “satellite” changes by a jump at the Stokes line (red) and becomes oscillatory at the anti-Stokes (AS) lines

Then

a) The real positive semi-axis is the Stokes line at which the power expansion (15) has all its terms positive and therefore maximally dominant with respect to an exponentially small additional component  $g_{SD}(z)$ ; the value of  $g_{SD}(z)$  changes almost by a jump when the positive semi-axis is crossed;

b) The subdominant component behaves like

$$g_{SD}(z) \propto \frac{e^{-z}}{z^{\beta-1}};$$

c) The imaginary semiaxes are the anti-Stokes lines where  $g_{SD}(z)$  becomes oscillatory and comparable to the power expansion, see Fig. 2;

d) Under certain assumptions about the properties of  $g(z)$  we have,

$$g(z) = \sum_{n=1}^{n^*(z,q)} \frac{c_n}{z^n} + R(z), \quad (17)$$

$$R(z) \approx e^{-z} \frac{\pi}{z^{\beta-1}} \left[ i \operatorname{erf} \left( \frac{y}{\sqrt{2x}} \right) + i\nu + \eta(z, q) \right]. \quad (18)$$

The upper limit of the sum is  $n^*(z, q) = \operatorname{Int} \{|z| + q\}$  where  $q$  is of the order unity and otherwise arbitrary. The error function  $\operatorname{erf}(\sigma) = \frac{2}{\sqrt{\pi}} \int_0^\sigma e^{-t^2} dt$  in (17) is close to 1 for  $|z|$  large and  $\arg z > \delta > 0$ , and to  $-1$  for  $\arg z < -\delta < 0$ ; on the real axis it is zero. The almost jump-like change of the subdominant component when the real axis is crossed, is the Stokes phenomenon. The small real correction

$$\eta(z, q) = \frac{2}{\sqrt{2\pi x}} \left( \operatorname{Fract} \{|z| + q\} + \beta - q - \frac{4}{3} - \frac{y^2}{6x} \right) e^{-y^2/2x}.$$

is significantly non-zero only close to the  $x$ -axis; its dependence on  $q$  compensates that of the sum in (17). The constant  $\nu$  must be deduced from additional information on the function  $g(z)$ . In particular, if  $g(z)$  is real on the real axis we must choose  $\nu = 0$ ; the asymptotics of  $g(z)$  contains then oscillatory components at both anti-Stokes lines.

Let us apply the method to the non-oscillatory part of the symplectic correlator. The terms of its expansion  $a_{n,S}/\varepsilon^n$  are all positive at the positive imaginary axis of  $\varepsilon$  which is the Stokes line where the power series is maximally dominant; the anti-Stokes lines are the real semiaxes of  $\varepsilon$ . The results above are applicable with  $\varepsilon = iz$ ,  $\beta = 2$ . According to its definition (1), the complex correlator must be real for positive imaginary  $\varepsilon$ . Therefore we must choose in (17)  $\nu = 0$  such that the oscillatory “Stokes satellite” of the power series must be present on both real semiaxes,

$$R(\varepsilon) \approx e^{i\varepsilon} \frac{\pi}{2\varepsilon}, \quad \varepsilon \rightarrow +\infty, \quad (19)$$

$$R(\varepsilon) \approx -e^{i\varepsilon} \frac{\pi}{2\varepsilon} = e^{i\varepsilon} \frac{\pi}{2|\varepsilon|}, \quad \varepsilon \rightarrow -\infty. \quad (20)$$

This is indeed the leading term in the oscillatory part of the symplectic correlator recovered by the Borel method. It is subdominant in the upper half-plane away from the real axis and experiences the erf –like approximate discontinuity at the positive imaginary axis.

It is instructive to investigate what happens if we choose to continue  $\log(1 - \tau)$  as  $\log(\tau - 1) \pm i\pi$  for  $\tau > 1$  in the form factor at the first stage of the Borel summation. The functions obtained by the inverse Fourier transform would then differ from the correct complex correlator by the additional terms

$$\mp \frac{i\pi}{2} \int_1^\infty d\tau e^{i\varepsilon\tau} \tau = \pm \frac{\pi}{2\varepsilon^2} (i + \varepsilon) e^{i\varepsilon} \sim \pm e^{i\varepsilon} \frac{\pi}{2\varepsilon}$$

They would cancel (19) at one of the real semiaxes and double its amplitude at the other one, i.e., exactly what we would get if we chose  $\nu = \pm 1$  in (17). Therefore the alternative choices of  $\nu$  are equivalent to different continuation of the result of the first Borel stage beyond the branch point in the  $\tau$  domain.

Finally let us convince ourselves that additional oscillatory components do not arise in the orthogonal case. The  $\varepsilon^{-1}$ –expansion with the coefficients  $a_{n,0}$  has its terms all positive on the negative imaginary semi-axis, i.e. in the non-physical half-plane of  $\varepsilon$  where (1) is inapplicable and the correlator need not be real. The parameter  $\beta = 2$  is the same as in the symplectic case but we must now set  $\varepsilon = -iz$  and the Stokes satellite now behaves like  $\sim e^{-i\varepsilon}/\varepsilon$ . Let us change the phase of  $\varepsilon$  from  $-\pi/2$  via 0 to positive values; if the satellite were present with a non-zero amplitude it would become exponentially large in the physical region  $\text{Im } \varepsilon > 0$ . This is forbidden, and we must choose thus in (17)  $\nu = -1$  which corresponds to absence of the exponential term in the sector  $-\pi/2 + \delta < \arg \varepsilon < \pi$ , in particular at the real positive semi-axis.

## VI. CONCLUSION

We studied the generating function, complex correlator and form factor of systems with spin 1/2. Expanding the generating function into a sum over periodic orbit quadruplets we showed that in the presence of spin, the terms of the expansion acquire an additional sign factor whose effect is to interchange the role of the numerator and the denominator of the generating function. As a result, the generating functions of the orthogonal and symplectic class turn out to be connected by a simple substitution of their arguments.

The periodic orbit expansion of the generating function supplemented by the Riemann-Siegel look-alike formula for the spectral determinants yields the complex correlator as combination of two asymptotic series in  $\varepsilon^{-1}$ , the second one multiplied by  $e^{i2\varepsilon}$ ; they are responsible for the form factor at small times and at times larger than the Heisenberg time. We demonstrate how the Borel summation reveals in the symplectic case one more oscillatory term  $\propto e^{i\varepsilon}$  associated with the logarithmic singularity of the form factor; the origin of that term is clarified by the Dingle-Berry method of smart summation of the asymptotic series as a display of the Stokes phenomenon. With the missing oscillatory term restored, complete equivalence of the correlation functions of chaotic systems with spin 1/2 and the Gaussian symplectic ensemble of RMT is reached.

The inherent ambiguity in restoration of a function from its asymptotic series is solved on the ground of reality of the energy eigenvalues. The same reason is at the heart of the Riemann-Siegel look-alike, such that existence of both oscillatory components of the complex correlator in the symplectic case can be traced to unitarity of the quantum mechanical evolution.

There are several possible further developments of the theory. An obvious generalization would be to consider parametric correlation in systems with spin 1/2 at times comparable with the Heisenberg time [27]. Away from the deep semiclassical limit, system-specific deviations from the universal behavior in systems with half-integer spin can be of physical interest.

## VII. ACKNOWLEDGEMENT

I want to thank Fritz Haake for continuous help and support in the years of my work in Essen and Duisburg; he read the paper before publication and made many important suggestions. The author is grateful to Martin Zirnbauer for providing his unpublished results on the generating functions of RMT, and Stefan Heusler, Sebastian Mueller, Thomas Guhr and Yan Fedorov for useful discussions. Financing by Sonderforschungsbereich TR12 is acknowledged.

## VIII. APPENDIX

### A. RMT complex correlator of the orthogonal and symplectic case

The complex correlators are conveniently expressed in terms of the functions,

$$f_{\pm}(z) = \int_z^{\infty} \frac{e^{\pm i(t-z)}}{t} dt;$$

The integral representations are applicable whenever the integral converges; for all  $z$  we have

$$f_{\pm}(z) \equiv e^{\mp iz} \left[ \pm i \frac{\pi}{2} - \text{Ci}(z) \mp i \text{Si}(z) \right]. \quad (21)$$

where Ci, Si are the integral sine and cosine. These functions are analytic in the plane of  $z$  with the cut along the negative real axis.

The complex correlator of the orthogonal case can be represented in terms of  $f_{\pm}$  as,

$$\begin{aligned} C_{\text{GOE}}(\varepsilon) &= 2 \int_0^{\infty} e^{i2\varepsilon\tau} K_{\text{GOE}}(\tau) d\tau = C_{\text{GOE}}^{(1)} + C_{\text{GOE}}^{(2)}, \\ C_{\text{GOE}}^{(1)} &= -\frac{1}{2\varepsilon^2} + \frac{1}{2\varepsilon^2} (i\varepsilon + 1) f_+(\varepsilon), \\ C_{\text{GOE}}^{(2)} &= -\frac{e^{i2\varepsilon}}{2\varepsilon^2} [(-i\varepsilon + 1) f_+(\varepsilon) - 1], \end{aligned} \quad (22)$$

while in the symplectic case

$$\begin{aligned} C_{\text{GSE}}(\varepsilon) &= 2 \int_0^{\infty} e^{i\varepsilon\tau} K_{\text{GSE}}(\tau) d\tau = C_{\text{GSE}}^{(1)} + C_{\text{GSE}}^{(2)} + C_{\text{GSE}}^{(3)}, \\ C_{\text{GSE}}^{(1)} &= -\frac{1}{2\varepsilon^2} + \frac{1}{2\varepsilon^2} (-i\varepsilon + 1) f_-(\varepsilon), \\ C_{\text{GSE}}^{(2)} &= -\frac{e^{2i\varepsilon}}{2\varepsilon^2} [(-i\varepsilon + 1) f_+(\varepsilon) - 1], \\ C_{\text{GSE}}^{(3)} &= \frac{\pi}{2\varepsilon^2} (\varepsilon + i) e^{i\varepsilon}. \end{aligned} \quad (23)$$

The components  $C^{(1)}$  in both cases have non-oscillatory asymptotic expansion in powers of  $\varepsilon^{-1}$  at the real positive axis while  $C^{(2)}$  oscillates like  $e^{i2\varepsilon}$ . That follows from the asymptotic representations of  $f_{\pm}(z)$ ,

$$f_{\pm}(z) \sim -\sum_{k=0}^{\infty} \frac{k!}{(\pm iz)^{k+1}} \equiv \sigma_{\pm}(z), \quad |z| \rightarrow \infty, \quad (24)$$

$$\begin{aligned} -\pi + \delta < \arg z \leq \pi - 0, & \quad (f_+); \\ -\pi + 0 \leq \arg z < \pi - \delta. & \quad (f_-). \end{aligned}$$

The Stokes line where the power expansion is dominant is  $z = -it$ ,  $t > 0$ , for  $f_+(z)$  and  $z = it$ ,  $t > 0$ , for  $f_-(z)$ ; the subdominant satellite,  $-2\pi i e^{-iz}$  for  $\sigma_+(z)$  and  $2\pi i e^{iz}$  for  $\sigma_-(z)$ , exists in the quadrant to the left of the respective Stokes line, see Fig. 3.

The form factors obtained from the complex correlators by the transformation (2) are

- GOE:

$$\begin{aligned} K(\tau) &= 2\tau - \tau \ln(1 + 2\tau), \quad \tau < 1; \\ K(\tau) &= 2 - \tau \ln \frac{2\tau + 1}{2\tau - 1}, \quad \tau > 1. \end{aligned}$$

- GSE:

$$\begin{aligned} K(\tau) &= \frac{\tau}{2} - \frac{\tau}{4} \ln |1 - \tau|, \quad \tau < 2; \\ K(\tau) &= 1, \quad \tau > 2. \end{aligned}$$






FIG. 3: Asymptotics of the functions  $f_+(\epsilon)$  (a) and  $f_-(\epsilon)$  (b): exponential term is present left of the Stokes line (red)

### B. Spin factor for quadruplets

Contribution of the quadruplet  $(AC)(BD)$  in the periodic orbit expansion of the generating function (4) contains in the symplectic case the spin factor  $\Xi_{AC,BD} \equiv \prod_{\gamma \in (AC)} \text{Tr} U_\gamma \prod_{\gamma' \in (BD)} \text{Tr} U_{\gamma'}$  which is product of traces of the periodic orbits composing the quadruplet. Non-vanishing contributions to the sum are created by the quadruplets such that the orbits of  $(BD)$  are constructed from pieces of the orbits in  $(AC)$  connected in different order and possibly traversed with a different sense. We assume ergodicity of the spin motion and independence of the spin evolution along different orbit pieces. Averaging is then done step by step by integration over the evolution matrices  $\mathcal{D}_l$  associated with the orbit pieces; the integration domain is the group  $SU_2$ . Possible outcomes of a single integration are summed up by the relations [21],

$$\int d\mathcal{D} \text{Tr} (\mathcal{A}\mathcal{D}\mathcal{B}\mathcal{D}) = -\frac{1}{2} \text{Tr} (\mathcal{A}\mathcal{B}^{-1}), \quad (25)$$

$$\int d\mathcal{D} \text{Tr} (\mathcal{A}\mathcal{D}\mathcal{B}\mathcal{D}^{-1}) = \frac{1}{2} \text{Tr} \mathcal{A} \text{Tr} \mathcal{B}, \quad (26)$$

$$\int d\mathcal{D} \text{Tr} (\mathcal{A}\mathcal{D}) \text{Tr} (\mathcal{B}\mathcal{D}) = \frac{1}{2} \text{Tr} (\mathcal{A}\mathcal{B}^{-1}). \quad (27)$$

Here  $\mathcal{A}, \mathcal{B}$  are any fixed  $SU_2$  matrices. Applying these rules, e. g., to the structure with  $V = 1, L = 2$  shown in Fig. 1, we obtain the result,

$$\langle \Xi_{AC,BD} \rangle = \int d\mathcal{D}_1 d\mathcal{D}_2 \text{Tr} (\mathcal{D}_1 \mathcal{D}_2) \text{Tr} (\mathcal{D}_1) \text{Tr} (\mathcal{D}_2) = \frac{1}{2},$$

which is opposite in sign compared with the Sieber-Richter pair [20].

We shall find the average of  $\Xi_{AC,BD}$  for all structures extending the inductive method of Bolte and Harrison from pairs of orbits to the pseudo-orbits quadruplets. We remind the essence of the method. Consider an orbit pair  $\gamma, \gamma'$  differing in  $V$  encounters with  $L$  encounter stretches and assume that averaging of the spin factor produces the factor  $C_{\gamma\gamma'} = (-1)^{L-V} / 2^{L-V}$ . Introduce one more orbit  $\gamma''$  differing from  $\gamma'$  by an additional 2-encounter such that the pair  $\gamma, \gamma''$  contains  $V'' = V + 1$  active encounters with  $L'' = L + 2$  stretches. It is then shown using the recurrence relations that  $C_{\gamma\gamma''} = -C_{\gamma\gamma'} / 2 = (-1)^{L''-V''} / 2^{L''-V''}$ . Starting from a pair without active encounters  $L = V = 0$  when the formula is true, and adding 2-encounters one by one we obtain that the result is true for an arbitrary number of 2-encounters. Finally, reconnection in any  $l$ -encounter with  $l > 2$  can be reduced to  $l - 1$  successive reconnections in 2-encounters in  $l - 1$  steps, with the factor  $(-1)^{L-V} / 2^{L-V}$  correct on all steps. Indeed, reconnection

in an  $l$ -encounter can be described by a permutation of  $l$  elements, however, any permutation can be represented as a chain of transpositions of just two elements.

Let us apply this method to the pseudo-orbit quadruplets. Let  $\Gamma = (A, C)$  be the initial pseudo-orbit pair, and  $\Gamma' = (B, D)$  its partner differing from  $\Gamma$  in  $V$  active encounters with  $L$  stretches. Let  $\Gamma''$  be a pseudo-orbit pair differing from  $\Gamma'$  by reconnection in a single additional 2-encounter whose stretches belong to some orbit  $\gamma$  in  $\Gamma$  and to  $\gamma'$  in  $\Gamma'$  where  $\gamma, \gamma'$  may differ in an arbitrary number of other encounters. The only new case to be considered is that of a 2-encounter with almost parallel stretches such that  $\gamma'$  breaks up after reconnection into two orbits  $\gamma_p''$  and  $\gamma_q''$  belonging to  $\Gamma''$ , see the example in Fig. 1.

The averaged spin factors for the quadruplets  $\Gamma\Gamma'$  and  $\Gamma\Gamma''$  can be written as multiple integrals over  $SU_2$  with the Haar measure,

$$C_{\Gamma\Gamma'} = \int d(\dots) \int d(\gamma) d(\gamma'),$$

$$C_{\Gamma\Gamma''} = \int d(\dots) \int d(\gamma) d(\gamma_p'') d(\gamma_q'')$$

Here  $d(\gamma), d(\gamma')$  denote  $SU_2$  integration over matrices associated with pieces of  $\gamma$  and  $\gamma'$  while  $d(\gamma_p'') d(\gamma_q'')$  indicate integration over matrices associated with pieces of the disconnected orbit pair in  $\Gamma''$ . Integration  $d(\dots)$  is over the remaining variables, same for  $\Gamma\Gamma'$  and  $\Gamma\Gamma''$ . In the way of induction, let us assume that

$$C_{\Gamma\Gamma'} = \frac{(-1)^{L-V+\nu_\Gamma-\nu_{\Gamma'}}}{2^{L-V}} \quad (28)$$

where  $\nu_\Gamma = \nu_A + \nu_C$  and  $\nu_{\Gamma'} = \nu_B + \nu_D$  is the number of periodic orbits in  $\Gamma$  and  $\Gamma'$ , respectively, and prove that the analogous formula will be true for  $C_{\Gamma\Gamma''}$ .

Consider Fig. 4 where the orbits  $\gamma, \gamma'$  and the orbit pair  $\gamma''$  are depicted. In  $\gamma, \gamma'$  we see a parallel crossing; however it is not switched between  $\gamma, \gamma'$ , i.e., it is inactive and not counted in  $V = V_{\Gamma\Gamma'}$ . On the other hand, it is activated in the pair  $\gamma\gamma''$  such that

$$V_{\Gamma\Gamma''} = V + 1, \quad L_{\Gamma\Gamma''} = L + 2, \quad \nu_{\Gamma''} = \nu_{\Gamma'} + 1.$$

In Fig. 4 the orbit pieces adjacent to the encounter and incorporating parts of its stretches are denoted  $a, b, c, d$ ; they are assumed to coincide in  $\gamma, \gamma'$ , i.e., do not contain any additional active encounters. The associated matrices  $\mathcal{D}_a$  etc. involved in the  $SU_2$  integration will be denoted by the same letters for compactness,  $a \equiv \mathcal{D}_a$  etc. (we hope that  $d$  as the integration variable will not be mixed with  $d$  as the differential!) The two links attached to the selected 2-encounter are denoted  $L_1, L_2$  in  $\gamma$  and  $L'_1, L'_2$  in  $\gamma'$ . Unlike  $a, b, c, d$ , the links  $L_i$  and  $L'_i$  need not coincide; indeed,  $L_i$  can contain any amount of encounters active in  $\Gamma\Gamma'$  such that  $L'_i$  can contain pieces of all orbits of the original quasi-orbit pair  $\Gamma$  other than  $\gamma$ .

The  $SU_2$  integrals can be written, with  $dL_1$  being a shorthand for  $d\mathcal{D}_{L_1}$ , etc.,

$$C_{\Gamma\Gamma'} = \int d(\dots) dL_1 dL_2 dL'_1 dL'_2 \int da db dc dd \operatorname{Tr}(L_1 ab L_2 cd) \operatorname{Tr}(L'_1 ab L'_2 cd);$$

$$C_{\Gamma\Gamma''} = \int d(\dots) dL_1 dL_2 dL'_1 dL'_2 \int da db dc dd \operatorname{Tr}(L_1 ab L_2 cd) \operatorname{Tr}(L'_1 ad) \operatorname{Tr}(L'_2 cb).$$

Let us transform the integrals over  $a, b, c, d$ . In  $C_{\Gamma\Gamma'}$  we can take the matrices  $x = ab$  and  $y = cd$  as the new integration variables writing

$$\begin{aligned} & \int da db dc dd \operatorname{Tr}(L_1 ab L_2 cd) \operatorname{Tr}(L'_1 ab L'_2 cd) \\ &= \int dx dy \operatorname{Tr}(L_1 x L_2 y) \operatorname{Tr}(L'_1 x L'_2 y) \\ &= \frac{1}{2} \int dy \operatorname{Tr} \left[ L_2 y L_1 \left( L'_1 \right)^{-1} y^{-1} \left( L'_2 \right)^{-1} \right] \\ &= \frac{1}{4} \operatorname{Tr} \left[ L_1 \left( L'_1 \right)^{-1} \right] \operatorname{Tr} \left[ L_2 \left( L'_2 \right)^{-1} \right]; \end{aligned} \quad (29)$$

The cyclic invariance of the trace and the relations (27) and (26) were used.




FIG. 4: A 2-encounter belongs to the orbits  $\gamma$  in the pseudo-orbit  $\Gamma$  and  $\gamma'$  in the partner pseudo-orbit  $\Gamma'$ . Reconnection in that 2-encounter divides  $\gamma'$  into  $\gamma''_1, \gamma''_2$  creating a new pseudo-orbit  $\Gamma''$

Now let us carry out similar transformations of the integral in  $C_{\Gamma''}$  introducing consecutively new integration

variables  $x = ad$ ,  $y = cb$ ,  $z = cd$ ,

$$\begin{aligned}
& \int da db dc dd \operatorname{Tr} (L_1 ab L_2 cd) \operatorname{Tr} (L'_1 ad) \operatorname{Tr} (L'_2 cb) \\
&= \int dx db dc dd \operatorname{Tr} (L_1 x d^{-1} b L_2 cd) \operatorname{Tr} (L'_1 x) \operatorname{Tr} (L'_2 cb) \\
&= \frac{1}{2} \int db dc dd \operatorname{Tr} \left[ d^{-1} b L_2 cd L_1 (L'_1)^{-1} \right] \operatorname{Tr} (L'_2 cb) \\
&= \frac{1}{2} \int dy dc dd \operatorname{Tr} \left[ d^{-1} c^{-1} y L_2 cd L_1 (L'_1)^{-1} \right] \operatorname{Tr} (L'_2 y) \\
&= \frac{1}{4} \int dc dd \operatorname{Tr} \left[ L_2 c d L_1 (L'_1)^{-1} d^{-1} c^{-1} (L'_2)^{-1} \right] \\
&= \frac{1}{4} \int dz \operatorname{Tr} \left[ L_1 (L'_1)^{-1} z (L'_2)^{-1} L_2 z^{-1} \right] \\
&= \frac{1}{8} \operatorname{Tr} \left[ L_1 (L'_1)^{-1} \right] \operatorname{Tr} \left[ L_2 (L'_2)^{-1} \right].
\end{aligned}$$

Comparing with (29) we see that,

$$C_{\Gamma\Gamma''} = \frac{1}{2} C_{\Gamma\Gamma'}$$

which agrees with (28) with  $L \rightarrow L'' = L + 2$ ,  $V \rightarrow V'' = V + 1$ ,  $\nu_{\Gamma'} \rightarrow \nu_{\Gamma''} = \nu_{\Gamma'} + 1$ ; therefore if (28) is true for the quadruplet  $\Gamma\Gamma'$  it will also hold for  $\Gamma\Gamma''$ .

Evidently, the reversed process when two orbits of  $\Gamma'$  merge into a single orbit in  $\Gamma''$  after reconnection in a 2–encounter, also agrees with (28); reconnection in an  $l$ –encounter with  $l > 2$  is reducible to a sequence of reconnections in 2–encounters. By repeated activation of the encounters resulting in joining and disjoining the orbits, we can construct any pseudo-orbit quadruplet out of an orbit pair for which (28) is known to be correct; hence by induction it is true for all quadruplets.

For the spin different from  $1/2$ , the Bolte-Harrison recurrence relations (25),(26),(27) differ by the replacement of 2 in the denominator by  $2S + 1$ ; for integer spins the sign in (25) is plus. Otherwise the reasoning remains unchanged with the result (28) replaced by

$$\begin{aligned}
C_{\Gamma\Gamma'} &= \frac{(-1)^{L-V+\nu_{\Gamma}-\nu_{\Gamma'}}}{(2S+1)^{L-V}}, \quad \text{half-integer } S; \\
C_{\Gamma\Gamma'} &= \frac{1}{(2S+1)^{L-V}}, \quad \text{integer } S.
\end{aligned} \tag{30}$$

### C. Generating functions of RMT. GOE-GSE duality

The generating function of GOE found by Zirnbauer [28] has a Weyl symmetric form,

$$Z_{\text{GOE}}(\hat{\varepsilon}) = F_{\text{GOE}}(\hat{\varepsilon}) + F_{\text{GOE}}(w(\hat{\varepsilon}))$$

where  $\hat{\varepsilon} = (\varepsilon_A \varepsilon_B \varepsilon_C \varepsilon_D)$ ;  $w$  interchanges  $C$  and  $D$ . Denoting  $\overline{ad} = \varepsilon_A - \varepsilon_D$  etc and assuming  $\operatorname{Im} \overline{ab} > 0$  we can write,

$$F_{\text{GOE}}(\hat{\varepsilon}) = e^{i\frac{1}{2}(\overline{ab}-\overline{cd})} \frac{\overline{ad}}{ab} \frac{\overline{cb}}{cd} \left[ 1 + \frac{1}{2} \frac{\overline{ac}}{ac} \frac{\overline{bd}}{bd} \frac{2+i\overline{cd}}{cd^2} f_+ \left( \frac{\overline{ab}}{2} \right) \right]$$

with  $f_+$  defined above, see (21).

The generating function of GSE is given in [28] as,

$$\begin{aligned} Z_{\text{GSE}}(\varepsilon) &= Z_{\text{GUE}}(\varepsilon) + \frac{\overline{ac} \overline{ad} \overline{bc} \overline{bd}}{ab \overline{cd}} \frac{1}{4} G_1\left(\frac{\overline{ab}}{2}\right) G_0\left(\frac{\overline{cd}}{2}\right), \\ G_1(z) &= i \int_1^\infty e^{izq} q dq = -\frac{e^{iz}}{z^2} (i+z), \\ G_0(x) &= \int_{-1}^1 \frac{\sin qx}{q} dq = \pi + \int_1^\infty \frac{e^{-iqx}}{iq} dq - \int_1^\infty \frac{e^{iqx}}{iq} dq \\ &= (\pi - ie^{-ix} f_+^*(x)) + ie^{ix} f_+(x) = G_0^{(1)} + G_0^{(2)}. \end{aligned}$$

Here  $Z_{\text{GUE}}(\varepsilon)$  is the generating function of the unitary ensemble; in the last line  $x = \overline{cd}/2$  is assumed real positive.

The GSE generating function can be transformed to a sum of two components connected by the Weyl substitution  $w$ . First let us write  $Z_{\text{GSE}}(\varepsilon) = F^{(1)} + F^{(2)}$  with

$$\begin{aligned} F^{(1)} &= e^{i\frac{(\overline{ab}-\overline{cd})}{2}} \frac{\overline{ad} \overline{cb}}{ab \overline{cd}} \left[ 1 + \frac{1}{2} \frac{\overline{ac} \overline{bd}}{ab^2} \frac{(2-i\overline{ab})}{ab^2} f_+^*\left(\frac{\overline{cd}}{2}\right) \right] \\ &\quad + e^{i\frac{\overline{ab}}{2}} \frac{\overline{ad} \overline{cb} \overline{ac} \overline{bd}}{2ab \overline{cd}} \frac{(2-i\overline{ab})}{ab^2} i\pi \end{aligned}$$

and

$$F^{(2)} = e^{i\frac{(\overline{ab}+\overline{cd})}{2}} \frac{\overline{ac} \overline{db}}{ab \overline{dc}} \left[ 1 + \frac{1}{2} \frac{\overline{ad} \overline{bc}}{ab^2} \frac{(2-i\overline{ab})}{ab^2} f_+\left(\frac{\overline{cd}}{2}\right) \right].$$

The part  $F^{(1)}$  proportional to  $e^{i\frac{(\overline{ab}-\overline{cd})}{2}}$  generates the non-oscillatory component of the symplectic correlator while the part proportional to  $e^{i\frac{\overline{ab}}{2}}$  generates the correlator term proportional to  $\propto e^{i\varepsilon}$ . The part  $F^{(2)} \propto e^{i\frac{(\overline{ab}+\overline{cd})}{2}}$  is responsible for the component of the correlator  $\propto e^{i2\varepsilon}$ .

Let us define the function  $f_+(-x)$  for real positive  $x$ , i.e., at the branch cut, as the average of the values of the analytic function  $f_+(z)$  at the lips of the cut,

$$f_+(-x) = \frac{f_+(-x+i0) + f_+(-x-i0)}{2}, \quad x > 0.$$

Taking into account that

$$\begin{aligned} f_+(-x+i0) &= f_+^*(x), \\ f_+(-x+i0) - f_+(-x-i0) &= -2\pi i e^{ix}, \end{aligned}$$

we have  $G_0(x) = -if_+(-x)e^{-ix} + if_+(x)e^{ix}$ : we have concealed  $\pi$  in the first summand. Now considering that the substitution  $w$  changes the sign of  $\overline{cd}$  we obtain

$$Z_{\text{GSE}}(\hat{\varepsilon}) = F_{\text{GSE}}(\hat{\varepsilon}) + F_{\text{GSE}}(w(\hat{\varepsilon})). \quad (31)$$

with

$$F_{\text{GSE}}(\hat{\varepsilon}) = F^{(1)} = e^{i\frac{(\overline{ab}-\overline{cd})}{2}} \frac{\overline{ad} \overline{cb}}{ab \overline{cd}} \left[ 1 + \frac{1}{2} \frac{\overline{ac} \overline{bd}}{ab^2} \frac{2-i\overline{ab}}{ab^2} f_+\left(-\frac{\overline{cd}}{2}\right) \right].$$

The duality relation now holds,

$$\begin{aligned} F_{\text{GSE}}(\varepsilon_A, \varepsilon_B, \varepsilon_C, \varepsilon_D) &= F_{\text{GOE}}(-\varepsilon_C, -\varepsilon_D, -\varepsilon_A, -\varepsilon_B), \\ \text{Re}(\varepsilon_A - \varepsilon_B) &> 0, \quad \text{Im}(\varepsilon_A - \varepsilon_B) = +i0, \\ \varepsilon_C - \varepsilon_D &> 0. \end{aligned}$$

It is instructive to compare the RMT function (31) with the semiclassical results. Taking (8) into account and using the semiclassical duality, we have the following equivalence in the high-energy limit,

$$F_{\text{GSE}}(\hat{\varepsilon}) \sim Z_S^{(1)}(\hat{\varepsilon}) + e^{i(\varepsilon_A - \varepsilon_B)/2} \frac{\overline{ad} \overline{cb} \overline{ac} \overline{bd}}{2\overline{ab} \overline{cd}} \frac{(2i + \overline{ab})}{\overline{ab}^2} \pi, \quad (32)$$

$$F_{\text{GSE}}(w(\hat{\varepsilon})) \sim Z_S^{(2)}(\hat{\varepsilon}) \equiv Z_S^{(1)}(w(\hat{\varepsilon})).$$

We stress that the asymptotics of  $F_{\text{GSE}}(w(\hat{\varepsilon}))$  is not simply the Weyl-transposed asymptotics of  $F_{\text{GSE}}(\hat{\varepsilon})$ . The reason is the Stokes phenomenon; the Weyl operation changes the sign of  $\overline{cd}$ , and whereas the asymptotics of  $f_+(\overline{cd}/2)$  is purely power-like,  $f_+(-\overline{cd}/2)$  contains an additional exponential summand, in accordance with

$$f_+(x) \sim \sigma_+(x), \quad f_+(-x) \sim i\pi e^{ix} + \sigma_+(-x),$$

$$x \rightarrow +\infty.$$

where  $\sigma_+$  is the asymptotic power series defined in (24).

- 
- [1] F. Haake. *Quantum Signatures of Chaos*. Springer Series in Synergetics. Springer, 2010.
  - [2] O. Bohigas, M. J. Giannoni, and C. Schmit. Characterization of chaotic quantum spectra and universality of level fluctuation laws. *Phys. Rev. Lett.*, 52:1–4, Jan 1984.
  - [3] Steven W. McDonald and Allan N. Kaufman. Spectrum and eigenfunctions for a Hamiltonian with stochastic trajectories. *Phys. Rev. Lett.*, 42:1189–1191, Apr 1979.
  - [4] G. Casati, I. Guarneri, and F. Valz-Gris. On the connection between quantization of nonintegrable systems and statistical theory of spectra. *Lettere al Nuovo Cimento*, 28:279, 1980.
  - [5] M. V. Berry. The bakerian lecture, 1987: Quantum chaology. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 413(1844):183–198, 1987.
  - [6] M. V. Berry. Semiclassical theory of spectral rigidity. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 400(1819):229–251, 1985.
  - [7] Martin Sieber and Klaus Richter. Correlations between periodic orbits and their rôle in spectral statistics. *Physica Scripta*, 2001(T90):128, 2001.
  - [8] Sebastian Müller, Stefan Heusler, Petr Braun, Fritz Haake, and Alexander Altland. Semiclassical foundation of universality in quantum chaos. *Phys. Rev. Lett.*, 93:014103, Jul 2004.
  - [9] Sebastian Müller, Stefan Heusler, Petr Braun, Fritz Haake, and Alexander Altland. Periodic-orbit theory of universality in quantum chaos. *Phys. Rev. E*, 72(4):046207, Oct 2005.
  - [10] E. B. Bogomolny and J. P. Keating. Gutzwiller’s trace formula and spectral statistics: Beyond the diagonal approximation. *Phys. Rev. Lett.*, 77(8):1472–1475, Aug 1996.
  - [11] Stefan Keppeler. Two-point correlations of the Gaussian symplectic ensemble from periodic orbits. *Journal of Physics A: Mathematical and General*, 33(50):L503, 2000.
  - [12] M V Berry and J P Keating. A rule for quantizing chaos? *Journal of Physics A: Mathematical and General*, 23(21):4839, 1990.
  - [13] J. P. Keating. Periodic orbit resummation and the quantization of chaos. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, 436(1896):99–108, 1992.
  - [14] M. V. Berry and J. P. Keating. A new asymptotic representation for  $\zeta(\frac{1}{2}+it)$  and quantum spectral determinants. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, 437(1899):151–173, 1992.
  - [15] Jonathan P Keating and Sebastian Müller. Resummation and the semiclassical theory of spectral statistics. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Science*, 463(2088):3241–3250, 2007.
  - [16] Stefan Heusler, Sebastian Müller, Alexander Altland, Petr Braun, and Fritz Haake. Periodic-orbit theory of level correlations. *Phys. Rev. Lett.*, 98(4):044103, Jan 2007.
  - [17] Sebastian Müller, Stefan Heusler, Alexander Altland, Petr Braun, and Fritz Haake. Periodic-orbit theory of universal level correlations in quantum chaos. *New Journal of Physics*, 11(10):103025, 2009.
  - [18] S. Keppeler. *Spinning Particles - Semiclassics and Spectral Statistics*. Springer Tracts in Modern Physics. Springer Berlin Heidelberg, 2010.
  - [19] Jens Bolte and Stefan Keppeler. Semiclassical form factor for chaotic systems with spin 1/2. *Journal of Physics A: Mathematical and General*, 32(50):8863, 1999.
  - [20] Stefan Heusler. The semiclassical origin of the logarithmic singularity in the symplectic form factor. *Journal of Physics A: Mathematical and General*, 34(34):L483, 2001.
  - [21] Jens Bolte and Jonathan Harrison. The spin contribution to the form factor of quantum graphs. *Journal of Physics A: Mathematical and General*, 36(27):L433, 2003.

- [22] Alexander Altland, Petr Braun, Fritz Haake, Stefan Heusler, Gerhard Knieper, and Sebastian Mueller. Near action-degenerate periodic-orbit bunches: A skeleton of chaos. 2009. arXiv:0906.4930v1 [nlin.CD].
- [23] W. Balser. *From divergent series to analytic functions*. Springer, 2010.
- [24] R. B. Dingle. *Asymptotic Expansions: Their Derivation and Interpretation*. Academic Press, New York and London, 1973.
- [25] M. V. Berry. Uniform asymptotic smoothing of Stokes's discontinuities. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 422(1862):7–21, 1989.
- [26] M. V. Berry and C. J. Howls. High orders of the Weyl expansion for quantum billiards: Resurgence of periodic orbits, and the Stokes phenomenon. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, 447(1931):527–555, 1994.
- [27] Taro Nagao and Keiji Saito. Semiclassical approach to parametric spectral correlation with spin 1/2. *Journal of Physics A: Mathematical and Theoretical*, 40(40):12055, 2007.
- [28] M Zirnbauer. Generating functions for GUE, GOE, GSE and crossover cases. Private communication, 2008.