S-ADIC VERSION OF MINKOWSKI'S GEOMETRY OF NUMBERS AND MAHLER'S COMPACTNESS CRITERION

DMITRY KLEINBOCK, RONGGANG SHI, AND GEORGE TOMANOV

ABSTRACT. In this note we give a detailed proof of certain results on geometry of numbers in the S-adic case. These results are well-known to experts, so the aim here is to provide a convenient reference for the people who need to use them.

1. Introduction

The space of unimodular lattices in \mathbb{R}^n $(n \geq 2)$ can be identified with the homogeneous space $X = \operatorname{SL}_n(\mathbb{Z}) \setminus \operatorname{SL}_n(\mathbb{R})$ via the correspondence $\mathbb{Z}^n g \leftrightarrow \operatorname{SL}_n(\mathbb{Z}) g$ where $g \in \operatorname{SL}_n(\mathbb{R})$. It is proved by Mahler [11] that a subset R of X is relatively compact if and only if nonzero elements of the corresponding unimodular lattices are separated from zero. This phenomenon is called Mahler's compactness criterion [2, Chapter V]. It has been very useful in dynamical approach to number theory; we refer the readers to survey papers [4],[5] and [7] and references there for details.

Let S be a finite nonempty set of places of a global field K. We assume S contains all the archimedean places if K is a number field. For each place v of K, let K_v be the completion of K at v. Let $K_S = \prod_{v \in S} K_v$ and

(1.1) $I_S = \{a \in K : a \text{ is integral in } K_v \text{ for every place } v \notin S\}.$

We consider K and hence I_S as subrings of K_S via natural embeddings $K \to K_v$. Then the homogeneous space $\operatorname{SL}_n(I_S) \setminus \operatorname{SL}_n(K_S)$ can be identified with a set of free discrete I_S -modules of rank n in K_S^n with fixed covolume. The connection between dynamics and number theory also spreads to the S-adic setting, where the corresponding version of Mahler's criterion plays an important role. The extension of Mahler's criterion to the S-adic case when K is a number field has already been used in several papers, and a proof for $K = \mathbb{Q}$ can be found in [9]. Moreover, a preliminary version [8] of the paper [9], published as a preprint of MPIM (Bonn), contains a proof of the S-adic Mahler's criterion for arbitrary number field K. When K is a function field with genus zero and S contains a single place of degree one, Mahler's criterion is proved in [6]. The general S-adic case is known to

²⁰⁰⁰ Mathematics Subject Classification. Primary 11H06; Secondary 22E40.

Key words and phrases. Successive minima, Mahler's criterion, Homogeneous space.

The first-named author was supported by NSF grant DMS-1600814. The second-named author is supported by Fundamental Research Funds for the Central Universities (Grant No. 20720160006). The third named author acknowledges partial support by IMI of BAS.

experts, but it is not easy to find a convenient reference. Here we provide a self-contained proof of an S-adic version of Mahler's criterion.

Theorem 1.1. Let $n \geq 2$. A set $R \subset \mathrm{SL}_n(I_S) \backslash \mathrm{SL}_n(K_S)$ is relatively compact if and only if the subset

$$\{\xi \in I_S^n g : \xi \neq 0, g \in \operatorname{SL}_n(K_S) \text{ and } \operatorname{SL}_n(I_S)g \in R\}$$

of K_S^n is separated from zero, i.e. this set has empty intersection with some open neighborhood of zero in K_S^n .

Our proof of Theorem 1.1 is based on an S-adic version of Minkowski's lemma of geometry of numbers. Let **vol** be the normalized Haar measure on the additive group K_S^n (see §2). For a discrete I_S -module Γ in K_S^n the covolume of Γ (denoted by $\mathbf{cov}(\Gamma)$) is the **vol** of a fundamental domain of Γ in K_S^n . Let $B_r(K_S^n)$ be the closed ball of radius r centered at zero in K_S^n with respect to the normalized norm (see §2). For each integer $1 \leq m \leq n$, the m-th minimum of a discrete I_S -module Γ is defined by

$$(1.2) \lambda_m(\Gamma) = \inf\{r > 0 : \dim_K(\operatorname{span}_K(B_r(K_S^n) \cap \Gamma)) \ge m\}.$$

Here span_K is the K linear span of a set and \dim_K is the dimension of a vector space over K. Similar notations are used when K is replaced by other rings. We remark here that if $K = \mathbb{Q}$ and S contains only the archimedean place, then we get the usual concept of successive minima of lattice points in \mathbb{R}^n .

For two nonnegative real numbers s and t the notation $s \approx t$ means $C^{-1}s \leq t \leq Cs$ for some constant $C \geq 1$. Let σ and τ be the number of real and complex places of K respectively. Let $\sharp S = \tau + \mathbf{card}(S)$ where \mathbf{card} denotes the cardinality of a set. The S-adic version of Minkowski's theorem on successive minima (see [12, Chapter IV, §1] for the usual case) is the following theorem.

Theorem 1.2. Let $n \geq 1$ and let $\Gamma \subset K_S^n$ be a discrete I_S -module with finite covolume. Then

$$(\lambda_1(\Gamma)\dots\lambda_n(\Gamma))^{\sharp S} \asymp \mathbf{cov}(\Gamma)$$

where the implied constants depend on K, S and n.

A refined version of Theorem 1.2 will be proved in Theorem 4.4 where the implied constants will be explicitly calculated. If K is a function field of genus zero and S consists of a single place of degree one, then Theorem 1.2 is established in [10]. The adelic versions of Theorem 1.2 are proved in [1] (resp. [13]) when K is a number field (resp. function field).

¹If K is function field we have $\sigma = \tau = 0$.

2. Preliminaries: Notations

Let K be a global field and let P be the set of places of K. Throughout this paper we fix a positive integer n and a finite nonempty set $S \subset P$ such that $S \supset P_0$ where P_0 (possibly empty) is the set of archimedean places of K.

For every $v \in P$ let K_v be the completion of K at v. The S-adic numbers and integers are defined as

$$K_S \stackrel{\mathrm{def}}{=} \prod_{v \in S} K_v \quad \text{and} \quad I_S \stackrel{\mathrm{def}}{=} \{x \in K : x \text{ is integral for all } v \in P \smallsetminus S\}$$

respectively. We consider K as a subring of K_S via the natural inclusions $K \to K_v$. For $v \in P$, let $|\cdot|_v$ be the normalized absolute value on K_v . If v is archimedean, we identify K_v with real or complex numbers where the usual absolute value is $|\cdot|_v$. If v is ultrametric then $|a|_v^{-1} = \mathbf{card}(I_v/aI_v)$ for all $a \in I_v$ where I_v is the ring of integers of K_v . For each ultrametric place $v \in P$ we fix a uniformizer ϖ_v (a generator of the maximal ideal of I_v) and take $q_v = |\varpi_v|_v^{-1}$. We define the absolute value and content for $x = (x_v)_{v \in S} \in K_S$ respectively by

$$|x| \stackrel{\text{def}}{=} \max_{v \in S} |x_v|_v$$
 and $\operatorname{\mathbf{cont}}(x) \stackrel{\text{def}}{=} \prod_{v \in S} |x_v|_v^{\varepsilon_v}$

where $\varepsilon_v = 2$ if $K_v = \mathbb{C}$ and $\varepsilon_v = 1$ otherwise.

The additive group K_S^n can be naturally identified with $\prod_{v \in S} K_v^n$ and we write every $\xi \in K_S^n$ as $(\xi_v)_{v \in S}$ according to this identification. More precisely, if $\xi = (x_1, \ldots, x_n)$ where $x_i = (x_{i,v})_{v \in S}$, then $\xi_v = (x_{1,v}, \ldots, x_{n,v})$. Similarly, the group $\operatorname{GL}_n(K_S)$ can be naturally identified with $\prod_{v \in S} \operatorname{GL}_n(K_v)$ and we write every $g \in \operatorname{GL}_n(K_S)$ as $(g_v)_{v \in S}$ according to this identification. The group $\operatorname{GL}_n(K_v)$ (resp. $\operatorname{GL}_n(K_S)$) acts on K_v^n (resp. K_S^n) by matrix multiplication from the right. Moreover, the action of $g \in \operatorname{GL}_n(K_S)$ on $\xi \in K_S^n$ is consistent with these identifications, that is, $\xi g = (\xi_v g_v)_{v \in S}$ under previous notations.

For $v \in P$ we take \mathbf{vol}_v to be the normalized Haar measure on K_v . For archimedean v, the measure \mathbf{vol}_v is the Lebesgue measure. If v is ultrametric, the measure satisfies $\mathbf{vol}_v(I_v^n) = 1$. It follows directly from definition that

$$\mathbf{vol}_v(aB) = |a|_v^{n\varepsilon_v} \mathbf{vol}_v(B)$$

for every $a \in K_v$ and any measurable subset B of K_v^n . We take the normalized Haar measure \mathbf{vol} on K_S^n to be the product measure. In the sequel we will abbreviate $d\mathbf{vol}(\xi)$ by $d\xi$ for the integration with respect to the volume measure. For a positive integer m, we use \mathbf{vol}_v^m and \mathbf{vol}^m to denote the normalized Haar measures on K_v^m and K_S^m respectively.

If v is archimedean we take $\|\cdot\|_v$ to be the Euclidean norm on K_v^n . If v is ultrametric we take $\|\cdot\|_v$ to be the sup norm with respect to coordinates,

that is

$$\|(a_1,\ldots,a_n)\|_v \stackrel{\text{def}}{=} \max_{1 \le i \le n} |a_i|_v \text{ where } a_i \in K_v.$$

We define the norm and content for $\xi \in K_S^n$ by

$$\|\xi\| \stackrel{\text{def}}{=} \max_{v \in S} \|\xi_v\|_v$$
 and $\operatorname{\mathbf{cont}}(\xi) \stackrel{\text{def}}{=} \prod_{v \in S} \|\xi_v\|_v^{\varepsilon_v}$.

For $a \in I_S^*$, where I_S^* is the group of multiplicatively invertible elements of I_S , we have that $\mathbf{cont}(a) = 1$ and $\mathbf{cont}(a\xi) = \mathbf{cont}(\xi)$ where $\xi \in K_S^n$. Also for every $g \in \mathrm{GL}_n(K_S)$ we have

(2.1)
$$d(\xi g) = \mathbf{cont}(\mathbf{det}(g)) d\xi,$$

where **det** is the determinant of a matrix.

The set of vectors in K_S^n (resp. K_v^n) with norm less than or equal to r is denoted by $B_r(K_S^n)$ (resp. $B_r(K_v^n)$). It can be checked directly that

$$(2.2) B_r(K_S^n) = \prod_{v \in S} B_r(K_v^n).$$

Let L be a free K_S -submodule of K_S^n with rank $m \leq n$. Then

$$(2.3) L = \prod_{v \in S} L_v,$$

where L_v is an m-dimensional subspace of K_v^n . There is a unique additive Haar measure \mathbf{vol}_L on L (resp. \mathbf{vol}_{L_v} on L_v) such that

$$\mathbf{vol}_L(L \cap B_1(K_S^n)) = \mathbf{vol}^m(B_1(K_S^m))$$
(resp.
$$\mathbf{vol}_{L_v}(L \cap B_1(K_v^n)) = \mathbf{vol}_v^m(B_1(K_v^m))$$
).

Moreover, the above definition, (2.2) and (2.3) imply

(2.4)
$$\mathbf{vol}_L = \prod_{v \in S} \mathbf{vol}_{L_v}.$$

In the case where $K = \mathbb{Q}$ and $\operatorname{card}(S) = 1$, the measure vol_L is the measure given by the inner product on L. Suppose $\xi = (\xi_v)_{v \in S}$ and $\xi_v \neq 0$; then the covolume of $I_S \xi$ (I_S -linear span of ξ) in $K_S \xi$ (K_S -linear span of ξ) with respect to $\operatorname{vol}_{K_S \xi}$ is equal to $\operatorname{cont}(\xi)$ multiplied by the covolume of I_S in K_S . The covolume of a discrete I_S -module Γ in K_S^n with respect to the induced measure $\operatorname{vol}_{K_S \Gamma}$ is called relative covolume of Γ and it is denoted by $\operatorname{cov}_{\Gamma}(\Gamma)$. The covolume of Γ with respect to vol is denoted by $\operatorname{cov}_{\Gamma}(\Gamma)$.

3. Discrete I_S -modules

Let $\Gamma \subset K_S^n$ be a discrete I_S -module. In this section, we use ideas of [9, §8] to study properties of Γ .

Lemma 3.1. Let $\Gamma \subset K_S^n$ be a discrete I_S -module and let $\xi_1, \ldots, \xi_m \in \Gamma$. The following statements are equivalent:

(1) ξ_1, \ldots, ξ_m are linearly independent over I_S ;

- (2) ξ_1, \ldots, ξ_m are linearly independent over K;
- (3) ξ_1, \ldots, ξ_m are linearly independent over K_S .

Proof. It suffices to show that (1) implies (3). We prove it by induction on m. Write $\xi_i = (\xi_{i,v})_{v \in S}$ as in §2. Suppose that ξ_1 is linearly dependent over K_S , then there exists $w \in S$ such that $\xi_{1,w} = 0$. According to the strong approximation theorem (see [3, Chapter II §15]), there is a sequence $\{c_i\}_{i\geq 1}$ of $I_S \setminus \{0\}$ such that $|c_i|_v \to 0$ as $i \to \infty$ for any $v \in S \setminus \{w\}$. Therefore $c_i\xi_1 \to 0$ which contradicts the assumption that Γ is discrete. This proves (3) in the case where m = 1.

Now suppose m > 1 and (1) implies (3) while m is replaced by m-1. By the case for m=1, we know $\xi_{1,v} \neq 0$ for every $v \in S$. So there exists $g \in \operatorname{GL}_n(K_S)$ such that $\xi_1 g = (1,0,\ldots,0)$. The right multiplication of g on K_S^n is a K_S linear isomorphism, so we can without loss of generality assume that $\xi_1 = (1,0,\ldots,0)$. Let $\varphi: K_S^n \to K_S^n/K_S\xi_1 \cong K_S^{n-1}$ be the natural quotient map. Since $I_S\xi_1$ is a cocompact lattice in $K_S\xi_1$ and $\Gamma \subset K_S^n$ is discrete, the module $\varphi(\Gamma)$ is discrete and $\varphi(\xi_2),\ldots,\varphi(\xi_m)$ are linearly independent over I_S . In view of the induction hypothesis, we have $\varphi(\xi_2),\ldots,\varphi(\xi_m)$ are linear independent over K_S . Therefore ξ_1,\ldots,ξ_m are linearly independent over K_S .

Remark 3.2. The implication (2) \Rightarrow (3) holds without assuming that ξ_1, \ldots, ξ_m belong to a discrete I_S -module, see [9, Lemma 7.1].

For a discrete I_S -module $\Gamma \subset K_S^n$ let $K\Gamma$ (resp. $K_S\Gamma$) be the K-linear (resp. K_S -linear) span of Γ in K_S^n . We call the dimension over K of $K\Gamma$ the rank of Γ . It follows from Lemma 3.1 that the rank of Γ is less than or equal to n and the equality holds if and only if Γ has finite covolume.

In the following lemma we prove a Gram-Schmidt orthogonalization process for ultrametric local fields.

Lemma 3.3. Let K_v be a ultrametric local field. For any K_v -linearly independent vectors $\xi_1, \ldots, \xi_m \in K_v^n$ there exist linearly independent vectors $\eta_1, \ldots, \eta_m \in K_v^n$ such that η_1, \ldots, η_r are in the K_v -linear span of ξ_1, \ldots, ξ_r for all $r \leq m$, and

$$(3.1) ||a_1\eta_1 + \dots + a_m\eta_m||_v = \max_{1 \le i \le m} |a_i|_v for all a_i \in K_v.$$

Remark 3.4. In the sequel we call a basis of $L_v \stackrel{\text{def}}{=} \mathbf{span}_{K_v} \{\xi_1, \dots, \xi_m\}$ which satisfies (3.1) an orthonormal basis of L_v . The map

$$\varphi: K_v^m \to L_v$$
 where $\varphi_v(a_1, \ldots, a_m) = a_1 \eta_1 + \cdots + a_m \eta_m$

is an isometric embedding sending \mathbf{vol}_v^m to \mathbf{vol}_{L_v} .

Proof. Contrary to the archimedean case, here we choose an entry with maximal absolute value for the corresponding vector. Write

$$\xi_i = (x_{i1}, \dots, x_{in}), \text{ where } x_{ij} \in K_v \text{ and } 1 \le i \le m.$$

First we choose $j_1 \leq n$ such that $\|\xi_1\|_v = |x_{1j_1}|_v$ and set $\eta_1 = x_{1j_1}^{-1}\xi_1$. Next we take $\eta'_2 = \xi_2 - x_{2j_1}\eta_1 = (y_{21}, \ldots, y_{2n})$. We choose $j_2 \leq n$ such that $\|\eta'_2\|_v = |y_{2j_2}|_v$ and set $\eta_2 = y_{2j_2}^{-1}\eta'_2$. In general after r steps we have r different integers $j_1 \ldots, j_r$ and unit norm vectors η_1, \ldots, η_r such that η_i has j_i -th entry 1 and j_s -th entry zero for s < i. We take

$$\eta'_{r+1} = \xi_{r+1} - x_{r+1,j_1} \eta_1 - \dots x_{r+1,j_r} \eta_r = (y_1,\dots,y_n)$$

and choose j_{r+1} such that $\|\eta'_{r+1}\|_v = |y_{j_{r+1}}|_v$. We set $\eta_{r+1} = y_{j_{r+1}}^{-1} \eta'_{r+1}$. Then it has j_{r+1} -th entry 1 and j_s -th entry 0 for s < r + 1. This induction process gives m unit norm vectors η_1, \ldots, η_m .

For
$$(a_1, \ldots, a_m) \in K_v^m$$
 let

$$k = \min\{1 \le r \le m : |a_{j_r}|_v = \max_{1 \le i \le m} |a_i|_v\}.$$

It is clear from the construction that

$$||a_1\eta_1 + \dots + a_m\eta_m||_v = |a_{j_k}|_v,$$

which proves (3.1).

The next lemma appeared as [9, Corollary 8.4] for $K = \mathbb{Q}$ and as [8, Corollary 5.8] for K a number field.

Lemma 3.5. Suppose Γ and Γ' are discrete I_S -modules in K_S^n with

$$(3.2) K\Gamma \cap K\Gamma' = \{0\}.$$

Then

(3.3)
$$\operatorname{cov}_{\mathbf{r}}(\Gamma + \Gamma') \leq \operatorname{cov}_{\mathbf{r}}(\Gamma)\operatorname{cov}_{\mathbf{r}}(\Gamma').$$

Proof. Let $L = K_S\Gamma$, $L' = K_S\Gamma'$ and L'' = L + L'. In view of (3.2), Lemma 3.3 implies that L'' is a direct sum of L and L'. The right (resp. left) hand side of (3.3) is the covolume of $\Gamma + \Gamma'$ with relative to $\mathbf{vol}_L \times \mathbf{vol}_{L'}$ (resp. $\mathbf{vol}_{L''}$). Let $L = \prod_{v \in S} L_v$ and $L' = \prod_{v \in S} L'_v$ according to (2.3). In view of (2.4), it suffices to prove that for each $v \in S$ there is a positive Haar measure set R_v of $L_v + L'_v$ such that

$$(3.4) \mathbf{vol}_{L_v + L'_v}(R_v) \le (\mathbf{vol}_{L_v} \times \mathbf{vol}_{L'_v})(R_v).$$

Let r and m be the rank of L and L'' respectively. For each $v \in S$ we choose an orthonormal basis $\xi_{1,v}, \ldots, \xi_{r,v}$ of L_v and an orthonormal basis $\xi_{r+1,v}, \ldots, \xi_{m,v}$ of L'_v . We will show that (3.4) holds for

$$R_v := \{a_1 \xi_{1,v} + \ldots + a_m \xi_{m,v} : a_i \in B_1(K_v)\}.$$

For all $v \in S$

$$(\mathbf{vol}_{L_v} \times \mathbf{vol}_{L'_v})(R_v) = \mathbf{vol}_v^1(B_1(K_v))^m.$$

If v is archimedean, then it is clear from Eulidean geometry (i.e. volume of parallelotope) that

$$\mathbf{vol}_{L_v+L_v'}(R_v) \leq \mathbf{vol}_v^1(B_1(K_v))^m = (\mathbf{vol}_{L_v} \times \mathbf{vol}_{L_v'})(R_v).$$

If v is ultrametric, we let $\eta_{1,v}, \ldots, \eta_{m,v}$ be an orthonormal basis of $L_v + L'_v$. Then Remark 3.4 implies that R_v is contained in

$$R'_v := \{(a_1, \dots, a_m) \in B_1(K_v^m) : a_1\eta_1 + \dots + a_m\eta_m\}.$$

Using Remark 3.4 again together with (3.5), we have

$$\mathbf{vol}_{L_v+L_v'}(R_v) \leq \mathbf{vol}_{L_v+L_v'}(R_v') = \mathbf{vol}_v^1 \big(B_1(K_v)\big)^m = (\mathbf{vol}_{L_v} \times \mathbf{vol}_{L_v'})(R_v).$$

4. Successive minima

The aim of this section is to prove Theorem 1.2.

Lemma 4.1. Let $\Gamma \subset K_S^n$ be a discrete I_S -module with finite covolume and let $R \subset K_S^n$ be a measurable subset. Then there exists $\xi \in K_S^n$ such that

(4.1)
$$\operatorname{card}((\xi + R) \cap \Gamma) \ge \operatorname{vol}(R)/\operatorname{cov}(\Gamma).$$

Proof. Let χ_R be the characteristic function of R, and let $F \subset K_S^n$ be a fundamental domain for Γ . Then

$$\int_{F} \mathbf{card} ((\xi + R) \cap \Gamma) d\xi = \int_{F} \sum_{\gamma \in \Gamma} \chi_{R}(\gamma - \xi) d\xi = \mathbf{vol}(R).$$

Therefore there exists $\xi \in F$ such that (4.1) holds.

Lemma 4.2. Let $\Gamma \subset K_S^n$ be a discrete I_S -module with finite covolume. Let R_1 be a centrally symmetric convex subset of $K_{P_0}^n$ and let R_2 be a closed additive subgroup of $K_{S \setminus P_0}^n$. Suppose $R \subset K_S^n$ is equal to $R_1 \times R_2$ with the natural identification of K_S^n with $K_{P_0}^n \times K_{S \setminus P_0}^n$. If $\mathbf{vol}(R) > 2^{n(\sigma+2\tau)}\mathbf{cov}(\Gamma)$, then R contains a nonzero element of Γ .

Proof. Let $R' = (\frac{1}{2}R_1) \times R_2$. It follows from the assumption on the $\mathbf{vol}(R)$ that $\mathbf{vol}(R') > \mathbf{cov}(\Gamma)$. According to Lemma 4.1, we can find two distinct elements $\gamma_1, \gamma_2 \in \Gamma$ and $\xi \in K_S^n$ such that $\gamma_i - \xi \in R'$ for i = 1, 2. Therefore the nonzero element $\gamma_1 - \gamma_2$ belongs to R.

Recall that $\lambda_m(\Gamma)$ $(1 \leq m \leq n)$ is the *m*-th minimum of a discrete I_{S} -module Γ , see (1.2). It follows directly from the definition that there exist K-linearly independent vectors $\xi_1, \ldots, \xi_n \in \Gamma$ with

$$\|\xi_m\| = \lambda_m \text{ for all } 1 \leq m \leq n.$$

Moreover, by Lemma 3.1 these vectors are also linearly independent over K_S .

According to Lemma 4.2 for any $0 < t < \lambda_1(\Gamma)$ we have

$$(4.2) \qquad \Big(\prod_{v \in S \setminus P_0} q_v^{-n}\Big) t^{n\sharp S} \mathbf{vol}\big(B_1(K_S^n)\big) \leq \mathbf{vol}\big(B_t(K_S^n)\big) \leq 2^{n(\sigma+2\tau)} \mathbf{cov}(\Gamma).$$

Since $B_1(\mathbb{R}^m)$ contains $\{(x_1,\ldots,x_m)\in\mathbb{R}^m:-m^{-1/2}\leq x_i\leq m^{-1/2}\}$, for archimedean $v\in P$ we have

(4.3)
$$\operatorname{vol}_{v}(B_{1}(K_{v}^{n})) \geq 2^{n\varepsilon_{v}} n^{-n\varepsilon_{v}/2}$$

where for complex place we use $B_1(\mathbb{C}^n) = B_1(\mathbb{R}^{2n})$. By (4.2), (4.3) and (2.2) we have

(4.4)
$$\lambda_1(\Gamma)^{n\sharp S} \le n^{n(\sigma+2\tau)/2} \Big(\prod_{v \in S \setminus P_0} q_v^n \Big) \mathbf{cov}(\Gamma).$$

Lemma 4.3. Let Γ be a discrete I_S -module with finite covolume. Suppose that $\xi_1, \ldots, \xi_n \in \Gamma$ are K-linearly independent vectors and $\|\xi_m\| = \lambda_m(\Gamma)$ for all $1 \leq m \leq n$. Then there exists $g \in GL_n(K_S)$ such that

(4.5)
$$\operatorname{cont}(\operatorname{det}(g)) = \prod_{i=1}^{n} \operatorname{cont}(\xi_i)^{-1},$$

and any nonzero vector of $\Gamma' \stackrel{\text{def}}{=} \Gamma g$ has norm greater than or equal to one.

Proof. Suppose that $\xi_i = (\xi_{i,v})_{v \in S}$ where $\xi_{i,v} \in K_v^n$ (the notation here is the same as $\S 2$). By Lemma 3.1, for every $v \in S$ the vectors $\xi_{1,v}, \ldots, \xi_{n,v}$ are K_v -linearly independent in K_v^n . Using Gram-Schmidt orthogonalization process (see Lemma 3.3 for the ultrametric case), for each $v \in S$ we can find an orthonormal basis $\eta_{1,v}, \ldots, \eta_{n,v}$ such that for every $1 \leq m \leq n$ the K_v -linear span of $\eta_{1,v}, \ldots, \eta_{m,v}$ is the same as that of $\xi_{1,v}, \ldots, \xi_{m,v}$. Let $b_i = (b_{i,v})_{v \in S} \in K_S$ $(1 \leq i \leq n)$ such that $|b_{i,v}|_v = ||\xi_{i,v}||_v$. It follows from the definition of content that

(4.6)
$$\operatorname{cont}(b_i) = \operatorname{cont}(\xi_i).$$

Since $\eta_i \stackrel{\text{def}}{=} (\eta_{i,v})_{v \in S}$ $(1 \leq i \leq n)$ is a K_S -basis of K_S^n , there is a unique $g \in GL_n(K_S)$ such that $\eta_i g = b_i^{-1} \eta_i$. We claim that this g satisfies the requirement of the lemma.

The equation (4.5) follows from

$$\operatorname{cont}(\det(g)) = \operatorname{cont}(b_1^{-1} \dots b_n^{-1}) = \prod_{i=1}^n \operatorname{cont}(b_i)^{-1} = \prod_{i=1}^n \operatorname{cont}(\xi_i)^{-1},$$

where in the last equality we use (4.6). For the other conclusion suppose that $\zeta = c_1 \eta_1 + \cdots + c_m \eta_m \in \Gamma' = \Gamma g$ where $c_i \in K_S$ and $c_m \neq 0$. We have

$$\zeta g^{-1} = c_1 b_1 \eta_1 + \dots + c_m b_m \eta_m \in \Gamma.$$

Since for every $v \in S$ the basis $\eta_{1,v}, \ldots, \eta_{n,v}$ is orthonormal, we have

(4.7)
$$\|\zeta g^{-1}\| \le \|\zeta\| \max_{1 \le i \le m} |b_i| = \|\zeta\| \cdot \lambda_m(\Gamma).$$

On the other hand for any $1 \leq j \leq m$, the K_S -linear span of η_1, \ldots, η_j is the same as that of ξ_1, \ldots, ξ_j . Since $c_m b_m \neq 0$, Lemma 3.1 implies that $\xi_1, \ldots, \xi_{m-1}, \zeta g^{-1}$ are K-linearly independent. Thus it follows from the

definition of the *m*-th minimum of $\Gamma = \Gamma' g^{-1}$ that $\|\zeta g^{-1}\| \geq \lambda_m(\Gamma)$. This estimate together with (4.7) imply $\|\zeta\| \geq 1$, which completes the proof. \square

Theorem 4.4. Let Γ be a discrete I_S -module with finite covolume. Let $\xi_1, \ldots, \xi_n \in \Gamma$ be K linearly independent vectors with $\|\xi_m\| = \lambda_m(\Gamma)$ for all $1 \leq m \leq n$. Then we have

$$(4.8) \quad \mathbf{cov}(I_S^n)^{-1}\mathbf{cov}(\Gamma) \le \prod_{i=1}^n \mathbf{cont}(\xi_i) \le n^{n(\sigma+2\tau)/2} \Big(\prod_{v \in S \setminus P_0} q_v^n\Big) \mathbf{cov}(\Gamma).$$

Proof. We first prove the upper bound of (4.8). Suppose that $\Gamma' = \Gamma g$, where $g \in GL_n(K_S)$, satisfies the conclusion of Lemma 4.3. Then $\lambda_1(\Gamma') \geq 1$. Applying (4.4) for Γ' we have

$$(4.9) 1 \leq \lambda(\Gamma')^{n\sharp S} \leq n^{n(\sigma+2\tau)/2} \Big(\prod_{v \in S \setminus P_0} q_v^n\Big) \mathbf{cov}(\Gamma').$$

On the other hand by (2.1) and (4.5)

(4.10)
$$\operatorname{cov}(\Gamma') = \operatorname{cov}(\Gamma) \cdot \operatorname{cont}(\operatorname{det}(g)) = \operatorname{cov}(\Gamma) \cdot \prod_{i=1}^{n} \operatorname{cont}(\xi_i)^{-1}.$$

The upper bound of (4.8) follows from (4.9) and (4.10).

Let Γ'' be the I_S -linear span of ξ_1, \ldots, ξ_n . Since Γ'' is a submodule of Γ , by Lemma 3.5 we get

$$\mathbf{cov}(\Gamma) \leq \mathbf{cov}(\Gamma'') \leq \prod_{i=1}^{n} \mathbf{cov}_{\mathbf{r}}(I_{S}\xi_{i}) = \mathbf{cov}(I_{S}^{n}) \cdot \prod_{i=1}^{n} \mathbf{cont}(\xi_{i}),$$

which implies lower bound of (4.8).

To prove Theorem 1.2 we need a balance between contents and norms of vectors in K_S^n . The following lemma is a generalization of [9, Lemma 8.6] and [8, Lemma 5.9], and the proof is the same.

Lemma 4.5. For any $\xi \in K_S^n$ with $\mathbf{cont}(\xi) \neq 0$, there exists $a \in I_S^*$ such that $||a\xi||^{\sharp S} \times \mathbf{cont}(\xi)$ where the implied constants depend on K and S.

Proof. Suppose that $S = \{v_1, \ldots, v_m\}$ where $m = \mathbf{card}(S)$. Let \mathbb{R}_+ be the multiplicative group of positive real numbers. We define a map

$$\varphi: K \to \mathbb{R}^m_+$$
 by $\varphi(a) = (|a|_{v_1}, \dots, |a|_{v_m}).$

Let

$$H = \{(r_1, \dots, r_m) \in \mathbb{R}_+^m : \prod_{i=1}^m r_i^{\varepsilon_{v_i}} = 1\}.$$

It follows from Dirichlet's unit theorem (see [3, Chapter II §18]) that the group $\varphi(I_S^*) \subset H$ is a cocompact lattice in H. So there exists $A \geq 1$ which depends on K and S such that for any $(r_1, \ldots, r_m) \in H$ we can find $a \in I_S^*$ with

$$(4.11) A^{-1} \le r_i |a|_{v_i} \le A.$$

Suppose $\xi = (\xi_v)_{v \in S}$. It follows from the definition that

$$(\|\xi_{v_1}\|_{v_1} \cdot \mathbf{cont}(\xi)^{-1/(m+\tau)}, \dots, \|\xi_{v_m}\|_{v_m} \cdot \mathbf{cont}(\xi)^{-1/(m+\tau)}) \in H.$$

By (4.11) one can find $a \in I_S^*$ such that for all $1 \le i \le m$

$$A^{-1} \le \|\xi_{v_i}\|_{v_i} \mathbf{cont}(\xi)^{-1/(m+\tau)} |a|_{v_i} \le A.$$

Therefore

$$A^{-m-\tau}\mathbf{cont}(\xi) \le ||a\xi||^{m+\tau} \le A^{m+\tau}\mathbf{cont}(\xi).$$

Proof of Theorem 1.2. Let $\xi_1, \ldots, \xi_n \in \Gamma$ be K-linearly independent vectors with $\|\xi_i\| = \lambda_i(\Gamma)$. By Theorem 4.4

(4.12)
$$\prod_{i=1}^{n} \mathbf{cont}(\xi_i) \approx \mathbf{cov}(\Gamma),$$

where the implied constants depend on K, S and n. The definitions of content and norm imply

(4.13)
$$\operatorname{cont}(\xi_i) \le \lambda_i(\Gamma)^{\sharp S}.$$

According to Lemma 4.5 there exists $a_1, \ldots, a_n \in I_S^*$ such that

$$(4.14) \qquad \qquad \mathbf{cont}(\xi_i) \gg \|a_i \xi_i\|^{\sharp S},$$

where the implied constant depends on K and S. Note that elements $a_1\xi_1,\ldots,a_n\xi_n\in\Gamma$ are linear independent over K. So the definition of successive minima implies

$$(4.15) \qquad \prod_{i=1}^{n} \lambda_i(\Gamma) \le \prod_{i=1}^{n} ||a_i \xi_i||.$$

Therefore the conclusion of Theorem 1.2 follows from (4.12), (4.13), (4.14) and (4.15).

5. Mahler's compactness criterion

Let $X = \operatorname{SL}_n(I_S) \backslash \operatorname{SL}_n(K_S)$. There is a one-to-one correspondence between X and

$$\{I_S^n g : g \in \operatorname{SL}_n(K_S)\}$$

via the map $SL_n(I_S)g \to I_S^n g$. In this section $\mathbf{e}_1, \dots, \mathbf{e}_n$ denotes the standard basis of K_S^n , i.e. \mathbf{e}_i has *i*-th entry 1 and other entries 0. Before proving Theorem 1.1 we need the following lemma. See [9, Corollary 8.6] for $K = \mathbb{Q}$ and [8, Corollary 5.1] for K a number field.

Lemma 5.1. Let M > 0. Then there are only finitely many I_S -submodules Γ of I_S^n such that $\mathbf{card}(I_S^n/\Gamma) \leq M$.

Proof. Let $\Gamma \subset I_S^n$ be an I_S -submodule with $\mathbf{card}(I_S^n/\Gamma) \leq M$. For every $1 \leq i \leq n$ there is an ideal J_i of I_S such that

$$I_S \mathbf{e}_i \cap \Gamma = J_i \mathbf{e}_i$$
 and $\mathbf{card}(I_S/J_i) \leq M$.

Therefore

$$J_1 \times \cdots \times J_n \subset \Gamma \subset I_S^n$$
.

Note that I_S is a Dedekind domain. It follows from the structure theory of ideals in I_S that there are only finitely many ideals J in I_S such that $\operatorname{card}(I_S/J) \leq M$. So the conclusion of the lemma holds.

Proof of Theorem 1.1. Let $\pi: \mathrm{SL}_n(K_S) \to X$ be the natural quotient map and let

(5.1)
$$r = \inf\{\|\xi g\| : \xi \in I_S^n, \ \xi \neq 0, \ g \in \operatorname{SL}_n(K_S), \ \pi(g) \in R\}.$$

Suppose R is relatively compact. There exists a relatively compact subset $F \subset \mathrm{SL}_n(K_S)$ with $\pi(F) = R$. Therefore there exists C > 0 such that

(5.2)
$$\|\xi g\| \le C\|\xi\|$$
 for every $\xi \in K_S^n$ and $g \in F$.

The discreteness of Γ and (5.2) imply r > 0.

Now we assume r > 0 and prove that R is relatively compact. Let $\{g_i\}_{i \geq 1}$ be a sequence in $\pi^{-1}(R)$. It suffices to show that there exists $g \in \mathrm{SL}_n(K_S)$ such that $\pi(g)$ is a limit point of a subsequence of $\{\pi(g_i)\}_{i \geq 1}$. By Theorem 1.2 there exists $C \geq 1$ such that for any free I_S -module $\Gamma \in R$ one has

$$(5.3) r \le \lambda_1(\Gamma) \le \lambda_n(\Gamma) \le C.$$

For every $i \geq 1$ let $\xi_1^{(i)}, \ldots, \xi_n^{(i)} \in I_S^n$ be K-linearly independent vectors such that $\|\xi_i^{(i)}g_i\|$ equals to the j-th minimum of $I_S^ng_i$. By (5.3) we have

(5.4)
$$\|\xi_{i}^{(i)}g_{i}\| \leq C \quad \forall i \geq 1 \text{ and } 1 \leq j \leq n.$$

Let

$$\Gamma_i = \mathbf{span}_{I_S} \{ \xi_j^{(i)} : 1 \le j \le n \}.$$

According to (5.3) and Theorem 1.2 there exists M > 0 such that

(5.5)
$$\operatorname{cov}(I_S^n) \le \operatorname{cov}(\Gamma_i) = \operatorname{cov}(\Gamma_i g_i) \le M \quad \forall i \ge 1.$$

By Lemma 5.1 and (5.5), the set $\{\Gamma_i: i \geq 1\}$ is finite. Therefore by possibly passing to a subsequence we may assume that there exists $h \in \operatorname{GL}_n(K)$ such that $\Gamma_i = I_S^n h$ for all $i \geq 1$. It follows that there is a sequence $\{f_i\}_{i \geq 1}$ in $\operatorname{GL}_n(I_S)$ such that $\mathbf{e}_j f_i h = \xi_j^{(i)}$ for all $i \geq 1$ and $1 \leq j \leq n$. By (5.4) there is a subsequence $\{g_{i_k}\}_{k \geq 1}$ of $\{g_i\}_{i \geq 1}$ and $g \in \operatorname{GL}_n(K_S)$ such that

$$(5.6) f_{i_k} h g_{i_k} \to g as k \to \infty.$$

Since **det** is continuous and I_S is discrete in K_S , for k sufficiently large we have $\det(f_{i_k}) = \det(f_{i_{k+1}}) \in I_S^*$. Therefore by possibly passing to a subsequence and multiplying the first row of h by some element of I_S^* , we assume without loss of generality that $f_{i_k} \in \operatorname{SL}_n(I_S)$ for all k.

The group $h^{-1}\operatorname{SL}_n(I_S)h\cap\operatorname{SL}_n(I_S)$ has finite index in $h^{-1}\operatorname{SL}_n(I_S)h$. So by possibly passing to a subsequence we can find $f \in h^{-1}\operatorname{SL}_n(I_S)h$ and a sequence $\{h_k\}_{k\geq 1}$ of $\operatorname{SL}_n(I_S)$ such that

(5.7)
$$h^{-1}f_{i_k}h = fh_k \quad \forall \ k \ge 1.$$

By (5.6) and (5.7) we have $hfh_kg_{i_k}\to g$ as $k\to\infty$. Therefore

$$h_k g_{i_k} \to f^{-1} h^{-1} g$$
 as $k \to \infty$.

Since $h_k \in \mathrm{SL}_d(I_S)$ we have $\pi(g_{i_k}) \to \pi(f^{-1}h^{-1}g)$ as $k \to \infty$.

REFERENCES

- [1] E. Bombieri and J. Vaaler, On Siegels lemma, Invent. Math. 73 (1983), 11–32.
- [2] J.W.S. Cassels, An introduction to the geometry of numbers, Die Grundlehren der mathematischen Wissenschaften, Band 99, Springer-Verlag, Berlin-New York, 1971.
- [3] J. Cassels and A. Frohlich, *Algebraic number theory*, Proceedings of the Brighton Conference, Academic Press, New York, 1968.
- [4] M. Einsiedler, Applications of measure rigidity of diagonal actions, in: Proceedings of the International Congress of Mathematicians, vol. III, 2010, 1740–1759
- [5] M. Einsiedler and E. Lindenstrauss, Diagonal flows on locally homogeneous spaces and number theory, in: Proceedings of the International Congress of Mathematicians, Vol. II, 2006, 1731–1759.
- [6] A. Ghosh, Metric Diophantine approximation over a local field of positive characteristic, J. Number Ttheory 124 (2007), 454–469.
- [7] D. Kleinbock, N. Shah, and A. Starkov, Dynamics of subgroup actions on homogeneous spaces of Lie groups and applications to number theory, in: Handbook of Dynamical Systems, Vol. 1A (2002), North-Holland, Amsterdam, 813–930.
- [8] D. Kleinbock and G. Tomanov, Flows on S-arithmetic homogeneous spaces and applications to metric Diophantine approximation, MPIM Preprint 2003-65, https://www.mpim-bonn.mpg.de/preblob/2234.
- [9] ______, Flows on S-arithmetic homogeneous spaces and applications to metric Diophantine approximation, Comment. Math. Helv'82 (2007), 519–581.
- [10] K. Mahler, An analogue to Minkowski's geometry of numbers in a field of series, Ann. Math. 42 (1941), 488–522.
- [11] ______, On lattice points in n-dimensional star bodies: I. Existence theorems, Proc. Roy. Soc. London, A 187 (1946), 151–187.
- [12] W. Schmidt, Diophantine approximation, Lecture Notes in Mathematics, vol. 785, Springer-Verlag, Berlin, 1980.
- [13] J.L. Thunder, Siegel's lemma for function fields, Mich. Math. J. 42 (1995), 147–162.

Department of Mathematics, Brandeis University, Waltham, MA 02454-9110, USA

 $E ext{-}mail\ address: kleinboc@brandeis.edu}$

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, PR CHINA

E-mail address: ronggang@xmu.edu.cn

Institut Girard Desargues, Université Claude Bernard-Lyon 1, 69622 Villeurbanne cedex, France

 $E ext{-}mail\ address: Georges.Tomanov@desargues.univ-lyon1.fr}$