

# Probabilist Set Inversion using a new framework for interval arithmetic

Abdel KENOUI\*

*SCORE, Scientific Consulting for Research and Engineering, Strasbourg, France*

Nicolas GOZE<sup>†</sup>

*CETE, Centre d'Etudes Techniques de l'Equipement, Strasbourg, France*

Michel GOZE<sup>‡</sup>

*LMIA, Laboratoire de Mathématiques, Informatique et Applications, Université de Haute-Alsace, Mulhouse, France*

(Dated: November 24, 2021)

In this paper, we present how to use an interval arithmetics framework based on free algebra construction, in order to build better defined inclusion function for interval semi-group and for its associated vector space. One introduces the  $\psi$ -algorithm, which performs set inversion of functions and exhibits some numerical examples developed with the *python* programming langage.

## I. INTRODUCTION

Many problems dealing with real numbers can be solved with intervals computations<sup>3,4,26-29</sup>, such as set inversion, functions optimization, linear algebra. Their key stone is the construction of the inclusion function from the natural one. Some approaches have developed methods to circumvent it with using boolean inclusion tests, series or limited expansions of the natural function where the derivatives are computed at a certain point of the intervals. Nevertheless, those transfers from real functions to functions defined on intervals are not systematic and not given by a formal process. This yields to the fact that the inclusion function definition has to be adapted to each problem with the risk to miss the primitive scope.

We propose here a clear and simple scheme to build inclusion functions from the natural one. It is based on the arithmetic developed by one of the author in his PhD thesis<sup>24</sup>, called in this paper free-algebra based arithmetic. This article reminds first the definition and characteristics of the intervals semi-group  $\mathbb{IR}$  and the construction of its associated vector space  $\overline{\mathbb{IR}}$ . We explain how to get an associative and distributive arithmetic of intervals by embedding the vector space into a free algebra. After that, one shows from an example how to generalize the inclusion function construction for the semi-group and the vector space. This permits to present the  $\psi$ -algorithm in order to perform set inversion which is based on this construction and probability calculations. We ends with numerical applications examples developed in *python*<sup>30</sup> programming langage for set inversion.

## II. THE BANACH SPACE $\mathbb{IR}$ AND THE ASSOCIATIVE ALGEBRA $\mathcal{A}_4$

An interval  $X$  is defined as a non-empty and connected set of real numbers. It can be represented as a point in the half-plane of  $\mathbb{R}^2$ ,  $\mathcal{P}_1 = \{(a, b) \in \mathbb{R}^2, a \leq b\}$ . The set  $\mathcal{P}_2 = \{(a, b) \in \mathbb{R}^2, a \geq b\}$  is the set of anti-intervals as shown on figure 1. One writes real numbers as intervals with same bounds,  $\forall a \in \mathbb{R}, a \equiv [a, a]$ . We denote by  $\mathbb{IR} = \mathcal{P}_1$  the set of intervals of  $\mathbb{R}$ . The semantical arithmetic operations on intervals, called *Minkowski or classical operations*, are defined such that the result of the corresponding operation on elements belonging to operand intervals belongs to the resulting interval. That is, if  $\diamond$  denotes one of the usual operations  $+$ ,  $-$ ,  $*$ ,  $/$ , we have, if  $X$  and  $Y$  are bounded intervals of  $\mathbb{R}$ ,

$$X \diamond Y = \{x \diamond y / x \in X, y \in Y\},$$

But  $\mathbb{IR}$  is only a semi-group for the addition and no subtraction can be defined for it.

### A. Modal interval arithmetic

An algebraic extension of the classical interval arithmetic, called generalized interval arithmetic<sup>3,4</sup> has been proposed first by M. Warmus<sup>1,2</sup>. It has been followed in the seventies by H. -J. Ortolof and E. Kaucher<sup>5-9</sup>. In this former interval arithmetic, the intervals form a group with respect to addition and a complete lattice with respect to inclusion. In order to adapt it to semantic problems, Gardenes et al. have developed an approach called modal interval arithmetics<sup>10-15</sup>. S. Markov and others investigate the relation between generalized intervals operations and Minkowski operations for classic intervals and propose the so-called directed interval arithmetic, in which Kaucher's generalized intervals can be viewed as classic intervals plus direction, hence the name directed interval arithmetic<sup>17,18</sup>. In this arithmetic framework, proper and improper intervals are considered as intervals with sign<sup>19</sup>. Interesting relations and developements for proper and improper intervals arithmetic and for applications can be found in litterature<sup>20-22</sup>.

Our approach<sup>24,25</sup>, that we remind below in this artice, is similar to the previous ones in the sense that intervals are extended to generalized intervals. However we use a construction which is more canonical and based on the semi-group completion into a a group, which permits then to build the associated real vector space, to get an analogy with directed interval arithmetic and to perform differential calculus and linear algebra naturally.

## B. New interval arithmetic

Let  $\overline{\mathbb{R}}$  be the  $\mathbb{R}$  completed group. An element of  $\overline{\mathbb{R}}$  is written  $\overline{(A, B)}$  where  $A, B \in \mathbb{R}$  and the equivalence relation being  $\overline{(A, B)} \sim \overline{(A', B')}$  if and only if  $A + B' = B + A'$ . For example  $\overline{([3, 4], [1, 4])} \sim \overline{([2, 0], [0, 0])} \equiv \overline{([2, 0], 0)}$ . Any equivalence class can be written as  $\overline{(X, 0)}$  or  $\overline{(0, X)}$  or  $\overline{(a, a)}$  with  $X \in \mathbb{R}$  or  $a \in \mathbb{R}$ . We denote  $\mathcal{X} \setminus \mathcal{Y}, \mathcal{X}, \mathcal{Y} \in \overline{\mathbb{R}}$ , the difference into the abelian group  $\overline{\mathbb{R}}$ . The external multiplication

$$\mathbb{R} \times \overline{\mathbb{R}} \longrightarrow \overline{\mathbb{R}} \quad (1)$$

given by

$$a \cdot \overline{(X, 0)} = \overline{(a \cdot X, 0)} \text{ if } a \geq 0 \quad (2)$$

$$= \overline{(0, -a \cdot X)} \text{ if } a \leq 0 \quad (3)$$

$$(4)$$

and

$$a \cdot \overline{(0, X)} = \overline{(0, a \cdot X)} \text{ if } a \geq 0 \quad (5)$$

$$= \overline{(-a \cdot X, 0)} \text{ if } a \leq 0 \quad (6)$$

$$(7)$$

provides  $\overline{\mathbb{R}}$  with a 2-dimensional  $\mathbb{R}$  vector space structure. A basis is given by the family  $\{\overline{([1, 1], 0)}, \overline{([0, 1], 0)}\}$ . The following map

$$\|\cdot\| : \overline{\mathbb{R}} \longrightarrow \mathbb{R}^+ \quad (8)$$

$$\overline{(X, 0)} \mapsto l(X) + |c(X)| \quad (9)$$

$$\overline{(0, X)} \mapsto l(X) + |c(X)| \quad (10)$$

$$(11)$$

with respectively  $l(X)$  and  $c(X)$  the width and the center of the interval  $X$ , is a norm. It is easy to prove<sup>24,25</sup> that  $\overline{\mathbb{R}}$  is isomorphic to  $\mathbb{R}^2$  which is complete, this yields to the fact that this norm endows  $\overline{\mathbb{R}}$  with a Banach space structure. Moreover, one can define an order relation  $<$  on  $\overline{\mathbb{R}}$  which is compatible with this norm :  $X, Y \in \overline{\mathbb{R}}, X < Y$  if

$$X \not\subset Y \text{ and } c(X) < c(Y) \quad (12)$$

or

$$X \subset Y \text{ and } l(X) < l(Y) \quad (13)$$

Due to the fact that every intervals can be compared to each others, it yields that this partial order relation can be extended to a total one.

In order to define a natural multiplication on  $\overline{\mathbb{R}}$ , we consider the following 4-dimensional associative algebra  $\mathcal{A}_4$  with basis :

$$\begin{cases} e_1 = (1, 1), \\ e_2 = (0, 1), \\ e_3 = (-1, 0), \\ e_4 = (-1, -1). \end{cases}$$

and product :

	$e_1$	$e_2$	$e_3$	$e_4$
$e_1$	$e_1$	$e_2$	$e_3$	$e_4$
$e_2$	$e_2$	$e_2$	$e_3$	$e_3$
$e_3$	$e_3$	$e_3$	$e_2$	$e_2$
$e_4$	$e_4$	$e_3$	$e_2$	$e_1$

(14)



product. If they are belonging both to  $\mathcal{P}_{12}$ , the result is greater than the Minkowski one. If one needs a result  $X \cdot Y$  closer to  $X \diamond Y$  or even equal to, one can embed  $\mathbb{IR}$  in higher dimension algebras, such as  $\mathcal{A}_5, \mathcal{A}_6, \mathcal{A}_7$ , and so on. This former one is sufficient to recover the distributivity and equality to Minkowski product<sup>24</sup>.

- It can be naturally extended to  $\overline{\mathbb{IR}}$ .
- The product satisfies the monotony property.
- With the canonical injection  $X \mapsto \overline{(X, 0)}$  from  $\mathbb{IR}$  to  $\overline{\mathbb{IR}}$ , one can consider that  $\mathbb{IR} \subset \overline{\mathbb{IR}}$ .




Figure 2. Interval embedding in the algebra  $\mathcal{A}_4$ .

### III. INCLUSION FUNCTIONS

It is necessary for some problems to extend the definition of a function defined for real numbers  $f : \mathbb{R} \rightarrow \mathbb{R}$  to function defined for intervals  $[f] : \mathbb{IR} \rightarrow \mathbb{IR}$  such as  $[f]([a, a]) = f(a)$  for any  $a \in \mathbb{R}$ . It will be convenient to have the same formal expression for  $f$  and  $[f]$ . Usually the lack of distributivity in Minkowski arithmetics doesn't give the possibility to get the same formal expressions. But with the arithmetic we have presented, there is no data dependency anymore and one can define easily inclusion functions from the natural one. For example, let's extend to intervals the real functions  $f_1(x) = x^2 - 2x + 1$ ,  $f_2(x) = (x - 1)^2$ ,  $f_3(x) = x(x - 2) + 1$ . Usually, with the Minkowski operations, the three expressions of this same function for the interval  $X = [3, 4]$  are  $[f]_1(X) = [2, 11]$ ,  $[f]_2(X) = [4, 9]$  and  $[f]_3(X) = [6, 12]$ . Data dependency occurs when the variable appears more than once in the function expression. The deep reason of that is the lack of distributivity in Minkowski arithmetics. But within the arithmetic developed in  $\mathcal{A}_4$  or higher dimension free algebras<sup>24,25</sup>, this problem vanishes. For example : with  $X = [3, 4]$  and since  $X \in \mathcal{P}_{11}$ ,

$$\varphi(X) = (3, 4 - 3, 0, 0) = (3, 1, 0, 0) = 3e_1 + e_2. \quad (19)$$

Since  $e_1 \equiv 1$ , and with means of product table (14) one has

$$\begin{aligned} \varphi([f]_1(X)) &= (3e_1 + e_2)^2 - 2(3e_1 + e_2) + 1 \\ &= 9e_1^2 + 2 \cdot 3e_1e_2 + e_2^2 - 2 \cdot 3e_1 - 2e_2 + 1 \\ &= 9e_1 + 6e_2 + e_2 - 6e_1 - 2e_2 + e_1 = 4e_1 + 5e_2 \\ &= \varphi([4, 9]), \end{aligned} \quad (20)$$

$$\begin{aligned} \varphi([f]_2(X)) &= (3e_1 + e_2 - 1)^2 \\ &= (2e_1 + e_2)^2 \\ &= 4e_1^2 + 4e_1e_2 + e_2^2 \\ &= 4e_1 + 4e_2 + e_2 \\ &= 4e_1 + 5e_2 \\ &= \varphi([4, 9]), \end{aligned} \quad (21)$$

and

$$\begin{aligned}
\varphi([f]_3(X)) &= (3e_1 + e_2) \cdot (3e_1 + e_2 - 2) + 1 \\
&= 9e_1 + 3e_1e_2 - 6e_1 + 3e_1e_2 + e_2^2 - 2e_2 + e_1 \\
&= 4e_1 + 3e_2 + 3e_2 + e_2 - 2e_2 \\
&= 4e_1 + 5e_2 \\
&= \varphi([4, 9]).
\end{aligned} \tag{22}$$

Thus,  $[f]_1(X) = [f]_2(X) = [f]_3(X) = [4, 9]$ .

On the other hand, the construction of the inclusion function depends on the type of problem one deals with. If one aims to perform set inversion for example, it has to be done in the semi-group  $\mathbb{IR}$  due to semantic definition of substraction. But, the substraction is not defined in  $\mathbb{IR}$ . This problem can be circumvented by replacing it with an addition and a multiplication with the interval  $e_4 = [-1, -1]$ . This maintains the associativity and distributivity of arithmetics and permits to introduce a semantical substraction. For example : if  $f(x) = x^2 - x = x(x - 1)$  for real numbers, one defines  $[f](X) = X^2 + e_4 \cdot X$ . One reminds the product  $[-1, -1] \cdot [a, b]$  is equal to  $[-b, -a]$ . Due to the fact that the arithmetics is now associative and distributive, one doesn't have data dependency anymore and  $[f](X) = X^2 + e_4 \cdot X = X \cdot (X + e_4)$ . The last term corresponds to the transfer of  $x(x - 1)$ . Division can be transferred to the semi-group in the same way by replacing  $\frac{1}{x} = x^{-1}$  with  $X^{e_4}$ .

Taylor polynomial expansions, differential calculus and linear algebra operations are defined only in a vector space. Therefore the transfer for the vector space is done directly. This permits to get infinitesimal intervals with the substraction and to compute derivatives. This is of course not allowed and not possible into the semi-group. From  $\mathbb{IR}$  to the vector space  $\mathbb{IR}$ ,  $f : x \mapsto -x$  is transferred to  $[f] : X \equiv (X, 0) \mapsto \setminus(X, 0) \equiv \setminus X$ . This means that  $[a, b]$  substraction is the anti-interval  $[-a, -b]$  addition. One of the most important consequence is that it is possible to transfer some functions directly to the intervals. For example, it is easy to prove analytically in  $\mathbb{IR}$  that  $\overline{\exp}(\overline{([a, b], 0)}) = (\overline{\exp(a)}, \overline{\exp(b)}, 0)$  with means of Taylor expansion.

#### IV. PROBABILIST SET INVERSION : $\psi$ -ALGORITHM

##### A. Flowchart

One presents an efficient set inversion method whose flowchart is very simple. One of the powerful application of interval calculus is the set inversion of a real-valued function defined on real numbers. It finds a huge number of applications in automation, optimization, resolution of algebraic and differential equations. The mathematical modelization of this problem is the following as shown on Fig.(3) : let's note  $f : \mathbb{R}^n \mapsto \mathbb{R}^p$  a function for a physical system, which is required to be surjective only,  $\mathcal{R} \subset \mathbb{R}^n$  the set of adjustments, and  $\mathcal{P} \subset \mathbb{R}^p$  the set of performance of a system. Set inversion consists of the computation of  $\mathcal{S} = f^{-1}(\mathcal{P}) \cap \mathcal{R}$ . For semantical reasons, one has to perform this set inversion within the semi-group  $\mathbb{IR}$ .

Some interesting and powerful methods using intervals have been developped those last years, especially SIVIA<sup>32</sup>, Set Inversion *Via* Interval Analysis. But the inclusion function being not well defined in the semi-group with the Minkowski arithmetic, SIVIA uses boolean inclusion tests and finds accepted, rejected and "uncertain" domains. With the algebraic arithmetic, one doesn't need boolean tests since the inclusion functions are well-defined. Thus, we propose the  $\psi$ -algorithm (Probabilist Set Inversion) inspired from SIVIA but without boolean tests and with a conditional probability calculation and domain bisections. This yields to accepted or rejected domains only. We are interested to compute the following conditional probability

$$\begin{aligned}
p(\mathcal{X}) &= p([f](\mathcal{X}) \subset \mathcal{P} \mid f(x) \in [f](\mathcal{X}), \forall x \in \mathcal{X}) \\
&= \frac{mes([f](\mathcal{X}) \cap \mathcal{P})}{mes([f](\mathcal{X}))} \\
&= \frac{mes(\mathcal{Y} \cap \mathcal{P})}{mes(\mathcal{Y})} = \frac{mes(\mathcal{I})}{mes(\mathcal{Y})}
\end{aligned} \tag{23}$$







Figure 3. Set inversion modelization with intervals

Figure 4.  $\psi$ -algorithm for  $f_1$ 

where  $mes$  is the Lebesgue measure in  $\mathbb{R}^p$  (length, surface, ...). If this probability equals 1 then the set is added to the list of solutions. If it is zero the set is rejected and removed from the list of interval candidates. If the probability is such as  $p(\mathcal{X}) \in ]0, 1[$ , then  $\mathcal{X}$  is bisected and  $\psi$ -algorithm applies the same procedure recursively for the resulting intervals until the size is lower than a fixed size resolution of the intervals or until the sets are accepted or rejected. Since  $\psi$ -algorithm creates sequences of decreasing intervals which are compact sets, it is obvious that  $\psi$ -algorithm converges to fixed points probabilities which are simply 0 and 1. In fact, we consider a sequence of compact sets  $\{K_n\}_{n \in \mathbb{N}}$  satisfying  $d(K_n) < d(K_{n+1})$  where  $d$  is the diameter of the compact set. If  $K_n \cap K_m = \emptyset$ , for any  $n \neq m$ , the sequence is convergent and the limit is the empty set. Then, one has just to consider the sequence of bisected sets.

## B. Numerical applications



We have developed a numerical library for *python* environnement<sup>30</sup> called yet *interval.lib*. It is a pure numerical implementation performing the basic arithmetics presented above<sup>24,25</sup>. This library aims to give simple and optimized routines to perform interval calculations based on the algebraic arithmetics. One gives in this section some numerical application examples of  $\psi$ -algorithm in order to illustrate how it can treat usual inversion problems and

Figure 5.  $\psi$ -algorithm for  $f_2$ Figure 6.  $\psi$ -algorithm for  $f_3$ 

build well-defined inclusion functions.

Let's define the non-linear functions  $f_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  with respectively adjustments and performances sets  $\mathcal{R}_i, \mathcal{P}_i$  :

$$\begin{aligned} f_1(x,y) &= (x^2 + y^2, x + y), \quad \mathcal{R}_1 = [-1, 2]^2, \quad \mathcal{P}_1 = [1, 2] \times [1, 4] \\ f_2(x,y) &= (x^2 - y^2, x \cdot y), \quad \mathcal{R}_2 = [-3, 3]^2, \quad \mathcal{P}_2 = [0, 4] \times [-2, 2] \\ f_3(x,y) &= (x^2 - y^2, \frac{y}{1+x}), \quad \mathcal{R}_3 = [0, 6] \times [0, 10], \quad \mathcal{P}_3 = [0, 5] \times [-4, 4]. \end{aligned}$$

Figure 7.  $\psi$ -algorithm for  $f_4$ Figure 8. Accepted and candidate intervals total ratios for numbers and surfaces according iterations in  $\psi$ -algorithm for  $f_4$  with  $10^{-4}$  spatial resolution.

Those examples have been chosen to give examples of addition, subtraction, product and division transfers from  $\mathbb{R}$  to  $\mathbb{IR}$ , and to exhibit the difference between the usual Minkowski arithmetic and the algebraic one<sup>24,25</sup>. The calculations with the  $\psi$ -algorithm are shown on figures 4, 5, 6 and 7. The convergence to 0 or 1 probabilities only, shows that inclusion functions are well constructed and that the interval arithmetic is robust. The last example

$$f_4(x, y) = (x^2 - y^2 \cdot \exp(x) + x \cdot \exp(y), x \cdot (x + y) - y^2), \mathcal{R}_4 = \mathcal{P}_4 = [-5, 5]^2 \quad (25)$$

presented on figure 7 shows clearly that the  $\psi$ -algorithm implemented in the algebraic arithmetic we use is not data dependant. The variables appear more than once in the formal expression of the function  $f_4$ . The CPU time for this inversion is about 255 seconds on a simple 1.67 Ghz Intel processor for a spatial surface resolution of  $10^{-4}$ . In figure 8, one computes for this resolution the number of solution intervals according to candidates ones and the total surfaces ratio of the same intervals. The strong correlation between the two curves characterizes the fact that the measures computations are well-defined because the probabilities are very close to the frequencies computed with  $\psi$ -algorithm independantly of the initial adjustments set. This confirms that the free algebra based arithmetic framework is relevant even for a semi-group  $\mathbb{IR}$  calculations example. Due to the bisection, the algorithm computational complexity is exponential according to the iterations  $N$ , and it is not improved compared to SIVIA one. In our scheme, computational time is defined as

$$T_{comp} = \mathcal{O}(N) = k \cdot 2^N. \quad (26)$$

However, if the native function is differentiable on  $\mathbb{R}^n$ , it is possible to define an adaptative mesh, with bisection spanned only in the space directions where the derivative magnitude is larger than a certain fixed value, because it is not useful to bisect in flat directions. This will obviously decrease the computational complexity constant  $k$ .

## V. CONCLUSION

We have presented in this article an heuristic way to transfer real functions to inclusion ones depending on the space needed, semi-group or vector space of intervals. This permits to define a simple but very efficient algorithm for set inversion inspired from SIVIA<sup>32</sup>, the  $\psi$ -algorithm, which uses a free algebra based arithmetic<sup>24,25</sup> and probability calculations. The convergence of this algorithm is assured for linear and non-linear problems. It offers several possibilities of applications, such as solving algebraic equations, differential equations, probability law of random variables calculations (discrete or continuous), topological analysis, numerical Lebesgue integrals computations, data analysis such as principal components analysis, and parameters identification.

## ACKNOWLEDGMENTS

We thank Michel Gondran, Luc Jaulin and Jean-François Osselin for useful and interesting discussions.

---

\* kenoufi@s-core.fr, abdel.kenoufi@uha.fr

† nicolas.goze@developpement-durable.gouv.fr

‡ michel.goze@uha.fr

<sup>1</sup> Warmus, M.: Calculus of Approximations. Bull. Acad. Pol. Sci., Cl. III, 4(5): 253-259, 1956.

<sup>2</sup> Warmus, M.: Approximations and Inequalities in the Calculus of Approximations: Classification of Approximate Numbers. Bull. Acad. Pol. Sci., Cl. III, 9(4): 241-245, 1961.

<sup>3</sup> Sunaga, T.: Theory of Interval Algebra and its Applications to Numerical Analysis. RAAG Memoirs, 2, 1958, pp. 29-46.

<sup>4</sup> Moore, R. E.: Interval Analysis. Prentice Hall, Englewood Cliffs, N. J., 1966.

<sup>5</sup> Ortolof, H. -J.: Eine Verallgemeinerung der Intervallarimetik. Gesellschaft fuer Mathematik und Datenverarbeitung, Bonn 11, 1969, pp. 1-71.

<sup>6</sup> Kaucher, E.: Ueber metrische und algebraische Eigenschaften einiger beim numerischen Rechnen auftretender Raume. Dissertation, Universitaet Karlsruhe, 1973.

<sup>7</sup> Kaucher, E.: Algebraische Erweiterungen der Intervallrechnung unter Erhaltung der Ordnungs und Verbandstrukturen Computing, Suppl. 1, 1977, pp. 65-79.

<sup>8</sup> Kaucher, E.: Ueber Eigenschaften und en der Anwendungsmoeglichkeiten der erweiterten Intervallrechnung und des hyperbolischen Fastkoerpers ueber R Computing, Suppl. 1, 1977, pp. 81-94.

<sup>9</sup> Kaucher, E.: Interval Analysis in the Extended Interval Space IR Computing, Suppl. 2, 1980, pp. 33-49.

<sup>10</sup> Gardenes, E.; Trepata, A.: Fundamentals of SIGLA, an Interval Computing System over the Completed Set of Intervals. Computing, 24, 1980, pp. 161-179.

<sup>11</sup> Gardenes, E.; Trepata, A.; Janer, J. M.: SIGLA-PL/1 Development and Applications. In: Nickel, K.: Interval Mathematics 1980, Academic Press, 1980, pp. 301-315.

- <sup>12</sup> Gardenes, E.; Trepát, A.; Janer, J. M.: Approaches to Simulation and to the Linear Problem in the SIGLA System. Freiburger Interval-Berichte 81/8, 1981, pp. 1-28.
- <sup>13</sup> Gardenes, E.; Trepát, A.; Mieglo H.: Present Perspective of the SIGLA Interval System. Freiburger Interval-Berichte 82/9, 1982, pp. 1-65.
- <sup>14</sup> Gardenes, E.; Mielgo, H.; Trepát, A.: Modal intervals: Reason and Ground Semantics, In: K. Nickel (ed.): Interval Mathematics 1985, Lecture Notes in Computer Science, Vol. 212, Springer-Verlag, Berlin, Heidelberg, 1986, pp. 27-35.
- <sup>15</sup> E. Gardenes, M.A. Sainz, L. Jorba, R. Calm, R. Estela, H. Mielgo, A. Trepát: Modal Intervals, Reliable Computing 7, 2001, pp. 77-111.
- <sup>16</sup> M. Markov. *Isomorphic Embeddings of Abstract Interval Systems*. Reliable Computing 3: 199–207, 1997.
- <sup>17</sup> Markov, S. M.: Extended Interval Arithmetic Involving Infinite Intervals. Mathematica Balkanica, New Series, 6, 3, 1992, pp. 269-304.
- <sup>18</sup> Markov, S. M.: On Directed Interval Arithmetic and its Applications. J. UCS, 1, 7, 1995, pp. 510-521.
- <sup>19</sup> Markov, S. M.: On the Foundations of Interval Arithmetic. In: Alefeld, G.; Frommer, A.; Lang, B. (Eds.): Scientific Computing and Validated Numerics Akademie Verlag, Berlin, pp. 507-513.
- <sup>20</sup> Dimitrova, N.; Markov, S. M.; Popova, E.: Extended Interval Arithmetics: New Results and Applications. In Atanassova, L.; Herzberger, J. (Eds.): Computer Arithmetic and Enclosure Methods. Elsevier Sci. Publishers B. V., 1992, pp. 225-232.
- <sup>21</sup> Popova, E. D.: Multiplication Distributivity of Proper and Improper Intervals. Reliable Computing 7, 2, 2001, pp.129-140.
- <sup>22</sup> Popova, E. D.: All about Generalized Interval Distributive Relations. I. Complete Proof of the Relations. Sofia, March 2000.
- <sup>23</sup> E. Popova: On the Efficiency of Interval Multiplication Algorithms. Proceedings of III-rd International Conference “Real Numbers and Computers”, Paris, April 27-29, 1998, 117-132.
- <sup>24</sup> Nicolas Goze, PhD Thesis, “*n*-ary algebras and interval arithmetics”, Université de Haute-Alsace, France, March 2011.
- <sup>25</sup> A. Kenoufi, N. Goze, E. Remm, M. Goze, “A new algebraic and arithmetic framework for interval computations”, arXiv:1107.3667, submitted to Reliable Computing, July 2011.
- <sup>26</sup> R.E. Moore, Interval Analysis I by R.E. Moore with C.T. Yang, LMSD-285875, September 1959, Lockheed Aircraft Corporation, Missiles and Space Division, Sunnyvale, California.
- <sup>27</sup> R.E. Moore, Interval Integrals by R.E. Moore, Wayman Strother and C.T. Yang, LMSD-703073, 1960 Lockheed Aircraft Corporation, Missiles and Space Division, Sunnyvale, California.
- <sup>28</sup> R.E. Moore, Ph.D. Thesis (Stanford, 1962).
- <sup>29</sup> R.E. Moore, A test for existence of solutions to nonlinear systems, SIAM J. Numer. Anal., 14 (4), 611–615, 1977).
- <sup>30</sup> <http://www.python.org>
- <sup>31</sup> Introduction to Interval Analysis (Ramon E. Moore, R. Baker Kearfott, and Michael J. Cloud), SIAM, Philadelphia, January, 2009.
- <sup>32</sup> L. Jaulin, M. Kieffer, O. Didrit and E. Walter. *Introduction to interval analysis*. SIAM. 2009 Applied Interval Analysis. Springer-Verlag, London, 2001.