Spectral determinants and zeta functions of Schrödinger operators on metric graphs

JM Harrison¹, K Kirsten¹ and C Texier^{2,3}

- ¹ Department of Mathematics, Baylor University, Waco, TX 76798, USA
- ² Univ. Paris Sud; CNRS; LPTMS, UMR 8626, Bât. 100, F-91405 Orsay, France
- ³ Univ. Paris Sud; CNRS; LPS, UMR 8502, Bât. 510, F-91405 Orsay, France

 $E-mail: \verb|jon_harrison@baylor.edu|, klaus_kirsten@baylor.edu|, christophe.texier@u-psud.fr\\$

Abstract. A derivation of the spectral determinant of the Schrödinger operator on a metric graph is presented where the local matching conditions at the vertices are of the general form classified according to the scheme of Kostrykin and Schrader. To formulate the spectral determinant we first derive the spectral zeta function of the Schrödinger operator using an appropriate secular equation. The result obtained for the spectral determinant is along the lines of the recent conjecture [32].

AMS classification scheme numbers: 34B45, 81Q10, 81Q35

Submitted to: J. Phys. A: Math. Gen.

1. Introduction

Quantum graphs have become a widely employed model for a range of phenomena in mathematical physics from Anderson localization and quantum chaos to problems in mesoscopic physics, microelectronics and the theory of waveguides, see e.g. [27, 29]. Much of the success for the analysis of these problems on graphs can be attributed to the existence of well behaved spectral functions. Consequently spectral determinants and trace formulae in graphs have become important problems widely studied in the literature [2, 3, 4, 5, 7, 9, 8, 10, 11, 13, 15, 18, 19, 23, 26, 30, 33]. Here we consider the Schödinger operator $-\Delta + V(x)$ on a metric graph where each bond of the B bonds is associated with an interval $[0, L_b], b = 1, \ldots, B$. Matching conditions at the vertices of the graph are parameterized by $2B \times 2B$ matrices A and B according to the scheme introduced by Kostrykin and Schrader [24], see Section 2. We also restrict the possible matching conditions to local matching conditions which both respect the graph topology, only relating values of the functions and their derivatives at the ends of intervals where they meet at a vertex, and are independent of the metric structure of the graph, namely the bond lengths $\{L_b\}_{b=1,\dots,B}$ and the energy. The matrices A and B can thus be considered to define the graph topology as they record how the bonds are connected.

In particular we will be concerned with the recent broad conjecture, introduced by one of the authors in [32], for the *spectral determinant* which is defined formally as

$$S(\gamma) = \det[\gamma - \Delta + V(x)] = \prod_{j=0}^{\infty} (\gamma + E_j) , \qquad (1)$$

where $0 < E_0 \le E_1 \le \ldots$ is the spectrum of the Schrödinger operator and γ is some spectral parameter; the condition on the positivity of the spectrum can be relaxed by accepting additional technicalities [21]. The first result for the spectral determinant of the Schrödinger operator on a graph with general vertex matching conditions was obtained by Desbois [9]; the method was, however, unable to determine the γ -independent prefactor of the determinant.

In order to regularize the formal definition of $S(\gamma)$ we employ the *spectral zeta* function,

$$\zeta(s) = \sum_{j=0}^{\infty} (\gamma + E_j)^{-s} . \tag{2}$$

The zeta function of the Schrödinger operator can be formulated by extending a technique introduced for graph Laplacians by some of the authors [17, 18]. Having derived the zeta function the regularized spectral determinant is then defined by $S(\gamma) = \exp(-\zeta'(0))$. More precisely we prove the following theorem along the lines of the conjecture.

Theorem 1. For the Schrödinger operator on a graph with local vertex matching conditions defined by a pair of matrices \mathbb{A} and \mathbb{B} , with $\mathbb{AB}^{\dagger} = \mathbb{BA}^{\dagger}$ and rank $(\mathbb{A}, \mathbb{B}) = 2B$, and potential functions $V_b(x_b) \in C^{\infty}$ for b = 1, ..., B the spectral determinant is

$$S(\gamma) = \left(\prod_{b=1}^{B} \frac{-2}{f_b'(L_b; \gamma)}\right) \frac{\det\left(\mathbb{A} + \mathbb{B}M(\gamma)\right)}{c_N \gamma^P} .$$

 $f_b'(x;\gamma)$ and $M(\gamma)$ are defined in terms of the solution to a boundary value problem on the interval $[0, L_b]$ described in Section 4. c_N is the coefficient of the leading order $t \to \infty$ asymptotic behavior of $\det(\mathbb{A} + \mathbb{B}M(t^2))$, determined in Section 5.1. Furthermore the exponent P characterizes the $t \to 0$ behavior of the determinant; P = 0 if $\det(\mathbb{A} + \mathbb{B}M(0)) \neq 0$, the generic case, otherwise $\det(\mathbb{A} + \mathbb{B}M(\gamma)) \sim \gamma^P$ as $\gamma \to 0$.

To complete the structure of the article, in Section 2 we define the terminology of the quantum graph model. Section 3 introduces the technique, followed subsequently, in the simple case of the Schrödinger operator on an interval with Dirichlet boundary conditions and a graph with Dirichlet conditions at the ends of every bond. To derive the zeta function of a general graph we start from a secular equation whose solutions are $\sqrt{E_j}$; an appropriate secular equation for the Schrödinger operator is identified in Section 4. In Section 5 we formulate the graph zeta function with general local matching conditions at the vertices. Finally, in Section 6, we apply the results for the zeta function to prove Theorem 1. The Appendix shows the derivation of the $t \to \infty$ asymptotics of the function $f'_b(x; t^2)$.

2. Graph model

A graph G is a collection of vertices $v=1,\ldots,V$ and bonds $b=1,\ldots,B$. Each bond connects a pair of vertices b=(v,w) and we denote by o(b)=v the initial vertex of b and t(b)=w the terminal vertex. $\overline{b}=(w,v)$ will denote the reversed bond with the initial and terminal vertices exchanged. We consider undirected graphs with B undirected bonds, so b and \overline{b} refer to the same physical bond but the availability of two labels for each bond will be notationally convenient. To be explicit the label b will refer to the bond with o(b) < t(b). m_v will denote the number of bonds meeting at a vertex v, the valency of v. To determine the valency b and \overline{b} refer to a single bond.

In order to define a metric graph we associate with each bond b an interval $[0, L_b]$ where L_b is the length of b and $x_b = 0$ at o(b) and $x_b = L_b$ at t(b). So x_b is the distance to a point in the interval measured from the initial vertex o(b) and we also use $x_{\overline{b}} = L_b - x_b$ for the distance to the same point measured from the terminal vertex of b. The total length of G is denoted by $\mathcal{L} = \sum_{b=1}^{B} L_b$. A function ψ on G is defined by specifying the set of functions $\{\psi_b(x_b)\}_{b=1,\dots,B}$ on the collection of intervals. The redundancy of notation enforces the relation $\psi_b(x_b) = \psi_{\overline{b}}(x_{\overline{b}}) = \psi_{\overline{b}}(L_b - x_b)$. The Hilbert space of G is consequently

$$\mathcal{H} = \bigoplus_{b=1}^{B} L^2([0, L_b]) . \tag{3}$$

Motivated by physical applications we consider here Schrödinger operators on the

metric graph. The eigenproblem on bond b is

$$\left(i\frac{\mathrm{d}}{\mathrm{d}x_b} + A_b\right)^2 \psi_b(x_b) + V_b(x_b)\psi_b(x_b) = k^2 \psi_b(x_b) . \tag{4}$$

The set $\{A_1, \ldots, A_B\}$ defines a vector potential on the graph constant on each bond.† For consistency $A_{\overline{b}} = -A_b$ as the direction is reversed when changing coordinate.

3. Dirichlet determinants

3.1. Laplace operator on a finite interval

We first analyse the determinant of the Laplace operator on a single finite interval [0, L] with Dirichlet boundary conditions, a *wire*. The spectrum of the wire is $E_j = (\frac{j\pi}{L})^2$ for $j \in \mathbb{N}^*$ and we can compute the zeta-function directly,

$$\zeta_{\text{Dir}}(s) = \sum_{j=1}^{\infty} E_j^{-s} = \left(\frac{L}{\pi}\right)^{2s} \zeta_R(2s) . \tag{5}$$

Differentiating,

$$\zeta_{\text{Dir}}'(0) = 2\zeta_R'(0) + 2\zeta_R(0)\log(L/\pi) = -\log 2 - \log L , \qquad (6)$$

where we have used $\zeta_R(0) = -\frac{1}{2}$ and $\zeta_R'(0) = -\frac{1}{2}\log(2\pi)$ [16]. We therefore obtain the spectral determinant for $\gamma = 0$,

$$S_{\text{Dir}}(0) = \det(-\Delta) = \exp\left(-\zeta'_{\text{Dir}}(0)\right) = 2L , \qquad (7)$$

which agrees with the formulation of the spectral determinant of the Laplace operator in [18, 20, 32].

3.2. Schrödinger operator on a finite interval

To compute the spectral determinant of the operator $-\frac{d^2}{dx^2} + V(x)$ for $x \in [0, L]$ with Dirichlet boundary conditions let us introduce two useful basis for the solutions of the differential equation

$$\left(-\frac{\mathrm{d}^2}{\mathrm{d}x^2} + V(x)\right)\psi(x) = k^2\psi(x). \tag{8}$$

The first basis shown to be useful in [8, 9] is: $f(x; -k^2)$ is a solution of (8) such that $f(0; -k^2) = 1$ and $f(L; -k^2) = 0$, with the second linearly independent solution denoted $\bar{f}(\bar{x}; -k^2)$ where $\bar{x} = L - x$ and \bar{f} satisfies $\bar{f}(0; -k^2) = 1$ and $\bar{f}(L; -k^2) = 0$. For a symmetric potential, V(x) = V(L - x) and we have $\bar{f}(x; -k^2) = f(x; -k^2)$. An alternative basis was employed in [14]: $u_k(x; -k^2)$ and $\bar{u}(\bar{x}; -k^2)$ are solutions satisfying $u(0; -k^2) = 0$ and $u'(0; -k^2) = 1$, where prime denotes derivation with respect to the

† A vector potential $A_b(x_b)$ that depends on the position on the bond can be made constant on each bond by a gauge transformation.

spatial coordinate x, i.e. first argument of the functions. The relation between the two basis is given by

$$\bar{u}(\bar{x}; -k^2) = -\frac{f(x; -k^2)}{f'(L; -k^2)}$$
 or $f(x; -k^2) = \frac{\bar{u}(\bar{x}; -k^2)}{\bar{u}(L; -k^2)}$. (9)

A secular equation whose solutions k are square roots of the eigenvalues of the Schrödinger operator is then

$$u(L; -k^2) = 0; (10)$$

note that the secular equation may be written in different forms, thanks to the relations $u(L;-k^2)=\bar{u}(L;-k^2)=-1/f'(L;-k^2)=-1/\bar{f}'(L;-k^2)$, that hold for an arbitrary potential. Note, however, that $u(x;\gamma)$ and $\bar{u}(x;\gamma)$ do not coincide in general (apart for a symmetric potential V(x)=V(L-x)), nevertheless $u(L;\gamma)=\bar{u}(L;\gamma)$ always holds. This may be shown by computing the Wronskian of the two solutions at the ends of the interval.

Example 1: For a free particle where V(x) = 0,

$$f(x; -k^2) = \bar{f}(x; -k^2) = \frac{\sin k(L-x)}{\sin kL}$$
 and $u(x; -k^2) = \bar{u}(x; -k^2) = \frac{1}{k}\sin kx$. (11)

The Dirichlet spectrum is then given by solutions of the secular equation

$$F(k) = u(L; -k^2) = \frac{-1}{f'(L; -k^2)} = \frac{1}{k} \sin kL = 0.$$
 (12)

Example 2: In the case of a linear potential $V(x) = \omega x$ we find

$$u(x; -k^{2}) = \frac{\pi}{\omega^{1/3}} \left[\operatorname{Ai}(-\omega^{-2/3}k^{2}) \operatorname{Bi}(\omega^{1/3}(x - k^{2}/\omega)) - \operatorname{Bi}(-\omega^{-2/3}k^{2}) \operatorname{Ai}(\omega^{1/3}(x - k^{2}/\omega)) \right] , \qquad (13)$$
$$\bar{u}(\bar{x}; -k^{2}) = \frac{\pi}{\omega^{1/3}} \left[\operatorname{Bi}(\omega^{1/3}(L - k^{2}/\omega)) \operatorname{Ai}(\omega^{1/3}(x - k^{2}/\omega)) - \operatorname{Ai}(\omega^{1/3}(L - k^{2}/\omega)) \operatorname{Bi}(\omega^{1/3}(x - k^{2}/\omega)) \right] , \qquad (14)$$

with the Airy functions Ai(z) and Bi(z) [1]. The Dirichlet spectrum is given by solving the secular equation

$$F(k) = \text{Ai}(-\omega^{-2/3}k^2)\text{Bi}(\omega^{1/3}(L - k^2/\omega)) - \text{Bi}(-\omega^{-2/3}k^2)\text{Ai}(\omega^{1/3}(L - k^2/\omega)) = 0. (15)$$

Example 3: Another interesting example is the case of a potential of the form $V(x) = \phi(x)^2 + \phi'(x)$. The Hamiltonian

$$H_S = -\frac{\mathrm{d}^2}{\mathrm{d}x^2} + \phi(x)^2 + \phi'(x) \tag{16}$$

describes so-called supersymmetric quantum mechanics [34]. In this case, exploiting the factorization $H_S = Q^{\dagger}Q$ with $Q = -\frac{d}{dx} + \phi(x)$, it is possible to construct explicitly two linearly independent solutions of the differential equation $H_S\psi = 0$. We may choose

 $\psi_0(x) = \exp \int_0^x \phi(y) \, dy$ and $\psi_1(x) = \psi_0(x) \int_x^L \frac{dy}{\psi_0(y)^2}$. We obtain the two useful solutions for k = 0:

$$u(x;0) = \psi_0(0)\psi_0(x) \int_0^x \frac{\mathrm{d}y}{\psi_0(y)^2} \quad \text{and} \quad \bar{u}(\bar{x};0) = \psi_0(L)\psi_0(x) \int_x^L \frac{\mathrm{d}y}{\psi_0(y)^2}$$
(17)

We now explain the general method followed in [18] and in the present article. A secular equation F(k) = 0 can be related to the zeta-function using the argument principle,

$$\zeta_{\text{Dir}}(s,\gamma) = \frac{1}{2i\pi} \int_{\mathcal{C}} (z^2 + \gamma)^{-s} \frac{\mathrm{d}}{\mathrm{d}z} \log (F(z)) \, \mathrm{d}z , \qquad (18)$$

where the contour wraps around the positive real axis enclosing the roots of F(z), the solutions of the secular equation, and avoiding any poles, Figure 1(a).




Figure 1. The contours used to evaluate the graph zeta function, (a) before, and (b) after, the contour deformation. The two branch cuts are represented by hashed lines.

In the present case the function $F(z) = u(L; -z^2)$ does not present poles. However, as there is some freedom in the choice of F(z), in the following section it is convenient to choose a function F for the general graph that will possess poles. Transforming the contour C to the imaginary axis z = it, Figure 1(b), we obtain

$$\zeta_{\text{Dir}}(s,\gamma) = \frac{\sin \pi s}{\pi} \int_{\sqrt{\gamma}}^{\infty} (t^2 - \gamma)^{-s} \, \frac{\mathrm{d}}{\mathrm{d}t} \log \left(u(L; t^2) \right) \, \mathrm{d}t \,, \tag{19}$$

where, assuming $F(0) \neq 0$, the segment between $i\sqrt{\gamma}$ and $-i\sqrt{\gamma}$ does not contribute to the integral as F(z) = F(-z) (which follows from the symmetry of the secular equation); comments about the case F(0) = 0 follow in Section 6. Analysis of the integral shows that this representation of the zeta-function is valid in the strip $1/2 < \Re s < 1$. The restriction to $\Re s > 1/2$ comes from the asymptotic behavior of $u(x; t^2)$ for $t \to \infty$. The asymptotics of the function $u(x; t^2)$ for $t \to \infty$ were evaluated in [14],

$$\log u(L; t^2) \underset{t \to \infty}{\sim} tL - \log 2t + O(t^0). \tag{20}$$

An explicit example of such an expansion is provided in the appendix, see Eq. (A.9). To obtain a representation valid for $s \to 0$ we subtract and add the first two terms in

the asymptotic expansion. Then upon integration we find,

$$\zeta_{\text{Dir}}(s,\gamma) = \frac{\sin \pi s}{\pi} \int_{\sqrt{\gamma}}^{\infty} (t^2 - \gamma)^{-s} \frac{\mathrm{d}}{\mathrm{d}t} \left[\log \left(u(L;t^2) \right) - tL + \log(2t) \right] \, \mathrm{d}t \\
+ L \frac{\Gamma(s - 1/2)}{2\sqrt{\pi}\Gamma(s)} \gamma^{\frac{1}{2} - s} - \frac{1}{2} \gamma^{-s} , \tag{21}$$

valid in the strip $-1/2 < \Re s < 1$. Differentiating

$$\zeta_{\text{Dir}}'(0,\gamma) = -\log\left(u(L;\gamma)\right) - \log(2\sqrt{\gamma}) + \frac{1}{2}\log\gamma = -\log\left(2u(L;\gamma)\right). \tag{22}$$

Consequently the zeta-regularized spectral determinant of a wire with Dirichlet boundary conditions is [22],

$$S_{\text{Dir}}(\gamma) = \exp\left(-\zeta'_{\text{Dir}}(0,\gamma)\right) = 2u(L;\gamma) = \frac{-2}{f'(L;\gamma)}.$$
 (23)

Example 1: In the absence of a potential we obtain

$$S_{\text{Dir}}(\gamma) = \frac{2\sinh\sqrt{\gamma}L}{\sqrt{\gamma}}.$$
 (24)

As $\gamma \to 0$ this expression agrees with (7).

Example 2: Using (13), the spectral determinant of $H = -\frac{d^2}{dx^2} + \omega x$ therefore is $S_{\text{Dir}}(\gamma) = \frac{2\pi}{\omega^{1/3}} \left[\text{Ai}(\omega^{-2/3}\gamma) \, \text{Bi}(\omega^{1/3}(L + \gamma/\omega)) - \text{Bi}(\omega^{-2/3}\gamma) \, \text{Ai}(\omega^{1/3}(L + \gamma/\omega)) \right] .$ (25)

Example 3: Using the expression (17) we can obtain an explicit formula for the spectral determinant of the supersymmetric Hamiltonian for vanishing spectral parameter:

$$S_{\text{Dir}}(0) = 2\,\psi_0(0)\psi_0(L)\int_0^L \frac{\mathrm{d}y}{\psi_0(y)^2}\,,\tag{26}$$

i.e.

$$\det\left(-\frac{\mathrm{d}^2}{\mathrm{d}x^2} + \phi^2 + \phi'\right) = 2e^{\int_0^L \phi(x) \, \mathrm{d}x} \int_0^L e^{-2\int_0^x \phi(y) \, \mathrm{d}y} \, \mathrm{d}x.$$
 (27)

Note that this explicit formula furnishes another way to obtain the determinant $\det(\gamma - \frac{d^2}{dx^2} + V)$ in terms of one solution among a family of solutions of a Riccati equation $\phi'(x) = \gamma - \phi(x)^2 + V(x)$. For example, in the free case, V(x) = 0, the Riccati problem is solved by $\phi(x) = \sqrt{\gamma} \tanh \sqrt{\gamma}(x + x_0)$: we can check that (27) leads to (24). Formula (27) may also be illustrated in the case $\phi(x) = \omega x$ (harmonic oscillator on a finite interval),

$$\det\left(-\frac{\mathrm{d}^2}{\mathrm{d}x^2} + \omega^2 x^2 + \omega\right) = \sqrt{\frac{\pi}{\omega}} e^{\frac{1}{2}\omega L^2} \operatorname{erf}(\sqrt{\omega}L), \qquad (28)$$

where $\operatorname{erf}(x)$ is the error function [16]. The limit $L \to \infty$ of (28) is singular and does not approach either $\sqrt{\pi/\omega}$ or $\sqrt{\pi/2\omega}$, the determinant of the operator defined on $\mathbb R$ or $\mathbb R_+$ respectively. This is not unexpected as eigenfunctions on the interval [0,L] will not be square integrable in the limit $L \to \infty$. Eigenfunctions of high enough energy always feel the presence of the boundary conditions at both ends of the interval and thus differ significantly from the infinite space situation.

3.3. Dirichlet determinant of a graph

We are now also in a position to consider the case of a graph with Dirichlet boundary conditions at every vertex. The Dirichlet conditions describe a graph where all the bonds are decoupled and consequently the spectrum is simply the union of the Dirichlet spectra of the individual intervals. Since the zeta-functions of each wire are additive and the determinants are multiplicative in this case we recover a result of [32]

$$S_{\text{Dir}}(\gamma) = \prod_{b=1}^{B} \frac{-2}{f_b'(L_b; \gamma)}.$$
 (29)

To compare with the free case, we use (11) to get $S_{\text{Dir}}(\gamma) = 2^B \gamma^{-B/2} \prod_{b=1}^B \sinh \sqrt{\gamma} L_b$. Taking the limit $\gamma \to 0$ we see (29) reduces to the explicit formula $S_{\text{Dir}}(0) = 2^B \prod_{b=1}^B L_b$ which agrees with (7).

4. Secular equation of a Schrödinger operator with general matching conditions

For a general graph let f_b be the solution of the boundary value problem on the interval $[0, L_b]$,

$$\left(-\frac{\mathrm{d}^2}{\mathrm{d}x_b^2} + V_b(x_b)\right) f_b(x_b; -k^2) = k^2 f_b(x_b; -k^2) , \qquad (30)$$

such that $f_b(0; -k^2) = 1$ and $f_b(L_b; -k^2) = 0$. $f_{\overline{b}}(x_{\overline{b}}; -k^2)$ will denote the complimentary solution on the same bond where the function vanishes at the initial vertex of b and is unity at the terminal vertex, i.e. $f_{\overline{b}}(0; -k^2) = 1$ and $f_{\overline{b}}(L_b; -k^2) = 0.$ ‡ f_b and $f_{\overline{b}}$ are a pair of linearly independent functions on the interval satisfying the eigenvalue problem. Hence the component of the wavefunction of the Schrödinger operator (4) with energy k^2 on bond b can then be written

$$\psi_b(x_b, -k^2) = c_b f_b(x_b; -k^2) e^{iA_b x_b} + c_{\overline{b}} f_{\overline{b}}(x_{\overline{b}}; -k^2) e^{iA_{\overline{b}} x_{\overline{b}}}.$$
(31)

Matching conditions at the graph vertices are specified by a pair of $2B \times 2B$ matrices $\mathbb A$ and $\mathbb B$ via

$$\mathbb{A}\boldsymbol{\psi} + \mathbb{B}\hat{\boldsymbol{\psi}} = \mathbf{0} , \qquad (32)$$

where

$$\psi = (\psi_1(0), \dots, \psi_B(0), \psi_1(L_1), \dots, \psi_B(L_B))^T,$$
(33)

$$\hat{\boldsymbol{\psi}} = (D_1 \psi_1(0), \dots, D_B \psi_B(0), D_{\overline{1}} \psi_1(L_1), \dots, D_{\overline{B}} \psi_B(L_B))^T . \tag{34}$$

 $D_b := \frac{\mathrm{d}}{\mathrm{d}x_b} - \mathrm{i}A_b$ is the covariant derivative and in (33) we consider ψ_b a function of x_b so $x_b = 0$ at o(b) and $x_b = L_b$ at t(b). $\hat{\psi}$ is therefore the vector of inward pointing covariant derivatives at the ends of the intervals.

The following theorem of Kostrykin and Schrader which classifies all matching conditions of self-adjoint realizations of the Laplace operator [24] also applies to the Schrödinger operator.

‡ Notice that it is convenient to consider $f_{\overline{b}}$ as a function of $x_{\overline{b}} = L_b - x_b$.

Theorem 2. The Laplace operator with matching conditions specified by \mathbb{A} and \mathbb{B} is self-adjoint if and only if (\mathbb{A}, \mathbb{B}) has maximal rank and $\mathbb{AB}^{\dagger} = \mathbb{BA}^{\dagger}$.

See [28] for an alternative unique classification scheme for general vertex matching conditions. Note that the general matching conditions we employ can be approximated by ornamenting the vertices of the graph with subgraphs whose vertices have δ -type matching conditions [6].

We consider only local matching conditions where the matrices \mathbb{A} and \mathbb{B} relate values of functions and their derivatives on the intervals where they meet at a vertex and where the matrices are independent of the metric structure of the graph, namely the bond lengths and the spectral parameter k. The restriction on the form of matching conditions is necessary in order to employ the argument principle later.

Representing the components of an energy eigenfunction on G following (31) we see that $\psi = (c_1, \ldots, c_B, c_{\overline{1}}, \ldots, c_{\overline{B}})^T$ and $\hat{\psi} = M(-k^2)\psi$ where $M(-k^2)$ is the $2B \times 2B$ matrix

$$M_{ab} = \delta_{a,b} f_b'(0; -k^2) - \delta_{a,\overline{b}} f_{\overline{b}}'(L_b; -k^2) e^{iA_b L_b} . \tag{35}$$

The vector ψ of coefficients of the two linearly independent solutions on each bond characterizes the wavefunction. Substituting in (32) we see that for ψ to define a wavefunction of energy k^2 satisfying the matching conditions we require

$$\det\left(\mathbb{A} + \mathbb{B}M(-k^2)\right) = 0 , \tag{36}$$

which defines a secular equation for the graph Schrödinger operator.

Example 1: For a free particle $V_b(x_b) = 0$ and we can write f_b explicitly,

$$f_b(x_b; -k^2) = \frac{\sin k(L_b - x_b)}{\sin kL_b}$$
, so $f'_b(x_b; -k^2) = \frac{-k\cos k(L_b - x_b)}{\sin kL_b}$. (37)

Consequently the matrix M takes the form

$$M(-k^2) = -k \begin{pmatrix} \cot k\mathbf{L} & -\csc k\mathbf{L} \\ -\csc k\mathbf{L} & \cot k\mathbf{L} \end{pmatrix} , \tag{38}$$

where $\cot k\mathbf{L} = \operatorname{diag}\{\cot kL_1, \dots, \cot kL_B\}$ and $\operatorname{csc} k\mathbf{L}$ is defined similarly.

5. Zeta function of a Schrödinger operator with general matching conditions

Let us define the function

$$F(z) := \det\left(\mathbb{A} + \mathbb{B}M(-z^2)\right) \tag{39}$$

so the secular equation (36) reads F(k) = 0. As previously the zeta function is written as a contour integral

$$\zeta(s,\gamma) = \frac{1}{2\pi i} \int_{\mathcal{C}} (z^2 + \gamma)^{-s} \frac{\mathrm{d}}{\mathrm{d}z} \log F(z) \,\mathrm{d}z , \qquad (40)$$

where the contour C encloses the zeros of F and avoids its poles, Figure 1(a). Making a contour transformation to the imaginary axis, Figure 1(b), the integral breaks into two parts

$$\zeta(s,\gamma) = \zeta_{\text{Im}}(s,\gamma) + \zeta_{\text{P}}(s,\gamma) , \qquad (41)$$

where we denote by ζ_{Im} the integral along the imaginary axis and ζ_{P} is the series that results from subtracting the residue $(z_0^2 + \gamma)^{-s}$ at each pole z_0 of F. Writing z = it the imaginary axis integral becomes

$$\zeta_{\text{Im}}(s,\gamma) = \frac{\sin \pi s}{\pi} \int_{\sqrt{\gamma}}^{\infty} (t^2 - \gamma)^{-s} \frac{\mathrm{d}}{\mathrm{d}t} \log F(\mathrm{i}t) \,\mathrm{d}t , \qquad (42)$$

which converges in the strip $1/2 < \Re s < 1$.

As poles of F lie at eigenvalues of the graph Schrödinger operator with Dirichlet boundary conditions we see that $\zeta_{\rm P}(s,\gamma)=\zeta_{\rm Dir}(s,\gamma)$, which we evaluated previously. It remains to formulate the imaginary axis integral so that the zeta function given in (42) is well defined to the left of the line $\Re s=1/2$.

5.1. Large t asymptotics of F(it)

To extend the zeta function to the left of the line $\Re s = 1/2$ we require the large t asymptotics of $F(it) = \det(\mathbb{A} + \mathbb{B}M(t^2))$. The $t \to \infty$ asymptotics of f'_b are analysed in the appendix where we find

$$f_b'(0;t^2) \sim -t + \sum_{j=1}^{\infty} s_{b,j}(0)t^{-j}$$
, (43)

and $f'_b(L_b; t^2)$ vanishes exponentially. The functions $s_{b,j}(x_b)$ are determined iteratively from the potential $V_b(x_b)$ via a recursion relation (A.4).

Using (43) we can write the expansion of $M(t^2)$ as $t \to \infty$ up to exponentially suppressed terms.

$$M_{ab} \sim \delta_{a,b}(-t + \sum_{j=1}^{\infty} s_{b,j}(0)t^{-j})$$
 (44)

Hence

$$F(it) \sim \det \left(\mathbb{A} + \mathbb{B}(-t\mathbf{I} + D(t)) \right)$$
 (45)

where

$$D(t) = \sum_{j=1}^{\infty} t^{-j} \operatorname{diag} \left\{ s_{1,j}(0), \dots, s_{B,j}(0), s_{\overline{1},j}(0), \dots, s_{\overline{B},j}(0) \right\} . \tag{46}$$

The expansion of F(it) fits with the free particle case investigated in [18] where, in the absence of a potential, $D(t) = \mathbf{0}$ and $F(it) \sim \det(\mathbb{A} - t\mathbb{B})$.

Using (45) and the recursion relation (A.4) the expansion of F(it) can be carried out to arbitrary order using a computer algebra package. In general the expansion will take the form

$$F(it) \sim \det\left(\mathbb{A} + \mathbb{B}(-t\mathbf{I} + D(t))\right) = \sum_{j=0}^{\infty} c_j t^{2B-j} , \qquad (47)$$

where $c_0 = \det \mathbb{B}$. We denote by c_N the first nonzero coefficient in the expansion of F(it), so N = 0 if $\det \mathbb{B} \neq 0$. To evaluate the zeta-regularized spectral determinant we need a representation of the zeta function valid about s = 0. To find this we only require the leading order in the expansion. Subtracting and adding the leading order behavior we obtain an expression for the imaginary axis integral valid in the strip $-1/2 < \Re s < 1$.

$$\zeta_{\text{Im}}(s,\gamma) = \frac{\sin \pi s}{\pi} \int_{\sqrt{\gamma}}^{\infty} (t^2 - \gamma)^{-s} \frac{d}{dt} \log (F(it)t^{N-2B}/c_N) dt + \frac{1}{2} (2B - N)\gamma^{-s} . \tag{48}$$

Correspondingly

$$\zeta_{\text{Im}}'(0,\gamma) = \log c_N - \log \left(F(i\sqrt{\gamma})\gamma^{\frac{N-2B}{2}} \right) - \frac{2B-N}{2} \log \gamma$$
$$= \log c_N - \log \det \left(\mathbb{A} + \mathbb{B}M(\gamma) \right) . \tag{49}$$

6. Zeta-regularized spectral determinant

Collecting the results so far we can now formulate the spectral determinant, $S(\gamma) = \exp(-\zeta'_{\text{Dir}}(0,\gamma) - \zeta'_{\text{Im}}(0,\gamma))$, of a graph with general local vertex matching conditions, we obtain,

$$S(\gamma) = \frac{S_{\text{Dir}}(\gamma)}{c_N} \det \left(\mathbb{A} + \mathbb{B}M(\gamma) \right) \quad \text{where} \quad S_{\text{Dir}}(\gamma) = \prod_{b=1}^B \frac{-2}{f_b'(L_b; \gamma)} . \tag{50}$$

This is the statement of Theorem 1. The γ -dependent part of this expression coincides with Desbois' result [9] obtained by a different approach. If we choose Dirichlet matching conditions at the graph vertices then $\mathbb{A} = \mathbf{I}_{2B}$ and \mathbb{B} is a matrix of zeros, see (32). Hence $c_{2B} = 1$, which is the only non-zero term in the large t expansion of F(it), and $\det(\mathbb{A} + \mathbb{B}M(\gamma)) = 1$, consequently (50) reduces to (29) as required.

F(z) vanishing at zero. If F(0)=0 the contour integral will pick up contributions associated with the behavior at z=0. While F(0) does not vanish generically, when it does vanish it is straightforward to compensate for this by adjusting the function F. Assume $F(z) \sim z^{2P}$ as $z \to 0$ for $P \in \mathbb{N}^*$.† We may define a new function $\tilde{F}(z) = F(z)/z^{2P}$. Then $\tilde{F}(z) = 0$ is a secular equation but \tilde{F} does not vanish at zero. Evaluating the zeta function using \tilde{F} the procedure is identical and we find,

$$S(\gamma) = \frac{S_{\text{Dir}}(\gamma)}{c_N \gamma^P} \det \left(\mathbb{A} + \mathbb{B}M(\gamma) \right) . \tag{51}$$

The same technique was used, for example, to evaluate the zeta function of the Laplacian on a star graph with Neumann like matching conditions in [18]. To determine P in a particular case one requires the asymptotic expansion of $f'_b(0, t^2)$ and $f'_b(L_b, t^2)$ as $t \to 0$. We now discuss the spectral determinant in two particular cases.

† The even power is due to the symmetry of F.

Functions continuous at the vertices. We consider local vertex matching conditions for which the wavefunction is continuous at the vertices, namely the δ -type conditions. At a vertex v the local δ -type matching conditions relate the functions and covariant derivatives at the ends of the intervals meeting at v. One explicit choice of $m_v \times m_v$ matrices to encode such matching conditions is

$$\mathbb{A}_{v} = \begin{pmatrix}
-\lambda_{v} & 0 & 0 & \cdots & 0 \\
-1 & 1 & 0 & \cdots & 0 \\
0 & -1 & 1 & \cdots & 0 \\
\vdots & \vdots & & & \vdots \\
0 & 0 & 0 & \cdots & 1
\end{pmatrix} \quad \text{and} \quad \mathbb{B}_{v} = \begin{pmatrix}
1 & 1 & \cdots & 1 & 1 \\
0 & 0 & \cdots & 0 & 0 \\
0 & 0 & \cdots & \cdots & 0 \\
\vdots & \vdots & & & \vdots \\
0 & 0 & \cdots & \cdots & 0
\end{pmatrix} \tag{52}$$

where λ_v is the strength of the coupling at v. We note that $\det(\mathbb{A}_v - t\mathbb{B}_v) = -(\lambda_v + m_v t)$. By relabeling the 2B ends of the intervals we may write the matrices \mathbb{A} and \mathbb{B} describing matching conditions on the whole graph so the matrices \mathbb{A}_v and \mathbb{B}_v for $v = 1, \ldots, V$ appear as blocks on the diagonal. We recall that the factor c_N appearing in the spectral determinant is the coefficient of the highest power of t in the expansion of $\det(\mathbb{A} + \mathbb{B}(-t\mathbf{I} + D(t))$, where D(t) is a diagonal matrix where each non-zero element is a series in inverse powers of t. Hence we see that for a graph with δ -type conditions at all the vertices and any potentials on the bonds

$$\det(\mathbb{A} + \mathbb{B}M(\gamma)) \underset{t \to \infty}{\sim} \det(\mathbb{A} + \mathbb{B}(-t\mathbf{I} + D(t)) \underset{t \to \infty}{\sim} \left(\prod_{v} -m_{v}\right) t^{V} + O(t^{V-1}). \tag{53}$$

Thus we find N = V and $c_V = \prod_v (-m_v)$. Consequently

$$S(\gamma) = \left(\prod_{b=1}^{B} \frac{-2}{f_b'(L_b; \gamma)}\right) \frac{\det(\mathbb{A} + \mathbb{B}M(\gamma))}{\prod_v(-m_v)},$$
 (54)

reproducing the result in [32]. Note that the choice of a wavefunction continuous at the vertices allows one to introduce vertex variables and simplify the spectral determinant by replacing $\det(\mathbb{A} + \mathbb{B}M(\gamma))$ by a matrix coupling vertices of smaller dimension [2, 9, 32].

Functions whose derivative is continuous at the vertices. Another interesting case of matching conditions, studied in [31, 32], are the so called δ'_s -type matching conditions corresponding to a wavefunction whose derivative is continuous at the vertices [12]. A possible choice of matching matrices at a vertex v is

$$\mathbb{A}_{v} = \begin{pmatrix}
1 & 1 & \cdots & 1 & 1 \\
0 & 0 & \cdots & 0 & 0 \\
0 & 0 & \cdots & \cdots & 0 \\
\vdots & \vdots & & & \vdots \\
0 & 0 & \cdots & \cdots & 0
\end{pmatrix} \quad \text{and} \quad \mathbb{B}_{v} = \begin{pmatrix}
-\mu_{v} & 0 & 0 & \cdots & 0 \\
-1 & 1 & 0 & \cdots & 0 \\
0 & -1 & 1 & \cdots & 0 \\
\vdots & \vdots & & & \vdots \\
0 & 0 & 0 & \cdots & 1
\end{pmatrix} .$$
(55)

So $\det(-\frac{1}{t}\mathbb{A}_v + \mathbb{B}_v) = -(\mu_v + \frac{1}{t}m_v)$, therefore

$$\det(\mathbb{A} + \mathbb{B}M(\gamma)) \underset{t \to \infty}{\sim} t^{2B-V} \prod_{v} (-m_v) \qquad \text{for } \mu_v = 0 \,\forall v \,. \tag{56}$$

The exponent N=2B-V and the coefficient $c_{2B-V}=\prod_v(-m_v)$. We again obtain (54).

Comparison with the conjecture of [32]. The zeta-regularized spectral determinant was obtained in [32] in the two particular cases considered above with a different method. Based on the observation that both sets of matching conditions led to Eq. (54) it was conjectured that this result is of greater generality (Eq. (61) of this reference). In light of the present work we now inject two observations.

- The matrices \mathbb{A} and \mathbb{B} describing boundary conditions are not unique. Any transformation $(\mathbb{A}, \mathbb{B}) \to (U\mathbb{A}, U\mathbb{B})$ where U is a matrix such that $\det U \neq 0$, leaves the nature of the matching conditions invariant. Therefore the general expression of the zeta-regularized spectral determinant should be invariant under such a transformation. It is clear that this is not the case for the form (54) conjectured in [32] whereas our main result Theorem 1 is invariant.
- The conjecture was based on a continuity argument with respect to variations of \mathbb{A} and \mathbb{B} . However, in general, c_N may depend on the particular choice of the matching conditions rather than simply the topology of the graph. In particular, it is not possible to rule out, a priori, that for a given choice of matching conditions and potentials a term in the series expansion (47) vanishes. We have seen in the previous pair of examples that the exponent N does indeed vary with the nature of the matching conditions.

Acknowledgments

The authors would like to thank Alain Comtet, Sebastian Eggers né Endres, Guglielmo Fucci, Frank Steiner and Yves Tourigny for stimulating discussions. KK is supported by National Science Foundation grant PHY-0757791. JMH was supported by the Baylor University summer sabbatical program.

Appendix. Large t asymptotics of $f'(x; t^2)$

 $f(x; t^2)$ is the solution of

$$\left(-\frac{\mathrm{d}^2}{\mathrm{d}x^2} + V(x) + t^2\right) f(x; t^2) = 0 \tag{A.1}$$

on the interval [0, L] with $f(0; t^2) = 1$ and $f(L; t^2) = 0$. To find the asymptotics of $f'(x; t^2)$ in the limit $t \to \infty$ we let $\phi_t(x)$ be a general solution of (A.1) and call $S_t(x) = \frac{d}{dx} \log \phi_t(x)$. S_t satisfies the differential equation

$$S'_t(x) = t^2 + V(x) - S_t^2(x)$$
 (A.2)

Consequently, in the limit $t \to \infty$, \mathcal{S}_t has an expansion of the form

$$S_t(x) = \sum_{j=-1}^{\infty} s_j(x) t^{-j} . \tag{A.3}$$

The functions $s_i(x)$ are determined by the recurrence relation

$$s_{j+1}(x) = \mp \frac{1}{2} \left(s'_j(x) + \sum_{k=0}^j s_k(x) s_{j-k}(x) \right)$$
(A.4)

where $s_{-1}(x) = \pm 1$, $s_0(x) = 0$ and $s_1(x) = \pm \frac{1}{2}V(x)$. The two alternative signs in the expansion produce exponentially growing and decaying solutions of (A.1). If we denote with $\mathcal{S}_t^{\pm}(x)$ the functions with $s_{-1}(x) = \pm 1$ respectively then we may write the large t behavior of $f(x; t^2)$ as a linear combination of the exponentially growing and decaying parts,

$$f(x;t^2) = A^+ \exp \int_0^x \mathcal{S}_t^+(u) \, \mathrm{d}u + A^- \exp \int_0^x \mathcal{S}_t^-(u) \, \mathrm{d}u \ . \tag{A.5}$$

Imposing the boundary conditions we require $A^- = 1 - A^+$ and as it turns out

$$A^{\pm} = \frac{\mp \exp \int_0^L \mathcal{S}_t^-(u) \, \mathrm{d}u}{\exp \int_0^L \mathcal{S}_t^+(u) \, \mathrm{d}u - \exp \int_0^L \mathcal{S}_t^-(u) \, \mathrm{d}u} . \tag{A.6}$$

Consequently differentiating (A.5) we find,

$$f'(x;t^{2}) = \frac{S_{t}^{-}(x) \exp(\int_{0}^{x} S_{t}^{-}(u) du + \int_{0}^{L} S_{t}^{+}(u) du) - S_{t}^{+}(x) \exp(\int_{0}^{x} S_{t}^{+}(u) du + \int_{0}^{L} S_{t}^{-}(u) du)}{\exp \int_{0}^{L} S_{t}^{+}(u) du - \exp \int_{0}^{L} S_{t}^{-}(u) du}.$$
(A.7)

We are interested in the $t \to \infty$ asymptotics of both $f'(0; t^2)$ and $f'(L; t^2)$. From (A.7) we see that at x = L, $f'(x; t^2)$ vanishes exponentially. At x = 0, up to exponentially suppressed terms, we have the asymptotic relation

$$f'(0;t^2) \sim \mathcal{S}_t^-(0) = -t + \sum_{j=1}^{\infty} s_j(0)t^{-j}$$
 (A.8)

Example: In the case of the linear potential $V(x) = \omega x$ considered earlier the first eight functions s_j in the expansion of \mathcal{S}_t^- constructed iteratively from the recurrence relation are shown bellow.

$$s_{-1}(x) = -1 s_0(x) = 0 s_1(x) = -\frac{\omega}{2}x$$

$$s_2(x) = -\frac{\omega}{2^2} s_3(x) = \frac{\omega^2}{2^3}x^2 s_4(x) = \frac{\omega^2}{2^2}x$$

$$s_5(x) = \frac{3\omega^2}{2^5} - \frac{\omega^3}{2^3}x^3 s_6(x) = -\frac{11\omega^3}{2^5}x^2$$

Consequently

$$f'(0;t^2) \sim -t - \frac{\omega}{2^2 t^2} + \frac{3\omega^2}{2^5 t^5} + \cdots$$
 (A.9)

References

- [1] M Abramowitz and IA Stegun, Handbook of mathematical functions with formulas, graphs, and mathematical tables, (Dover 1965)
- [2] E Akkermans, A Comtet, J Desbois, G Montambaux and C Texier "Spectral determinant on quantum graphs," 2000 Ann. Phys. **284** 10–51
- [3] G Berkolaiko and JP Keating, "Two-point spectral correlations for star graphs," 1999 J. Phys. A 32 7827–7841
- [4] J Bolte and S Endres, "The trace formula for quantum graphs with general self adjoint boundary conditions," 2009 Ann. Henri Poincare 10 189-223 arXiv:0805.3111
- [5] R Carlson, "Eigenvalue problems on directed graphs," 1999 Trans. Amer. Math. Soc. 99 4069–4088
- [6] T Cheon, P Exner and O Turek, "Approximation of a general singular vertex coupling in quantum graphs," 2010 Ann. Phys. 325 548–578
- [7] A Comtet, J Desbois and C Texier, "Functionals of the Brownian motion, localization and metric graphs," 2005 J. Phys. A: Math. Gen. 38 R341–R383
- [8] J Desbois, "Spectral determinant of Schrödinger operators on graphs," 2000 J. Phys. A: Math. Gen. 33 L63–L67
- [9] J Desbois, "Spectral determinant on graphs with generalized boundary conditions," 2001 Eur. Phys. J. B 24 261–266
- [10] S Endres and F Steiner, "The Berry-Keating operator on $L^2(\mathbb{R}_>, \mathrm{d}x)$ and compact quantum graphs with general self-adjoint realizations," 2010 J. Phys. A: Math. Theor. 43 095204 arXiv:0912.3183
- [11] S Endres and F Steiner, "A simple infinite quantum graph," 2009 (Preprint) www.uni-ulm.de/fileadmin/website_uni_ulm/nawi.inst.260/paper/09/tp09-1.pdf
- [12] P Exner, "Lattice Kronig-Penney models," 1995 Phys. Rev. Lett. 74 3503
- [13] L Friedlander, "Determinant of the Schrödinger operator on a metric graph" In: G. Berkolaiko, R. Carlson, S. A. Fulling, and P. Kuchment (Editors) Quantum Graphs and Their Applications, Contemporary Mathematics, 415, (AMS 2006) 151–160
- [14] G Fucci, K Kirsten and P Morales, "Pistons modeled by potentials" In: SD Odintsov, D Saez-Gomez, and S Xambo-Descamps (Eds) Cosmology, Quantum Vacuum and Zeta Functions: In Honour of Emilio Elizalde, (Springer 2011) 313 arXiv:1106.0731
- [15] S Gnutzmann and U Smilansky, "Quantum graphs: Applications to quantum chaos and universal spectral statistics," 2006 Advances in Physics 55 5 527-625
- [16] IS Gradshteyn and IM Ryzhik, *Table of integrals, series and products*, fifth edition (Academic Press 1994)
- [17] JM Harrison and K Kirsten, "Vacuum energy, spectral determinant and heat kernel asymptotics of graph Laplacians with general vertex matching conditions," in KA Milton and M Bordag (Eds) Quantum field theory under the influence of external conditions (QFEXT09), World Scientific, 2010, 421–425 arXiv:0912.0036
- [18] JM Harrison and K Kirsten, "Zeta functions of quantum graphs," 2011 J. Phys. A: Math. Theor. 44 235301 arXiv:0911.2509
- [19] JP Keating, J Marklof and B Winn, "Value distribution of the eigenfunctions and spectral determinants of quantum star graphs," 2003 Comm. Math. Phys. 241, 421–452
- [20] K Kirsten and P Loya, "Computation of determinants using contour integrals," 2008 Am. J. Phys. 76 60–64 arXiv:0707.3755v1
- [21] K Kirsten and AJ McKane, "Functional determinants for general Sturm-Liouville problems," 2004 J. Phys. A: Math. Gen. 37 4649–4670
- [22] K Kirsten and AJ McKane, "Functional determinants in the presence of zero modes," in KA Milton (Ed), Quantum Field theory under the influence of external conditions, Rinton Press, 2004, 146–151
- [23] V Kostrykin, J Potthoff and R Schrader, "Heat kernels on metric graphs and a trace formula,"

- in F Germinet and PD Hislop (Eds), Adventures in mathematics, Contemporary Mathematics 447, AMS, 2007, 175–198
- [24] V Kostrykin and R Schrader, "Kirchhoff's rule for quantum wires," 1999 J. Phys. A: Math. Gen. 32 595–630
- [25] T Kottos and U Smilansky, "Quantum chaos on graphs," 1997 Phys. Rev. Lett. 79 4794–4797
- [26] T Kottos and U Smilansky, "Periodic orbit theory and spectral statistics for quantum graphs," 1999 Ann. Phys. 274 76–124
- [27] P Kuchment, "Graph models of wave propogation in thin structures," 2002 Waves Random Media 12 R1–24
- [28] P Kuchment, "Quantum graphs: I Some basic structures," 2004 Waves Random Media 14 S107–28
- [29] P Kuchment, "Quantum graphs: an introduction and a brief survey," in P Exner, JP Keating, P Kuchment, T Sunada, and A Teplyaev (Eds), Analysis on Graphs and Its Applications Proceedings of Symposia in Pure Mathematics, 77, AMS, 2008, 291–312
- [30] J-P Roth, "Le spectre du Laplacien sur un graphe," in G Mokobodzki and D Pinchon (Eds), *Théorie du potentiel*, Lecture Notes in Maths, **1096**, Springer, Berlin, 1984, 521–539
- [31] C Texier, "On the spectrum of the Laplace operator of metric graphs attached at a vertex Spectral determinant approach," 2008 J. Phys. A: Math. Theor. 41, 085207
- [32] C Texier, "ζ-regularised spectral determinants on metric graphs," 2010 J. Phys. A: Math. Theor. 43 425203 arXiv:1006.2163
- [33] C Texier and G Montambaux, "Quantum oscillations in mesoscopic rings and anomalous diffusion," 2005 J. Phys. A: Math. Gen. 38 3455-71
- [34] E Witten, "Dynamical breaking of supersymmetry," 1981 Nucl. Phys. B 188 513