

# Solution of the momentum-space Schrödinger equation for bound states of the $N$ -dimensional Coulomb problem (revisited)

Radosław Szmytkowski\*

Atomic Physics Division, Department of Atomic Physics and Luminescence,  
Faculty of Applied Physics and Mathematics, Gdańsk University of Technology,  
Narutowicza 11/12, 80–233 Gdańsk, Poland

January 28, 2019

## Abstract

The Schrödinger–Coulomb Sturmian problem in  $\mathbb{R}^N$ ,  $N \geq 2$ , is considered in the momentum representation. An integral formula for the Gegenbauer polynomials, found recently by Cohl [arXiv:1105.2735], is used to separate out angular variables and reduce an integral Sturmian eigenvalue equation in  $\mathbb{R}^N$  to a Fredholm one on  $\mathbb{R}_+$ . A kernel of the latter equation contains the Legendre function of the second kind. A symmetric Poisson-type series expansion of that function into products of the Gegenbauer polynomials, established by Ossicini [Boll. Un. Mat. Ital. 7 (1952) 315], is then used to determine the Schrödinger–Coulomb Sturmian eigenvalues and associated momentum-space eigenfunctions. Finally, a relationship existing between solutions to the Sturmian problem and solutions to a (physically more interesting) energy eigenvalue problem is exploited to find the Schrödinger–Coulomb bound-state energy levels in  $\mathbb{R}^N$ , together with explicit representations of the associated normalized momentum-space Schrödinger–Coulomb Hamiltonian eigenfunctions.

**KEY WORDS:** Coulomb problem; momentum representation; integral equations; Gegenbauer polynomials; Legendre functions

**PACS:** 03.65.Ge, 02.30.Gp, 02.30.Rz

## 1 Introduction

A closed-form expression for the bound-state hydrogenic wave functions in the momentum representation in  $\mathbb{R}^3$  was first obtained in 1929 by Podolsky and Pauling [1], who succeeded to carry out a Fourier transform of the position-space Coulomb eigenfunctions, found three years earlier by Schrödinger [2]. Guided by the results of Ref. [1], in 1932 Hylleraas [3] derived a second-order partial differential equation obeyed by the momentum-space Coulomb wave functions. In 1933, Elsasser [4] attacked the problem in the spirit of Ref. [1], but with the Fourier transform done with the aid of the complex contour-integration technique. A completely different approach to the subject was presented in 1935 by Fock [5]. He showed that the momentum-space Schrödinger–Coulomb wave equation in  $\mathbb{R}^3$ , known to be an *integral* one, might be transformed into a form which appeared to be identical with an equation satisfied on the unit hypersphere  $\mathbb{S}^3$  by the hyperspherical harmonics. This fact led Fock to the conclusion that for bound states the symmetry group of the quantum-mechanical Coulomb problem in  $\mathbb{R}^3$  is  $O(4)$ . That thread was pursued further a year later by Bargmann [6]. Since then, the problem was revisited and discussed in various

---

\*Email: radek@mif.pg.gda.pl

aspects by a number of authors. Space limitations allow us to mention here only several representative papers [7–16], as well as relevant reviews and monographs [17–24], where references to other pertinent works may be found in abundance.

In this paper, we present a method which allows one to determine bound-state energy levels and associated Hamiltonian eigenfunctions of the Schrödinger–Coulomb problem in the  $N$ -dimensional ( $N \geq 2$ ) Euclidean momentum space, without making any use of the  $O(N+1)$  dynamical symmetry specific to the problem. We proceed along the following steps. First, we replace the integral energy eigenproblem by a related Sturmian integral eigenvalue equation, in which energy is fixed and the role of the eigenparameter is played by the Coulomb potential strength. Starting with the next movement, our approach seems to be markedly different from what may be found in the existing literature on the subject. We reduce the aforementioned Sturmian integral equation in  $\mathbb{R}^N$  to a Fredholm one on  $\mathbb{R}_+$ , exploiting a recent result from the theory of Gegenbauer polynomials due to Cohl [25]. A kernel in the latter one-dimensional equation contains the Legendre function of the second kind. We solve that integral equation using a Poisson-type expansion of the aforementioned Legendre function in Gegenbauer polynomials, derived long ago by Ossicini [26], but now buried deeply in the mathematical literature and little known. Finally, we use a relationship existing between solutions to the Sturmian problem and to the energy eigenvalue problem, and determine the Schrödinger–Coulomb bound-state energy levels in  $\mathbb{R}^N$ , together with explicit representations of the associated normalized momentum-space Schrödinger–Coulomb Hamiltonian eigenfunctions.

## 2 Integral equation for the momentum-space Schrödinger–Coulomb wave functions

In the position representation, the time-independent Schrödinger equation for a nonrelativistic spinless particle moving in the  $N$ -dimensional ( $N \geq 2$ ) Euclidean space  $\mathbb{R}^N$ , in the field of a force derivable from the scalar local potential  $V(\mathbf{r})$ , is

$$-\frac{\hbar^2}{2m}\nabla^2\Psi(\mathbf{r}) + V(\mathbf{r})\Psi(\mathbf{r}) = E\Psi(\mathbf{r}). \quad (2.1)$$

The counterpart equation in the momentum representation is

$$\left(\frac{p^2}{2m} - E\right)\Phi(\mathbf{p}) + \frac{1}{(2\pi\hbar)^{N/2}} \int_{\mathbb{R}^N} d^N\mathbf{p}' U(\mathbf{p} - \mathbf{p}')\Phi(\mathbf{p}') = 0, \quad (2.2)$$

where  $\Phi(\mathbf{p})$  and  $U(\mathbf{p})$  are the Fourier transforms of  $\Psi(\mathbf{r})$  and  $V(\mathbf{r})$ , respectively. In the case of the attractive Coulomb potential

$$V(\mathbf{r}) = -\frac{Ze^2}{(4\pi\epsilon_0)r} \quad (Z > 0), \quad (2.3)$$

which concerns us here,  $U(\mathbf{p})$  may be shown (cf. Appendix) to be

$$U(\mathbf{p}) = -\frac{Ze^2}{(4\pi\epsilon_0)} \frac{(2\hbar)^{N/2-1}\Gamma\left(\frac{N-1}{2}\right)}{\sqrt{\pi}} \frac{1}{p^{N-1}}, \quad (2.4)$$

so that Eq. (2.2) becomes

$$\left(\frac{p^2}{2m} - E\right)\Phi(\mathbf{p}) = \frac{Ze^2}{(4\pi\epsilon_0)} \frac{\Gamma\left(\frac{N-1}{2}\right)}{2\pi^{(N+1)/2}\hbar} \int_{\mathbb{R}^N} d^N\mathbf{p}' \frac{\Phi(\mathbf{p}')}{|\mathbf{p}' - \mathbf{p}|^{N-1}}. \quad (2.5)$$

We aim at finding solutions to the above equation which are associated with, yet unknown, *negative* (bound-state) energy eigenvalues  $E$ .

### 3 Solution of the integral equation for the momentum-space Schrödinger–Coulomb Sturmian functions

#### 3.1 Preliminaries

Instead of attacking Eq. (2.5) directly, it is easier to consider first the associated Sturmian eigenproblem

$$\left(\frac{p^2}{2m} - E\right) \Sigma(E, \mathbf{p}) = \lambda \frac{Ze^2}{(4\pi\epsilon_0)} \frac{\Gamma\left(\frac{N-1}{2}\right)}{2\pi^{(N+1)/2}\hbar} \int_{\mathbb{R}^N} d^N \mathbf{p}' \frac{\Sigma(E, \mathbf{p}')}{|\mathbf{p}' - \mathbf{p}|^{N-1}}, \quad (3.1)$$

in which  $\lambda \equiv \lambda(E)$  is an eigenvalue, whereas  $E < 0$  is a *fixed* parameter. It is evident that once the  $\lambda$ -spectrum ( $\{\lambda_\alpha\}$ ) for Eq. (3.1) is determined, the sought negative energy eigenvalues to the original problem (2.5) are roots to the algebraic equations

$$\lambda_\alpha(E) = 1 \quad (E < 0). \quad (3.2)$$

If  $E_{\alpha\beta} < 0$  is a particular solution to Eq. (3.2) (the second subscript at  $E$  serves to distinguish between different roots to Eq. (3.2), if there are more such roots than one), the corresponding negative-energy momentum-space Coulomb wave function  $\Phi_{\alpha\beta\gamma}(\mathbf{p})$  is

$$\Phi_{\alpha\beta\gamma}(\mathbf{p}) = A \Sigma_{\alpha\gamma}(E_{\alpha\beta}, \mathbf{p}), \quad (3.3)$$

$A$  being an arbitrary non-zero constant (the second subscript at  $\Sigma$  appears if the eigenvalue  $\lambda_\alpha$  is degenerate).

Before proceeding further, we shall rewrite Eq. (3.1) in a slightly more compact form. To this end, we define

$$q = \sqrt{-2mE}, \quad q_B = \frac{\hbar}{a_B} \quad (3.4)$$

(here  $a_B = (4\pi\epsilon_0)\hbar^2/me^2$  is the Bohr radius). Use of Eq. (3.4) transforms Eq. (3.1) into

$$(p^2 + q^2)\Sigma(E, \mathbf{p}) = \lambda Z q_B \frac{\Gamma\left(\frac{N-1}{2}\right)}{\pi^{(N+1)/2}} \int_{\mathbb{R}^N} d^N \mathbf{p}' \frac{\Sigma(E, \mathbf{p}')}{|\mathbf{p}' - \mathbf{p}|^{N-1}}. \quad (3.5)$$

#### 3.2 Reduction of Eq. (3.5) to the radial form

We try whether Eq. (3.5) has solutions in the form

$$\Sigma_{l\eta}(E, \mathbf{p}) = F_l(E, p) Y_{l\eta}^{(N-1)}(\mathbf{n}_p). \quad (3.6)$$

Here  $Y_{l\eta}^{(N-1)}(\mathbf{n}_p)$ , with  $l \in \mathbb{N}$  and  $\eta \in \{1, 2, \dots, d_l^{(N-1)}\}$ , where

$$d_l^{(N-1)} = (2l + N - 2) \frac{(l + N - 3)!}{l!(N - 2)!}, \quad (3.7)$$

are the hyperspherical harmonics [21, 22], that are orthonormal on the unit hypersphere  $\mathbb{S}^{N-1}$  in the sense of

$$\oint_{\mathbb{S}^{N-1}} d^{N-1} \mathbf{n}_p Y_{l\eta}^{(N-1)*}(\mathbf{n}_p) Y_{l'\eta'}^{(N-1)}(\mathbf{n}_p) = \delta_{ll'} \delta_{\eta\eta'}. \quad (3.8)$$

We insert Eq. (3.6) into Eq. (3.5), multiply the resulting equation by  $Y_{l\eta}^{(N-1)*}(\mathbf{n}_p)$ , and then sum over  $\eta$ . Using the addition theorem [22, Eqs. (3.80), (3.86) and (3.40)]

$$\sum_{\eta=1}^{d_l^{(N-1)}} Y_{l\eta}^{(N-1)*}(\mathbf{n}_p) Y_{l\eta}^{(N-1)}(\mathbf{n}'_p) = \frac{d_l^{(N-1)}}{S_{N-1}} \frac{C_l^{(N/2-1)}(\mathbf{n}_p \cdot \mathbf{n}'_p)}{C_l^{(N/2-1)}(1)}, \quad (3.9)$$

where  $C_n^{(\alpha)}(\xi)$  is the Gegenbauer polynomial and  $S_{N-1}$  is the surface area of  $\mathbb{S}^{N-1}$ , we obtain

$$(p^2 + q^2)F_l(E, p) = \lambda Z q_B \frac{\Gamma\left(\frac{N-1}{2}\right)}{2^{(N-1)/2} \pi^{(N+1)/2} C_l^{(N/2-1)}(1) p^{(N-1)/2}} \int_0^\infty dp' p'^{(N-1)/2} F_l(E, p') \\ \times \oint_{\mathbb{S}^{N-1}} d^{N-1} \mathbf{n}'_p \frac{C_l^{(N/2-1)}(\mathbf{n}_p \cdot \mathbf{n}'_p)}{\left(\frac{p^2 + p'^2}{2pp'} - \mathbf{n}_p \cdot \mathbf{n}'_p\right)^{(N-1)/2}}. \quad (3.10)$$

It is evident that the angular integral appearing in Eq. (3.10) may be expressed as

$$\oint_{\mathbb{S}^{N-1}} d^{N-1} \mathbf{n}'_p \frac{C_l^{(N/2-1)}(\mathbf{n}_p \cdot \mathbf{n}'_p)}{\left(\frac{p^2 + p'^2}{2pp'} - \mathbf{n}_p \cdot \mathbf{n}'_p\right)^{(N-1)/2}} = S_{N-2} \int_{-1}^1 d\xi (1-\xi^2)^{(N-3)/2} \frac{C_l^{(N/2-1)}(\xi)}{\left(\frac{p^2 + p'^2}{2pp'} - \xi\right)^{(N-1)/2}}. \quad (3.11)$$

However, it has been recently shown by Cohl [25] that

$$\int_{-1}^1 d\xi \frac{(1-\xi^2)^{\alpha-1/2} C_n^{(\alpha)}(\xi)}{(z-\xi)^{\kappa+1/2}} = e^{i\pi(\alpha-\kappa)} 2^{\alpha+1/2} C_n^{(\alpha)}(1) \frac{\Gamma(\alpha + \frac{1}{2})}{\Gamma(\kappa + \frac{1}{2})} (z^2 - 1)^{(\alpha-\kappa)/2} \mathfrak{Q}_{n+\alpha-1/2}^{\kappa-\alpha}(z) \\ (\text{Re } \alpha > -\frac{1}{2}, \kappa \in \mathbb{C}, n \in \mathbb{N}, z \in \mathbb{C} \setminus (-\infty, 1]), \quad (3.12)$$

where  $\mathfrak{Q}_\nu^\mu(z)$  is the associated Legendre function of the second kind. It follows from Eq. (3.12) that

$$\int_{-1}^1 d\xi \frac{(1-\xi^2)^{\alpha-1/2} C_n^{(\alpha)}(\xi)}{(z-\xi)^{\alpha+1/2}} = 2^{\alpha+1/2} C_n^{(\alpha)}(1) \mathfrak{Q}_{n+\alpha-1/2}(z) \\ (\text{Re } \alpha > -\frac{1}{2}, n \in \mathbb{N}, z \in \mathbb{C} \setminus (-\infty, 1]) \quad (3.13)$$

( $\mathfrak{Q}_\nu(z) \equiv \mathfrak{Q}_\nu^0(z)$  is the Legendre function of the second kind), and consequently

$$\oint_{\mathbb{S}^{N-1}} d^{N-1} \mathbf{n}'_p \frac{C_l^{(N/2-1)}(\mathbf{n}_p \cdot \mathbf{n}'_p)}{\left(\frac{p^2 + p'^2}{2pp'} - \mathbf{n}_p \cdot \mathbf{n}'_p\right)^{(N-1)/2}} = 2^{(N-1)/2} S_{N-2} C_l^{(N/2-1)}(1) \mathfrak{Q}_{l+(N-3)/2}\left(\frac{p^2 + p'^2}{2pp'}\right). \quad (3.14)$$

Plugging Eq. (3.14) into Eq. (3.10) and making use of the relation

$$S_{N-2} = \frac{(N-1)\pi^{(N-1)/2}}{\Gamma\left(\frac{N+1}{2}\right)}, \quad (3.15)$$

we arrive at the following integral equation for the radial function  $F_l(E, p)$ :

$$(p^2 + q^2)F_l(E, p) = \lambda \frac{2Z q_B}{\pi p^{(N-1)/2}} \int_0^\infty dp' p'^{(N-1)/2} \mathfrak{Q}_{l+(N-3)/2}\left(\frac{p^2 + p'^2}{2pp'}\right) F_l(E, p') \quad (3.16)$$

(for  $N = 3$  and  $\lambda = 1$ , Eq. (3.16) coincides, as it should, with Eq. (8.7) in Ref. [27]). After some straightforward rearrangements, this may be further cast into the form

$$[\sqrt{p^2 + q^2} p^{(N-1)/2} F_l(E, p)] \\ = \lambda \frac{2Z q_B}{\pi} \int_0^\infty dp' \frac{\mathfrak{Q}_{l+(N-3)/2}\left(\frac{p^2 + p'^2}{2pp'}\right)}{\sqrt{p^2 + q^2} \sqrt{p'^2 + q^2}} [\sqrt{p'^2 + q^2} p'^{(N-1)/2} F_l(E, p')], \quad (3.17)$$

the kernel in the above equation being real and manifestly symmetric.

### 3.3 Intermezzo on the spectral theory for Fredholm integral equations

Before we attack Eq. (3.17), we have to recall some very basic facts from the spectral theory for integral equations. If the kernel  $M(p, p')$  in the homogeneous Fredholm equation of the second kind

$$f(p) = \lambda \int_0^\infty dp' M(p, p') f(p') \quad (3.18)$$

is real and symmetric, and if its  $\lambda$ -spectrum (known to be real, in view of the aforementioned assumptions about the kernel) is purely discrete, the following spectral expansion of  $M(p, p')$  holds:

$$M(p, p') = \sum_k \frac{f_k(p) f_k(p')}{\lambda_k}, \quad (3.19)$$

provided the eigenfunctions  $\{f_k(p)\}$  are orthonormal in the sense of

$$\int_0^\infty dp f_k(p) f_{k'}(p) = \delta_{kk'}. \quad (3.20)$$

This implies that if we cope with some particular equation of the form (3.18), and if, in whatever manner and from whichever premises, we infer that the kernel  $M(p, p')$  possesses the series representation

$$M(p, p') = \sum_k \varepsilon_k g_k(p) g_k(p') \quad (\varepsilon_k = \pm 1), \quad (3.21)$$

the functions  $\{g_k(p)\}$  being orthogonal,

$$\int_0^\infty dp g_k(p) g_{k'}(p) = \|g_k\|^2 \delta_{kk'}, \quad (3.22)$$

we may immediately deduce that the eigenvalues for Eq. (3.18) are

$$\lambda_k = \varepsilon_k \|g_k\|^{-2}, \quad (3.23)$$

whereas the associated normalized eigenfunctions are

$$f_k(p) = \frac{g_k(p)}{\|g_k\|} = \sqrt{\varepsilon_k \lambda_k} g_k(p) \quad (3.24)$$

(observe that  $\varepsilon_k^{-1} = \varepsilon_k$ ).

### 3.4 Solution of the radial integral equation (3.17)

We return to Eq. (3.17). It is evident that it falls into the category (3.18), with

$$f_l(p) = \sqrt{p^2 + q^2} p^{(N-1)/2} F_l(E, p) \quad (3.25)$$

(here the subscript  $l$  at  $f$  corresponds to the one at  $F$ ) and

$$M_l(p, p') = \frac{2Zq_B}{\pi} \frac{\Omega_{l+(N-3)/2} \left( \frac{p^2 + p'^2}{2pp'} \right)}{\sqrt{p^2 + q^2} \sqrt{p'^2 + q^2}}. \quad (3.26)$$

An expansion of the kernel (3.26) in the symmetric form (3.21) may be derived from the following Poisson-type identity:

$$\begin{aligned} \Omega_\nu \left( \frac{1 - 2h\xi\xi' + h^2}{2h\sqrt{(1-\xi^2)(1-\xi'^2)}} \right) &= 2^{2\nu+1} [\Gamma(\nu+1)]^2 (1-\xi^2)^{(\nu+1)/2} (1-\xi'^2)^{(\nu+1)/2} \\ &\times \sum_{n_r=0}^{\infty} \frac{n_r!}{\Gamma(n_r+2\nu+2)} h^{n_r+\nu+1} C_{n_r}^{(\nu+1)}(\xi) C_{n_r}^{(\nu+1)}(\xi') \\ &(|h| \leq 1, -1 \leq \xi, \xi' \leq 1), \quad (3.27) \end{aligned}$$

discovered several decades ago by Ossicini [26, Eq. (11)] (there is a misprint in the formula provided originally by Ossicini — on its left-hand side  $[\Gamma(\lambda)]$  should be replaced by  $[\Gamma(\lambda)]^2$ ; moreover, one should be aware that in Ref. [28, Eq. (47.6.11)] the formula in question was reprinted with several errors!). For  $h = 1$ , Eq. (3.27) becomes

$$\begin{aligned} \Omega_\nu \left( \frac{1 - \xi\xi'}{\sqrt{(1 - \xi^2)(1 - \xi'^2)}} \right) &= 2^{2\nu+1} [\Gamma(\nu + 1)]^2 (1 - \xi^2)^{(\nu+1)/2} (1 - \xi'^2)^{(\nu+1)/2} \\ &\times \sum_{n_r=0}^{\infty} \frac{n_r!}{\Gamma(n_r + 2\nu + 2)} C_{n_r}^{(\nu+1)}(\xi) C_{n_r}^{(\nu+1)}(\xi') \\ &(-1 \leq \xi, \xi' \leq 1), \end{aligned} \quad (3.28)$$

from which, with the substitutions

$$\xi = \frac{q^2 - p^2}{q^2 + p^2}, \quad \xi' = \frac{q'^2 - p'^2}{q'^2 + p'^2}, \quad (3.29)$$

we infer that

$$\begin{aligned} \Omega_\nu \left( \frac{p^2 + p'^2}{2pp'} \right) &= 2^{2\nu+1} [\Gamma(\nu + 1)]^2 \left( \frac{2qp}{q^2 + p^2} \right)^{\nu+1} \left( \frac{2qp'}{q'^2 + p'^2} \right)^{\nu+1} \\ &\times \sum_{n_r=0}^{\infty} \frac{n_r!}{\Gamma(n_r + 2\nu + 2)} C_{n_r}^{(\nu+1)} \left( \frac{q^2 - p^2}{q^2 + p^2} \right) C_{n_r}^{(\nu+1)} \left( \frac{q'^2 - p'^2}{q'^2 + p'^2} \right) \end{aligned} \quad (3.30)$$

(the particular reason for which we have denoted the summation index as  $n_r$  will become clear shortly). Combining Eqs. (3.26) and (3.30), we see that the kernel in Eq. (3.17) may be written as

$$M_l(p, p') = \sum_{n_r=0}^{\infty} g_{n_r, l}(p) g_{n_r, l}(p') \quad (\Rightarrow \forall n_r \in \mathbb{N} : \varepsilon_{n_r, l} = 1), \quad (3.31)$$

with

$$g_{n_r, l}(p) = \Gamma \left( l + \frac{N-1}{2} \right) \sqrt{\frac{Z q_B n_r!}{\pi (n_r + 2l + N - 2)!}} \frac{(4qp)^{l+(N-1)/2}}{(q^2 + p^2)^{l+N/2}} C_{n_r}^{(l+(N-1)/2)} \left( \frac{q^2 - p^2}{q^2 + p^2} \right). \quad (3.32)$$

In the next step, we evaluate the integral  $\int_0^\infty dp g_{n_r, l}(p) g_{n'_r, l}(p)$ , in order to verify whether the functions (3.32) form an orthogonal set [as they should, cf. Eq. (3.22)]. Exploiting the known [29, Eq. (7.313)] integral formula

$$\int_{-1}^1 d\xi (1 - \xi^2)^{\alpha-1/2} C_{n_r}^{(\alpha)}(\xi) C_{n'_r}^{(\alpha)}(\xi) = \frac{\pi \Gamma(n_r + 2\alpha)}{2^{2\alpha-1} n_r! (n_r + \alpha) [\Gamma(\alpha)]^2} \delta_{n_r, n'_r} \quad (\text{Re } \alpha > -\frac{1}{2}, \alpha \neq 0), \quad (3.33)$$

we find that

$$\int_0^\infty dp g_{n_r, l}(p) g_{n'_r, l}(p) = \frac{Z}{n_r + l + \frac{N-1}{2}} \frac{q_B}{q} \delta_{n_r, n'_r}, \quad (3.34)$$

i.e., the functions (3.32) are indeed orthogonal, the norm of  $g_{n_r, l}(p)$  being

$$\|g_{n_r, l}\| = \sqrt{\frac{Z}{n_r + l + \frac{N-1}{2}} \frac{q_B}{q}}. \quad (3.35)$$

Hence, applying Eqs. (3.23) and (3.24) we infer that the spectrum of eigenvalues in Eq. (3.17) is given by

$$\lambda_{n_r, l} = \frac{n_r + l + \frac{N-1}{2}}{Z} \frac{q}{q_B} \quad (n_r \in \mathbb{N}), \quad (3.36)$$

whereas the associated orthonormal eigenfunctions are

$$f_{n_r l}(E, p) = \Gamma\left(l + \frac{N-1}{2}\right) \sqrt{\frac{qn_r!(n_r + l + \frac{N-1}{2})}{\pi(n_r + 2l + N - 2)!}} \\ \times \frac{(4qp)^{l+(N-1)/2}}{(q^2 + p^2)^{l+N/2}} C_{n_r}^{(l+(N-1)/2)} \left(\frac{q^2 - p^2}{q^2 + p^2}\right). \quad (3.37)$$

Consequently (cf. Eq. (3.25)), the sought radial momentum-space Coulomb Sturmian functions are found to be

$$F_{n_r l}(E, p) = 2^{N-1} \Gamma\left(l + \frac{N-1}{2}\right) \sqrt{\frac{n_r!(n_r + l + \frac{N-1}{2})}{\pi(n_r + 2l + N - 2)!}} \\ \times q^{N/2} \frac{(4qp)^l}{(q^2 + p^2)^{l+(N+1)/2}} C_{n_r}^{(l+(N-1)/2)} \left(\frac{q^2 - p^2}{q^2 + p^2}\right) \quad (3.38)$$

(it is seen that  $n_r$  is simply the radial quantum number). They obey the weighted orthonormality relation

$$\int_0^\infty dp p^{N-1} (p^2 + q^2) F_{n_r l}(E, p) F_{n_r' l'}(E, p) = \delta_{n_r n_r'} \quad (3.39)$$

and are standardized to be positive in the immediate vicinity of  $p = 0$ .

### 3.5 Some properties of solutions to Eq. (3.5)

Once the radial integral equation (3.16) is solved, we may return to the  $N$ -dimensional problem (3.5).

It follows from Eq. (3.36) that the degree of degeneracy of the eigenvalues in Eq. (3.5) is higher than  $d_l^{(N-1)}$ , as  $\lambda_{n_r l}$  depends on the quantum numbers  $n_r$  and  $l$  through their sum. To distinguish between different eigenvalues, we introduce the principal quantum number

$$n = n_r + l + 1, \quad (3.40)$$

in terms of which we have

$$\lambda_{n_r l} \equiv \lambda_n = \frac{n + \frac{N-3}{2}}{Z} \frac{q}{q_B}. \quad (3.41)$$

The degree of degeneracy of the particular eigenvalue  $\lambda_n$  obviously is

$$D_n^{(N)} = \sum_{l=0}^{n-1} d_l^{(N-1)}. \quad (3.42)$$

In view of Eq. (3.7), this may be rewritten as

$$D_n^{(N)} = 2 \sum_{l=0}^{n-1} \binom{l+N-2}{N-2} - \sum_{l=0}^{n-1} \binom{l+N-3}{N-3}. \quad (3.43)$$

The sums in Eq. (3.43) are elementary; we have [29, Eq. (0.15.1)]

$$\sum_{k=0}^n \binom{k+m}{m} = \binom{n+m+1}{m+1}, \quad (3.44)$$

and consequently

$$D_n^{(N)} = (2n + N - 3) \frac{(n + N - 3)!}{(n - 1)!(N - 1)!}. \quad (3.45)$$

Comparison of Eqs. (3.45) and (3.7) shows that it holds that

$$D_n^{(N)} = d_{n-1}^{(N)}. \quad (3.46)$$

(The relation in Eq. (3.46) is, of course, a symptom of the  $O(N+1)$  dynamical symmetry of the  $N$ -dimensional negative-energy Schrödinger–Coulomb problem.)

Degenerate Sturmian eigenfunctions associated with the characteristic numbers (3.41), labeled with the principal rather than the radial quantum number, are

$$\Sigma_{nl\eta}(E, \mathbf{p}) = F_{n-l-1, l}(E, p) Y_{l\eta}^{(N-1)}(\mathbf{n}_p). \quad (3.47)$$

It is implied by Eqs. (3.8) and (3.39) that these functions obey the weighted orthonormality relation

$$\int_{\mathbb{R}^N} d^N \mathbf{p} (p^2 + q^2) \Sigma_{nl\eta}^*(E, \mathbf{p}) \Sigma_{n'l'\eta'}(E, \mathbf{p}) = \delta_{nn'} \delta_{ll'} \delta_{\eta\eta'}. \quad (3.48)$$

It remains to consider the completeness problem for the Sturmian set  $\{\Sigma_{nl\eta}(E, \mathbf{p})\}$ . It is known that the functions  $\{(1 - \xi^2)^{\alpha/2-1/4} C_{n_r}^{(\alpha)}(\xi)\}$ , with  $\text{Re } \alpha > -\frac{1}{2}$ , form a complete set in the functional space  $L^2((-1, 1))$ , the corresponding closure relation being

$$\frac{2^{2\alpha-1} [\Gamma(\alpha)]^2}{\pi} \sum_{n_r=0}^{\infty} \frac{n_r! (n_r + \alpha)}{\Gamma(n_r + 2\alpha)} C_{n_r}^{(\alpha)}(\xi) C_{n_r}^{(\alpha)}(\xi') = \frac{\delta(\xi - \xi')}{[(1 - \xi^2)(1 - \xi'^2)]^{\alpha/2-1/4}} \quad (-1 < \xi, \xi' < 1, \text{Re } \alpha > -\frac{1}{2}), \quad (3.49)$$

with  $\delta(\xi - \xi')$  denoting the Dirac delta distribution. We set in the above equation  $\alpha = l + (N-1)/2$ , plug Eq. (3.29), combine the result with Eq. (3.38), and then exploit the identity

$$\delta(\xi - \xi') = \frac{(p^2 + q^2)(p'^2 + q^2)}{4\sqrt{pp'} q^2} \delta(p - p'), \quad (3.50)$$

which follows from Eq. (3.29) and the well-known properties of the Dirac delta. This chain of operations transforms Eq. (3.49) into the closure relation for the radial Sturmians:

$$\sum_{n_r=0}^{\infty} F_{n_r l}(E, p) F_{n_r l}(E, p') = \frac{\delta(p - p')}{(pp')^{(N-1)/2} \sqrt{(p^2 + q^2)(p'^2 + q^2)}}. \quad (3.51)$$

Next, we have the closure relation

$$\sum_{l=0}^{\infty} d_l^{(N-1)} \sum_{\eta=1}^{d_l^{(N-1)}} Y_{l\eta}^{(N-1)}(\mathbf{n}_p) Y_{l\eta}^{(N-1)*}(\mathbf{n}'_p) = \delta^{(N-1)}(\mathbf{n}_p - \mathbf{n}'_p), \quad (3.52)$$

which reflects the fact that the hyperspherical harmonics form a complete set in the functional space  $L^2(\mathbb{S}^{N-1})$ . We insert Eqs. (3.51) and (3.52) into the right-hand side of the identity

$$\sum_{n=1}^{\infty} \sum_{l=0}^{n-1} \sum_{\eta=1}^{d_l^{(N-1)}} \Sigma_{nl\eta}(E, \mathbf{p}) \Sigma_{nl\eta}^*(E, \mathbf{p}') = \sum_{l=0}^{\infty} \sum_{\eta=1}^{d_l^{(N-1)}} Y_{l\eta}^{(N-1)}(\mathbf{n}_p) Y_{l\eta}^{(N-1)*}(\mathbf{n}'_p) \sum_{n_r=0}^{\infty} F_{n_r l}(E, p) F_{n_r l}(E, p') \quad (3.53)$$

and make use of the obvious fact that

$$\delta^{(N)}(\mathbf{p} - \mathbf{p}') = \frac{\delta(p - p') \delta^{(N-1)}(\mathbf{n}_p - \mathbf{n}'_p)}{(pp')^{(N-1)/2}}. \quad (3.54)$$

This leads us to the symmetric closure relation

$$\sum_{n=1}^{\infty} \sum_{l=0}^{n-1} \sum_{\eta=1}^{d_l^{(N-1)}} \Sigma_{nl\eta}(E, \mathbf{p}) \Sigma_{nl\eta}^*(E, \mathbf{p}') = \frac{\delta^{(N)}(\mathbf{p} - \mathbf{p}')}{\sqrt{(p^2 + q^2)(p'^2 + q^2)}}. \quad (3.55)$$

We have thus proved that the Sturmian functions  $\{\Sigma_{nl\eta}(E, \mathbf{p})\}$  form a complete orthonormal set in the functional space  $L^2_{p^2+q^2}(\mathbb{R}^N)$ ; equivalently, the functions  $\{\sqrt{p^2 + q^2} \Sigma_{nl\eta}(E, \mathbf{p})\}$  form a complete orthonormal set in the functional space  $L^2(\mathbb{R}^N)$ .



## 4 Determination of bound-state eigensolutions to the momentum-space Schrödinger–Coulomb wave equation (2.5)

We are now ready to determine bound-state eigensolutions to Eq. (2.5), following the procedure outlined at the beginning of Sec. 3.1. At first, we substitute  $\lambda_n$ , as given in Eq. (3.41), for  $\lambda_\alpha(E)$  in Eq. (3.2), then use Eq. (3.4), and solve the resulting equation for  $E$ . For each particular  $n$ , there is just one solution to that equation, which we shall denote as  $E_n$ ; it is

$$E_n = -\frac{Z^2 e^2}{2\left(n + \frac{N-3}{2}\right)^2 (4\pi\epsilon_0)a_B}. \quad (4.1)$$

Corresponding eigenfunctions, in accordance with Eqs. (3.3) and (3.6), are

$$\Phi_{nl\eta}(\mathbf{p}) = \mathcal{F}_{nl}(p) Y_{l\eta}^{(N-1)}(\mathbf{n}_p), \quad (4.2)$$

with

$$\mathcal{F}_{nl}(p) = A F_{n-l-1,l}(E_n, p). \quad (4.3)$$

Equation (2.5) shows that  $E_n^{-1}$  is an eigenvalue of the Hermitian (in fact — real and symmetric) kernel

$$\frac{pp'}{2m} \delta^{(N)}(\mathbf{p} - \mathbf{p}') - \frac{Ze^2}{(4\pi\epsilon_0)} \frac{\Gamma\left(\frac{N-1}{2}\right)}{2\pi^{(N+1)/2}\hbar} |\mathbf{p} - \mathbf{p}'|^{-(N-1)}.$$

From this and from Eqs. (4.2) and (3.8), one routinely deduces that the functions  $\{\Phi_{nl\eta}(\mathbf{p})\}$  are orthogonal in the sense of

$$\int_{\mathbb{R}^N} d^N \mathbf{p} \Phi_{nl\eta}^*(\mathbf{p}) \Phi_{n'l'\eta'}(\mathbf{p}) \sim \delta_{nn'} \delta_{ll'} \delta_{\eta\eta'}. \quad (4.4)$$

If, additionally, we require the functions under consideration are normalized to unity under the same scalar product that is used in the above equation, and if we choose the constant  $A$  in Eq. (4.3) to be real, we obtain the constraint

$$A^2 \int_0^\infty dp p^{N-1} [F_{n-l-1,l}(E_n, p)]^2 = 1. \quad (4.5)$$

To evaluate the integral in Eq. (4.5), we use Eq. (3.38). This yields

$$\begin{aligned} \int_0^\infty dp p^{N-1} [F_{n-l-1,l}(E_n, p)]^2 &= 2^{2l+N-3} \left[ \Gamma\left(l + \frac{N-1}{2}\right) \right]^2 \frac{(n-l-1)! \left(n + \frac{N-3}{2}\right)}{\pi(n+l+N-3)!} q_n^{-2} \\ &\times \int_{-1}^1 d\xi_n (1 - \xi_n^2)^{l+N/2-1} (1 + \xi_n) \left[ C_{n-l-1}^{(l+(N-1)/2)}(\xi_n) \right]^2, \end{aligned} \quad (4.6)$$

with

$$q_n = \frac{Zq_B}{n + \frac{N-3}{2}}. \quad (4.7)$$

It is evident that in the integrand on the right-hand side of Eq. (4.6) one may replace the factor  $1 + \xi_n$  by 1 without altering the value of the integral. Applying then the formula (3.33) and imposing a further restriction on  $A$  to be positive, we obtain

$$A = \sqrt{2} q_n. \quad (4.8)$$

Hence, we conclude that the bound-state momentum-space eigenfunctions of the Schrödinger–Coulomb problem in  $\mathbb{R}^N$ ,  $N \geq 2$ , associated with the energy levels (4.1) and orthonormal in the sense of

$$\int_{\mathbb{R}^N} d^N \mathbf{p} \Phi_{nl\eta}^*(\mathbf{p}) \Phi_{n'l'\eta'}(\mathbf{p}) = \delta_{nn'} \delta_{ll'} \delta_{\eta\eta'}, \quad (4.9)$$

may be chosen in the form

$$\begin{aligned}\Phi_{nl\eta}(\mathbf{p}) &= 2^{N-1/2}\Gamma\left(l + \frac{N-1}{2}\right)\sqrt{\frac{(n-l-1)!\left(n + \frac{N-3}{2}\right)}{\pi(n+l+N-3)!}} \\ &\times q_n^{N/2+1}\frac{(4q_n p)^l}{(q_n^2 + p^2)^{l+(N+1)/2}}C_{n-l-1}^{(l+(N-1)/2)}\left(\frac{q_n^2 - p^2}{q_n^2 + p^2}\right)Y_{l\eta}^{(N-1)}(\mathbf{n}_p).\end{aligned}\quad (4.10)$$

The degeneracy of the energy level  $E_n$  is, obviously, the same as that of the Sturmian eigenvalue  $\lambda_n$ , and is given by Eq. (3.45).

With Eqs. (4.1) and (4.10) being obtained, our task to determine bound-state eigensolutions to the Schrödinger–Coulomb problem in the  $N$ -dimensional Euclidean momentum space is accomplished.

## Acknowledgments

I thank Dr. Howard S. Cohl for drawing my attention to his work [25].

## Appendix: The Fourier transform of $V(\mathbf{r}) = Ar^{-\nu}$ , $0 < \nu < N$ , in $\mathbb{R}^N$

In this Appendix, we shall follow closely Ref. [30, Sec. 5.9]. We wish to evaluate the Fourier transform

$$U(\mathbf{p}) = \frac{1}{(2\pi\hbar)^{N/2}}\int_{\mathbb{R}^N}d^N\mathbf{r}e^{-i\mathbf{p}\cdot\mathbf{r}/\hbar}V(\mathbf{r}),\quad (A.1)$$

where

$$V(\mathbf{r}) = \frac{A}{r^\nu}\quad (0 < \nu < N).\quad (A.2)$$

To accomplish the task, at first we observe that

$$\int_0^\infty d\xi \xi^{\nu/2-1}e^{-\xi r^2} = r^{-\nu}\int_0^\infty d\eta \eta^{\nu/2-1}e^{-\eta} = \Gamma\left(\frac{\nu}{2}\right)r^{-\nu},\quad (A.3)$$

hence, it follows that

$$r^{-\nu} = \frac{1}{\Gamma\left(\frac{\nu}{2}\right)}\int_0^\infty d\xi \xi^{\nu/2-1}e^{-\xi r^2}.\quad (A.4)$$

If we plug Eq. (A.4) into Eq. (A.1), we obtain

$$\begin{aligned}U(\mathbf{p}) &= \frac{A}{(2\pi\hbar)^{N/2}\Gamma\left(\frac{\nu}{2}\right)}\int_{\mathbb{R}^N}d^N\mathbf{r}e^{-i\mathbf{p}\cdot\mathbf{r}/\hbar}\int_0^\infty d\xi \xi^{\nu/2-1}e^{-\xi r^2} \\ &= \frac{A}{(2\pi\hbar)^{N/2}\Gamma\left(\frac{\nu}{2}\right)}\int_0^\infty d\xi \xi^{\nu/2-1}\int_{\mathbb{R}^N}d^N\mathbf{r}e^{-i\mathbf{p}\cdot\mathbf{r}/\hbar}e^{-\xi r^2}.\end{aligned}\quad (A.5)$$

The integral over  $\mathbf{r}$  is readily found to be

$$\int_{\mathbb{R}^N}d^N\mathbf{r}e^{-i\mathbf{p}\cdot\mathbf{r}/\hbar}e^{-\xi r^2} = \left(\frac{\pi}{\xi}\right)^{N/2}e^{-p^2/4\hbar^2\xi},\quad (A.6)$$

so that Eq. (A.5) becomes

$$U(\mathbf{p}) = \frac{A}{(2\hbar)^{N/2}\Gamma\left(\frac{\nu}{2}\right)}\int_0^\infty d\xi \xi^{(\nu-N)/2-1}e^{-p^2/4\hbar^2\xi},\quad (A.7)$$

and further, after switching to the integration variable  $\eta = p^2/4\hbar^2\xi$ ,

$$U(\mathbf{p}) = A \frac{(2\hbar)^{N/2-\nu}}{\Gamma(\frac{\nu}{2})} \frac{1}{p^{N-\nu}} \int_0^\infty d\eta \eta^{(N-\nu)/2-1} e^{-\eta}. \quad (\text{A.8})$$

The remaining integral is elementary and we eventually arrive at

$$U(\mathbf{p}) = A \frac{(2\hbar)^{N/2-\nu} \Gamma(\frac{N-\nu}{2})}{\Gamma(\frac{\nu}{2})} \frac{1}{p^{N-\nu}}. \quad (\text{A.9})$$

In the case of the Coulomb potential

$$V(\mathbf{r}) = -\frac{Ze^2}{(4\pi\epsilon_0)r}, \quad (\text{A.10})$$

from Eq. (A.9) we find that

$$U(\mathbf{p}) = -\frac{Ze^2}{(4\pi\epsilon_0)} \frac{(2\hbar)^{N/2-1} \Gamma(\frac{N-1}{2})}{\sqrt{\pi}} \frac{1}{p^{N-1}}. \quad (\text{A.11})$$

## References

- [1] B. Podolsky, L. Pauling, The momentum distribution in hydrogen-like atoms, *Phys. Rev.* 34 (1929) 109
- [2] E. Schrödinger, Quantisierung als Eigenwertproblem (Erste Mitteilung), *Ann. Phys. (Leipzig)* 79 (1926) 361 [English translation: E. Schrödinger, Quantisation as a Problem of Proper Values (Part I), in: E. Schrödinger, *Collected Papers on Wave Mechanics* (AMS Chelsea, Providence, RI, 2001) p. 1]
- [3] E. A. Hylleraas, Die Wellengleichung des Keplerproblems im Impulsraume, *Z. Phys.* 74 (1932) 216
- [4] W. Elsasser, Atomare Wellenfunktion im Impulsraum, *Z. Phys.* 81 (1933) 332
- [5] V. Fock, Zur Theorie des Wasserstoffatoms, *Z. Phys.* 98 (1935) 145 [English translations: V. Fock, Hydrogen atom and non-Euclidean geometry, in: V. A. Fock, *Selected Works. Quantum Mechanics and Quantum Field Theory*, edited by L. D. Faddeev, L. A. Khal'fin, I. V. Komarov (Chapman & Hall/CRC, Boca Raton, FL, 2004) p. 369; V. Fock, On the theory of the hydrogen atom, in: *Dynamical Groups and Spectrum Generating Algebras*, vol. 1, A. Bohm, Y. Ne'eman, A. O. Barut (eds.) (World Scientific, Singapore, 1988) p. 411; V. Fock, On the theory of the hydrogen atom, in: S. F. Singer, *Linearity, Symmetry, and Prediction in the Hydrogen Atom* (Springer, New York, 2005) p. 286]
- [6] V. Bargmann, Zur Theorie des Wasserstoffatoms, *Z. Phys.* 99 (1936) 576 [English translation: V. Bargmann, On the theory of the hydrogen atom, in: *Dynamical Groups and Spectrum Generating Algebras*, vol. 1, A. Bohm, Y. Ne'eman, A. O. Barut (eds.) (World Scientific, Singapore, 1988) p. 411]
- [7] T. Shibuya, C. E. Wulfman, The Kepler problem in two-dimensional momentum space, *Am. J. Phys.* 33 (1965) 570
- [8] J. J. Klein, Eigenfunctions of the hydrogen atom in momentum space, *Am. J. Phys.* 34 (1966) 1039
- [9] R. Aldrovandi, J. A. C. Alcaras, P. Leal Ferreira, Accidental degeneracy of the Coulomb interaction, *Am. J. Phys.* 35 (1967) 520

- [10] E. V. Ivash, A momentum representation treatment of the hydrogen atom problem, *Am. J. Phys.* 40 (1972) 1095
- [11] O. L. de Lange, R. E. Raab, An operator solution for the hydrogen atom with application to the momentum representation, *Am. J. Phys.* 55 (1987) 913
- [12] J. D. Hey, On the momentum representation of hydrogenic wave functions: some properties and an application, *Am. J. Phys.* 61 (1993) 28
- [13] B. R. Holstein, Quantum mechanics in momentum space: The Coulomb system, *Am. J. Phys.* 63 (1995) 710
- [14] W. Dittrich, The Coulomb Green's function in two dimensions, *Am. J. Phys.* 67 (1999) 768
- [15] A. V. Meremianin, J.-M. Rost, Multipole expansions and Fock symmetry of the hydrogen atom, *J. Phys. A* 39 (2006) 12427
- [16] D.-H. Lin, Solution of Coulomb system in momentum space, *Ann. Phys. (NY)* 323 (2008) 317
- [17] M. Bander, C. Itzykson, Group theory and the hydrogen atom (I), *Rev. Mod. Phys.* 38 (1966) 330
- [18] V. S. Popov, On a "hidden" symmetry of a hydrogen atom, in: *High Energy Physics and Theory of Elementary Particles (Naukova Dumka, Kiev, 1967)* (in Russian) p. 702
- [19] J. C. Y. Chen, A. C. Chen, Nonrelativistic off-shell two-body Coulomb amplitudes, *Adv. At. Mol. Phys.* 8 (1972) 71
- [20] H. van Haeringen, *Charged-Particle Interactions. Theory and Formulas* (Coulomb Press, Leyden, 1985)
- [21] J. Avery, *Hyperspherical Harmonics. Applications in Quantum Theory* (Kluwer, Dordrecht, 1989)
- [22] J. Avery, *Hyperspherical Harmonics and Generalized Sturmians* (Kluwer, Dordrecht, 2000)
- [23] V. Aquilanti, S. Cavalli, C. Coletti, D. Di Domenico, G. Grossi, Hyperspherical harmonics as Sturmian orbitals in momentum space: a systematic approach to the few-body Coulomb problem, *Int. Rev. Phys. Chem.* 20 (2001) 673
- [24] J. Avery, Harmonic polynomials, hyperspherical harmonics, and Sturmians, in: *Fundamental World of Quantum Chemistry, vol. I*, E. J. Brändas, E. S. Kryachko (eds.) (Kluwer, Dordrecht, 2003) p. 261
- [25] H. S. Cohl, On a generalization of the generating function for Gegenbauer polynomials, [arXiv:1105.2735](https://arxiv.org/abs/1105.2735)
- [26] A. Ossicini, Funzione generatrice dei prodotti di due polinomi ultrasferici, *Boll. Un. Mat. Ital.* 7 (1952) 315
- [27] H. A. Bethe, E. E. Salpeter, *Quantum Mechanics of One- and Two-electron Atoms* (Springer, Berlin, 1957)
- [28] E. R. Hansen, *A Table of Series and Products* (Prentice-Hall, Englewood Cliffs, NJ, 1975)
- [29] I. S. Gradshteyn, I. M. Ryzhik, *Table of Integrals, Series, and Products*, 7th ed. (Elsevier, Amsterdam, 2007)
- [30] E. Lieb, M. Loss, *Analysis*, 2nd ed. (American Mathematical Society, Providence, RI, 2001)