

SIMPLE CUBIC RANDOM-SITE PERCOLATION THRESHOLDS FOR COMPLEX NEIGHBOURHOODS

Lukasz Kurzawski and Krzysztof Malarz*
AGH University of Science and Technology,
Faculty of Physics and Applied Computer Science,
al. Mickiewicza 30, 30-059 Krakow, Poland.
e-mail: malarz@agh.edu.pl

(Received June 3, 2019)

In this communication with computer simulation we evaluate simple cubic random-site percolation thresholds for neighbourhoods including the nearest-neighbours (NN), the second-nearest neighbours (2NN) and the third-nearest neighbours (3NN). Our estimations base on finite-size scaling analysis of the percolation probability vs. site occupation probability plots. The Hoshen–Kopelman algorithm has been applied for cluster labelling. The calculated thresholds are $0.137 \dots$, $0.142 \dots$, $0.097 \dots$, $0.199 \dots$, $0.103 \dots$, $0.245 \dots$ for (NN + 2NN), (NN + 3NN), (NN + 2NN + 3NN), 2NN, (2NN + 3NN), 3NN neighbourhoods, respectively. In contrast to the results obtained for a square lattice the calculated percolation thresholds decrease monotonically with the site coordination number z , at least for inspected neighbourhoods.

Keywords: site percolation; percolation thresholds; computer simulations

1. Introduction

Percolation [1,2] is a mathematical description of a geometrical phase transition. This allows for purely theoretical studies of all phenomena occurring near the critical point with computer experiments solely or, sometimes, even analytically [3]. The mixture of occupied and empty sites of given lattice may exhibit some features of real physical systems. Among typical applications of the percolation theory one may find material science [4], immunology [5] or forest fires problems [6] and studies of liquids moving in porous media [7], etc. [8,9].

Generally speaking, the percolation theory deals with statistical properties of the clusters of occupied nodes (site percolation) or occupied edges (bond percolation) for a given graph, network or regular lattice. In the site percolation problem, the cluster is defined as a group of the occupied lattice vertexes which belong to the site neighbourhoods. When each site is occupied with some probability p there is a critical probability

* <http://home.agh.edu.pl/malarz/>

of sites occupation p_c above which the cluster spanning through the whole system appears for the first time [8,9]. This special probability is called percolation threshold p_c and it separates two phases (in the language of material science a conductor and an isolator). The value of percolation threshold p_c depends on kind of percolation (site/bond), lattice/graph/network topology and assumed sites neighbourhoods. In the simplest case only the nearest-neighbours constitute the neighbourhoods (von Neumann's neighbourhood) or the nearest-neighbours and second-nearest neighbours are considered (Moore's neighbourhood).

In the vicinity of the phase transition the quantity A describing the system follows a scaling relation

$$A \propto L^\alpha \cdot f(x^\beta), \quad (1)$$

where L describes the linear size of the system, $f(x)$ is a scaling function and x is dimensionless scaling parameter. For physical systems x usually plays the role of reduced temperature $(T - T_C)/T_C$, where T_C stands for critical temperature. The α and β parameters are universal exponents which—in the first approximation—do not depend on system details (kind of order/disorder phenomenon, lattice shape, site or bond percolation, etc.) but only on the system dimensionality [8, 10]. However, for α and β calculations the precise value of T_C is required. For geometrical model of the phase transition percolation threshold p_c plays the role of critical temperature.

In this communication we evaluate with computer simulations [11] the random-site simple cubic percolation thresholds for neighbourhoods including the nearest-neighbours (NN), the second-nearest neighbours (2NN) and the third-nearest neighbours (3NN). Our estimations base on finite size scaling analysis [8,10] of the percolation probability vs. site occupation probability plots. The Hoshen–Kopelman algorithm [12] has been applied for cluster labelling. The calculated thresholds p_c are $0.137 \dots$, $0.142 \dots$, $0.097 \dots$, $0.199 \dots$, $0.103 \dots$, $0.245 \dots$ for (NN + 2NN), (NN + 3NN), (NN + 2NN + 3NN), 2NN, (2NN + 3NN), 3NN neighbourhoods, respectively. In contrast to the results obtained for a square lattice [13–15] the calculated percolation thresholds decrease monotonically with the site coordination number z , at least for inspected neighbourhoods.




Fig. 1: Basic neighbourhoods for simple cubic lattice. The NN neighbourhood is often referred to as von Neumann's neighbourhood while combination of (NN+2NN) is called Moore's neighbourhood.

2. Calculations

We use Hoshen–Kopelman algorithm [12] for occupied sites labelling. In the Hoshen–Kopelman scheme each site has one label: all sites in given cluster have the same labels and different clusters have assigned different labels.

Examples of percolation probability P vs. sites occupation probability p for various neighbourhoods and for various lattice linear sizes L are presented in Fig. 2. We use finite-size scaling analysis to determine p_c numerically. As for finite systems phase transition is never sharp we observe it when for increasing lattice sizes L the $P(p)$ curves become more and more steep and intersect at a common point corresponding also to the case of $L \rightarrow \infty$ [16]. The common cross-point predicts percolation threshold p_c . Such strategy was successfully applied for many system description where phase transition may be observed including percolation [17], Ising model [18], majority-vote models [19] or opinion dynamics [20].

Basing on $P(p)$ dependence for various L we look for an interval of the length $\Delta p = 10^{-3}$ where curves for $L = 63$ and 100 cross each other. For example for 2NN case this interval is $(0.199, 0.200)$. We assume, that these three significant figures of the left border of this interval approximate percolation threshold p_c , i.e. these three digits may be treated as exact.

3. Results

The $P(p)$ dependencies for $L = 22, 63$ and 100 are presented in Fig. 2. The evaluated percolation thresholds p_c for various neighbourhoods are collected in Tab. 1. To check the accuracy of our estimations we evaluated the percolation threshold for von Neumann’s neighbourhood as well. The obtained value $p_c(\text{NN}) = 0.311\dots$ agrees finely with the results of extensive numerical simulations [21].

Table 1: Simple-cubic lattice random-site percolation thresholds p_c for various neighbourhoods constructed with basic neighbourhoods NN, 2NN and 3NN.

neighbourhood	z	p_c
NN	6	0.311...
2NN	12	0.199...
3NN	8	0.245...
NN+2NN	18	0.137...
NN+3NN	14	0.142...
2NN+3NN	20	0.103...
NN+2NN+3NN	26	0.097...

These p_c values decrease with sites coordination number z as shown in Fig. 3. The $p_c(z)$ dependencies may be fitted nicely by two straight lines in a logarithmic plot, namely: $p_c(z) \propto z^{-\gamma}$ with $\gamma = -0.919 \pm 0.025$ and $\gamma = -0.927 \pm 0.005$.




Fig. 2: Percolation probability P vs. site occupation probability p for various neighbourhoods in three dimensions for three lattice sizes L . The results have been averaged over N clusters realisations.




Fig. 3: Percolation thresholds p_c vs. sites coordination number z for inspected three-dimensional neighbourhoods. The straight lines are the least squares fits of $p_c(z) \propto z^{-\gamma}$ to the experimental data.

4. Conclusions

In this communication for the first time we evaluate the random-site percolation thresholds for the simple cubic lattice with neighbourhoods for which sites from the first, the second and the third coordination shells were included (see Tab. 1). The obtained thresholds p_c decrease monotonically with sites coordination number z in two series according to the power law $p_c \propto z^{-\gamma}$, both with exponent $\gamma \approx 0.92$.

In contrast to the results obtained for a square lattice [15] the calculated percolation thresholds decrease monotonically with the site coordination number z , at least for inspected neighbourhoods.

The obtained results may be helpful in studies of the universal formulae [22] for percolation thresholds p_c dependence on sites coordination number z .

Finally, we propose to name (NN+2NN+3NN) neighbourhood in simple cubic lattice *the Rubik's neighbourhood* as it is identical with the famous Rubik's cube [23] — a very popular logical puzzle in early 80's.

Acknowledgements

Supported by the Polish Ministry of Science and Higher Education and its grants for scientific research. The numerical calculations were carried out in the Academic Computer Centre CYFRONET-AGH (grant No. MEiN/SGI3700/AGH/024/2006).

REFERENCES

- [1] S. R. Broadbent, J. M. Hammersley: *Proc. Cambridge Phil. Soc.* **53**, 629 (1957);
H. L. Frisch, J. M. Hammersley, D. J. A. Welsh: *Phys. Rev.* **126**, 949 (1962).
- [2] M. F. Sykes, M. Glen: *J. Phys. A* **9**, 87 (1976);
M. F. Sykes, D. S. Gaunt, M. Glen: *ibid.* **9**, 97 (1976);
M. F. Sykes, D. S. Gaunt, M. Glen: *ibid.* **9**, 715 (1976);
M. F. Sykes, D. S. Gaunt, M. Glen: *ibid.* **9**, 725 (1976);
D. S. Gaunt, M. F. Sykes: *ibid.* **9**, 1109 (1976).
- [3] J. C. Wierman, S. P. Naor, R. Cheng: *Phys. Rev. E* **72**, 066116 (2005);
A. Rosowsky: *Eur. Phys. J. B* **15**, 77 (2000).
- [4] B. I. Halperin and D. J. Bergman: *Physica B* **405**, 2908-2914 (2010);
P. R. Shearing, D. J. L. Brett, N. P. Brandon: *International Materials Reviews* **55**, 347-363 (2010);
J. Silva, R. Simoes, S. Lanceros-Mendez, R. Vaia: *EPL* **93**, 6 (2011).
- [5] S. U. Suzuki, A. Sasaki *J. Theor. Biol.* **276**, 117-125 (2011);
J. Lindquist, Junling Ma, P. van den Driessche, et al: *J. Math. Biol.* **62**, 143-164 (2011);
E. N. Naumova, J. Gorski, Yu. N. Naumov: *Annales Zoologici Fennici* **45**, 369-384 (2008);
W. Floyd, L. Kay, M. Shapiro: *Bull. Math. Biol.* **70**, 713-727 (2008).
- [6] G. Camelo-Neto, S. Coutinho: *J. Stat. Mech.*, P06018 (2011);
N. Guisoni, E. S. Loscar, E. V. Albano: *Phys. Rev. E* **83**, 011125 (2011);
A. Simeoni, P. Salinesi, F. Morandini: *Int. J. Wildland Fire* **20**, 625-632 (2011);
K. Malarz, S. Kaczanowska, K. Kulakowski: *Int. J. Mod. Phys. C* **13**, 1017 (2002).
- [7] Y. Amiaz, S. Sorek, Y. Enzel, et al: *Water Resources Research* **47**, W10513 (2011);
S. F. Bolandtaba, A. Skauge: *Transport in Porous Media* **89**, 357-382 (2011);
V. V. Mourzenko, J.-F. Thovert, P. M. Adler: *Phys. Rev. E* **84**, 036307 (2011).
- [8] D. Stauffer, A. Aharony: *Introduction to Percolation Theory*, Taylor and Francis, London 1994;
- [9] H. Kesten: *Percolation Theory for Mathematicians*, Birkhauser, Boston 1982;
M. Sahimi: *Applications of Percolation Theory*, Taylor and Francis, London 1994;
D. Stauffer: *Physica A* **242**, 1 (1997).
- [10] D. P. Landau, K. Binder: *A Guide to Monte Carlo Simulation in Statistical Physics*, Cambridge University Press, Cambridge 2005.
- [11] L. Kurzawski: *M.Sc. Thesis*, AGH University of Science and Technology, Kraków 2011.
- [12] J. Hoshen, R. Kopelman: *Phys. Rev. B* **14**, 3428 (1976).
- [13] K. Malarz, S. Galam: *Phys. Rev. E* **71**, 016125 (2005).
- [14] S. Galam, K. Malarz: *Phys. Rev. E* **72**, 027103 (2005).
- [15] M. Majewski, K. Malarz: *Acta Phys. Pol. B* **38**, 2191 (2007).
- [16] V. Privman (Ed.): *Finite size scaling and numerical simulation of statistical systems*, World Scientific, Singapore 1990.

- [17] M. E. J. Newman, R. M. Ziff: *Phys. Rev. E* **85**, 016706 (2001).
- [18] F. W. S. Lima, J. Mostowicz, K. Malarz: *Commun. Comput. Phys.* **10**, 912 (2011).
- [19] J. C. Santos, F. W. S. Lima, K. Malarz: *Physica A* **390**, 359 (2011);
F. W. S. Lima, A. O. Sousa, M. A. Sumuorc: *Physica A* **387**, 3503-3510 (2008).
F. W. S. Lima, K. Malarz: *Int. J. Mod. Phys. C* **17**, 1273 (2006);
- [20] D. Stauffer, A. O. Sousa, S. Moss de Oliveira: *Int. J. Mod. Phys. C* **11**, 1239 (2000);
A. A. Moreira, J. S. Andrade Jr., D. Stauffer: *Int. J. Mod. Phys. C* **12**, 39 (2001);
A. O. Sousa: *Physica A* **348**, 701 (2005).
- [21] P. Grassberger: *J. Phys. A* **25**, 5867-5888 (1992);
C. D. Lorenz, R. M. Ziff: *J. Phys. A* **31**, 8147-8157 (1998);
N. Jan, D. Stauffer: *Int. J. Mod. Phys. C* **9**, 341-347 (1998);
P. N. Ballesteros, L. A. Fernandez, V. Martín-Mayor, et al: *J. Phys. A* **32**, 1-13 (1999);
Youjin Deng, H. W. J. Blöte: *Phys. Rev. E* **72**, 016126 (2005);
M. Acharyya, D. Stauffer: *Int. J. Mod. Phys. C* **9**, 643-647 (2009);
J. Škvor, I. Nezbeda: *Phys. Rev. E* **79**, 041141 (2009).
- [22] M. Sahimi, B. D. Hughes, L. E. Scriven, T. Davis: *J. Phys. A* **16**, L67 (1983);
S. Galam, A. Mauger: *J. Appl. Phys.* **75**, 5526 (1994);
S. Galam, A. Mauger: *Physica A* **205**, 502 (1994);
S. C. van der Marck: *Phys. Rev. E* **55**, 1228 (1997).
- [23] <http://www.rubiks.com/>