Segre maps and entanglement for multipartite systems of indistinguishable particles

Janusz Grabowski*

Faculty of Mathematics and Natural Sciences, College of Sciences Cardinal Stefan Wyszyński University, Wóycickiego 1/3, 01-938 Warszawa, Poland

Marek Kuś[†]

Center for Theoretical Physics, Polish Academy of Sciences, Aleja Lotników 32/46, 02-668 Warszawa, Poland

Giuseppe Marmo[‡]

Dipartimento di Scienze Fisiche, Università "Federico II" di Napoli and Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, Complesso Universitario di Monte Sant Angelo, Via Cintia, I-80126 Napoli, Italy

April 11, 2018

Abstract

We elaborate the concept of entanglement for multipartite system with bosonic and fermionic constituents and its generalization to systems with arbitrary parastatistics. The entanglement is characterized in terms of generalized Segre maps, supplementing thus an algebraic approach to the problem by a more geometric point of view.

Key words: entanglement, tensor product, symmetry group, Bose statistics, Fermi statistics, parastatistics, Young diagram, Segre map.

PACS: 03.65.Aa, 03.67.Mn, 02.10.Xm.

MSC 2000: 81P40, 81P16 (Primary); 15A69, 81R05 (Secondary).

1 Introduction

The possibility of identifying subsystems states in a given total state of a composite quantum system goes under the name of *separability*. In the case of pure states such a possibility is guaranteed if the composite state takes the form of the tensor product of subsystems states.

On the other hand, with the advent of Quantum Field Theory, we have identified elementary particles which are either bosons or fermions. As a matter of fact, according to the spin-statistics theorem all particles are either bosons or fermions. The difference is that a state is unchanged by the interchange of two identical bosons, whereas it changes the sign under the interchange of two identical fermions. The characterization of fermionic states contains already the lack of the factorization of the total state of the

^{*}email: jagrab@impan.pl

[†]email: marek.kus@cft.edu.pl

[‡]email: marmo@na.infn.it

composite system. According to usual wisdom, this would always imply the presence of an entanglement. In our opinion this state of affairs cannot be maintained, so there is a need of a refinement of the notion of entanglement that describes better the situation when we are dealing with bosons and fermions or even with 'parabosons' or 'parafermions' arising from potentially meaningful parastatistics [1, 2].

In [3] we analyzed a concept of entanglement for a multipartite system with bosonic and fermionic constituents in purely algebraic way using the the representation theory of the underlying symmetry groups. Correlation properties of indistinguishable particles become relevant when subsystems are no longer separated by macroscopic distances, like e.g. in quantum gates based on quantum dots, where they are confined to the same spatial regions [4]. In our approach to bosons and fermions we adopted the concept of entanglement put forward in [4, 5] for fermionic systems and extended in [6, 7] in a natural way to bosonic ones¹.

The problem of quantifying and measuring entanglement in systems of many indistinguishable particles remains a topic of vivid investigations from different points of view. Like in the case of distinguishable particles, also nonclassical correlations of indistinguishable particles can be studied with the use of various characteristics, e.g., a suitably adapted concept of the entropy of a state [14], [15] or by mapping the Fock spaces onto spaces of qubits [16], [17], elaborating thus a concept of the so called mode entanglement proposed earlier by Zanardi [18], [19]. Another area of the current theoretical research on indistinguishable particles concentrates around problems of identifying and measuring the entanglement in realistic experimental circumstances [20], [21]. An exhaustive exposition of these aspects of the theory, out of the scope of the present paper, is contained in a recent review article by Tichy, Mintert, and Buchleitner [22].

Our approach appeared to be sufficiently general to define entanglement also for systems with an arbitrary *parastatistics* in a consistent and unified way. For pure states we defined the S-rank, generalizing the notion of the Schmidt rank for distinguishable particles and playing an analogous role in the characterization of the degree of entanglement among particles with arbitrary exchange symmetry (parastatistics).

In the algebraic geometry, a canonical embedding of the product $\mathbb{C}P^{n-1} \times \mathbb{C}P^{m-1}$ of complex projective spaces into $\mathbb{C}P^{nm-1}$ is known under the name the *Segre embedding* (or the *Segre map*). In the quantum mechanical context, the complex projective space $\mathbb{C}P^{n-1}$ represents pure states in the Hilbert space \mathbb{C}^n , and $\mathbb{C}P^{nm-1}$ represent pure states in $\mathbb{C}^n \otimes \mathbb{C}^m$, so that the Segre embedding gives us a geometrical description of separable pure states and, as shown in [8, 9], this description can be extended also to mixed states.

In the present paper we give a geometric description of the entanglement for systems with arbitrary symmetry (with respect to exchanging of subsystems) in terms of *generalized Segre embeddings* associated with particular parastatistics. This description is complementary to the one presented in [3] in terms of the S-rank. For systems with arbitrary exchange symmetries, unlike for the systems of distinguishable particles, the spaces of states are not, in general, projectivizations of the full tensor products of the underlying Hilbert spaces of subsystems, but rather some parts of them. We show in the following how to extend properly the concept of the Segre embedding to achieve a geometric description analogous to that for distinguishable particles. This approach uses a unifying mathematical framework based on the representation theory and strongly suggesting certain concepts of the separability, thus of the entanglement, in the case of indistinguishable particles. For physicists, this approach may be viewed as being too mathematical and abstract, but in our opinion it covers exactly the logic structure of the notion of entanglement for systems of particles with some symmetries.

In the next section we shortly review the relevant concepts of composite systems of distinguishable particles, their description in terms of the classical Segre maps, as well as entanglement measures for systems of distinguishable particles. In sections 3 and 4 we give a brief review of the algebraic description of the entanglement for bosons and fermions in terms of the S-rank of tensors presented in [3]. The main results are contained in Sections 5-7 where we construct the Segre maps, first for bosons and fermions and, finally, for systems with an arbitrary parastatistics. The constructions are based on Theorem 6.1 which relates simple tensor of a given parastatistics with the corresponding Young diagram.

¹For discussion of a slightly different treatment of bosons in [10, 11, 12, 13] see the Introduction and Conclusion sections of [3].

2 Composite systems, separability, and entanglement

Let \mathcal{H} be a Hilbert space with a Hermitian product $\langle \cdot | \cdot \rangle$, $gl(\mathcal{H})$ be the vector space of complex linear operators on \mathcal{H} , $GL(\mathcal{H})$ be the group of invertible operators from $gl(\mathcal{H})$, and $U(\mathcal{H})$ be its subgroup of unitary operators on \mathcal{H} . For simplicity, we will assume that \mathcal{H} is finite-dimensional, say $\dim(\mathcal{H}) = n$, but a major part of our work remains valid also for Hilbert spaces of infinite dimensions. Note only that in the infinite dimensions the corresponding tensor product $\mathcal{H}_1 \otimes \mathcal{H}_2$ is the tensor product in the category of Hilbert spaces, i.e., corresponding to the Hilbert-Schmidt norm.

With $u(\mathcal{H})$ we will denote the Lie algebra of the Lie group $U(\mathcal{H})$ consisting of anti-Hermitian operators, while $u^*(\mathcal{H})$ will denote its dual interpreted as the Euclidean space of Hermitian operators with the scalar product

$$\langle A, B \rangle_{u^*} = \frac{1}{2} \operatorname{Tr}(AB).$$
 (1)

The space of non-negatively defined operators from $gl(\mathcal{H})$, i.e. of those $\rho \in gl(\mathcal{H})$ which can be written in the form $\rho = T^{\dagger}T$ for a certain $T \in gl(\mathcal{H})$, we denote as $\mathcal{P}(\mathcal{H})$. It is a convex cone in $gl(\mathcal{H})$ and the set of *density states* $\mathcal{D}(\mathcal{H})$ is distinguished in $\mathcal{P}(\mathcal{H})$ by the normalizing condition $\operatorname{Tr}(\rho) = 1$. We will regard $\mathcal{P}(\mathcal{H})$ and $\mathcal{D}(\mathcal{H})$ as embedded in $u^*(\mathcal{H})$, so that the space $\mathcal{D}(\mathcal{H})$ of density states is a convex set in the affine hyperplane in $u^*(\mathcal{H})$ determined by the equation $\operatorname{Tr}(\rho) = 1$. As the difference of two Hermitian operators of trace 1 (the vector connection two points in the affine hyperplane) is a Hermitian operator of trace 0, the model vector spaces of this affine hyperplane is therefore canonically identified with the space of Hermitian operators with the trace 0.

Denote the set of all operators from $\mathcal{D}(\mathcal{H})$ of rank k with $\mathcal{D}^k(\mathcal{H})$. It is well known that the set of extreme points of $\mathcal{D}(\mathcal{H})$ coincides with the set $\mathcal{D}^1(\mathcal{H})$ of *pure states*, i.e. the set of one-dimensional orthogonal projectors $\rho_x = |x\rangle\langle x|, ||x|| = 1$. Hence, every element of $\mathcal{D}(\mathcal{H})$ is a convex combination of points from $\mathcal{D}^1(\mathcal{H})$. The space $\mathcal{D}^1(\mathcal{H})$ of all pure states can be identified with the complex projective space $\mathbb{P}\mathcal{H} \simeq \mathbb{C}P^{n-1}$ via the projection

$$\mathcal{H} \setminus \{0\} \ni x \mapsto \rho_x = \frac{|x\rangle\langle x|}{\|x\|^2} \in \mathcal{D}^1(\mathcal{H})$$
(2)

which identifies the points of the orbits of the action in \mathcal{H} of the multiplicative group $\mathbb{C} \setminus \{0\}$ by complex homotheties. It is well known that the complex projective space $\mathbb{P}\mathcal{H} = \mathcal{D}^1(\mathcal{H})$ is canonically a Kähler manifold. The symplectic structure on $\mathcal{D}^1(\mathcal{H}) \subset u^*(\mathcal{H})$ is the canonical symplectic structure of an $U(\mathcal{H})$ -coadjoint orbit, and the metric, called the *Fubini-Study metric* is just the metric induced from the embedding of $\mathcal{D}^1(\mathcal{H})$ into the Euclidean space $u^*(\mathcal{H})$. This is the best known compact Kähler manifold in the algebraic geometry.

Suppose now that our Hilbert space has a fixed decomposition into the tensor product of two Hilbert spaces, $\mathcal{H} = \mathcal{H}^1 \otimes \mathcal{H}^2$. This additional input is crucial in studying composite quantum systems. Observe first that the tensor product map

$$\otimes: \mathcal{H}^1 \times \mathcal{H}^2 \to \mathcal{H} = \mathcal{H}^1 \otimes \mathcal{H}^2 \tag{3}$$

associates the product of rays with a ray, so it induces a canonical embedding on the level of complex projective spaces,

$$\operatorname{Seg}: \mathbb{P}\mathcal{H}^1 \times \mathbb{P}\mathcal{H}^2 \to \mathbb{P}\mathcal{H} = \mathbb{P}(\mathcal{H}^1 \otimes \mathcal{H}^2),$$
(4)

$$(|x^{1}\rangle\langle x^{1}|, |x^{2}\rangle\langle x^{2}|) \quad \mapsto \quad |x^{1}\otimes x^{2}\rangle\langle x^{1}\otimes x^{2}|, \quad ||x^{1}|| = ||x^{2}|| = 1.$$

$$(5)$$

This embedding of the product of complex projective spaces into the projective space of the tensor product is called in the literature the *Segre embedding* [24]. Note that the elements of the image $Seg(\mathbb{P}\mathcal{H}^1 \times \mathbb{P}\mathcal{H}^2)$ in $\mathbb{P}\mathcal{H} = \mathbb{P}(\mathcal{H}^1 \otimes \mathcal{H}^2)$ are usually called *separable pure states* (separable with respect to the decomposition $\mathcal{H} = \mathcal{H}^1 \otimes \mathcal{H}^2$).

The Segre embedding is related to the (external) tensor product of the basic representations of the unitary groups $U(\mathcal{H}^1)$ and $U(\mathcal{H}^2)$, i.e. with the representation of the direct product group in $\mathcal{H} = \mathcal{H}^1 \otimes \mathcal{H}^2$,

$$U(\mathcal{H}^1) \times U(\mathcal{H}^2) \ni (T_1, T_2) \quad \mapsto \quad T_1 \otimes T_2 \in U(\mathcal{H}) = U(\mathcal{H}^1 \otimes \mathcal{H}^2),$$

$$(T_1 \otimes T_2)(x^1 \otimes x^2) \quad = \quad T_1(x^1) \otimes T_2(x^2).$$

Note that $T_1 \otimes T_2$ is unitary, since the Hermitian product in \mathcal{H} is related to the Hermitian products in \mathcal{H}^1 and \mathcal{H}^2 by

$$\langle x^1 \otimes x^2, y^1 \otimes y^2 \rangle_{\mathcal{H}} = \langle x^1, y^1 \rangle_{\mathcal{H}^1} \cdot \langle x^2, y^2 \rangle_{\mathcal{H}^2}.$$
 (6)

The above group embedding gives rise to the corresponding embedding of Lie algebras or, by our identification, of their duals which, with some abuse of notation, we will denote by

Seg: $u^*(\mathcal{H}^1) \times u^*(\mathcal{H}^2) \to u^*(\mathcal{H}), \quad (A, B) \mapsto A \otimes B.$ (7)

The original Segre embedding is just the latter map reduced to pure states. In fact, a stronger result holds true [8, 9].

Proposition 2.1. The embedding (7) maps $\mathcal{D}^k(\mathcal{H}^1) \times \mathcal{D}^l(\mathcal{H}^2)$ into $\mathcal{D}^{kl}(\mathcal{H})$.

Let us denote the image $\text{Seg}(\mathcal{D}^1(\mathcal{H}^1) \times \mathcal{D}^1(\mathcal{H}^2))$, i.e. the set of separable pure states, with $\mathcal{S}^1(\mathcal{H})$, and its convex hull, $conv(\mathcal{S}^1(\mathcal{H}))$, i.e. the set of all mixed separable states in $u^*(\mathcal{H})$, with $\mathcal{S}(\mathcal{H})$. The states from

$$\mathcal{E}(\mathcal{H}) = \mathcal{D}(\mathcal{H}) \setminus \mathcal{S}(\mathcal{H}),$$

i.e. those which are not separable, are called *entangled states*. It is well known (see e.g. [8]) that $S^1(\mathcal{H})$ is exactly the set of extremal points of $S(\mathcal{H})$. What we have just presented is a very simple geometric interpretation of separability and entanglement.

The entangled states play an important role in quantum computing and one of main problems is to decide effectively whether a given composite state is entangled or not. An abstract measurement of entanglement can be based on the following observation (see also [23])

Let *E* be the set of all extreme points of a compact convex set *K* in a finite-dimensional real vector space *V*, and let E_0 be a compact subset of *E* with the convex hull $K_0 = \operatorname{conv}(E_0) \subset K$. For every non-negative function $f: E \to \mathbf{R}_+$ define its extension (*convex roof*) $f_K: K \to \mathbf{R}_+$ by

$$f_K(x) = \inf_{x = \sum t_i \alpha_i} \sum t_i f(\alpha_i), \tag{8}$$

where the *infimum* is taken with respect to all expressions of x in the form of convex combinations of points from E. Recall that, according to Krein-Milman theorem, K is the convex hull of its extreme points.

Proposition 2.2. For every non-negative continuous function $f : E \to \mathbf{R}_+$ which vanishes exactly on E_0 , the function f_K is convex on K and vanishes exactly on K_0

An immediate consequence is the following (cf. [8, 9]).

Corollary 2.1. Let $\mathcal{H} = \mathcal{H}^1 \otimes \mathcal{H}^2$ and let $F : \mathcal{D}^1(\mathcal{H}) \to \mathbf{R}_+$ be a continuous function which vanishes exactly on on the set $\mathcal{S}^1(\mathcal{H})$ of separable pure states. Then,

$$\mu = F_{\mathcal{D}(\mathcal{H})} : \mathcal{D}(\mathcal{H}) \to \mathbf{R}_+$$

is a measure of entanglement, i.e. μ is convex and $\mu(\rho) = 0$ if and only if the (mixed) density state ρ is separable. Moreover, if f is taken $U(\mathcal{H}^1) \times U(\mathcal{H}^2)$ -invariant, then μ is $U(\mathcal{H}^1) \times U(\mathcal{H}^2)$ -invariant.

Remark 2.1. In the terminology of [23], the convex roof function $F_{\mathcal{D}(\mathcal{H})}$ is *entanglement monotone* if F is entanglement monotone on pure states.

3 Tensor algebras, fermions, and bosons

To describe some properties of systems composed of indistinguishable particles and to fix the notation, let us start with introducing corresponding tensor algebras associated with a Hilbert space \mathcal{H} .

In the tensor power $\mathcal{H}^{\otimes k} = \underbrace{\mathcal{H} \otimes \cdots \otimes \mathcal{H}}_{k-\text{times}}$, we distinguish the subspaces: $\mathcal{H}^{\vee k} = \underbrace{\mathcal{H} \vee \cdots \vee \mathcal{H}}_{k-\text{times}}$ of totally symmetric tensors and $\mathcal{H}^{\wedge k} = \underbrace{\mathcal{H} \wedge \cdots \wedge \mathcal{H}}_{k-\text{times}}$ of totally antisymmetric ones, together with the symmetrization, $\pi_k^{\vee} : \mathcal{H}^{\otimes k} \to \mathcal{H}^{\vee k}$, and antisymmetrization, $\pi_k^{\wedge} : \mathcal{H}^{\otimes k} \to \mathcal{H}^{\wedge k}$, projectors:

$$\pi_k^{\vee}(f_1 \otimes \cdots \otimes f_k) = \frac{1}{k!} \sum_{\sigma \in S_k} f_{\sigma(1)} \otimes \cdots \otimes f_{\sigma(k)}, \tag{9}$$

$$\pi_k^{\wedge}(f_1 \otimes \cdots \otimes f_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (-1)^{\sigma} f_{\sigma(1)} \otimes \cdots \otimes f_{\sigma(k)}.$$
(10)

Here, S_k is the group of all permutations $\sigma : \{1, \ldots, k\} \to \{1, \ldots, k\}$, and $(-1)^{\sigma}$ denotes the sign of the permutation σ . Note that with every permutation $\sigma \in S_k$ there is a canonically associated unitary operator U_{σ} on $\mathcal{H}^{\otimes k}$ defined by

$$U_{\sigma}(f_1 \otimes \cdots \otimes f_k) = f_{\sigma(1)} \otimes \cdots \otimes f_{\sigma(k)},$$

so that the map $\sigma \mapsto U_{\sigma}$ is an injective unitary representation of S_k in $\mathcal{H}^{\otimes k}$. We will write simply σ instead of U_{σ} if no misunderstanding is possible. Symmetric and skew-symmetric tensors are characterized in terms of this unitary action by $\sigma(v) = v$ and $\sigma(v) = (-1)^{\sigma}v$, respectively, for all $\sigma \in S_k$

We put, by convention, $\mathcal{H}^{\otimes 0} = \mathcal{H}^{\vee 0} = \mathcal{H}^{\wedge 0} = \mathbb{C}$. It is well known that the obvious structure of a unital graded associative algebra on the graded space $\mathcal{H}^{\otimes} = \bigotimes_{k=0}^{\infty} \mathcal{H}^{\otimes k}$ (the *tensor algebra*) induces canonical unital graded associative algebra structures on the spaces $\mathcal{H}^{\vee} = \bigoplus_{k=0}^{\infty} \mathcal{H}^{\vee k}$ (called the *bosonic Fock space*) and $\mathcal{H}^{\wedge} = \bigoplus_{k=0}^{\infty} \mathcal{H}^{\wedge k}$ (called the *fermionic Fock space*) of symmetric and antisymmetric tensors. This simply means that we have associative multiplications

$$v_1 \vee v_2 = \pi^{\vee}(v_1 \otimes v_2),\tag{11}$$

$$w_1 \wedge w_2 = \pi^{\wedge}(w_1 \otimes w_2), \qquad (12)$$

where

$$\pi^{\vee} = \bigoplus_{k=0}^{\infty} \pi_k^{\vee} : \mathcal{H}^{\otimes} \to \mathcal{H}^{\vee}, \tag{13}$$

and

$$\pi^{\wedge} = \bigoplus_{k=0}^{\infty} \pi_k^{\wedge} : \mathcal{H}^{\otimes} \to \mathcal{H}^{\wedge}, \tag{14}$$

are the symmetrization and antisymmetrization projections. Moreover, these multiplications respect the grading, i.e. $v_1 \vee v_2 \in \mathcal{H}^{\vee(k+l)}$ if $v_1 \in \mathcal{H}^{\vee k}$, $v_2 \in \mathcal{H}^{\vee l}$, and $w_1 \wedge w_2 \in \mathcal{H}^{\wedge(k+l)}$ if $w_1 \in \mathcal{H}^{\wedge k}$, $w_2 \in \mathcal{H}^{\wedge l}$. Note also that the multiplication in \mathcal{H}^{\vee} is commutative, $v_1 \vee v_2 = v_2 \vee v_1$, and the multiplication in \mathcal{H}^{\wedge} is graded commutative, $w_1 \wedge w_2 = (-1)^{k_1 \cdot k_2} w_2 \wedge w_1$, for $w_i \in \mathcal{H}^{\wedge k_i}$.

It is well known that the symmetric tensor algebra \mathcal{H}^{\vee} can be canonically identified with the algebra $Pol(\mathcal{H})$ of polynomial functions on \mathcal{H} . Indeed, any $f \in \mathcal{H}$ can be identified with the linear function x_f on \mathcal{H} by means of the Hermitian product: $x_f(y) = \langle f | y \rangle$. We must stress, however, that the identification $f \mapsto x_f$ is anti-linear. This can be extended to an anti-linear isomorphism of commutative algebras in which $f_1 \vee \cdots \vee f_k$ corresponds to the homogenous polynomial $x_{f_1} \cdots x_{f_k}$. Similarly, one identifies \mathcal{H}^{\wedge} with the Grassmann algebra $Grass(\mathcal{H})$ of polynomial (super)functions on \mathcal{H} . Here, however, with $f \in \mathcal{H}$ we associate a linear function ξ_f on \mathcal{H} regarded as and odd function: $\xi_f \xi_{f'} = -\xi_{f'} \xi_f$. In the language of supergeometry one speaks about the purely odd vector space $\Pi \mathcal{H}$ obtained from \mathcal{H} by changing the parity.

If we fix a basis e_1, \ldots, e_n in \mathcal{H} and associate with its elements even linear functions x_1, \ldots, x_n on \mathcal{H} , and odd linear functions ξ_1, \ldots, ξ_n on $\Pi \mathcal{H}$, then \mathcal{H}^{\vee} can be identified with the algebra of complex polynomials in n commuting variables, $\mathcal{H}^{\vee} \simeq \mathbb{C}[x_1, \ldots, x_n]$. Similarly, $\mathcal{H}^{\wedge} \simeq \mathbb{C}[\xi_1, \ldots, \xi_n]$, i.e., \mathcal{H}^{\wedge} can be identified with the algebra of complex Grassmann polynomials in n anticommuting variables. The

subspaces $\mathcal{H}^{\vee k}$ and $\mathcal{H}^{\wedge k}$ correspond to homogenous polynomials of degree k. It is straightforward that homogeneous polynomials $x_1^{k_1} \cdots x_n^{k_n}$, with $k_1 + \cdots + k_n = k$, form a basis of $\mathcal{H}^{\vee k}$, while homogeneous Grassmann polynomials $\xi_{i_1} \wedge \cdots \wedge \xi_{i_k}$, with $1 \leq i_1 < i_2 < \cdots < i_k \leq n$, form a basis of $\mathcal{H}^{\wedge k}$. In consequence, dim $\mathcal{H}^{\vee k} = \binom{n+k-1}{k}$ and dim $\mathcal{H}^{\wedge k} = \binom{n}{k}$, so the gradation in the fermionic Fock space is finite-dimensional (for a finite-dimensional \mathcal{H}).

Note that any basis $\{e_1, \ldots, e_n\}$ in \mathcal{H} induces a basis $\{e_{i_1} \otimes e_{i_2} \otimes \cdots \otimes e_{i_k} \mid i_1, \ldots, i_k \in \{1, \ldots, n\}\}$ in $\mathcal{H}^{\otimes k}$. Therefore, any $u \in \mathcal{H}^{\otimes k}$ can be uniquely written as a linear combination

$$u = \sum_{i_1,\dots,i_k=1}^n u^{i_1\dots i_k} e_{i_1} \otimes \dots \otimes e_{i_k}.$$
(15)

If $u \in \mathcal{H}^{\vee k}$, then the tensor coefficients $u^{i_1 \dots i_k}$ are totally symmetric and, after applying the symmetrization projection to (15), we get

$$u = \sum_{i_1, \dots, i_k=1}^n u^{i_1 \dots i_k} e_{i_1} \vee \dots \vee e_{i_k}.$$
 (16)

Similarly, if $u \in \mathcal{H}^{\wedge k}$, the tensor coefficients $u^{i_1 \dots i_k}$ are totally antisymmetric and

$$u = \sum_{i_1, \dots, i_k=1}^n u^{i_1 \dots i_k} e_{i_1} \wedge \dots \wedge e_{i_k}.$$
 (17)

We will refer to the coefficients $u^{i_1...i_k}$ as to the *coefficients of* u *in the basis* $\{e_1, \ldots, e_n\}$.

The Hermitian product in \mathcal{H} has an obvious extension to a Hermitian product in $\mathcal{H}^{\otimes k}$,

$$\langle f_1 \otimes \cdots \otimes f_k | g_1 \otimes \cdots \otimes g_k \rangle = \prod_{i=1}^k \langle f_i | g_i \rangle,$$
 (18)

and viewing symmetric and antisymmetric tensors as canonically embedded in the tensor algebra, we find the corresponding Hermitian products in $\mathcal{H}^{\vee k}$ and $\mathcal{H}^{\wedge k}$.

For $f_1, \ldots, f_k \in \mathcal{H}$ and $g_1, \ldots, g_k \in \mathcal{H}$, we get

$$\langle f_1 \vee \dots \vee f_k | g_1 \vee \dots \vee g_k \rangle = \frac{1}{(k!)^2} \sum_{\sigma, \tau \in S_k} \prod_{i=1}^k \langle f_{\sigma(i)} | g_{\tau(i)} \rangle = \frac{1}{k!} \operatorname{per}(\langle f_i | g_j \rangle).$$
(19)

Here, $\frac{1}{k!} \sum_{\tau \in S_k} \prod_{i=1}^k a_{i\tau(i)} = \text{per}(a_{ij})$ is the permanent of the matrix $A = (a_{ij})$. Similarly,

$$\langle f_1 \wedge \dots \wedge f_k | g_1 \wedge \dots \wedge g_k \rangle = \frac{1}{k!} \det(\langle f_i | g_j \rangle).$$
 (20)

These Hermitian products can be generalized to certain 'pairings' (contractions or inner products) between $\mathcal{H}^{\vee k}$ and $\mathcal{H}^{\vee l}$ on one hand, and $\mathcal{H}^{\wedge k}$ and $\mathcal{H}^{\wedge l}$ on the other, $l \leq k$. For the standard simple tensors $f = f_1 \otimes \cdots \otimes f_k \in \mathcal{H}^{\otimes k}$ and $g = g_1 \otimes \cdots \otimes g_l \in \mathcal{H}^{\otimes l}$, we just put

$$i_g f = \langle f_1 \otimes \cdots \otimes f_l | g_1 \otimes \cdots \otimes g_l \rangle f_{l+1} \otimes \cdots \otimes f_k$$

and extend it by linearity to all tensors. It is easy to see now that, if $v = f_1 \vee \cdots \vee f_k \in \mathcal{H}^{\vee k} \subset \mathcal{H}^{\otimes k}$ and $\nu = g_1 \vee \cdots \vee g_l \in \mathcal{H}^{\vee l} \subset \mathcal{H}^{\otimes l}$, then $\iota_{\nu} v \in \mathcal{H}^{\vee (k-l)}$.

Similarly, $\iota_{\omega} w \in \mathcal{H}^{\wedge (k-l)}$, if $w \in \mathcal{H}^{\wedge k} \subset \mathcal{H}^{\otimes k}$ and $\omega \in \mathcal{H}^{\wedge l} \subset \mathcal{H}^{\otimes l}$. Explicitly,

$$i_{g_{1}\vee\cdots\vee g_{l}}f_{1}\vee\cdots\vee f_{k} = \frac{1}{k!\,l!}\sum_{\substack{\sigma\in S_{k}\\\tau\in S_{l}}}\prod_{j=1}^{l}\langle f_{\sigma(j)}|g_{\tau(j)}\rangle f_{\sigma(l+1)}\otimes\cdots\otimes f_{\sigma(k)}$$
$$= \frac{(k-l)!}{k!}\sum_{\substack{S\in S(l,k-l)\\\tau\in S_{l}}}\prod_{j=1}^{l}\langle f_{S(j)}|g_{\tau(j)}\rangle f_{S(l+1)}\vee\cdots\vee f_{S(k)},$$
(21)

where S(l, k-l) denotes the group of all (l, k-l) shuffles. Recall that a permutation τ in S_{p+q} is a (p,q) shuffle if $\tau(1) < \cdots < \tau(p)$ and $\tau(p+1) < \cdots < \tau(p+q)$.

For skew-symmetric tensors,

$$i_{g_1 \wedge \dots \wedge g_l} f_1 \wedge \dots \wedge f_k = \frac{1}{k! \, l!} \sum_{\substack{\sigma \in S_k \\ \tau \in S_l}} (-1)^{\sigma} (-1)^{\tau} \prod_{j=1}^l \langle f_{\sigma(j)} | g_{\tau(j)} \rangle f_{\sigma(l+1)} \otimes \dots \otimes f_{\sigma(k)}$$
$$= \frac{(k-l)!}{k!} \sum_{\substack{S \in S(l,k-l) \\ \tau \in S_l}} (-1)^{\sigma} (-1)^{\tau} \prod_{j=1}^l \langle f_{S(j)} | g_{\tau(j)} \rangle f_{S(l+1)} \wedge \dots \wedge f_{S(k)}.$$
(22)

In particular,

$$i_{g_1 \vee \cdots \vee g_k} f_1 \vee \cdots \vee f_k = \langle f_1 \vee \cdots \vee f_k | g_1 \vee \cdots \vee g_k \rangle, \tag{23}$$

and

$$i_{g_1 \wedge \dots \wedge g_k} f_1 \wedge \dots \wedge f_k = \langle f_1 \wedge \dots \wedge f_k | g_1 \wedge \dots \wedge g_k \rangle.$$
⁽²⁴⁾

Moreover,

$$i_{g_1 \vee \dots \vee g_{k-1}} f_1 \vee \dots \vee f_k = \frac{1}{k!} \sum_{j=1}^k \langle f_1 \vee \overset{j}{\overset{\vee}{\dots}} \vee f_k | g_1 \vee \dots \vee g_{k-1} \rangle f_j , \qquad (25)$$

and

$$g_1 \wedge \dots \wedge g_{k-1} f_1 \wedge \dots \wedge f_k = \frac{1}{k!} \sum_{j=1}^k (-1)^{k-j} \langle f_1 \wedge \cdots \rangle \langle f_k | g_1 \wedge \dots \wedge g_{k-1} \rangle f_j, \qquad (26)$$

where $\stackrel{j}{\vee}$ stands for the omission.

ı

4 The S-rank and entanglement for multipartite Bose and Fermi systems

There are many concepts of a rank of a tensor used in describing its complexity. One of the simplest and most natural is the one based on the inner product operators defined in the previous section. This rank, called in [3] the *S*-rank and used there to define the entanglement for systems of indistinguishable particles, is a natural generalization of the *Schmidt rank* of 2-tensors.

Definition 4.1. Let $u \in \mathcal{H}^{\otimes k}$. By the *S*-rank of u, we understand the maximum of dimensions of the linear spaces $i_{\mathcal{H}}^{k-1}\sigma(u)$, for $\sigma \in S_k$, which are the images of the contraction maps

$$\mathcal{H}^{\otimes (k-1)} \ni \nu \mapsto \imath_{\nu} \sigma(u) \in \mathcal{H}.$$
(27)

Non-zero tensors of minimal S-rank in $\mathcal{H}^{\otimes k}$ (resp., $\mathcal{H}^{\vee k}$, $\mathcal{H}^{\wedge k}$) we will call simple (resp., simple symmetric, simple antisymmetric).

Note that he above definition has its natural counterpart for distinguishable particles, so tensors from $\mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_k$. We just do the contractions with tensors from $\mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_{k-1}$ and the corresponding permutations. If particles are identical, $\mathcal{H}_i = \mathcal{H}$, and indistinguishable, e.g. the tensors are symmetric or skew-symmetric, we can skip using permutations. In other words, for $u \in \mathcal{H}^{\vee k}$ (resp., $u \in \mathcal{H}^{\wedge k}$), the S-rank of u equals the dimension of the linear space which is the image of the contraction map,

$$\mathcal{H}^{\vee(k-1)} \ni \nu \mapsto \imath_{\nu} u \in \mathcal{H},\tag{28}$$

(resp.,

$$\mathcal{H}^{\wedge(k-1)} \ni \nu \mapsto \imath_{\nu} u \in \mathcal{H}). \tag{29}$$

Theorem 4.1. ([3])

(a) The minimal possible S-rank of a non-zero tensor $u \in \mathcal{H}^{\otimes k}$ equals 1. A tensor $u \in \mathcal{H}^{\otimes k}$ is of S-rank 1 if and only if u is decomposable, i.e., it can be written in the form

$$u = f_1 \otimes \dots \otimes f_k, \quad f_i \in \mathcal{H}, \quad f_i \neq 0.$$
(30)

Such tensors span $\mathcal{H}^{\otimes k}$.

(b) The minimal possible S-rank of a non-zero tensor $v \in \mathcal{H}^{\vee k}$ equals 1. A tensor $v \in \mathcal{H}^{\vee k}$ is of S-rank 1 if and only if v can be written in the form

$$v = f \lor \dots \lor f, \quad f \in \mathcal{H}, \quad f \neq 0.$$
(31)

Such tensors span $\mathcal{H}^{\vee k}$.

(c) The minimal possible S-rank of a non-zero tensor $w \in \mathcal{H}^{\wedge k}$ equals k. A tensor $w \in \mathcal{H}^{\wedge k}$ is of S-rank k if and only if w can be written in the form

$$w = f_1 \wedge \dots \wedge f_k,\tag{32}$$

where $f_1, \ldots, f_k \in \mathcal{H}$ are linearly independent. Such tensors span $\mathcal{H}^{\wedge k}$.

In particular, the S-rank is 1 for simple and simple symmetric tensors and it is k for simple antisymmetric tensors from $\mathcal{H}^{\wedge k}$. Simple tensors have the form (30), simple symmetric tensors have the form (31), and simple antisymmetric tensors have the form (32).

Using the concept of simple tensors we can define simple (non-entangled or separable) and entangled pure states for multipartite systems of bosons and fermions.

Definition 4.2.

- (a) A pure state ρ_x on $\mathcal{H}^{\vee k}$ (resp., on $\mathcal{H}^{\wedge k}$), $\rho_x = \frac{|x \rangle \langle x|}{||x||^2}$, with $x \in \mathcal{H}^{\vee k}$ (resp., $x \in \mathcal{H}^{\wedge k}$), $x \neq 0$, is called a bosonic (resp., fermionic) simple (or non-entangled) pure state if x is a simple symmetric (resp., antisymmetric) tensor. If x is not simple symmetric (resp., antisymmetric), we call ρ_x a bosonic (resp., fermionic) entangled state.
- (b) A mixed state ρ on $\mathcal{H}^{\vee k}$ (resp., on $\mathcal{H}^{\wedge k}$) we call *bosonic* (resp., *fermionic*) *simple* (or *non-entangled*) *mixed state* if it can be written as a convex combination of bosonic (resp., fermionic) simple pure states. In the other case, ρ is called *bosonic* (resp., *fermionic*) *entangled mixed state*.

According to Theorem 4.1, bosonic simple pure k-states are of the form

$$|e \lor \cdots \lor e\rangle\langle e \lor \cdots \lor e|$$

for unit vectors $e \in \mathcal{H}$, and fermionic simple pure k-states are of the form

$$k!|e_1 \wedge \cdots \wedge e_k\rangle\langle e_1 \wedge \cdots \wedge e_k|$$

for orthonormal systems e_1, \ldots, e_k in \mathcal{H} .

Fixing a base in \mathcal{H} results in defining coefficients $[u^{i_1...i_k}]$ of $u \in \mathcal{H}^{\otimes k}$. Formulae characterizing simple tensors, thus simple pure states, can be written in terms of quadratic equations with respect to these coefficients as follows. The corresponding characterization of entangled pure states are obtained by negation of the latter.

Theorem 4.2. ([3])

(a) The pure state ρ_u , associated with a tensor $u = [u^{i_1 \dots i_k}] \in \mathcal{H}^{\otimes k}$, is entangled if and only if there exist $i_1, \dots, i_k, j_1, \dots, j_k$, and $s = 1, \dots, k$ such that

$$u^{i_1...i_s...i_k} u^{j_1...j_s...j_k} \neq u^{i_1...j_s...i_k} u^{j_1...i_s...j_k}.$$
(33)

(b) The bosonic pure state ρ_v , associated with a symmetric tensor $v = [v^{i_1 \dots i_k}] \in \mathcal{H}^{\vee k}$, is bosonic entangled if and only if there exist $i_1, \dots, i_k, j_1, \dots, j_k$, such that

$$v^{i_1\dots i_{k-1}i_k}v^{j_1\dots j_{k-1}j_k} \neq v^{i_1\dots i_{k-1}j_k}v^{j_1\dots j_{k-1}i_k}.$$
(34)

(c) The fermionic pure state ρ_w , associated with an antisymmetric tensor $w = [w^{i_1 \dots i_k}] \in \mathcal{H}^{\wedge k}$, is fermionic entangled if and only if there exist $i_1, \dots, i_{k+1}, j_1, \dots, j_{k-1}$ such that

$$w^{[i_1\dots i_k}w^{i_{k+1}]j_1\dots j_{k-1}} \neq 0,$$
(35)

where the left-hand side is the antisymmetrization of $w^{i_1...i_k}w^{i_{k+1}j_1...j_{k-1}}$ with respect to the indices i_1, \ldots, i_{k+1} .

Note that the opposite to (35), $w^{[i_1...i_k}w^{i_{k+1}]j_1...j_{k-1}} = 0$, are sometimes called the *Plücker relations*.

Example 4.1. Assume that deal with qubit systems, and $|0\rangle$, $|1\rangle$ is an orthonormal basis in \mathcal{H} . The tensor $u = |0\rangle \otimes |0\rangle$ has the S-rank 1:

$$\imath_{a|0\rangle+b|1\rangle}\sigma(u)=\imath_{a|0\rangle+b|1\rangle}(u)=(a\langle 0|+b\langle 1|)|0\rangle|0\rangle=a|0\rangle$$

which is the 1-dimensional space spanned by $|0\rangle$, while the tensor $u_{\pm} = |0\rangle \otimes |1\rangle \pm |1\rangle \otimes |0\rangle$ has the S-rank 2:

$$\begin{split} u_{a|0\rangle+b|1\rangle}\sigma(u) &= \pm u_{a|0\rangle+b|1\rangle}u_{\pm} = \\ &\pm (a\langle 0|+b\langle 1|)|0\rangle|1\rangle \pm (a\langle 0|+b\langle 1|)|1\rangle|0\rangle = \pm a|1\rangle \pm b|0\rangle \,. \end{split}$$

Example 4.2. For the GHZ-states $|GHZ_k\rangle$ and W-states $|W_k\rangle$ the S-rank is 2 independently on $k \ge 2$. Indeed, it is clear that contractions of

$$|GHZ_k\rangle = \frac{1}{\sqrt{2}} \left(|0\rangle^{\otimes k} + |1\rangle^{\otimes k} \right)$$

with (k-1)-tensors give all linear combinations of $|0\rangle$ and $|1\rangle$. The same is true for

$$|W_k\rangle = \frac{1}{\sqrt{k}} \left(|0\cdots 01\rangle + |0\cdots 10\rangle + \cdots + |1\cdots 00\rangle\right).$$

On the other hand, the S-rank cannot exceed 2 for qubit systems.

Remark 4.1. The S-rank does not distinguish between $|GHZ_k\rangle$ and $|W_k\rangle$. Note however that we can slightly generalize our notion of the S-rank including also contractions $\iota_{\nu} u$ with shorter tensors, i.e. tensors $\nu \in \mathcal{H}^{\otimes (k-l)}$ with 0 < l < k. We have not insist on this generalization in order to avoid additional technical complications. The simple version of the S-rank is sufficient for distinguishing simple tensors. This extended version could be useful in measuring the entanglement.

Example 4.3. If $|i_1\rangle$ and $|i_2\rangle$ are orthonormal sets in \mathcal{H}_1 and \mathcal{H}_2 , then the S-rank of $u\sum_{i=1}^r \lambda_i |i_1\rangle \otimes |i_2\rangle$ is r, as $\iota_{\mathcal{H}_1} u$ is spanned by $|i_2\rangle$, $i = 1, \ldots, r$, and $\iota_{\mathcal{H}_2} \sigma(u)$ for the transposition $\sigma(f \otimes g) = g \otimes f$ is spanned by $|i_1\rangle$, $i = 1, \ldots, r$. In other words the S-rank equals the Schmidt rank in this case, so the S-rank is a natural generalization of the latter.

5 Entanglement and Segre maps for Bose and Fermi statistics

Similarly to the case of distinguishable particles (see [8, 9]), the sets of all bosonic simple pure states (resp., fermionic simple pure states) can be described as the images of certain maps defined on the products of projective Hilbert spaces, the generalized Segre maps, as follows.

Consider first the standard Segre embedding Seg_k induced by the tensor product map:

where $\mathcal{H}_{\circ} = \mathcal{H} \setminus \{0\}.$

It is clear that the analogous map, $\operatorname{Seg}_k^{\vee}$, for the Bose statistics should be

where $x^k = x \vee \cdots \vee x = x \otimes \cdots \otimes x$ (k-factors), and for the Fermi statistics:

where $\mathcal{H}_{\circ}^{\times k}$ denotes

$$\mathcal{H}^{\times k} \setminus \{(x_1, \dots, x_k) : x_1 \wedge \dots \wedge x_k = 0\}$$

and $(\mathbb{P}\mathcal{H})^{\times k}_{\circ}$ is

$$(\mathbb{P}\mathcal{H})^{\times k} \setminus \{(\rho_{x_1}, \dots, \rho_{x_k}) : x_1 \wedge \dots \wedge x_k = 0\}.$$

Note that the condition $x_1 \wedge \cdots \wedge x_k \neq 0$ does not depend on the choice of the vectors x_1, \ldots, x_k in their projective classes and means that $\rho_{x_1}, \ldots, \rho_{x_k}$ do not lie in a common projective hyperspace. The subset $\mathcal{H}_{\circ}^{\times k}$ (resp., $(\mathbb{P}\mathcal{H})_{\circ}^{\times k}$) is open and dense in $\mathcal{H}^{\times k}$ (resp., $(\mathbb{P}\mathcal{H})^{\times k}$). The following is an immediate consequence of Theorem 4.1.

Theorem 5.1. A bosonic (fermionic) pure state $\rho \in \mathbb{P}(\mathcal{H}^{\vee k})$ (resp., $\rho \in \mathbb{P}(\mathcal{H}^{\wedge k})$) is entangled if and only if it lies outside the range of the Segre map

$$\operatorname{Seg}_{k}^{\vee} : \mathbb{P}\mathcal{H} \to \mathbb{P}(\mathcal{H}^{\vee k}) \quad (\operatorname{resp.}, \quad \operatorname{Seg}_{k}^{\wedge} : (\mathbb{P}\mathcal{H})_{\circ}^{\times k} \to \mathbb{P}(\mathcal{H}^{\wedge k})).$$

A mixed bosonic (fermionic) state is entangled if and only if it lies outside the convex hull of the range of the corresponding Segre map.

6 Entanglement for generalized parastatistics

Our approach to the entanglement of composite systems for identical particles is so general and natural that it allows for an immediate implications also for generalized parastatistics. Parastatistics were introduced by Green [1] as a refinement of the spin-statistics connection introduced by Pauli [25]. Green was motivated by a two-pages paper by Wigner addressing the connection between equations of motion and the commutation relations (Wigner's problem) [26, 27]. The context of Green's paper is Quantum Field Theory, while most of the applications which have been proposed deal with thermodynamical aspects, in particular with the calculation of the partition function. In this paper we are concerned with parastatistics only to show that our proposed scheme applies to all the situations where states of composite systems are by construction not factorizable and are associated with representations of the permutation group acting on the tensor product of states of the subsystems. The reader who wants to know more about parastatistics could read [2, 28, 29].

Observe first that simple tensors of length 1 in $\tilde{\mathcal{H}} = \mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_k$ form an orbit of the group $U(\mathcal{H}_1) \times \cdots \times U(\mathcal{H}_k)$ acting on $\tilde{\mathcal{H}}$ in the obvious way. In fact, each such tensor can be written as $e_1^1 \otimes \cdots \otimes e_1^k$

for certain choice of orthonormal bases $e_1^j, \ldots, e_{n_j}^j$ in \mathcal{H}_j , $j = 1, \ldots, k$. This means that simple tensors are just vectors of highest (or lowest, depending on the convention) weight of the compact Lie group $U(\mathcal{H}_1) \times \cdots \times U(\mathcal{H}_k)$ relative to some choice of a maximal torus and Borel subgroups. If indistinguishable particles are concerned, the symmetric and antisymmetric tensors form particular irreducible parts of the 'diagonal' representation of the compact group $U(\mathcal{H})$ in the Hilbert space $\mathcal{H}^{\otimes k}$, defined by

$$U(x_1 \otimes \dots \otimes x_k) = U(x_1) \otimes \dots \otimes U(x_k).$$
(39)

Recall that we identify the symmetry group S_k with the group of certain unitary operators on the Hilbert space \mathcal{H}^k in the obvious way,

$$\sigma(x_1 \otimes \cdots \otimes x_k) = x_{\sigma(1)} \otimes \cdots \otimes x_{\sigma(k)}.$$

Note that the operators of S_k intertwine the unitary action of $U(\mathcal{H})$. In the cases of the symmetric and antisymmetric tensors, we speak about the Bose and Fermi statistics, respectively. But, for k > 2, there are other irreducible parts of the representation (39) associated with invariant subspaces of the S_k action. We shall call them *(generalized) parastatistics*. Any of these k-parastatistics (i.e. any irreducible subspace of the tensor product $\mathcal{H}^{\otimes k}$) is associated with a Young tableau α with k-boxes (chambers) as follows (see e.g. [30, 31, 32]).

Consider partitions of $k: k = \lambda_1 + \cdots + \lambda_r$, where $\lambda_1 \ge \cdots \ge \lambda_r \ge 1$. To a partition $\lambda = (\lambda_1, \ldots, \lambda_r)$ is associated a Young diagram (sometimes called a Young frame or a Ferrers diagram) with λ_i boxes in the *i*th row, the rows of boxes lined up on the left. Define a tableau on a given Young diagram to be a numbering of the boxes by the integers $1, \ldots, k$, and denote with Y_{λ} the set of all such Young tableaux. Finally, put Y(k) to be the set of all Young tableaux with k boxes. Given a tableau $\alpha \in Y(k)$ define two subgroups in the symmetry group S_k :

$$P = P_{\alpha} = \{ \sigma \in S_k : \sigma \text{ preserves each row of } \alpha \}$$

and

 $Q = Q_{\alpha} = \{ \tau \in S_k : \tau \text{ preserves each column of } \alpha \}.$

In the space of linear operators on $\mathcal{H}^{\otimes k}$ we introduce two operators associated with these subgroups:

$$a_{\alpha} = \sum_{\tau \in Q} (-1)^{\tau} \tau, \quad b_{\alpha} = \sum_{\sigma \in P} \sigma.$$
(40)

Finally, we define the Young symmetrizer

$$c_{\alpha} = a_{\alpha} \circ b_{\alpha} = \sum_{\sigma \in P, \ \tau \in Q} (-1)^{\tau} \tau \circ \sigma \,.$$

$$\tag{41}$$

It is well known (see, e.g., [30, 31, 32]) that $\pi^{\alpha} = \frac{1}{\mu(\alpha)}c_{\alpha}$, for some non-zero rational number $\mu(\alpha)$, is an orthogonal projector and that the image \mathcal{H}^{α} of c_{α} is an irreducible subrepresentation of $U(\mathcal{H})$, i.e. the *parastatistics associated with* α . As a matter of fact, these representations for Young tableaux on the same Young diagram are equivalent, so that the constant $\mu(\alpha)$ depends only on the Young diagram λ of α and does not depend on the enumeration of boxes. Hence, $\mu(\alpha) = \mu(\lambda)$ and this number is related to the multiplicity $m(\lambda)$ of the corresponding irreducible representation in $\mathcal{H}^{\otimes k}$ by $\mu(\lambda) \cdot m(\lambda) = k!$. For a given Young diagram (partition) λ , the map

$$\epsilon_{\lambda} = \frac{1}{\mu(\lambda)^2} \sum_{\alpha \in Y_{\lambda}} c_{\alpha} \tag{42}$$

is an orthogonal projection, called the *central Young symmetrizer*, onto the invariant subspace being the sum of all copies of the irreducible representations equivalent to that with a parastatistics from Y_{λ} .

The symmetrization π^{\vee} (antisymmetrization π^{\wedge}) projection corresponds to a Young tableau with just one row (one column) and arbitrary enumeration. It is well known that any irreducible representation \mathcal{H}^{α} of $U(\mathcal{H})$ contains cyclic vectors which are of highest weight relative to some choice of a maximal torus and Borel subgroups in $U(\mathcal{H})$. We will call them α -simple tensors or simple tensors in \mathcal{H}^{α} . Note that such vectors can be viewed as generalized coherent states [33]. They can be also regarded as the 'most classical' states with respect to their correlation properties [34]. These are exactly the tensors associated with simple (non-entangled) pure states for composite systems of particles with (generalized) parastatistics. This is because α -simple tensors represent the minimal amount of quantum correlations for tensors in \mathcal{H}^{α} , namely the quantum correlations forced directly by the particular parastatistics.

- **Example 6.1.** (a) For k = 2 we have just the obvious splitting of $\mathcal{H}^{\otimes 2}$ into symmetric and antisymmetric tensors: $\mathcal{H}^{\wedge 2} \oplus \mathcal{H}^{\vee 2}$.
- (b) For k = 3, besides symmetric and antisymmetric tensors associated with the Young tableaux

$$\alpha_0 = \boxed{1 \ 2 \ 3}$$
 and $\alpha_3 = \boxed{2 \ 3}$

we have two additional irreducible parts associated with the Young tableaux

$$\alpha_1 = \boxed{\frac{1}{3}} \quad and \quad \alpha_2 = \boxed{\frac{1}{2}}, \tag{43}$$

namely

$$\mathcal{H}^{\otimes 3} = \mathcal{H}^{\wedge 3} \oplus \mathcal{H}^{\alpha_1} \oplus \mathcal{H}^{\alpha_2} \oplus \mathcal{H}^{\vee 3} \,. \tag{44}$$

Since $P_{\alpha_1} = \{id, (1,2)\}$ and $Q_{\alpha_1} = \{id, (1,3)\}$, we get $a_{\alpha_1} = id - (1,3)$ and $b_{\alpha_1} = id + (1,2)$, so that

$$c_{\alpha_1} = a_{\alpha_1} \circ b_{\alpha_1} = (id - (1,3)) \circ (id + (1,2)) = id + (1,2) - (1,3) - (123).$$

As the multiplicity of the representation is 2, we have $\mu(\alpha_1) = 3!/2 = 3$ and the projection

$$\pi^{\alpha_1}: \mathcal{H}^{\otimes 3} \to \mathcal{H}^{\alpha_1},\tag{45}$$

takes the form

$$\pi^{\alpha_1}(x_1 \otimes x_2 \otimes x_3) = \frac{1}{3}(x_1 \otimes x_2 \otimes x_3 + x_2 \otimes x_1 \otimes x_3 - x_3 \otimes x_2 \otimes x_1 - x_3 \otimes x_1 \otimes x_2).$$
(46)

Similarly,

$$\pi^{\alpha_2}: \mathcal{H}^{\otimes 3} \to \mathcal{H}^{\alpha_2},\tag{47}$$

$$\pi^{\alpha_2}(x_1 \otimes x_2 \otimes x_3) = \frac{1}{3}(x_1 \otimes x_2 \otimes x_3 + x_3 \otimes x_2 \otimes x_1 - x_2 \otimes x_1 \otimes x_3 - x_2 \otimes x_3 \otimes x_1).$$
(48)

The simple tensors (the highest weight vectors) in \mathcal{H}^{α_1} can be written as

$$v_{\lambda}^{\alpha_1} = \lambda(e_1 \otimes e_1 \otimes e_2 - e_2 \otimes e_1 \otimes e_1), \tag{49}$$

for certain choice of an orthonormal basis e_i in \mathcal{H} and $\lambda \neq 0$. Analogously, the simple tensors in \mathcal{H}^{α_2} , in turn, take the form

$$v_{\lambda}^{\alpha_2} = \lambda(e_1 \otimes e_2 \otimes e_1 - e_2 \otimes e_1 \otimes e_1).$$
(50)

For dim $(\mathcal{H}) = 3$, the simple tensors of length 1 form an orbit of the unitary group $U(\mathcal{H})$ of the (real) dimension 7 in \mathcal{H}^{α_1} and \mathcal{H}^{α_2} . The simple symmetric tensors of length 1 form an orbit of the dimension 5, and the simple antisymmetric ones (of length 1) – an orbit of the dimension 1. The dimensions of the irreducible representations are: dim $(\mathcal{H}^{\wedge 3}) = 1$, dim $(\mathcal{H}^{\vee 3}) = 10$, dim $(\mathcal{H}^{\alpha_1}) = \dim(\mathcal{H}^{\alpha_2}) = 8$.

A fundamental observation is that α -simple tensors can also be characterized in terms of the S-rank.

Theorem 6.1. A tensor $v \in \mathcal{H}^{\alpha}$, $\alpha \in Y(k)$, is simple if and only if it has the minimal S-rank among non-zero tensors from \mathcal{H}^{α} . This minimal S-rank equals the number r of rows in the corresponding Young diagram and the simple tensor reads as

$$v = \pi^{\alpha} \left(e_{\alpha(1)} \otimes \dots \otimes e_{\alpha(k)} \right) \,, \tag{51}$$

where e_1, \ldots, e_r are some linearly independent vectors in \mathcal{H} and $\alpha(i)$ is the number of the row in which the box with the number i appears in the tableaux α . In other words, the tensor

$$E_{\alpha} = e_{\alpha(1)} \otimes \dots \otimes e_{\alpha(k)} \tag{52}$$

is the tensor product of k vectors from the sequence $E = (e_1, \ldots, e_r)$ obtained by putting e_j in the places indicated by the number of the boxes in the *j*th row.

Proof. Assume that v is of the form (51). Passing to the complexification $GL(\mathcal{H})$ of $U(\mathcal{H})$ and using a basis e_1, \ldots, e_n extending the linear independent family $e_1, \ldots, e_r \in \mathcal{H}$ to identify $GL(\mathcal{H})$ with $GL(n; \mathbb{C})$, we easily see that E_{α} , thus v, is an eigenvector for any diagonal matrix and is killed by any upper-triangular matrix. Moreover, $v \neq 0$, hence v is a vector of highest weight. Indeed, by definition $b_{\alpha}(E_{\alpha})$ is non-zero and proportional to E_{α} , so that $v = t \cdot a_{\alpha}(E_{\alpha})$ for a non-zero constant t. It is now easy to see that $\{\tau(E_{\alpha}) : \tau \in Q\}$ is a family of linearly independent tensors in $\mathcal{H}^{\otimes k}$, so that $v = \sum_{\tau \in Q} (-1)^{\tau} \tau(E_{\alpha})$ is non-zero. As tensors from $\{\tau(E_{\alpha}) : \tau \in Q\}$ are linearly independent and have one of e_1, \ldots, e_r as the last factor, the S-rank of v is at least r. On the other hand, v is composed from tensor products of e_1, \ldots, e_r only, so the S-rank is at most r.

Conversely, let $v \in \mathcal{H}^{\alpha}$, $v \neq 0$. Without loss of generality we can assume that the numbers in the first column of the Young tableaux α are $k - r + 1, \ldots, k$. Since $\epsilon_{\lambda}(v) = v$, we have

$$v = \frac{1}{\mu(\lambda)^2} \sum_{\varsigma \in S_r} (-1)^{\varsigma} \varsigma \left(\sum_{\tau \in Q'} (-1)^{\tau} \tau(b_{\alpha}(v)) \right) ,$$

where

 $Q' = \{ \tau \in Q : \tau \text{ is identical on the first column of } \alpha \}$

and S_r is the permutation group of $\{k - r + 1, \ldots, k\}$. This means that v is skew-symmetric with respect to the last r positions, so the contractions $i_{\nu}v$, with $\nu \in (\mathcal{H})^{\otimes (k-1)}$, can be written as contractions of a skew-symmetric r-tensor, thus they span a vector space of dimension $\geq r$ (see Theorem 4.1 (c)). If this dimension is exactly r, then v can be written as a combination of linearly independent tensor products $e_{s(1)} \otimes \cdots \otimes e_{s(k)}$ of vectors from $\{e_1, \ldots, e_r\}$. Since the tensor is symmetric with respect to permutations preserving rows and skew-symmetric with respect to permutations of the first column, each tensor product $e_{s(1)} \otimes \cdots \otimes e_{s(k)}$ in this combination should satisfy $e_{s(i)} = e_{s(j)}$ if i and j are in the same row of α . Hence, v is proportional to $\pi^{\alpha}(E_{\alpha})$.

Let $\mathcal{H}^{\alpha} \subset \mathcal{H}^{\otimes k}$ denotes the irreducible component of the tensor representation of the unitary group $U(\mathcal{H})$ in $\mathcal{H}^{\otimes k}$ associated with a Young diagram $\alpha \in Y(k)$.

Definition 6.1.

- (a) We say that a pure state ρ_v on $\mathcal{H}^{\otimes k}$ obeys a parastatistics $\alpha \in Y(k)$ (is a pure α -state for short) if $v \in \mathcal{H}^{\alpha}$, i.e. ρ is a pure state on the Hilbert space \mathcal{H}^{α} .
- (b) A pure state ρ on $\mathcal{H}^{\otimes k}$ obeying a parastatistics α is called a *simple pure state for the parastatistics* α (*simple pure* α -*state* for short) if ρ is represented by an α -simple tensor in \mathcal{H}^{α} . If ρ is not simple α -state, we call it an *entangled pure* α -*state*.
- (c) A mixed state ρ on \mathcal{H}^{α} we call a simple (mixed) state for the parastatistics α (simple α -state for short), if it can be written as a convex combination of simple pure α -states. In the other case, ρ is called an *entangled mixed* α -state.

7 Segre maps for generalized parastatistics

In general, for an arbitrary parastatistics (Young tableau) $\alpha \in Y(k)$ with the partition (Young diagram) $\lambda = (\lambda_1, \ldots, \lambda_r)$, we define the generalized Segre map Seg^{α} (α -Segre map) as a map Seg^{α} : $(\mathbb{P}\mathcal{H})^{\times r}_{\circ} \to \mathbb{P}(\mathcal{H}^{\alpha})$ described as follows.

Let us consider first the map

$$i_{\alpha}: \mathcal{H}^{\times r} \to \mathcal{H}^{\otimes k}, \quad (x_1, \dots, x_r) \mapsto x_{\alpha(1)} \otimes \dots \otimes x_{\alpha(k)},$$

where $\alpha(i)$ is the number of the row in which the box with the number *i* appears in the tableaux α . In other words, we make a tensor product of *k* vectors from $\{x_1, \ldots, x_r\}$ by putting x_j in the places indicated by the number of the boxes in the *j*th row. For instance, the Young tableaux from Example 6.1 give $i_{\alpha_1}(x_1, x_2) = x_1 \otimes x_1 \otimes x_2$ and $i_{\alpha_2}(x_1, x_2) = x_1 \otimes x_2 \otimes x_1$. It is clear that $i_{\alpha}(x_1, \ldots, x_r)$ is an eigenvector of b_{α} .

The Segre map $\operatorname{Seg}^{\alpha}$ associates with $(\rho_{x_1}, \ldots, \rho_{x_r}) \in (\mathbb{P}\mathcal{H})^{\times r}_{\circ}$ the pure state $\rho_{\pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})}$ in \mathcal{H}^{α} , as shows the following diagram:

Note that $\pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})$ is proportional to the antisymmetrization of the tensor $x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)}$ and that the construction is correct, since $\pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})$ is non-zero if and only if $x_1 \wedge \cdots \wedge x_r \neq 0$, and its projective class is uniquely determined by the projective classes of x_1, \ldots, x_r . Note also that we can always take x_1, \ldots, x_r orthogonal, say, $x_1 = e_1, \ldots, x_r = e_r$, since the antisymmetrization kills the part of x_i which is the orthogonal projection of x_i onto the linear subspace spanned by the rest of the vectors x_j . Now, according to Theorem 6.1, $\pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})$ is α -simple, so that $\rho_{\pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})}$ is a simple pure α -state. Moreover, each simple pure α -state is of this form and for symmetric and antisymmetric tensors this construction agrees with (37) and (38). We therefore get the following.

Theorem 7.1. A pure α -state $\rho \in \mathbb{P}(\mathcal{H}^{\alpha})$ is an entangled α -state if and only if it is not in the range of the Segre map

$$\operatorname{Seg}^{\alpha} : (\mathbb{P}\mathcal{H})^{\times r}_{\circ} \to \mathbb{P}(\mathcal{H}^{\alpha}).$$

The set $S(\mathcal{H}^{\alpha})$ of mixed non-entangled α -states is the convex hull of the range of the normalized α -Segre map,

$$\mathcal{S}(\mathcal{H}^{\alpha}) = conv \left(\operatorname{Seg}^{\alpha} \left(\mathcal{P}(\mathcal{H})_{\alpha}^{\times r} \right) \right) \,,$$

and mixed entangled α -states are exactly members of

$$\mathcal{D}(\mathcal{H}^{\alpha})\setminus \mathcal{S}(\mathcal{H}^{\alpha})$$
.

Let us observe that

$$\|\pi^{\alpha}(x_{\alpha(1)}\otimes\cdots\otimes x_{\alpha(k)})\|^{2}\cdot\rho_{\pi^{\alpha}(x_{\alpha(1)}\otimes\cdots\otimes x_{\alpha(k)})}=\pi^{\alpha}\circ\left(\rho_{x_{\alpha(1)}}\otimes\cdots\otimes \rho_{x_{\alpha(k)}}\right)\circ\pi^{\alpha},$$

as operators on $\mathcal{H}^{\otimes k}$. Indeed, since π^{α} is an orthogonal projection, the left-hand side equals

$$|\pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})\rangle \langle \pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)})|(y) = \pi^{\alpha}(x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)}) \langle x_{\alpha(1)} \otimes \cdots \otimes x_{\alpha(k)}|\pi_{\alpha}(y)\rangle = (\pi^{\alpha} \circ (\rho_{x_{\alpha(1)}} \otimes \cdots \otimes \rho_{x_{\alpha(k)}}) \circ \pi^{\alpha})(y).$$

This suggests to look at the map (the big α -Segre map):

$$\widetilde{\operatorname{Seg}}^{\alpha}: (\mathfrak{u}^*(\mathcal{H}))^{\times r} \to \mathfrak{u}^*(\mathcal{H}^{\alpha}) \subset \mathfrak{u}^*(\mathcal{H}^{\otimes k})\,,$$

where $\mathfrak{u}^*(\mathcal{H})$ denotes the (real) vector space of selfadjoint operators on \mathcal{H} , defined by

$$\widetilde{\operatorname{Seg}}^{\alpha}(u_1,\ldots,u_r) = \pi_{\alpha} \circ (u_{\alpha(1)} \otimes \cdots \otimes u_{\alpha(k)}) \circ \pi_{\alpha}$$

The big α -Segre map is a natural generalization of the map (7). A closer study of the big Segre maps we postpone to a separate paper.

8 Conclusions

The presented geometric description, in terms of Segre maps, of entanglement properties for systems with arbitrary statistics parallels our previous algebraic approach to such systems based on the concept of Srank of a tensor. It puts on equal and unifying footing systems of distinguishable particles, for which both algebraic and geometric descriptions were known, and systems with indistinguishable particles. What is more, this description provides an explicit form of simple (separable) pure states for an arbitrary parastatistics an effective procedures to check the simplicity in the bosonic and the fermionic case. Such procedures for arbitrary parastatistics are not known for us and can be the subject of forthcoming papers. Also the problem of the decomposition of the total algebra of operators, associated with the S-rank, is an interesting and open question.

Acknowledgments

Research of the first two authors was financed by the Polish Ministry of Science and Higher Education under the grant No. N N202 090239. G. Marmo would like to acknowledge the support provided by the Santander/UCIIIM chair of Excellence programme 2011-2012. We are also indebted to referees for their useful comments and suggestions.

References

- [1] H. S. Green. A generalized method of field quantization. Phys. Rev., 90:270?273, 1953.
- [2] Y. Ohnuki and S. Kamefuchi. Quantum Field Theory and Parastatistics. Springer, Berlin (1982).
- [3] J. Grabowski, M. Kuś, and G. Marmo. Entanglement for multipartite systems of indistinguishable particles. J. Phys. A: Math. Gen. 44:175302 (21pp), 2011.
- [4] J. Schliemann, D. Loss, and A. H. MacDonald. Double-occupancy errors, adiabaticity, and entanglement of spin qubits in quantum dots. *Phys. Rev. B*, 63(8):085311, 2001.
- [5] J. Schliemann, J. I. Cirac, M. Kuś, M. Lewenstein, and D. Loss. Quantum correlations in twofermion systems. *Phys. Rev. A*, 64:022303, 2001.
- [6] K. Eckert, J. Schliemann, D. Bruß, and M Lewenstein. Quantum correlations in systems of identical particles. Ann. Phys., 299:88–127, 2002.
- [7] R. Paškauskas and L. You. Quantum correlations in two-boson wave functions. *Phys. Rev. A*, 64(4):042310, 2001.
- [8] J. Grabowski, M. Kuś, and G. Marmo. Geometry of quantum systems: density states and entanglement. J. Phys. A: Math. Gen. 38:10217–10244, 2005.
- [9] J. Grabowski, M. Kuś, and G. Marmo. Symmetries, group actions, and entanglement. Open Sys. Information Dyn., 13:343–362, 2006.
- [10] Y. S. Li, B. Zeng, X. S. Liu, and G. L. Long. Entanglement in a two-identical-particle system. *Phys. Rev. A*, 64(5):054302, 2001.
- [11] G. Ghirardi, L. Marinatto, and T. Weber. Entanglement and Properties of Composite Quantum Systems: A Conceptual and Mathematical Analysis. *Journal of Statistical Physics*, 108(1):49–122, 2002.
- [12] G. Ghirardi and L. Marinatto. General criterion for the entanglement of two indistinguishable particles. *Phys. Rev. A*, 70(1):012109, 2004.
- [13] G. Ghirardi and L. Marinatto. Identical particles and entanglement. Optics and Spectroscopy, 99(3):386–390, 2005.

- [14] F. Buscemi, P. Bordone, and A. Bertoni. Linear entropy as an entanglement measure in two-fermion systems. *Phys. Rev. A* 75:032301, 2007.
- [15] A. R. Plastino, D. Manzano, and J. S. Dehesa. Separability Criteria and Entanglement Measures for Pure States of N Identical Fermions. *Europhys. Lett.* 86:20005, 2009.
- [16] B. Lari, P. Durganandini, and P. S. Joag. Multipartite entanglement in fermionic systems via a geometric measure. *Phys. Rev. A* 82:062302, 2010.
- [17] F. Buscemi and P. Bordone A measure of tripartite entanglement in bosonic and fermionic systems. *Phys. Rev. A* 84:022303, 2011.
- [18] P. Zanardi. Quantum entanglement in fermionic lattices. Phys. Rev. A 65:042101(R), 2002.
- [19] P. Zanardi and and X. Wang Fermionic entanglement in itinerant systems. J. Phys. A 35:7947–7959, 2002.
- [20] M. R. Dowling, A. C. Doherty, and H. M. Wiseman. Entanglement of indistinguishable particles in condensed matter physics *Phys. Rev. A* 73:052323, 2006.
- [21] T. Sasaki, T. Ichikawa, and I. Tsutsui. Entanglement of indistinguishable particles. Phys. Rev. A 83:012113, 2011.
- [22] M. C. Tichy, F. Mintert, and A. Buchleitner Essential entanglement for atomic and molecular physics. J. Phys. B: At. Mol. Opt. Phys. 44:192001, 2011.
- [23] G. Vidal. Entanglement monotones. J. Mod. Opt. 47:355-376, 2000.
- [24] R. Hartshorne. Algebraic geometry. Springer, 1977, Sect. IV.2.
- [25] W. Pauli. On the connection between spin and statistics Phys. Rev. 58:716–728, 1940.
- [26] E. P. Wigner. Do the equations of motion determine the quantum mechanical commutation relations? Phys. Rev. 77:711–712, 1950.
- [27] V. I. Man'ko, G. Marmo, E. C. G. Sudarshan, and F. Zaccaria. Wigner's Problem and Alternative Commutation Relations for Quantum Mechanics Int. J. Mod. Phys. B 11:1281-1296, 1997.
- [28] A. M. L. Messiah, O. W. Greenberg. Symmetrization postulate and its experimental foundation. *Phys. Rev.(2)* 136:B248-B267, 1964.
- [29] R. Y. Cusson. Examples of parastatistics Ann. Physics 55:22-40, 1969.
- [30] W. Fulton and J. Harris. Representation Theory. A First Course. Springer Verlag, 1991.
- [31] W. Fulton Young Tableaux. Cambridge University Press, 1997.
- [32] D. P. Zhelobenko. Compact Lie Groups and Their Representations. Amer. Math. Soc., 1973 (Translated from Russian).
- [33] A. Perelomov. Generalized coherent states and their applications. Springer, Heidelberg, 1986.
- [34] M. Kuś and I. Bengtsson. 'Classical' quantum states. Phys. Rev. A 80:022319, 2009.