

Relaxation in finite and isolated classical systems: an extension of Onsager's regression hypothesis

Marcus V. S. Bonança*

Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, 09210-170, Santo André, SP, Brazil

(Dated: May 21, 2019)

In order to derive the reciprocity relations, Onsager formulated a relation between thermal equilibrium fluctuations and relaxation widely known as regression hypothesis. It is shown in the present work how such relation can be extended to finite and isolated classical systems. This extension is derived from the fluctuation-dissipation theorem for the microcanonical ensemble. The results are exemplified with a nonintegrable system in order to motivate possible applications to dynamical systems and statistical mechanics of finite systems.

PACS numbers: 05.70.Ln, 05.40.-a, 05.45.-a

In his seminal work on the reciprocity relations[1], Onsager formulated an hypothesis about the decay of thermal equilibrium fluctuations that was essential in his derivation. From this hypothesis, Onsager was able to relate the relaxation of nonequilibrium macroscopic observables, obtained from phenomenological equations, to the decay rate of equilibrium fluctuations of those quantities. Such relation between those two apparently different phenomena is however quite expected from the point of view of the fluctuation-dissipation theorem derived [2] 20 years after Onsager's work. Indeed, since the derivation of the reciprocity relations from linear response theory, Onsager's regression hypothesis is understood as a different but equivalent statement of the fluctuation-dissipation theorem.

On the other hand, Onsager's reciprocity relations have already been extended to far from equilibrium conditions [3] in different contexts [4] where nonlinear effects are taken into account. Those extensions are based on fluctuation theorems which quite often are derived from the so-called "chaotic hypothesis"[5]. Hence the fluctuation theorems play the role of linear response theory beyond near equilibrium conditions and have been also applied to understand the nonequilibrium behavior of systems far from the thermodynamic limit [6].

Onsager's original work and its extensions mentioned above always focus on systems in contact with reservoirs. Here however we intend to study the near equilibrium behavior of isolated systems when they are finite and standard linear response theory cannot be applied. The relaxation to equilibrium of finite and almost isolated quantum systems has been studied experimentally using cold atoms [7] and has motivated several theoretical works which try to understand it and discuss the controversies related to the role played by the nonintegrability in this process [8–10]. For classical systems under such constraints, we show here that a simple extension of the fluctuation-dissipation theorem [11] shows very clearly the importance of the dynamics in the relaxation to equi-

librium within a description that is essentially Onsager's regression hypothesis extended to this new situation.

We start presenting the usual regression hypothesis expressed in mathematical terms. Let us consider a system described by the following Hamiltonian

$$H(\lambda_o + d\lambda) = H(\lambda_o) + d\lambda \left. \frac{\partial H}{\partial \lambda} \right|_{\lambda=\lambda_o}. \quad (1)$$

When the value of the parameter λ is suddenly switched from $\lambda_1 = \lambda_o + d\lambda$ to λ_o at $t = t_o$, the nonequilibrium average value \bar{B} of an observable B of the system evolves as [12]

$$\bar{B}(t - t_o) = \langle B \rangle_{\lambda_o} - \frac{d\lambda}{k_B T} C_{\lambda_1}(t - t_o), \quad (2)$$

for $t > t_o$, where $\langle \cdot \rangle_{\lambda_o}$ denotes the equilibrium average value when $\lambda = \lambda_o$, k_B is the Boltzmann constant, T is the temperature and

$$C_{\lambda_1}(t) = \langle \delta A(0) \delta B(t) \rangle_{\lambda_1} \quad (3)$$

is the correlation function with $\delta X(t) = X(t) - \langle X \rangle_{\lambda_1}$ and $A = (\partial H / \partial \lambda)|_{\lambda=\lambda_o}$.

Equation (2) is Onsager's regression hypothesis expressed mathematically. It states that relaxation of \bar{B} to the equilibrium value is possible as long as the correlation function $C_{\lambda_1}(t)$ decays to zero. On the other hand, if the relaxation process can be described by some sort of phenomenological equation, one obtains the decay of $C_{\lambda_1}(t)$ from (2). Since Eq. (2) is derived in the usual context of the canonical ensemble, the decay of the correlation function is interpreted as a consequence of the heat bath influence.

We will now derive (2) in a different context, namely, when the system is isolated and finite, i.e. there is no heat bath and the number of degrees of freedom is such that the system is not in the thermodynamic limit. We consider first the system in equilibrium under $H(\lambda_o + d\lambda)$. When t is equal to t_o , λ_1 is suddenly switched to λ_o . The system then relax to an new equilibrium state. Assuming that the system was not far from the final equilibrium state, a linear response calculation describes the

* marcus.bonanca@ufabc.edu.br

relaxation process. Although the situation requires the microcanonical ensemble, the fundamental equations of linear response theory do not rely on any particular ensemble [13]. For $t > t_o$ the Hamiltonian is $H(\lambda_o)$ and the situation can be stated as follows: a system, whose Hamiltonian is initially given by (1), is in equilibrium when at $t = t_o$ the generalized force $d\lambda$ is suddenly removed and $H(\lambda_o + d\lambda) \rightarrow H(\lambda_o)$. Therefore, from linear response theory, one obtains [13]

$$\bar{B}(t - t_o) - \langle B \rangle_{\lambda_1} = d\lambda \int_{-\infty}^t ds \phi_{BA}(E, t - s) \Theta(s - t_o), \quad (4)$$

where $\Theta(x)$ is the step function and $\langle B \rangle_{\lambda_1}$ is the following microcanonical average over the phase space points (\mathbf{q}, \mathbf{p})

$$\langle B \rangle_{\lambda_1} = \int d\mathbf{q} d\mathbf{p} \rho_{\lambda_1}(\mathbf{q}, \mathbf{p}) B(\mathbf{q}, \mathbf{p}), \quad (5)$$

with the distribution function ρ_{λ_1} given by

$$\rho_{\lambda_1}(\mathbf{q}, \mathbf{p}) = \frac{\delta[E - H(\mathbf{q}, \mathbf{p}, \lambda_1)]}{Z_{\lambda_1}(E)}, \quad (6)$$

where $Z_{\lambda_1}(E) = \int d\mathbf{q} d\mathbf{p} \delta[E - H(\mathbf{q}, \mathbf{p}, \lambda_1)]$.

The response function $\phi_{BA}(E, t - s)$ is [13]

$$\phi_{BA}(E, t - s) = \langle \{\delta A(s), \delta B(t)\} \rangle_{\lambda_1}, \quad (7)$$

where $\{\cdot, \cdot\}$ is the Poisson bracket, $\delta X(t) = X(\mathbf{q}(t), \mathbf{p}(t)) - \langle X \rangle_{\lambda_1}$, $(\mathbf{q}(t), \mathbf{p}(t))$ is the solution of Hamilton's equations of motion for N degrees of freedom, B is any observable and $A = (\partial H / \partial \lambda)|_{\lambda=\lambda_o}$.

The fluctuation-dissipation theorem [11] yields then

$$\tilde{F}_{BA}(z, \omega) = \frac{i}{z\omega} \tilde{\chi}_{BA}(z, \omega), \quad (8)$$

where

$$\tilde{\chi}_{BA}(z, \omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\tau \int_0^{\infty} dE e^{-i\omega\tau + Ez} [Z_{\lambda}(E) \phi_{BA}(E, \tau)], \quad (9)$$

$$\tilde{F}_{BA}(z, \omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\tau \int_0^{\infty} dE e^{-i\omega\tau + Ez} [Z_{\lambda}(E) C_{BA}(E, \tau)], \quad (10)$$

and $C_{BA}(E, \tau)$, with $\tau = t - s$, is the following correlation function

$$C_{BA}(E, t - s) = \langle \delta A(s) \delta B(t) \rangle_{\lambda}, \quad (11)$$

which differs from (3) because of the microcanonical average.

Thus, Eq.(8) leads to

$$\phi_{BA}(E, \tau) = -\frac{1}{Z_{\lambda}(E)} \frac{\partial^2}{\partial \tau \partial E} [Z_{\lambda}(E) C_{BA}(E, \tau)]. \quad (12)$$

The integral in (4) can be written as

$$\int_{-\infty}^t ds \phi_{BA}(t - s) \Theta(s - t_o) = \int_0^{t-t_o} d\tau \phi_{BA}(\tau), \quad (13)$$

and from (12) and (13) one obtains

$$\int_0^{t-t_o} d\tau \phi_{BA}(\tau) = -\frac{1}{Z_{\lambda}} \frac{\partial}{\partial E} [Z_{\lambda} C_{BA}(\tau)] \Big|_0^{t-t_o} \quad (14)$$

The dependence on E was omitted for convenience.

Therefore, the relaxation of B to the new equilibrium state is described by the following expression for $t > t_o$

$$\bar{B}(t - t_o) = \langle B \rangle_{\lambda_o} - d\lambda \frac{1}{Z_{\lambda_1}} \frac{\partial}{\partial E} [Z_{\lambda_1} C_{BA}(t - t_o)] \quad (15)$$

where

$$\langle B \rangle_{\lambda_o} = \langle B \rangle_{\lambda_1} + d\lambda \frac{1}{Z_{\lambda_1}} \frac{\partial}{\partial E} [Z_{\lambda_1} C_{BA}(0)], \quad (16)$$

since $\lim_{\tau \rightarrow \infty} C_{BA}(\tau) = 0$ is assumed.

Analogously to Eq.(2), the relaxation to equilibrium of B is ruled by a correlation function related to it and the relaxation rate is given in terms of the decay rate of equilibrium fluctuations. Thus, the physical contents of Eq.(15) allows us to interpret it as an extension of Onsager's regression hypothesis to the context of finite and isolated classical systems.

Despite of the analogy with (2), Eq.(15) shows that the relaxation rate of \bar{B} (as well as the decay rate of $C_{BA}(E, t)$) is given only by the statistical properties of the dynamics produced by $H(\lambda_o)$. Since the system is isolated, there is no influence of external thermal fluctuations on the decay of $C_{BA}(E, t)$ or \bar{B} as in (2). Therefore, one might ask for which kind of dynamics relaxation occurs. For integrable systems, whose motion is quasi-periodic, $C_{BA}(E, t)$ would decay only for $N \rightarrow \infty$. Nevertheless, it is well-known that $C_{BA}(E, t)$ decays for chaotic systems [14–16]. For nonintegrable systems, the complete spectrum of behaviors, from quasi-periodic to chaotic, could be approximately obtained.

In order to motivate possible applications of Eq.(15) to both low-dimensional dynamical systems and statistical mechanics of finite systems, we will consider relaxation in a system described by the following Hamiltonian [17]

$$H = \frac{p_1^2}{2} + \frac{p_2^2}{2} + \frac{q_1^2 q_2^2}{2} + \lambda \frac{(q_1^4 + q_2^4)}{4}, \quad (17)$$

which is integrable only for $\lambda = 1$. It can be verified through Poincaré surface of sections that the motion generated by (17) gets less and less regular as λ decreases. If on one hand the relaxation process is very well defined for low-dimensional hyperbolic systems, on




FIG. 1. Correlation function $C_{AA}(E', t)$ for $A = (q_1^4 + q_2^4)/4$ and $E' = 2.5$. The solid black lines were obtained numerically for 4×10^5 initial conditions. The dashed red lines are the fitting of $\mathcal{A}'e^{-\alpha't} \cos(\omega't)$. (a) $\lambda_1 = 0.1$, $\mathcal{A}' = 21.5$, $\alpha' = 0.20$ and $\omega' = 1.7$. (b) $\lambda_1 = 0.12$, $\mathcal{A}' = 14.5$, $\alpha' = 0.17$ and $\omega' = 1.8$.

the other hand most of the realistic models used in statistical mechanics of classical systems are nonintegrable. For a nonintegrable systems with few degrees of freedom as (17), relaxation may or may not happen and one has to find numerically the range of parameters where it occurs. In our application of (15) to this case, we have chosen $A = (\partial H / \partial \lambda)|_{\lambda=\lambda_o} = (q_1^4 + q_2^4)/4$ and $B = A$. There is a small range of values of λ ($0.9 \lesssim \lambda \lesssim 0.13$) where $C_{BA}(\tau)$ can be fitted by the expression $\mathcal{A}e^{-\alpha\tau} \cos(\omega\tau)$ (see Fig.(1)). It is not our aim here to find the exact functional form of the correlation function. Instead we want to find out whether a simple description of it (as the one just written above) is enough to describe the relaxation process approximately.

The dynamics given by (17) is scalable with energy, i.e. the Hamilton equations remain invariant under a transformation $(q'_{1,2}, p'_{1,2}, t') \rightarrow (q_{1,2}, p_{1,2}, t)$ given by the equations

$$\frac{q_{1,2}}{q'_{1,2}} = \left(\frac{E}{E'}\right)^{1/4}, \quad \frac{p_{1,2}}{p'_{1,2}} = \left(\frac{E}{E'}\right)^{1/4}, \quad \frac{t}{t'} = \left(\frac{E'}{E}\right)^{1/4}, \quad (18)$$




FIG. 2. Relaxation of \bar{B} for $B = (q_1^4 + q_2^4)/4$ and initial energy $E = 5.0$. The solid black lines were obtained numerically for 4×10^5 initial conditions. The dashed red lines are the predictions of Eq.(15) with C_{AA} obtained from Fig.(1) and the scaling property. (a) $\lambda_1 = 0.1$ switched to $\lambda_o = 0.12$ and the value of $\langle B \rangle_{\lambda_1} = 7.54$ obtained numerically for 4×10^5 initial conditions with $E = 5.0$ and $\lambda_1 = 0.1$. (b) $\lambda_1 = 0.12$ switched to $\lambda_o = 0.1$ and the value of $\langle B \rangle_{\lambda_1} = 6.55$ obtained numerically for 4×10^5 initial conditions with $E = 5.0$ and $\lambda_1 = 0.12$.

From this property of (17) yields $\mathcal{A}/\mathcal{A}' = (E/E')^2$ since $(q_1^4 + q_2^4)^2 \propto (E/E')^2$, $\alpha/\alpha' = (E/E')^{1/4}$ and $\omega/\omega' = (E/E')^{1/4}$. One can also verify that $Z_\lambda(E) \propto E^{1/2}$. Therefore, the scaling with energy allows us to perform the derivative in (15) taking a certain value E' as reference and obtaining $C_{BA}(E, t)$ from the parameters of $C_{BA}(E', t)$, keeping of course the same functional form and the same value of λ .

All the numerical results were obtained from the integration of the equations of motion of (17) using a fourth-order symplectic integrator [18]. In Fig.(1), it is shown the numerical results for $C_{BA}(E', t)$. Although it is clear that the fitting is not excellent, one obtains afterwards a good agreement between numerical and analytical results for the relaxation of \bar{B} . The analytical results should indeed be called semi-analytical since both $\langle B \rangle_{\lambda_1}$ and C_{BA} were obtained numerically. In Fig.(2), $\bar{B}(t-t_o)$ is the result of

the average over several $B(t-t_o)$ obtained from the time evolution of initial conditions distributed over an energy surface with $E = 5.0$. In Fig.(2a) a sudden switching of $\lambda_1 = 0.1$ to $\lambda_o = 0.12$ at t_o leads to the relaxation of \bar{B} to a new equilibrium value. Although amplitude and frequency of oscillations are not correctly described, the relaxation time and the value of $\langle B \rangle_{\lambda_o}$ are well predicted. In Fig.(2b), the comparison between numerical and analytical results for \bar{B} is shown for $\lambda_1 = 0.12$ switched to $\lambda_o = 0.1$. As before, amplitude and frequency of oscillations are roughly described and the value predicted for $\langle B \rangle_{\lambda_o}$ is not as good as in Fig.(2a). The relaxation time however is still in good agreement with the numerical results.

In conclusion, we have derived an extension of Onsager's regression hypothesis from linear response theory

when the system of interest is isolated and finite. Although thermal fluctuations induced by a heat bath are absent in this context, the expression obtained relates, as usual, relaxation to equilibrium fluctuations. Hence the new feature is that the decay of correlations is given only by the intrinsic dynamics of the system. The relaxation of a nonintegrable systems with two degrees of freedom has illustrated the result. The outlook is to extend the present result to the quantum regime.

ACKNOWLEDGMENTS

The author acknowledges support of UFABC. The author is also grateful to M. de Koning and R. Venegeroles for valuable discussions and suggestions.

-
- [1] L. Onsager, Phys. Rev. **37**, 405 (1931); **38**, 2265 (1931).
 - [2] H. B. Callen and T. A. Welton, Phys. Rev. **83**, 34 (1951).
 - [3] G. Gallavotti, Phys. Rev. Lett. **77**, 4334 (1996).
 - [4] D. Andrieux and P. Gaspard, J. Stat. Mech. **2007**, P02006 (2007).
 - [5] G. Gallavotti and E. G. D. Cohen, Phys. Rev. Lett. **74**, 2594 (1995); J. Stat. Phys. **80**, 931 (1995).
 - [6] D. Collin, F. Ritort, C. Jarzynski, S. B. Smith, J. I. Tinoco, and C. Bustamante, Nature **437**, 231 (2005).
 - [7] T. Kinoshita, T. Wenger, and D. S. Weiss, Nature **440**, 900 (2006).
 - [8] M. Rigol, V. Dunjko, and M. Olshanii, Nature **452**, 854 (2008).
 - [9] A. V. Ponomarev, S. Denisov, and P. Hänggi, Phys. Rev. Lett. **106**, 180601 (2011).
 - [10] C. Gogolin, M. P. Müller, and J. Eisert, Phys. Rev. Lett. **106**, 040401 (2011).
 - [11] M. V. S. Bonança, Phys. Rev. E **78**, 031107 (2008).
 - [12] M. E. Tuckerman, *Statistical Mechanics: Theory and Molecular Simulation* (Oxford University Press, New York, 2010).
 - [13] R. Kubo, M. Toda, and N. Hashitsume, *Statistical Physics II* (Springer-Verlag, Berlin, 1985).
 - [14] D. Ruelle, Phys. Rev. Lett. **56**, 405 (1986); J. Stat. Phys. **44**, 281 (1986).
 - [15] P. L. Garrido and G. Gallavotti, J. Stat. Phys. **76**, 549 (1994).
 - [16] R. Artuso, G. Casati, and I. Guarneri, J. Stat. Phys. **83**, 145 (1996).
 - [17] A. Carnegie and I. C. Percival, J. Phys. A: Math. Gen. **17**, 801 (1984).
 - [18] E. Forest and R. D. Ruth, Physica D **43**, 105 (1990).