

# The Terwilliger algebra of Odd graphs

Qian Kong Benjian Lv Kaishun Wang\*

*Sch. Math. Sci. & Lab. Math. Com. Sys., Beijing Normal University, Beijing, 100875, China*

## Abstract

In [The Terwilliger algebra of the Johnson schemes, Discrete Mathematics 307 (2007) 1621–1635], Levstein and Maldonado computed the Terwilliger algebra of the Johnson scheme  $J(n, m)$  when  $3m \leq n$ . The distance- $m$  graph of  $J(2m + 1, m)$  is the Odd graph  $O_{m+1}$ . In this paper, we determine the Terwilliger algebra of  $O_{m+1}$  and give its basis.

*AMS classification:* 05E30

*Key words:* Terwilliger algebra; Odd graph

## 1 Introduction

Suppose  $\Gamma = (X, R)$  denotes a simple connected graph with diameter  $D$ . For each  $i \in \{0, 1, \dots, D\}$ , let  $\Gamma_i(x) = \{y \in X \mid \partial(x, y) = i\}$ , where  $\partial(x, y)$  is the distance between  $x$  and  $y$ . Define  $E_i^* = E_i^*(x)$  to be the diagonal matrix in  $\text{Mat}_X(\mathbb{C})$  with  $yy$ -entry

$$(E_i^*)_{yy} = \begin{cases} 1, & \text{if } y \in \Gamma_i(x), \\ 0, & \text{otherwise.} \end{cases}$$

The *Terwilliger algebra*  $\mathcal{T}(x)$  of  $\Gamma$  with respect to a given vertex  $x$  is the subalgebra of  $\text{Mat}_X(\mathbb{C})$  generated by the adjacency matrix  $A$  of  $\Gamma$  and  $E_0^*, E_1^*, \dots, E_D^*$ .

Terwilliger [10] initiated the study of the Terwilliger algebra of association schemes, which has been used to study (almost) bipartite  $P$ - and  $Q$ -polynomial association schemes [2, 3], 2-homogeneous bipartite distance-regular graphs [4], Hypercubes [5], Hamming graphs [7], Johnson graphs [8], incidence graphs of Johnson geometry [6] and so on.

Let  $\Omega$  be a set of cardinality  $2m + 1$  and let  $\binom{\Omega}{i}$  denote the set of all  $i$ -subsets of  $\Omega$ . The Odd graph  $O_{m+1}$  is the graph whose vertex set is the set  $X = \binom{\Omega}{m}$ , where two vertices are adjacent if they are disjoint. Levstein and Maldonado [8] determined the Terwilliger algebra of the Johnson graph  $J(n, m)$  when  $3m \leq n$ . Observe  $O_{m+1}$  is the distance- $m$  graph of the Johnson graph  $J(2m + 1, m)$ , and they have the same Terwilliger algebra. In this paper we shall determine the Terwilliger algebra of  $O_{m+1}$  (Theorem 3.5), give one of its bases (Proposition 3.6) and compute its dimension (Corollary 3.7).

---

\*Corresponding author.

E-mail addresses: kongqian@mail.bnu.edu.cn (Qian Kong); benjian@mail.bnu.edu.cn (Benjian Lv); wangks@bnu.edu.cn (Kaishun Wang)

## 2 Intersection matrix

In this section we first introduce the intersection matrix, then discover the relationship between the adjacency matrix of the Odd graph  $O_{m+1}$  and the intersection matrices.

Since  $O_{m+1}$  is distance-transitive with diameter  $m$  (cf. [1]), the isomorphism class of  $\mathcal{T}(x)$  is independent of the choice of  $x$ , denoted by  $\mathcal{T} := \mathcal{T}(x)$ .

Let  $V$  be a set of cardinality  $v$ . Let  $H_{i,j}^l(v)$  be a binary matrix with rows indexed by  $\binom{V}{i}$  and columns indexed by  $\binom{V}{j}$ , whose  $yz$ -entry is defined by

$$(H_{i,j}^l(v))_{yz} = \begin{cases} 1, & \text{if } |y \cap z| = l, \\ 0, & \text{otherwise.} \end{cases}$$

This matrix is a class of *intersection matrices*. Observe that  $H_{i,j}^l(v) \neq 0$  if and only if  $\max(0, i+j-v) \leq l \leq \min(i, j)$ . We adopt the convention that  $H_{i,j}^l(v) = 0$  for any integer  $l$  such that  $l < 0$  or  $l > \min(i, j)$ . From [9, Proposition 4], we have

$$H_{i,j}^l(v)H_{j,k}^s(v) = \sum_{g=0}^{\min(i,k)} \sum_{h=0}^g \binom{g}{h} \binom{i-g}{l-h} \binom{k-g}{s-h} \binom{v+g-i-k}{j+h-l-s} H_{i,k}^g(v). \quad (1)$$

In particular,

$$H_{i,j}^l(v)H_{j,k}^0(v) = \sum_{s=\max(0, i+j+k-l-v)}^{\min(i-l, k)} \binom{i-s}{l} \binom{v+s-i-k}{j-l} H_{i,k}^s(v). \quad (2)$$

**Lemma 2.1** *Let  $\Gamma$  be the Odd graph  $O_{m+1}$  with the adjacency matrix  $A$ , and let  $A_{i,j}$  be the submatrix of  $A$  with rows indexed by  $\Gamma_i(x)$  and columns indexed by  $\Gamma_j(x)$ . Then*

$$A_{i,j} = 0 \quad (0 \leq i \leq j \leq m, \quad i \neq j-1 \text{ or } i = j \neq m), \quad (3)$$

$$A_{2i, 2i+1} = H_{m-i, i}^0(m) \otimes H_{i, m-i}^0(m+1) \quad (0 \leq i \leq \lceil \frac{m}{2} \rceil - 1), \quad (4)$$

$$A_{2i+1, 2i+2} = H_{i, m-i-1}^0(m) \otimes H_{m-i, i+1}^0(m+1) \quad (0 \leq i \leq \lfloor \frac{m}{2} \rfloor - 1), \quad (5)$$

$$A_{m, m} = H_{\lfloor \frac{m}{2} \rfloor, \lfloor \frac{m}{2} \rfloor}^0(m) \otimes H_{\lceil \frac{m}{2} \rceil, \lceil \frac{m}{2} \rceil}^0(m+1), \quad (6)$$

where “ $\otimes$ ” denotes the Kronecker product of matrices.

**Proof.** Since  $O_{m+1}$  is almost bipartite, (3) is directed.

Pick  $y \in \Gamma_{2i}(x)$ ,  $z \in \Gamma_{2i+1}(x)$ . Note that  $\partial(x, y) = 2i$  if and only if  $|x \cap y| = m - i$ ;  $\partial(x, z) = 2i + 1$  if and only if  $|x \cap z| = i$ . Then  $|x \cap y| = m - i$  and  $|x \cap z| = i$ . Suppose  $y = \alpha_{m-i}\beta_i := \alpha_{m-i} \cup \beta_i$ ,  $z = \alpha'_i\beta'_{m-i}$ , where  $\alpha_{m-i} \in \binom{x}{m-i}$  and  $\beta_i \in \binom{\Omega \setminus x}{i}$ , while  $\alpha'_i \in \binom{x}{i}$  and  $\beta'_{m-i} \in \binom{\Omega \setminus x}{m-i}$ . Then

$$(A_{2i, 2i+1})_{yz} = (H_{m-i, i}^0(m) \otimes H_{i, m-i}^0(m+1))_{yz} = \begin{cases} 1, & \text{if } \alpha_{m-i} \cap \alpha'_i = \emptyset \text{ and } \beta_i \cap \beta'_{m-i} = \emptyset, \\ 0, & \text{otherwise,} \end{cases}$$

which leads to (4).

Similarly, (5) (6) hold.  $\square$

### 3 The Terwilliger algebra

In this section we fix  $x \in \binom{\Omega}{m}$ , then consider the Terwilliger algebra  $\mathcal{T} = \mathcal{T}(x)$  of  $O_{m+1}$ .

For  $0 \leq i, j \leq m$ , any matrix  $M$  indexed by elements in  $\Gamma_i(x) \times \Gamma_j(x)$  can be embedded into  $\text{Mat}_X(\mathbb{C})$  by

$$L(M)_{\Gamma_p(x) \times \Gamma_q(x)} = \begin{cases} M, & \text{if } p = i \text{ and } q = j, \\ 0, & \text{otherwise.} \end{cases}$$

Write  $G_{i,j}(v) = \{g \mid \max(0, i + j - v) \leq g \leq \min(i, j)\}$ . Let

$$\mathcal{M} = \bigoplus_{p,q=0}^m L(\mathcal{M}_{p,q}), \quad (7)$$

where  $L(\mathcal{M}_{p,q}) = \{L(M) \mid M \in \mathcal{M}_{p,q}\}$ , and

$$\mathcal{M}_{2i,2j} = \text{Span}\{H_{m-i,m-j}^l(m) \otimes H_{i,j}^s(m+1), l \in G_{m-i,m-j}(m), s \in G_{i,j}(m+1)\}, \quad (8)$$

$$\mathcal{M}_{2i,2j+1} = \text{Span}\{H_{m-i,j}^l(m) \otimes H_{i,m-j}^s(m+1), l \in G_{m-i,j}(m), s \in G_{i,m-j}(m+1)\}, \quad (9)$$

$$\mathcal{M}_{2i+1,2j} = \text{Span}\{H_{i,m-j}^l(m) \otimes H_{m-i,j}^s(m+1), l \in G_{i,m-j}(m), s \in G_{m-i,j}(m+1)\}, \quad (10)$$

$$\mathcal{M}_{2i+1,2j+1} = \text{Span}\{H_{i,j}^l(m) \otimes H_{m-i,m-j}^s(m+1), l \in G_{i,j}(m), s \in G_{m-i,m-j}(m+1)\}. \quad (11)$$

Note that  $\mathcal{M}$  is a vector space. By (1) we have  $\mathcal{M}$  is an algebra. Next we shall prove  $\mathcal{T} = \mathcal{M}$ .

**Lemma 3.1** *The Terwilliger algebra  $\mathcal{T}$  is a subalgebra of  $\mathcal{M}$ .*

**Proof.** By Lemma 2.1 we have  $A \in \mathcal{M}$ . Since

$$E_{2i}^* = L(H_{m-i,m-i}^{m-i}(m) \otimes H_{i,i}^i(m+1)) \in \mathcal{M}, \quad 0 \leq i \leq \lfloor \frac{m}{2} \rfloor,$$

$$E_{2i+1}^* = L(H_{i,i}^i(m) \otimes H_{m-i,m-i}^{m-i}(m+1)) \in \mathcal{M}, \quad 0 \leq i \leq \lceil \frac{m}{2} \rceil - 1,$$

we have  $\mathcal{T} \subseteq \mathcal{M}$ . □

For  $0 \leq i, j \leq m$ , let  $\mathcal{T}_{i,j} = \{M_{i,j} \mid M \in \mathcal{T}\}$ , where  $M_{i,j}$  is the submatrix of  $M$  with rows indexed by  $\Gamma_i(x)$  and columns indexed by  $\Gamma_j(x)$ . Since  $\mathcal{T}$  is an algebra, each  $\mathcal{T}_{i,j}$  is a linear space. From  $\mathcal{T}E_j^*\mathcal{T} \subseteq \mathcal{T}$  we obtain  $(\mathcal{T}E_j^*\mathcal{T})_{i,k} \subseteq \mathcal{T}_{i,k}$ , which implies that

$$\mathcal{T}_{i,j}\mathcal{T}_{j,k} \subseteq \mathcal{T}_{i,k}. \quad (12)$$

From  $A, E_i^* \in \mathcal{T}$ , we have  $AE_{i_2}^*AE_{i_3}^* \cdots AE_{i_{n-1}}^*A \in \mathcal{T}$ , which follows that

$$A_{i_1,i_2}A_{i_2,i_3} \cdots A_{i_{n-2},i_{n-1}}A_{i_{n-1},i_n} \in \mathcal{T}_{i_1,i_n}. \quad (13)$$

**Lemma 3.2** *For  $i \leq j \leq \lceil \frac{m}{2} \rceil - 1$  and  $0 \leq l \leq i$ , we have*

$$H_{i,j}^l(m) \otimes H_{m-i,m-j}^{m-j}(m+1) \in \mathcal{T}_{2i+1,2j+1}.$$

**Proof.** We use induction on  $l$ .

By (13), for  $i < j$  we have  $A_{2i+1,2i+2}A_{2i+2,2i+3} \cdots A_{2j,2j+1} \in \mathcal{T}_{2i+1,2j+1}$ , which yields that

$$H_{i,j}^i(m) \otimes H_{m-i,m-j}^{m-j}(m+1) \in \mathcal{T}_{2i+1,2j+1}. \quad (14)$$

When  $i = j$  we pick  $I_{(i)}^{(m)} \otimes I_{(m-i)}^{(m+1)} \in \mathcal{T}_{2i+1,2j+1}$ , which also satisfies (14).

Assume that  $H_{i,j}^g(m) \otimes H_{m-i,m-j}^{m-j}(m+1) \in \mathcal{T}_{2i+1,2j+1}$  for  $g \geq l$ . By (12) and (13), for  $2j+1 < m$  we obtain

$$(H_{i,j}^l(m) \otimes H_{m-i,m-j}^{m-j}(m+1))A_{2j+1,2j+2}A_{2j+2,2j+1} \in \mathcal{T}_{2i+1,2j+1}\mathcal{T}_{2j+1,2j+1} \subseteq \mathcal{T}_{2i+1,2j+1},$$

and for  $2j+1 = m$  we have

$$(H_{i,j}^l(m) \otimes H_{m-i,m-j}^{m-j}(m+1))A_{2j+1,2j+1}^2 \in \mathcal{T}_{2i+1,2j+1}.$$

Then we get

$$(a_1H_{i,j}^{l-1}(m) + a_2H_{i,j}^l(m) + a_3H_{i,j}^{l+1}(m)) \otimes H_{m-i,m-j}^{m-j}(m+1) \in \mathcal{T}_{2i+1,2j+1},$$

where  $a_1, a_2$  and  $a_3$  are some positive integers. It follows that  $H_{i,j}^{l-1}(m) \otimes H_{m-i,m-j}^{m-j}(m+1) \in \mathcal{T}_{2i+1,2j+1}$ . Hence the conclusion is obtained by induction.  $\square$

**Lemma 3.3** For  $i+1 \leq j \leq \lfloor \frac{m}{2} \rfloor$  and  $0 \leq l \leq i$ , we have

$$H_{i,m-j}^l(m) \otimes H_{m-i,j}^{j-i-1}(m+1) \in \mathcal{T}_{2i+1,2j}.$$

**Proof.** By (13) we have  $A_{2i+1,2i+2}A_{2i+2,2i+3} \cdots A_{2j-1,2j} \in \mathcal{T}_{2i+1,2j}$  for  $i+1 \leq j$ , i.e.,

$$H_{i,m-j}^0(m) \otimes H_{m-i,j}^{j-i-1}(m+1) \in \mathcal{T}_{2i+1,2j}.$$

Assume that  $H_{i,m-j}^g(m) \otimes H_{m-i,j}^{j-i-1}(m+1) \in \mathcal{T}_{2i+1,2j}$  for  $g \leq l$ . Then by (12) and (13), we obtain

$$(H_{i,m-j}^l(m) \otimes H_{m-i,j}^{j-i-1}(m+1))A_{2j,2j-1}A_{2j-1,2j} \in \mathcal{T}_{2i+1,2j},$$

which gives

$$(b_1H_{i,m-j}^{l-1}(m) + b_2H_{i,m-j}^l(m) + b_3H_{i,m-j}^{l+1}(m)) \otimes H_{m-i,j}^{j-i-1}(m+1) \in \mathcal{T}_{2i+1,2j},$$

where  $b_1, b_2$  and  $b_3$  are some positive integers. Thus  $H_{i,m-j}^{l-1}(m) \otimes H_{m-i,j}^{j-i-1}(m+1) \in \mathcal{T}_{2i+1,2j}$  and the conclusion is valid by induction.  $\square$

**Lemma 3.4** The algebra  $\mathcal{M}$  is a subalgebra of  $\mathcal{T}$ .

**Proof.** In order to prove this result, we only need to show that  $\mathcal{M}_{p,q} \subseteq \mathcal{T}_{p,q}$  for  $0 \leq p, q \leq m$ . Write  $\mathcal{M}_{p,q}^t = \{M^t \mid M \in \mathcal{M}_{p,q}\}$  and  $\mathcal{T}_{p,q}^t = \{M^t \mid M \in \mathcal{T}_{p,q}\}$ . Since  $\mathcal{M}_{q,p} = \mathcal{M}_{p,q}^t$  and  $\mathcal{T}_{q,p} = \mathcal{T}_{p,q}^t$ , it suffices to prove  $\mathcal{M}_{p,q} \subseteq \mathcal{T}_{p,q}$  for  $p \leq q$ . We use induction on  $p$ .

**Step 1.** Show  $\mathcal{M}_{0,q} \subseteq \mathcal{T}_{0,q}$  ( $0 \leq q \leq m$ ).

According to (8) (9), we get

$$\mathcal{M}_{0,2j} = \text{Span}\{H_{m,m-j}^{m-j}(m) \otimes H_{0,j}^0(m+1)\} \quad (0 \leq j \leq \lfloor \frac{m}{2} \rfloor),$$

and

$$\mathcal{M}_{0,2j+1} = \text{Span}\{H_{m,j}^j(m) \otimes H_{0,m-j}^0(m+1)\} \quad (0 \leq j \leq \lceil \frac{m}{2} \rceil - 1).$$

By Lemma 2.1 and (2), we have

$$A_{0,1}A_{1,2} \cdots A_{2j-1,2j} = c_1 H_{m,m-j}^{m-j}(m) \otimes H_{0,j}^0(m+1)$$

and

$$A_{0,1}A_{1,2} \cdots A_{2j,2j+1} = c_2 H_{m,j}^j(m) \otimes H_{0,m-j}^0(m+1),$$

where  $c_1, c_2$  are some positive integers. Then by (13) we have  $\mathcal{M}_{0,2j} \subseteq \mathcal{T}_{0,2j}$  and  $\mathcal{M}_{0,2j+1} \subseteq \mathcal{T}_{0,2j+1}$ .

**Step 2.** Assume that  $\mathcal{M}_{p,q} \subseteq \mathcal{T}_{p,q}$  for  $p \leq 2i$ . We will show that  $\mathcal{M}_{2i+1,q} \subseteq \mathcal{T}_{2i+1,q}$  and  $\mathcal{M}_{2i+2,q} \subseteq \mathcal{T}_{2i+2,q}$ .

**Step 2.1.** Show  $\mathcal{M}_{2i+1,q} \subseteq \mathcal{T}_{2i+1,q}$  ( $2i+1 \leq q \leq m$ ).

**Case 1.**  $q = 2j+1$  ( $i \leq j \leq \lceil \frac{m}{2} \rceil - 1$ ).

By inductive hypothesis we have

$$H_{m-i,j}^l(m) \otimes H_{i,m-j}^s(m+1) \in \mathcal{M}_{2i,2j+1} \subseteq \mathcal{T}_{2i,2j+1}, \quad l \in G_{m-i,j}(m), \quad s \in G_{i,m-j}(m+1).$$

Since  $A_{2i+1,2i} = A_{2i,2i+1}^t = H_{i,m-i}^0(m) \otimes H_{m-i,i}^0(m+1) \in \mathcal{T}_{2i+1,2i}$ , by (12) we have

$$(H_{i,m-i}^0(m) \otimes H_{m-i,i}^0(m+1))(H_{m-i,j}^l(m) \otimes H_{i,m-j}^s(m+1)) \in \mathcal{T}_{2i+1,2i} \mathcal{T}_{2i,2j+1} \subseteq \mathcal{T}_{2i+1,2j+1}.$$

From (2) we obtain

$$H_{i,j}^{j-l}(m) \otimes ((s+1)H_{m-i,m-j}^{m-j-s-1}(m+1) + (i-s+1)H_{m-i,m-j}^{m-j-s}(m+1)) \in \mathcal{T}_{2i+1,2j+1}. \quad (15)$$

Since  $l \in G_{m-i,j}(m)$  and  $s \in G_{i,m-j}(m+1)$ , from (15) and Lemma 3.2 we get

$$H_{i,j}^{l'}(m) \otimes H_{m-i,m-j}^{s'}(m+1) \in \mathcal{T}_{2i+1,2j+1}, \quad l' \in G_{i,j}(m), \quad s' \in G_{m-i,m-j}(m+1),$$

which implies  $\mathcal{M}_{2i+1,2j+1} \subseteq \mathcal{T}_{2i+1,2j+1}$ .

**Case 2.**  $q = 2j$  ( $i+1 \leq j \leq \lfloor \frac{m}{2} \rfloor$ ).

By inductive hypothesis, we have

$$H_{m-i,m-j}^l(m) \otimes H_{i,j}^s(m+1) \in \mathcal{M}_{2i,2j} \subseteq \mathcal{T}_{2i,2j}, \quad l \in G_{m-i,m-j}(m), \quad s \in G_{i,j}(m+1).$$

Thus by (13) we obtain

$$A_{2i+1,2i}(H_{m-i,m-j}^l(m) \otimes H_{i,j}^s(m+1)) \in \mathcal{T}_{2i+1,2i} \mathcal{T}_{2i,2j} \subseteq \mathcal{T}_{2i+1,2j}.$$

From (2), we have

$$H_{i,m-j}^{m-j-l}(m) \otimes ((s+1)H_{m-i,j}^{j-s-1}(m+1) + (i-s+1)H_{m-i,j}^{j-s}(m+1)) \in \mathcal{T}_{2i+1,2j}. \quad (16)$$

Since  $l \in G_{m-i,m-j}(m)$  and  $s \in G_{i,j}(m+1)$ , by (16) and Lemma 3.3 we have

$$H_{i,m-j}^{l''}(m) \otimes H_{m-i,j}^{s''}(m+1) \in \mathcal{T}_{2i+1,2j}, \quad l'' \in G_{i,m-j}(m), \quad s'' \in G_{m-i,j}(m+1),$$

which yields  $\mathcal{M}_{2i+1,2j} \subseteq \mathcal{T}_{2i+1,2j}$ .

**Step 2.2.** Show  $\mathcal{M}_{2i+2,q} \subseteq \mathcal{T}_{2i+2,q}$  ( $2i+2 \leq q \leq m$ ).

The proof of this step is similar to that of Step 2.1 and we omit it here.

Hence the desired result follows by induction.  $\square$

**Theorem 3.5** *Let  $\mathcal{T}$  be the Terwilliger algebra of the Odd graph  $O_{m+1}$  and  $\mathcal{M}$  be the algebra defined in (7). Then  $\mathcal{T} = \mathcal{M}$ .*

**Proof.** Combining Lemma 3.1 and Lemma 3.4, the desired result follows.  $\square$

Since the generating matrices of each vector space in (8)-(11) are linearly independent, we have the following result.

**Proposition 3.6** *The Terwilliger algebra  $\mathcal{T}$  of the Odd graph  $O_{m+1}$  has a basis:*

$$\{L(H_{m-i,m-j}^l(m) \otimes H_{i,j}^s(m+1)), l \in G_{m-i,m-j}(m), s \in G_{i,j}(m+1)\}_{i,j=0}^m.$$

**Corollary 3.7** *The dimension of  $\mathcal{T}$  is  $\binom{m+4}{4}$ .*

**Proof.** By Proposition 3.6 we get

$$\begin{aligned} & \dim \mathcal{T} \\ &= \sum_{i,j=0}^m |G_{m-i,m-j}(m)| |G_{i,j}(m+1)| \\ &= \sum_{i,j=0}^m (\min(m-i, m-j) - \max(0, m-i-j) + 1)(\min(i, j) - \max(0, i+j-m-1) + 1). \end{aligned}$$

By zigzag calculation, the desired result follows.  $\square$

## Acknowledgement

This research is partially supported by NSF of China, NCET-08-0052, and the Fundamental Research Funds for the Central Universities of China.

## References

- [1] A.E. Brouwer, A.M. Cohen and A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, Heidelberg, 1989.
- [2] J.S. Caughman IV, The Terwilliger algebra for bipartite  $P$ - and  $Q$ -polynomial association schemes, Discrete Math. 196 (1999) 65–95.
- [3] J.S. Caughman, M.S. MacLean, P. Terwilliger, The Terwilliger algebra of an almost-bipartite  $P$ - and  $Q$ -polynomial association scheme, Discrete Math. 292 (2005) 17–44.
- [4] B. Curtin, The Terwilliger algebra of a 2-homogeneous bipartite distance-regular graph, J. Combin. Theory Ser. A 81 (2001) 125–141.
- [5] J. Go, The Terwilliger algebra of the hypercube, Europ. J. Combin. 23 (2002) 399–429.
- [6] Q. Kong, B. Lv and K. Wang, The Terwilliger algebra of the incidence graph of Johnson geometry, arXiv: 1111.1369 [math.CO].
- [7] F. Levstein, C. Maldonado, D. Penazzi, The Terwilliger algebra of a Hamming scheme  $H(d, q)$ , Europ. J. Combin. 27 (2006) 1–10.

- [8] F. Levstein, C. Maldonado, The Terwilliger algebra of the Johnson schemes, *Discrete Mathematics* 307 (2007) 1621–1635.
- [9] M. Mohammad-Noori, N. Ghareghani, E. Ghorbani, Intersection matrices and the Johnson scheme, arXiv: 0902.4367v3 [math.CO].
- [10] P. Terwilliger, The subconstituent algebra of an association scheme I, *J. Algebr. Comb.* 1 (1992) 363–388; The subconstituent algebra of an association scheme II, *J. Algebr. Comb.* 2 (1993) 73–103; The subconstituent algebra of an association scheme III, *J. Algebr. Comb.* 2 (1993) 177–210.