

PROOF OF THE PRUNING FRONT CONJECTURE FOR CERTAIN HÉNON PARAMETERS

VALENTÍN MENDOZA

ABSTRACT. The *Pruning Front Conjecture* is proved for an open set of Hénon parameters far from unimodal. More specifically, for an open subset of Hénon parameter space, consisting of two connected components one of which intersects the area-preserving locus, it is shown that the associated Hénon maps are prunings of the horseshoe. In particular, their dynamics is a subshift of the two-sided two-shift.

1. INTRODUCTION

Pruning was introduced by Cvitanović [5] to describe the dynamics of the Hénon (or Lozi) family. The main idea of pruning is to describe Hénon maps as *partially formed* horseshoes or, put another way, as horseshoes from which part of the dynamics was destroyed or *pruned away*. Roughly, the *Pruning Front Conjecture (PFC)* states just that: every map in the Hénon family is a pruning of the horseshoe, which is proved here for an open set of parameter values far from the degenerate unimodal world. More precisely, it is shown that there are two open topological disks in parameter space, one of which intersects the area-preserving locus (and contains parameters previously studied by Davis, MacKay and Sannami [7]), whose associated Hénon maps are non-trivial prunings of the horseshoe.

Pruning can be described as a way to give a topological description of the dynamics of surface homeomorphisms and the mechanism of creation or destruction of chaos in 2-dimensions. A means to formalize this idea, due to de Carvalho¹, is to introduce *pruning isotopies* which destroy pieces of the dynamics of a surface homeomorphism f_0 in a controlled way [8]. Here *to destroy dynamics* means to convert non-wandering points into wandering points, and *in a controlled way* means that this destruction occurs in regions which we can define explicitly in terms of f_0 . A precise formulation of this is as follows:

Definition 1.1. Let D be a domain of the plane, $f_0: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ a homeomorphism of the plane. Let $\mathbb{P} := \cup_{i \in \mathbb{Z}} f_0^i(D)$ be the saturation of D under f_0 . An isotopy $f_t: f_0 \simeq f_1$ is a *pruning isotopy* if:

- (1) $\text{Supp}(f_t) \subset \mathbb{P}$, and
- (2) $\text{NW}(f_1) = \text{NW}(f_0) \setminus \mathbb{P}$,

where $\text{NW}(f)$ denotes the non-wandering set of f .

In [8] conditions on D were introduced to ensure the existence of pruning isotopies; a disk D satisfying these conditions is called a *pruning disk*.

Date: Dec 3, 2011.

2010 Mathematics Subject Classification. Primary .

Key words and phrases. Hénon maps, pruning fronts, symbolic dynamics.

Research supported by FAPESP grant 2007/55771-0.

¹A different way to formalize pruning for the Lozi family was introduced by Ishii in [13] which enabled him to prove the PFC for that family.

Theorem 1.2 (Pruning theorem). *Given a pruning disk D for a homeomorphism f_0 , there exists a pruning isotopy associated to D .*

In words, a pruning disk D for a homeomorphism f_0 is a topological disk for which it is possible to destroy all of the orbits of f_0 which enter D , while leaving other orbits untouched.

The homeomorphism f_1 is called *the pruning homeomorphism of f_0 associated to the disk D* . It is important to see that pruning theory gives models to understand the topological dynamics of homeomorphisms. If we start with a homeomorphism f_0 we can construct the *pruning family of f_0* , $\mathcal{P}(f_0)$, defined to be the closure of the family of all the homeomorphisms obtained applying the pruning theorem to f_0 a finite number of times. Depending on f_0 , the pruning family $\mathcal{P}(f_0)$ contains infinitely many different models of dynamics. For example, if $f_0 = F$ is the Smale horseshoe it can be proved that $\mathcal{P}(F)$ has uncountably many topological models. See [8] for these facts.

In this paper, we use pruning to study the Hénon family [10] which is the family of diffeomorphisms of the plane defined by

$$H_{a,b}(x, y) = (a - x^2 - by, x), \quad a, b \in \mathbb{R}, b > 0.$$

Our work is related to the *Pruning Front Conjecture (PFC)*; which states that each Hénon map can be understood as a partially formed horseshoe, i.e., what is obtained from the horseshoe after pruning some orbits:

Pruning Front Conjecture (PFC). *Up to semiconjugacy, the real Hénon family is contained in $\mathcal{P}(F)$, that is, for any choice of real parameter values a, b , the Hénon map $H_{a,b}$ belongs to $\mathcal{P}(F)$, up to semiconjugacy.*

Many authors have found numerical evidence that for certain choices of parameters (a, b) , the non-wandering set of $H_{a,b}$ is a subset of the Smale horseshoe, e.g. [6, 7]. In particular, Davis, Mackay and Sannami [7] give parameter values $(a_1, b_1) = (5.4, 1)$ where the non-wandering set of $H_{a,b}$ seems to be a subshift of finite type. On other hand, in [1] Arai showed a rigorous computational method to prove hyperbolicity for certain parameter values in the Hénon family. These parameter values contain (a_1, b_1) above and $(a_2, b_2) = (2.25, 0.25)$.

We use the above facts to prove the PFC in some open neighborhoods around (a_1, b_1) and (a_2, b_2) . Our main theorem is the following.

Theorem 1.3. *There exists an open set $A = A_1 \cup A_2$, where A_1 and A_2 are open topological disks in the real parameter plane, such that if $(a, b) \in A$ then $H_{a,b}$ is topologically conjugate to a pruning homeomorphism of the horseshoe.*

In §2 we will define which pruning disks we are going to use. In §3, we present some concepts and results for the complex Hénon family and using Arai's theorem [2], we show Theorem 1.3.

2. THE HORSESHOE, PRUNING THEORY AND SHIFT AUTOMORPHISMS

2.1. The Smale Horseshoe. In this section we recall the definition of the Smale horseshoe and set the notation to be used. The Smale horseshoe F is a hyperbolic diffeomorphism of the plane whose dynamics in its non-wandering set is conjugate to the shift σ in $\Sigma_2 := \{0, 1\}^{\mathbb{Z}}$.

Consider the square $Q = [-1, 1] \times [-1, 1]$ joined to two semi-disk of radii 1/2 and centered in $(-1, 0)$ and $(1, 0)$. Foliate the square Q with horizontal unstable leaves and vertical stable leaves, and begin by choosing the action of F on Q as depicted in Figure




FIGURE 1. Dynamics of F .

1. We require that F should stretch the unstable leaves uniformly and contract the stable leaves uniformly by factors λ and μ , respectively, with $\lambda > 2$ and $\mu < 1/2$.

Then it is possible to prove that the non-wandering set $NW(F)$ is a Cantor set in the plane together with a fixed point x in the left semi-circle. Making a homotopy we can suppose that x is the point ∞ so the non wandering set is only the Cantor set. We also suppose that if $z \in NW(F)$ then $W^s(z) \cap Q$ is a vertical segment and $W^u(z) \cap Q$ is a horizontal segment.

The conjugacy $\vartheta : NW(F) \rightarrow \Sigma_2$ between the horseshoe F and the shift σ is chosen to be usual one, that is, for any $z = (x, y)$ in $NW(F)$ the symbol sequence associated is $\vartheta(z) = (s_i)_{i \in \mathbb{Z}}$ where:

$$(2.1) \quad s_i := \begin{cases} 0 & \text{if } \pi_1(F^i(z)) < 0 \\ 1 & \text{if } \pi_1(F^i(z)) > 0 \end{cases}$$

and π_1 is the projection on the first coordinate $\pi_1(z) := x$. In the following, we identify one point $z \in NW(F)$ by its *code* or symbolic representation $h(z)$ in Σ_2 . When writing elements $s \in \Sigma_2$, it is common to juxtapose a point between s_{-1} and s_0 to indicate the origin of the sequence.

2.2. Pruning. We now present the main definitions and results from [8]. We adapt them to the needs of this paper and avoid technical details will not be used here.

In [8, Section 1] a pruning disk is defined to be an open topological disk D , whose closure is a closed topological disk, its boundary can be written as $\partial D = C \cup E$, where C and E are arcs which meet only at their endpoints, and satisfy the dynamical conditions below plus some technical conditions.

A disk D is a *pruning disk* if

$$(2.2) \quad F^n(C) \cap D = \emptyset \text{ and } F^{-n}(E) \cap D = \emptyset, \forall n \geq 1$$

and

$$(2.3) \quad \lim_{n \rightarrow \infty} \text{diam}(F^n(C)) = 0 \text{ and } \lim_{n \rightarrow \infty} \text{diam}(F^{-n}(E)) = 0.$$

For our purposes, C and E will always be segments of stable, unstable manifolds, respectively, and the definition is made using the concept of homoclinic pruning disk below.




FIGURE 2. Homoclinic disk.

Definition 2.1. A *homoclinic pruning disk* is a open topological disk $D \subset \mathbb{R}^2$ if D satisfies (2.2) and (2.3) and C and E satisfy:

- (i) $E \subset W^u(0^\infty)$ and $C \subset W^s(0^\infty)$,
- (ii) there is an open set G with $\overline{G} \subset D$ and $D \cap \text{NW}(F) \subset G$.

Observe that the conditions ensure that the stable and unstable manifolds of all points in $\text{NW}(F)$ do not accumulate internally on ∂D .

Example 2.2. Let D be the homoclinic disk defined by the homoclinic points $p_0 = 0^\infty 1111 \cdot 10110^\infty$ and $p_1 = 0^\infty 1110 \cdot 10110^\infty$. See Figure 2. In this case, we need three forward iterates of p_0 and p_1 to see that, $\forall n > 0$, the segment $F^n(C)$ does not intersect D , and three backward iterates to see that $F^{-n}(E)$ does not intersects D .

We define *pruning region associated to D* to be the open set $\mathbb{P} := \cup_{i=-\infty}^{\infty} F^i(D)$. The set $\mathbb{P}_s := \vartheta(\mathbb{P} \cap \text{NW}(F))$ is called the *symbolic pruning region*.

Now let $f_0 := F$ be the horseshoe. Then it follows from [8] that:

Theorem 2.3 (Pruning Theorem). *If D is a homoclinic pruning disk, then there exists a pruning isotopy $f_t: f_0 \simeq f_1$ of f_0 .*

It follows that the dynamics of the pruning map f_1 is what remains after removing the pruning region. Symbolically this is:

Lemma 2.4. *The restriction $\vartheta_s = \vartheta|_{\text{NW}(f_1)}$ is a conjugacy between $f_1|_{\text{NW}(f_1)}$ and $\sigma|_{\Sigma_2 \setminus \mathbb{P}_s}$.*

Because we are considering only homoclinic disks as in Definition 2.1, the subshifts obtained by this procedure will always be of finite type. More general subshifts can be obtained considering general pruning disks but they will not occur here.

The *pruning family $\mathcal{P}(F)$* is defined to be the closure, in the C^0 -topology, of the set of all the prunings of F . It can be shown that $\mathcal{P}(F)$ contains uncountably many different models of dynamics and the Pruning Front Conjecture states that \mathcal{P} contains enough models to describe *all* Hénon maps.

2.3. Pruning automorphisms. In this section certain automorphisms of the shift will be discussed. They will be called pruning automorphisms.

Given $N \in \mathbb{N}$ and $s \in \{0, 1\}$ we set $s^N := \underbrace{s_i s_i \dots s_i}_{N\text{-times}}$ and $s^0 := \emptyset$ as the empty word.

Define the homoclinic disk $D_{N,M}$ of the horseshoe F to be the disk that contains the blocks corresponding to the symbolic sequences $0^N 1 \cdot 010^M$ and $0^N 1 \cdot 110^M$. See Figure 3. Then the disk $D_{N,M}$ is bounded by the stable and unstable segments passing through the homoclinic points $p_0 = 0^\infty 110^{N-1} 1 \cdot 110^{M-1} 110^\infty$ and $p_1 = 0^\infty 110^{N-1} 1 \cdot 010^{M-1} 110^\infty$.




FIGURE 3. Pruning disk $D_{N,M}$ with $N = 1$ and $M = 2$.

Proposition 2.1. $D_{N,M}$ is a pruning disk, except when $N = 0$ and $M = 0$; $M = 0$ and $N = 1$; $N = 1$ and $M = 1$; or, $M = 1$ and $N = 0$.

The proof consists of following the orbits of C and E just as in figure 2. Note that $F(D_{N,M})$ is positioned similarly to the pruning disk of that figure.

Let $f_{N,M}$ be the pruning homeomorphism associated to $D_{N,M}$ as in theorem 2.3.

An automorphism of the shift is a shift commuting homeomorphism from Σ_2 to Σ_2 . The set of all the automorphisms of the shift is denoted by $\text{Aut}(\Sigma_2)$, and $\text{Fix}(\rho)$ denote the set of fixed points of a automorphism ρ .

The definition below associates an automorphism of the shift to a pruning disk $D_{N,M}$. These are called *pruning automorphisms*.

Definition 2.5. Let $\rho_{N,M} \in \text{Aut}(\Sigma_2)$ be the shift automorphism that interchanges the sequences $0^N 1010^M$ and $0^N 1110^M$:

$$(2.4) \quad (\rho_{N,M}(s))_i = \begin{cases} 0 & \text{if } s_{i-N-1} \dots s_i \dots s_{i+M+1} = 0^N 1110^M \\ 1 & \text{if } s_{i-N-1} \dots s_i \dots s_{i+M+1} = 0^N 1010^M \\ s_i & \text{otherwise} \end{cases}$$

In figure 3 this exchanges points in D to the left and right of the vertical center line, among many possible other symmetries. The automorphism $\rho_{N,M}$ is called *pruning automorphism associated to the disk $D_{N,M}$* .

By the above definition, $\Sigma_2 \setminus \mathbb{P}_s = \text{Fix}(\rho_{N,M})$ where \mathbb{P}_s is the symbolic pruning front associated to $D_{N,M}$. Hence, by lemma 2.4, $\vartheta_s: \text{NW}(f_{N,M}) \rightarrow \text{Fix}(\rho_{N,M})$ is a conjugacy between $f_{N,M}$ restricted to $\text{NW}(F) \setminus \mathbb{P}$ and the shift σ restricted to $\text{Fix}(\rho_{N,M})$. Therefore, in $\text{NW}(f_{N,M})$, we have the following equation:

$$(2.5) \quad \vartheta_s \circ f_{N,M} = \sigma \circ \vartheta_s.$$

3. THE HÉNON FAMILY

3.1. The horseshoe locus. The Hénon family has an extension, also denoted $H_{a,b}$, to \mathbb{C}^2 , with $(a,b) \in \mathbb{C} \times \mathbb{C}^*$. This has been investigated by many authors. See, e.g., [3, 4, 11, 12]. Here we state some definitions and results of real and complex Hénon maps.

Define

$$(3.1) \quad K_{a,b}^{\mathbb{C}} = \{p \in \mathbb{C}^2 \mid \{H_{a,b}^n(p)\}_{n \in \mathbb{Z}} \text{ is bounded} \}$$

to be the set of bounded orbits and $K_{a,b}^{\mathbb{R}} := K_{a,b}^{\mathbb{C}} \cap \mathbb{R}^2$. Let $\mathcal{H}^{\mathbb{C}}$ be the set of $(a,b) \in \mathbb{C} \times \mathbb{C}^*$ such that the restriction of $H_{a,b}$ to $K_{a,b}^{\mathbb{C}}$ is topologically conjugate to the full 2-shift (σ, Σ_2) . The set $\mathcal{H}^{\mathbb{C}}$ is called the *horseshoe locus*. Likewise, let $\mathcal{H}^{\mathbb{R}}$ be the set of $(a,b) \in \mathbb{R}^2$ such that the restriction of $H_{a,b}$ to $K_{a,b}^{\mathbb{R}}$ is topologically conjugate to the full 2-shift.

Devaney and Nitecki showed that $\mathcal{H}^{\mathbb{R}} \neq \emptyset$ and contains the set

$$(3.2) \quad \text{DN} := \{(a,b) \in \mathbb{R}^2 \mid a > (5 + 2\sqrt{5})(1 + |b|)^2/4, b \neq 0\};$$

they also prove that if $(a,b) \in \text{EMP} := \{(a,c) \in \mathbb{R}^2 : a < -(1 + |b|)^2/4\}$ then $K_{a,b}^{\mathbb{R}} = \emptyset$ (See [9]).

Hubbard and Oberste-Vorth ([14]) showed that $\mathcal{H}^{\mathbb{C}}$ contains the set

$$(3.3) \quad \text{HOV} := \{(a,b) \in \mathbb{C}^2 : |a| > 2(1 + |b|)^2, b \neq 0\}$$

Observe that $\text{DN} \subset \text{HOV}$.

In [2], Arai shows that there are parameters values in $(a,b) \in \mathcal{H}^{\mathbb{C}} \cap \mathbb{R}^2$ such that $K^{\mathbb{R}} \neq \emptyset$ but $H_{a,b}$ restricted to $K^{\mathbb{R}}$ is not conjugate to a full horseshoe. Those were called *parameters of type-3*. Also, in [4], Bedford and Smillie exhibited a way to join distinct connected components of the *real horseshoe locus*, $\mathcal{H}^{\mathbb{C}} \cap \mathbb{R}^2$, using loops in $\mathcal{H}^{\mathbb{C}}$. This can be accomplished by loops starting in DN. Let $\mathcal{H}_0^{\mathbb{C}}$ be the connected component of $\mathcal{H}^{\mathbb{C}}$ that contains HOV and take a point $(a_0, b_0) \in \text{DN} \subset \mathcal{H}_0^{\mathbb{C}}$. Then there exists a conjugacy $h_0 : K_{a_0, b_0}^{\mathbb{C}} \rightarrow \Sigma_2$. Let $\gamma(t)$ be a loop in $\mathcal{H}_0^{\mathbb{C}}$ with basepoint in (a_0, b_0) . By structural stability there are conjugacies h_t between $K_{\gamma(t)}^{\mathbb{C}}$ and Σ_2 . Since $\gamma(0) = \gamma(1) = (a_0, b_0)$, we can define an automorphism of the 2-shift

$$(3.4) \quad \rho(\gamma) := h_1 \circ h_0^{-1} \in \text{Aut}(\Sigma_2).$$

Thus, ρ sends a loop γ to an automorphism in $\text{Aut}(\Sigma_2)$ which only depends on the homotopy class of the loop, $[\gamma]$. Therefore, the map:

$$(3.5) \quad \rho : \pi_1(\mathcal{H}_0^{\mathbb{C}}, (a_0, b_0)) \rightarrow \text{Aut}(\Sigma_2)$$

given by $[\gamma] \rightarrow \rho(\gamma)$ is a group homomorphism. The map ρ is called the *monodromy homomorphism*.

Some results about the horseshoe locus were obtained by Arai in [2]. His main result relates to the loops in $\mathcal{H}_0^{\mathbb{C}}$ which pass through real parameter values (a,b) .

Theorem 3.1 (Arai). *If $(a,b) \in \mathcal{H}_0^{\mathbb{C}} \cap \mathbb{R}^2$, then there exists a loop $\gamma := \bar{\alpha}^{-1} \cdot \alpha$, where α is a path in $\mathcal{H}_0^{\mathbb{C}}$ starting in (a_0, b_0) and ending in (a,b) (so that $\gamma(1/2) = (a,b)$), such that*

- (i) the image of $K_{a,b}^{\mathbb{R}}$ under $h_{1/2}$ is exactly $\text{Fix}(\rho(\gamma))$, and
- (ii) this identification is dynamical, i.e.

$$(3.6) \quad \sigma \circ h_{1/2} = h_{1/2} \circ H_{a,b}.$$

Arai's proof uses the fact that a Hénon map $H_{a,b}$ is conjugate to $H_{\bar{a},\bar{b}}$ by the complex conjugation $\varphi(x, y) := (\bar{x}, \bar{y})$.

It is natural to ask what automorphisms in the theorem above are realized by real Hénon maps. In the next section, we will observe that among those automorphisms are some pruning automorphisms.

4. THE APPLICATION

Now we prove a theorem that allows one to show that a parameter pair (a, b) corresponds to a pruning homeomorphism. Theorem 1.3 follows as a direct application using the rigorous numerical work of Arai [1, 2].

Definition 4.1. Let γ be a closed curve in \mathbb{C}^2 and $\bar{\gamma} := \varphi(\gamma)$. In the following we suppose $\gamma(0) \in \text{DN}$. We say that γ is *symmetric* if $\bar{\gamma} = \gamma^{-1}$, that is, $\bar{\gamma}(t) = \gamma(1-t)$, $\forall t \in [0, 1]$. When γ is symmetric it follows that $\gamma(1/2) \in \mathbb{R}^2$.

Equation (2.5) gives us a connection between a pruning homeomorphism and the possible real dynamics of $H_{a,b}$ when $(a, b) \in \mathcal{H}_0^{\mathbb{C}} \cap \mathbb{R}^2$ in the following way:

Theorem 4.2. *Let $\rho_{N,M}$ be a pruning automorphism of the shift. Suppose that there exists a symmetric curve $\gamma \in \mathcal{H}_0^{\mathbb{C}}$ such that $\rho(\gamma) = \rho_{N,M}$ and define $(a, b) = \gamma(1/2)$. Then there exists a conjugacy between the Hénon map $H_{a,b}$, restricted to $K_{a,b}^{\mathbb{R}}$, and the pruning homeomorphism $f_{N,M}$ restricted to $\text{NW}(f_{N,M})$.*

Proof. Let h_t be the conjugacies between $H_{a,b}$ and σ restricted to $K_{\gamma(t)}^{\mathbb{C}}$ and Σ_2 , respectively. Then $\rho_{N,M} = h_1 \circ (h_0)^{-1}$ by definition. Let $\alpha := \gamma|_{[0,1/2]}$. Then $\gamma_{N,M} = \bar{\alpha}^{-1} \cdot \alpha$ and by theorem 3.1 of Arai, $h_{1/2}$ is a conjugacy between $H_{a,b}$, restricted to $K_{a,b}^{\mathbb{R}}$, and σ , restricted to $\text{Fix}(\rho_{N,M})$. By equation (2.5), the desired conjugacy is $(\vartheta_s)^{-1} \circ h_{1/2}$. \square

Now we prove the Theorem 1.3.

Proof of Theorem 1.3. Let $I_1 = [5.3125, 5.46875] \times \{1\}$ and take $(a, b) \in I_1$. By [1, Theorem 1.1], these parameter values correspond to hyperbolic Hénon maps. By structural stability there exists a connected open set A_1 , with $I_1 \subset A_1$, such that all its maps are conjugate. By [2, Theorem 4.], there exists a symmetric curve $\gamma_1 \in \mathcal{H}_0^{\mathbb{C}}$ such that $\gamma_1(1/2) = (a, b)$ and $\rho(\gamma_1) = \rho_{2,2}$. Hence, by Theorem 4.2, $H_{a,b}$, restricted to $K_{a,b}^{\mathbb{R}}$, is topologically conjugate to the pruning homeomorphism $f_{2,2}$ associated to the pruning disk $D_{2,2}$ and restricted to $\text{NW}(f_{2,2})$.

Let $I_2 = [2.21875, 2.296875] \times \{0.25\}$. The open set A_2 is obtained considering the values $(a, b) \in I_2$. The proof follows the same procedure. In this case $H_{a,b}$, restricted to $K_{a,b}^{\mathbb{R}}$, is topologically conjugate to the pruning homeomorphism $f_{0,2}$ associated to the pruning disk $D_{0,2}$ and restricted to $\text{NW}(f_{0,2})$. \square

5. FURTHER PARAMETERS VALUES

The technique above can be used to prove further cases of PFC provided the symmetric paths as in Theorem 4.2 are found. There is numerical evidence that many other parameter regions fall into this proof scheme but this has not been rigorously verified. For instance, some parameter values are the following with their respective pruning disks.

Parameter values (a, b)	Pruning disks
$(3.5, 0.55)$	$D_{2,3}$
$(2.766, 0.4)$	$D_{0,3}$ and $D_{1,2}$
$(2.887, 0.4)$	$D_{1,3}$ and $D_{2,2}$
$(2.345, 0.19)$	$D_{0,3}$




FIGURE 4. Homoclinic disks conjectured for $H_{2.812,0.4}$.

We conjecture that all these parameters are in the horseshoe locus, because, numerically, it seems that there are paths in $\mathcal{H}^{\mathbb{C}}$ joining these parameters values with points in DN : these paths can be found using the program **SaddleDrop**² and drawing the complex unstable manifold of the fixed point 0^{∞} .

As an example, consider the parameter values $(a, b) = (2.812, 0.4)$. The pruning disks conjectured for this map are $D_{1,2}$ and the homoclinic disk D' defined by the homoclinic orbits $p_0 = 0^{\infty}10111 \cdot 100110^{\infty}$ and $p_1 = 0^{\infty}10110 \cdot 100110^{\infty}$ (See Figure 4). To justify that $(2.812, 0.4) \in \mathcal{H}^{\mathbb{C}}$, we take 9 points in the complex parameter space which, at least numerically, seem to belong to a path joining $(2.812, 0.4)$ and the parameter values $(3.149, 0.4) \in DN$. To see that this path is entirely contained in $\mathcal{H}^{\mathbb{C}}$, draw (using SaddleDrop) the intersection of the set $K_{a,b}$ of bounded orbits with the complex unstable manifold of 0^{∞} . These are shown in Figures 5 - 13. Observing these figures one see what seems to be a Cantor set varying continuously in \mathbb{C}^2 . It suggests that these values are in the horseshoe locus.

6. ACKNOWLEDGEMENT

I am grateful to A. de Carvalho for showing me how to prune and for many suggestions to improve this work. I would like to thank FAPESP for their support, and Peter Hazard

²SaddleDrop is a program created by J. Hubbard and K. Papadantonakis to study the complex Hénon family. It can be found in <http://www.math.cornell.edu/~dynamics/SD/index.html>




FIGURE 5. $a \approx 2.8187 + 0.0119i$ and $b = 0.4$.




FIGURE 6. $a \approx 2.8255 - 0.032i$ and $b = 0.4$.




FIGURE 7. $a \approx 2.8187 - 0.079i$ and $b = 0.4$.




FIGURE 8. $a \approx 2.8187 - 0.095i$ and $b = 0.4$.




FIGURE 9. $a \approx 2.8931 - 0.0959i$ and $b = 0.4$.




FIGURE 10. $a \approx 3.00 - 0.0919i$ and $b = 0.4$.




FIGURE 11. $a \approx 3.1432 - 0.0919i$ and $b = 0.4$




FIGURE 12. $a \approx 3.1432 - 0.048i$ and $b = 0.4$.




FIGURE 13. $a \approx 3.149 - 0.00399i$ and $b = 0.4$.

for helping me prepare this manuscript. I acknowledge the hospitality of the Instituto de Matemática e Estatística IME-USP where this work was carried out.

REFERENCES

- [1] Arai, Z.: *On hyperbolic plateaus of the Hénon map*. Experiment. Math. **16(2)**, 181–188 (2007).
- [2] Arai, Z.: *On loops in the hyperbolic locus of the complex Hénon maps and their monodromies*. ArXiv:math.DS/07042978v1, (2007).
- [3] Bedford, E. and Smillie, J.: *Real polynomial diffeomorphisms with maximal entropy: Tangencies*. Ann. of Math (2). **160(1)**, 1–26 (2004).
- [4] Bedford, E. and Smillie, J.: *The Hénon family: the complex horseshoe locus and real parameter space*. In *Complex dynamics*, volume 396 of *Contemp. Math.*, 21–36. Amer. Math. Soc., Providence, RI, (2006).
- [5] Cvitanović, P.: *Periodic orbits as the skeleton of classical and quantum chaos*. Phys. D, **51(1-3)**, 138–151 (1991).
- [6] Cvitanović, P. and Gunaratne, G. and Procaccia, I.: *Topological and metric properties of Hénon-type strange attractors*. Phys. Rev. A (3), **38(3)**, 1503–1520 (1998).
- [7] Davis, M.J. and Mackay, R.S. and Sannami, A.: *Markov shifts in the Hénon family*. Phys. D, **52(2-3)**, 171–178 (1991).
- [8] de Carvalho, A.: *Pruning fronts and the formation of horseshoes*. Ergodic Theory and Dynam. Systems. **19(4)**, 851–894 (1999).
- [9] Devaney, R. and Nitecki, Z.: *Shift automorphisms in the Hénon mapping*. Comm. Math. Phys., **67(2)**, 137–146 (1979).
- [10] Hénon, M.: *A two dimensional mapping with a strange attractor*. Comm. Math. Phys. **50(1)**, 67–77 (1976).
- [11] Hubbard, J. and Oberste-Vorth, R.: *Hénon mappings in the complex domain. I. The global topology of dynamical systems*. Inst. Hautes tudes Sci. Publ. Math. **79**, 5–46 (1994).
- [12] Hubbard, J. and Oberste-Vorth, R.: *Hénon mappings in the complex domain. II. Projective and inductive limits of polynomials*. In *Real and complex dynamical systems(Hilerod, 1993)*, volume 464 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, 89–132. Kluwer Acad. Publ., Dordrecht (1995).
- [13] Ishii, Y.: *Towards a kneading theory for Lozi mappings. I. A solution of the pruning front conjecture and the first tangency problem*. Nonlinearity. **10(3)**, 731–747 (1997).
- [14] Oberste-Vorth, R.: *Complex horseshoes and the dynamics of mappings of two complex variables*. PhD thesis. Cornell University, (1987).

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO. RUA DO MATÃO 1010, CIDADE UNIVERSITARIA, CEP 05508-090, BRAZIL.

E-mail address: jvalentm@ime.usp.br