

Non-thermal escape of molecular hydrogen from Mars

M. Gacesa^{1,2}, P. Zhang¹, and V. Kharchenko^{1,2}

We present a detailed theoretical analysis of a non-thermal escape of molecular hydrogen from Mars induced by collisions with hot atomic oxygen from martian corona. To accurately describe the energy transfer in $O + H_2(v, j)$ collisions, we performed extensive quantum-mechanical calculations of state-to-state elastic, inelastic, and reactive cross sections. The escape flux of H_2 molecules was evaluated using a simplified 1D column model of the martian atmosphere with realistic densities of atmospheric gases and hot oxygen production rates for the low solar activity conditions. An average density of the non-thermal escape flux of H_2 of $1.9 \times 10^5 \text{ cm}^{-2}\text{s}^{-1}$ was obtained considering energetic O atoms produced in dissociative recombinations of O_2^+ ions. Predicted rovibrational distribution of the escaping H_2 was found to contain a significant fraction of higher rotational states. While the non-thermal escape rate was found to be lower than Jeans flux for H_2 molecules, the non-thermal escape rates of HD and D_2 are significantly higher than their respective Jeans rates. The accurate values of non-thermal escape fluxes of different molecular isotopes of H_2 may be important in analyses of evolution of the martian atmosphere. The described molecular ejection mechanism is general and expected to contribute to atmospheric escape of H_2 and other light molecules from planets, satellites, and exoplanetary bodies.

1. Introduction

The interaction of the martian atmosphere with the solar radiation and interplanetary plasma results in its evaporation due to thermal (Jeans) escape and a number of non-thermal mechanisms. The absence of the intrinsic magnetic field on Mars and its low gravitational potential make the martian atmosphere particularly susceptible to erosion [Acuna *et al.*, 1998]. The current low atmospheric pressure can be explained well by extrapolating the escape rates to a geological time frame while accounting for a change of the solar activity in the past [Chassefière and Leblanc, 2004]. The present day degradation of the martian atmosphere occurs mainly via non-thermal escape processes induced by ion charge-exchange, sputtering and ionospheric outflows driven by solar wind [Chassefière and Leblanc, 2004; Johnson *et al.*, 2008]. The dissociative recombination (DR) of O_2^+ is a major source of hot O atoms in the upper atmosphere of Mars, responsible for the escape of oxygen and formation of martian hot corona [Ip, 1988; Fox, 1993; Krest'yanikova and Shematovich, 2005].

¹Institute for Theoretical Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA.

²Department of Physics, University of Connecticut, Storrs, CT, USA.

Nascent hot O atoms collide with thermal constituents of the martian atmosphere and eject them from the planetary gravitational field, if a sufficient kinetic energy transfer occurs. Suprathermal neutral oxygen was shown to be important for analyses of Mars' corona and the non-thermal escape of neutral atoms [Krest'yanikova and Shematovich, 2005; Krest'yanikova and Shematovich, 2006; Fox and Hać, 2009]. Recent calculations of the non-thermal He escape from Mars carried out with accurate energy transfer parameters predicted a significant He escape flux induced by hot O atoms [Bovino *et al.*, 2011].

In this Letter we explore collisional ejection of molecules from the martian atmosphere. Specifically, we report the results of a quantum-mechanical study of the energy transfer from the hot O to H_2 molecules and their subsequent escape. Significant computational difficulties arise from the fact that molecular internal rotational and vibrational degrees of freedom can be excited in collisions. In addition, the reactive pathway leading to the production of OH molecules is energetically permitted. To account for the increased complexity, we have computed the cross sections for $O(^3P) + H_2$ reactive collision using fully quantum-mechanical approach. Kinetic theory was used to calculate the rate of energy transfer, as well as distributions of excited rotational and vibrational (RV) states of the recoiled H_2 molecules. The total escape flux of H_2 from Mars and RV distributions of escaping molecules have been evaluated for the low solar activity conditions. Also, we have estimated the non-thermal escape fluxes of HD and D_2 and compared them to the corresponding Jeans escape rates. Finally, the dependence of the molecular ejection fluxes on the gravitational escape threshold is analyzed for conditions present on other planets, satellites, and exoplanets.

2. Cross Sections and Energy Transfer

The DR of O_2^+ with electrons proceeds via five possible dissociation pathways, producing $O(^3P)$, $O(^1D)$, and $O(^1S)$ [Guberman, 1988; Fox and Hać, 2009]. Energetic metastable $O(^1D)$ atoms decay via spontaneous emission and quenching in collisions with atmospheric gases into $O(^3P)$ atoms [Kharchenko *et al.*, 2005]. The cross sections for $O(^3P)$ and $O(^1D)$ colliding with He were found to be very similar [Bovino *et al.*, 2011]. For simplicity, we assumed a similar behavior for $O(^3P) + H_2$ and $O(^1D) + H_2$ elastic collisions.

To describe the collision ejection of H_2 molecules, we have calculated elastic and inelastic cross sections for the center-of-mass (CM) collision energies from 0.01 to 4.5 eV. The quantum scattering code ABC [Skouteris *et al.*, 2000], that can treat elastic, inelastic, and open reactive channels of the OH production, $O(^3P) + H_2(v, j) \rightarrow OH(v'', j'') + H$, where (v, j) and (v'', j'') indicate initial and final RV levels of H_2 and OH, was used to solve the time-independent coupled-channel Schrödinger equation in Delves hyperspherical coordinates. In addition, for high collision energies, the elastic and inelastic cross sections were calculated using the MOLSCAT [Hutson, 1994] code to ensure the convergence of the nonreactive channels¹. Extensive numerical convergence tests were carried out for the both codes.




Figure 1. Elastic and momentum transfer cross sections for H₂($v = 0, j = 0, 1, 2$) + O collisions. The momentum transfer cross section shown is thermally averaged over the first three rotational states. *Inset:* Total inelastic cross sections $\sigma_{vj, v'j'}^{\text{inel}}(E) = \sum_{v', j'} \sigma_{vj, v'j'}(E)$ for H₂($v = 0, j = 0, 1, 2$) + O \rightarrow H₂($v' = 0, j'$) + O, and $j \neq j'$. The reactive cross section for H₂($v = 0, j = 0$) + O \rightarrow OH + H is also shown and compared with the experimental results (black squares) [Garton, 2003].

The O(³P)+H₂(v, j) interaction was described using two lowest potential energy surfaces, Rogers' LEPS ³A'' [Rogers *et al.*, 2000] and Brandão's BMS1 ³A' [Brandão *et al.*, 2004]. Partial cross sections for initial and final rotational levels j and j' were constructed as a statistically weighted sum of the independently calculated cross sections for the two potential surfaces, where both ³A'' and ³A' contribute a weight factor of 1/3 [Balakrishnan, 2004]. Elastic, inelastic, and momentum transfer partial cross sections for oxygen colliding with the hydrogen molecule in three energetically lowest rotational states are given in Figure 1. We compared our reactive cross sections for OH production to the previously published results [Balakrishnan, 2003, 2004; Braunstein *et al.*, 2004; Wei-keh Wu, 2010] and found them to be in close agreement within the available energy range.

To determine the energy transfer rate from the suprathermal oxygen to atmospheric H₂ and find its escape rate, we used kinetic theory with a quantum description of internal molecular structure and realistic anisotropic cross sections. Since the reactive cross sections are an order of magnitude smaller than the elastic cross sections (Figure 1), and the more massive OH molecule has a considerably higher escape threshold than H₂, we neglected it in this study. However, note that a small fraction of produced OH molecules may be sufficiently energetic to escape. The transferred energy from the energetic projectile O to the frozen target H₂ in the laboratory frame (LF) can be expressed as [Johnson, 1982]

$$T_{v', j'} = \frac{m_{\text{O}} m_{\text{H}_2}}{(m_{\text{O}} + m_{\text{H}_2})^2} (1 + \gamma_{v', j'} - 2\sqrt{\gamma_{v', j'}} \cos \theta) E, \quad (1)$$

where m_{O} and m_{H_2} are masses of O and H₂, respectively, E is the collision energy in the LF, $\gamma_{v', j'} = \epsilon_{v', j'}/\epsilon$, is the ratio of $\epsilon_{v', j'}$ and ϵ , the CM translational kinetic energies after and before the collision, respectively, and θ is the scattering angle in the CM frame. The energies $\epsilon_{v', j'}$ were calculated quantum mechanically for the two triplet potential surfaces. Eq. (1) takes into account that the energy transferred to H₂ molecules is spent on increasing their translational kinetic energy and exciting their internal RV degrees of freedom.

The fraction of energized H₂ molecules capable of escaping can be calculated as

$$\Gamma_{vj}^{v'j'}(E) = \frac{\int_{\theta_{\text{min}}}^{\pi} Q_{vj, v'j'}(\theta) (1 - \cos \theta) \sin \theta d\theta}{\int_0^{\pi} Q_{vj, v'j'}(\theta) (1 - \cos \theta) \sin \theta d\theta}, \quad (2)$$

where $Q_{vj, v'j'}(\theta)$ is the differential cross section for scattering of H₂ in the initial (v, j) into the final (v', j') state. The critical angle θ_{min} was determined from the condition that the translational part of the transferred energy $T_{v', j'}$ is equal to the minimum energy required for H₂ to escape from Mars, $E_{\text{esc}} = 0.26$ eV. An alternative description of the escape process could be constructed by performing Monte Carlo simulations with accurate quantum cross sections for angular distributions of the recoiled H₂ molecules.

Momentum transfer cross sections $\sigma_{vj, v'j'}^{\text{mt}}$ for inelastic collisions were calculated using [Parker and Pack, 1978]

$$\sigma_{vj, v'j'}^{\text{mt}} = 2\pi \int_0^{\pi} d\theta \sin \theta (1 - \sqrt{1 - \gamma_{v', j'} \cos \theta}) Q_{vj, v'j'}(\theta). \quad (3)$$

3. Flux and Distribution of Escaping H₂

Jeans escape and collisions with hot oxygen are the major mechanisms that contribute to the escape of neutral H₂ molecules and their isotopes from the martian atmosphere. Both processes are strongly dependent on the temperature and density of upper layers of the martian atmosphere. The temperature of the exosphere (above the altitude of about 180 km), T_{exo} , is approximately constant and estimated to be between 240 and 280 K, depending on the solar activity and gas density profiles [Krasnopolsky, 2010; Fox and Hać, 2009; Fox, 2003]. To obtain a conservative estimate of the non-thermal flux of escaping H₂, we considered the $T_{\text{exo}} = 240$ K, corresponding to the low solar activity. Furthermore, we assumed a thermal distribution of the initial rotational states of H₂, where more than 95 % of the total population is distributed between its first three rotational states, $j = 0, 1, 2$, with corresponding population fractions equal to 0.31, 0.46, and 0.19, respectively. Using Eqs. (2,3) we have calculated the values of the thermally averaged momentum transfer cross sections and the fractions $\Gamma_{vj}^{v'j'}(E)$ of rovibrationally excited H₂, sufficiently energetic to escape from Mars (Figure 2). The fraction $\Gamma_{vj}^{v'j'}(E)$ becomes significant at collision energies greater than 0.7 eV for H₂($j' = 0, 2$). Although higher rotational states require increasingly larger projectile energies, *e.g.* H₂($j' = 16$) can escape for $E > 1.65$ eV, their fraction in the RV distribution of the escaping molecules also becomes larger. Note that the initial population of higher vibrational levels of H₂ at the exobase is negligible.

Since $\Gamma_{vj}^{v'j'}(E)$ depends only on the energy transfer efficiency and the escape energy threshold, it can be easily generalized to different astronomical objects. We illustrate this for two hypothetical planets, the first corresponding in size and mass to Earth and the second to the extrasolar super-earth Kepler-10b (3.3 Earth masses) [Batalha *et al.*, 2011]. Note that only very energetic H₂, mostly in higher excited rotational levels, is able to escape (Figure 2).




Figure 2. *Top:* State-to-state momentum transfer cross sections for the H₂(v', j'). Curves for the final rotational levels, $j' = 0, 2, \dots, 16$ (labeled on graph), collisionally excited from the ground state of H₂, are shown. *Bottom:* Fraction $\Gamma_{v_j}^{v'j'}$ of the recoiled H₂($v' = 0, j'$) with energies greater than the escape energy for Mars. The escaping fractions for Earth for $j' = 0, 4$ (dashed blue) and Kepler-10b super-earth for $j' = 0, 8$ (light green) are given for comparison.

A simple estimate of the total escape flux of H₂(v', j') can be obtained from the exobase approximation [Fox, 2003; Chassefière and Leblanc, 2004; Krasnopolsky, 2010], using the density of exospheric H₂, and fractions $\Gamma_{v_j}^{v'j'}(E)$ calculated above. However, such an approach neglects the hot O production below the exobase and loss of the upward flux in atmospheric collisions, resulting in a large uncertainty in the computed flux.

We constructed a more realistic 1D model of escape, analogous to the one used to describe the escape of He atoms [Bovino et al., 2011]. In our model the explicit consideration of energy transfer collisions is combined with the altitude-dependent rate of production of hot O atoms, $f(E, h)$ via DR channels [Guberman, 1988; Petrigiani et al., 2005; Bovino et al., 2011]. In addition, we estimated the extinction of fluxes of suprathermal O and H₂ due to collisions with thermal atmospheric gases. All calculations were performed for low solar activity. We used the rate of production of hot O below 400 km by Fox and Hać [2009] and smoothly interpolated it to the rate given by Krasnopolsky and Gladstone [1996] at higher altitudes.

The volume production rate of escaping hot H₂(v', j') can be expressed as

$$P_{v',j'}(h) = \frac{1}{2} \int_0^\infty dE T_{H_2}(h, E) n_{H_2}(h) \Gamma_{v_j}^{v'j'}(E) \sigma_{v_j, v'j'}^{mt}(E) \times \int_{h_2^{\min}}^h dh_2 f(E, h_2) T_O(h_2, h, E), \quad (4)$$

with the transparency factors T_{H_2} and T_O defined as

$$T_{H_2}(h, E) = \exp \left[- \int_h^{h_{\max}} dh' \sum_i \sigma_{H_2, i}^{mt}(E) n_i(h') \right] \\ T_O(h_2, h, E) = \exp \left[- \int_{h_2}^h dh' \sum_i \sigma_{O, i}^{mt}(E) n_i(h') \right]. \quad (5)$$




Figure 3. Altitude profile of the volume production rate $P_{v',j'}(h)$ of the non-thermal flux of H₂ molecules escaping from Mars. The most significant rates with respect to the initial and final rotational states, $j = 0$ (solid), $j = 1$ (dashed) and $j = 2$ (dotted), and $j' = 0 - 10$, are shown. The curves are denoted as jj' .

The transparency factor T_{H_2} is equal to the escape probability of hot H₂(v', j') produced in collisions with the incident hot O of energy E at the altitude h_2 . The second transparency factor T_O is defined as the probability that the hot O atoms, produced at the altitude h_2 , reach the altitude h without the energy loss in collisions with other atmospheric constituents. The quantity $n_{H_2}(h) \Gamma_{v_j}^{v'j'}(E) \sigma_{v_j, v'j'}^{mt}(E)$ is the inverse mean free path for O+H₂(v, j) collisions, resulting in the energy transfer greater than the H₂ escape threshold. The prefactor 1/2 indicates that, in our simplified 1D model, approximately half of the nascent energetic atoms and recoiled H₂ molecules are scattered towards the planet and cannot escape regardless of the energy transferred. Summations of the flux loss of H₂ and O in collisions with the i -th atmospheric gas of density $n_i(h)$ and momentum transfer cross section $\sigma_{H_2, i}^{mt}(E)$ and $\sigma_{O, i}^{mt}(E)$, respectively, included major constituents of the martian upper atmosphere: CO₂, CO, N₂, O₂, H₂, H, Ar, and He. The momentum transfer cross sections for H-H₂ [Krstic and Schultz, 1999], Ar-H₂ [Udus et al., 2005], H₂-H₂ [Phelps, 1990] were used from the literature. Since no data were available in the required energy range, we used approximate mass-scaled cross sections for He-H₂ (from Ar-H₂), N₂-H₂, O₂-H₂, CO-H₂ (from O-H₂), and CO₂-H₂ (from O-N₂ [Balakrishnan et al., 1998]).

Using Eq. (4) we have calculated volume production rate of the escaping H₂($v' = 0, j'$) molecules induced in H₂($v = 0, j = 0 - 2$) + O collisions for a range of altitudes from $h_{\min} = 130$ km to $h_{\max} = 800$ km (Figure 3). Note that, by symmetry arguments, for a homonuclear H₂ only $\Delta j = 0, 2, 4, \dots$ transitions are allowed [Cohen-Tannoudji et al., 1986]. The resulting escape rates of H₂ molecules are the largest for the elastic collisions, followed by the three times smaller rates for the first two excited rotational states, $j' = 2$ and $j' = 4$. The rates remain significant for the final rotational levels up to $j' = 10$.




Figure 4. *Top:* Collisional and thermal total escape rate of H₂($v' = 0, j'$) for the first 16 rotational states j' . *Bottom:* The same as above for H₂, HD, and D₂.

The altitude profile of the production rate of H₂ capable of escaping is similar to the production rate profile of He [Bovino *et al.*, 2011]. This was expected, since the escape of both species is driven by collisions with the nascent fast O atoms, produced mostly below 150 km for the considered atmospheric and solar conditions. The calculated altitude profile can be used to compute the non-thermal escape flux of H₂ molecules and estimate the accuracy of the exobase approximation. We calculated $\phi_{j'}$, the non-thermal flux for H₂($v' = 0, j'$) molecules as

$$\phi_{j'} = \int_{h_{\min}}^{h_{\max}} P_{v'j'}(h)dh. \quad (6)$$

Total collisional and thermal fluxes were calculated as sums over all rotational levels and found to be $1.9 \times 10^5 \text{ cm}^{-2} \text{ s}^{-1}$ and $1.1 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$, respectively. A comparison of Jeans and non-thermal rates of escape of H₂($v' = 0, j'$) molecules, escaping in different rotational states j' from martian day-side, is given in Figure 4. To simplify the calculation we assumed the average solar conditions and neglected the latitude dependence of the production rate of hot O atoms. Jeans rate is about eight times greater than the non-thermal rate of the escaping H₂ for the lowest three rotational states, while for $j' > 3$ the latter starts to dominate. The distinct character of the two RV distributions is a clear signature of different physical escape mechanisms.

While, in case of H₂, thermal rate is almost an order of magnitude higher than the collisionally-induced rate of escape, relative importance of the two processes changes for heavier isotopologues, namely HD and D₂ (Table 1). A similar scaling of the collisional and thermal escape fractions can be expected for molecular escape from more massive astronomical objects.

Table 1. Total collisionally-induced escape rates of H₂, HD, and D₂ from the martian atmosphere.

	H ₂	HD	D ₂
Jeans escape rate (s ⁻¹)	1.1×10^6	2.7	3.3×10^{-6}
Non-thermal escape rate (s ⁻¹)	1.9×10^5	74	0.03

4. Conclusions

We find that the collisionally-induced outflow of H₂ molecules and their heavier isotopes contributes to the evolution of the martian atmosphere. Namely, the escape rate of molecular hydrogen induced by collisions with hot oxygen from the martian atmosphere was calculated and found to be about six times smaller than the corresponding Jeans escape rate for the low solar activity. For heavier molecules, the collisional escape will dominate over thermal, as we have illustrated in case of HD and D₂ isotope molecules. In fact, the described process of molecular ejection induced by collisions may be one of the most important escape mechanisms of HD and D₂ from Mars. Consequently, the calculated escape fluxes of H₂ isotopologues could be important in analyses of the H/D ratio on Mars and evolution of water in martian history.

The described mechanism of molecular escape, where collisions provides sufficient translational energy to exceed the escape threshold and simultaneously excite internal molecular degrees of freedom, is rather general. It could be used to evaluate non-thermal escape fluxes and RV distributions of heavier molecules, such as CO, N₂, or CH₄, from Mars, Solar system bodies, and exoplanets. For the escape flux induced by O atoms produced in DR the upper limit on the mass of the escaping molecule is about 30 *u* on Mars. Similarly, we estimate that the non-thermal escape of H₂ from a planetary atmosphere is possible for planetary masses up to about 3.4 Earth masses. A number of solar system bodies as well as the lightest currently confirmed exoplanets belong in that mass range. These limits do not include other non-thermal sources of hot atoms.

The escaping H₂ molecules exhibit a characteristic internal energy distribution, with a significant fraction of populated higher rotational states. Since H₂ molecules do not have a permanent electric dipole moment, they decay to the ground state mainly via collisions with the background gases present in an extended planetary corona. This is true for all escaping molecules that do not have their permanent dipole moment. It could be possible to indirectly detect the presence of H₂ or other rotationally excited light molecules in the extended martian corona from a careful analysis of the collision rates and abundancies of the excited coronal species. Finally, a significant amount of rovibrationally excited H₂ molecules remain in the martian atmosphere after colliding with hot O atoms. The cross sections and energy transfer parameters presented in this study can be used to determine non-thermal translational and rovibrational distributions of hot H₂ gas in the upper atmosphere of Mars.

Acknowledgments. We are grateful to D. Wang, A. Kupfermann, and J. Brandão for providing Fortran subroutines for constructing potential energy surfaces, and to N. Lewkow for reading and suggestions. M.G. and V.K. were supported by NASA grants NNX09AF13G and NNX10AB88G.

Notes

1. A detailed description of the scattering calculations and resulting cross sections for the two lowest potential energy surfaces will be published elsewhere.

References

- Acuna, M. H., *et al.* (1998), Magnetic Field and Plasma Observations at Mars: Initial Results of, *Science*, 279, 1676–+, doi:10.1126/science.279.5357.1676.

- Balakrishnan, N. (2003), Quantum mechanical investigation of the O+H₂ → OH+H reaction, *J. Chem. Phys.*, *119*, 195–199, doi:10.1063/1.1576532.
- Balakrishnan, N. (2004), Quantum calculations of the O(³P)+H₂ → OH+H reaction, *J. Chem. Phys.*, *121*, 6346–6352, doi:10.1063/1.1789871.
- Balakrishnan, N., V. Kharchenko, and A. Dalgarno (1998), Slowing of energetic O(³P) atoms in collisions with N₂, *J. Geophys. Res.*, *103*, 23,393–23,398, doi:10.1029/98JA02198.
- Batalha, N. M., et al. (2011), Kepler’s first rocky planet: Kepler-10b, *The Astrophysical Journal*, *729*(1), 27.
- Bovino, S., P. Zhang, F. A. Gianturco, A. Dalgarno, and V. Kharchenko (2011), Energy transfer in O collisions with He isotopes and Helium escape from Mars, *Geophys. Res. Lett.*, *38*, L02,203, doi:10.1029/2010GL045763.
- Brandão, J., C. Mogo, and B. C. Silva (2004), Potential energy surface for H₂O(³A'') from accurate ab initio data with inclusion of long-range interactions, *J. Chem. Phys.*, *121*, 8861–8868, doi:10.1063/1.1802434.
- Braunstein, M., S. Adler-Golden, B. Maiti, and G. C. Schatz (2004), Quantum and classical studies of the O(³P)+H₂(*v* = 0–3, *j* = 0) → OH+H reaction using benchmark potential surfaces, *J. Chem. Phys.*, *120*, 4316–4323, doi:10.1063/1.1642580.
- Chassefière, E., and F. Leblanc (2004), Mars atmospheric escape and evolution; interaction with the solar wind, *Planet. Space Sci.*, *52*, 1039–1058, doi:10.1016/j.pss.2004.07.002.
- Cohen-Tannoudji, C., B. Diu, and F. Laloe (1986), *Quantum Mechanics, Volume 2*.
- Fox, J. L. (1993), On the escape of oxygen and hydrogen from Mars, *Geophys. Res. Lett.*, *20*, 1747–1750, doi:10.1029/93GL01118.
- Fox, J. L. (2003), Effect of H₂ on the Martian ionosphere: Implications for atmospheric evolution, *J. Geophys. Res.*, *108*, 1223, doi:10.1029/2001JA000203.
- Fox, J. L., and A. B. Hać (2009), Photochemical escape of oxygen from Mars: A comparison of the exobase approximation to a Monte Carlo method, *Icarus*, *204*, 527–544, doi:10.1016/j.icarus.2009.07.005.
- Garton, et al.(2003), A crossed molecular beams study of the O(³P)+H₂ reaction: Comparison of excitation function with accurate quantum reactive scattering calculations, *J. Chem. Phys.*, *118*, 1585–1588, doi:10.1063/1.1539043.
- Guberman, S. L. (1988), The production of O(¹D) from dissociative recombination of O₂⁺, *Planet. Space Sci.*, *36*, 47–53, doi:10.1016/0032-0633(88)90145-6.
- Hutson, G. S., J. M. (1994), Molscat version 14, collaborative computational project 6 (daresbury laboratory: Uk engineering and phys. sci. res. council).
- Ip, W.-H. (1988), On a hot oxygen corona of Mars, *Icarus*, *76*, 135–145, doi:10.1016/0019-1035(88)90146-7.
- Johnson, R. E. (1982), *Introduction to atomic and molecular collisions*, Plenum Press, 1982, New York.
- Johnson, R. E., M. R. Combi, J. L. Fox, W.-H. Ip, F. Leblanc, M. A. McGrath, V. I. Shematovich, D. F. Strobel, and J. H. Waite (2008), Exospheres and Atmospheric Escape, *Space Sci. Rev.*, *139*, 355–397, doi:10.1007/s11214-008-9415-3.
- Kharchenko, V., A. Dalgarno, and J. L. Fox (2005), Thermospheric distribution of fast O(¹D) atoms, *J. Geophys. Res.*, *110*, A12,305, doi:10.1029/2005JA011232.
- Krasnopolsky, V. A. (2010), Solar activity variations of thermospheric temperatures on Mars and a problem of CO in the lower atmosphere, *Icarus*, *207*, 638–647, doi:10.1016/j.icarus.2009.12.036.
- Krasnopolsky, V. A., and G. R. Gladstone (1996), Helium on Mars: EUVE and PHOBOS data and implications for Mars’ evolution, *J. Geophys. Res.*, *101*, 15,765–15,772, doi:10.1029/96JA01080.
- Krest’yanikova, M. A., and V. I. Shematovich (2005), Stochastic models of hot planetary and satellite coronas: a photochemical source of hot Oxygen in the upper atmosphere of Mars, *Sol. Syst. Res.*, *39*, 22–32, doi:10.1007/s11208-005-0002-9.
- Krestyanikova, M. A., and V. I. Shematovich (2006), Stochastic models of hot planetary and satellite coronas: A hot oxygen corona of Mars, *Sol. Syst. Res.*, *40*, 384–392, doi:10.1134/S0038094606050030.
- Krstic, P. S., and D. R. Schultz (1999), Elastic and vibrationally inelastic slow collisions: H + H₂, H⁺ + H₂, *J. Phys. B*, *32*, 2415–2431, doi:10.1088/0953-4075/32/10/310.
- Parker, G. A., and R. T. Pack (1978), Rotationally and vibrationally inelastic scattering in the rotational IOS approximation. Ultrasimple calculation of total (differential, integral, and transport) cross sections for nonspherical molecules, *J. Chem. Phys.*, *68*, 1585–1601, doi:10.1063/1.435927.
- Petrignani, A., F. Hellberg, R. D. Thomas, M. Larsson, P. C. Cosby, and W. J. van der Zande (2005), Electron energy-dependent product state distributions in the dissociative recombination of O₂⁺, *J. Chem. Phys.*, *122*(23), 234,311–+, doi:10.1063/1.1937388.
- Phelps, A. V. (1990), Cross Sections and Swarm Coefficients for H⁺, H₂⁺, H₃⁺, H, H₂, and H⁻ in H₂ for Energies from 0.1 eV to 10 keV, *J. Phys. Chem. Ref. Data*, *19*, 653–675, doi:10.1063/1.555858.
- Rogers, S., D. Wang, A. Kuppermann, and S. Walch (2000), Chemically accurate ab initio potential energy surfaces for the lowest 3a and 3a electronically adiabatic states of o(3p) + h₂, *J. Phys. Chem. A*, *104*(11), 2308–2325, doi:10.1021/jp992985g.
- Skouteris, D., J. F. Castillo, and D. E. Manolopoulos (2000), ABC: a quantum reactive scattering program, *Comp. Phys. Comm.*, *133*, 128–135, doi:10.1016/S0010-4655(00)00167-3.
- Uudus, N., S. Magaki, and N. Balakrishnan (2005), Quantum mechanical investigation of rovibrational relaxation of H₂ and D₂ by collisions with Ar atoms, *J. Chem. Phys.*, *122*(2), 024,304–+, doi:10.1063/1.1829976.
- Wei-keh Wu, V. (2010), Stereodynamics of O(³P)+H₂ at Scattering Energies of 0.5, 0.75, and 1.0 eV, *Chin. J. Chem. Phys.*, *23*, 149–154, doi:10.1088/1674-0068/23/02/149-154.