

# COMPACT ACTIONS AND UNIQUENESS OF THE GROUP MEASURE SPACE DECOMPOSITION OF $\text{II}_1$ FACTORS

ADRIAN IOANA

ABSTRACT. We prove that any  $\text{II}_1$  factor  $L^\infty(X) \rtimes \Gamma$  arising from a compact, free, ergodic, probability measure preserving action of a countable group  $\Gamma$  with positive first  $\ell^2$ -Betti number, has a unique group measure space Cartan subalgebra, up to unitary conjugacy.

## 1. INTRODUCTION

The *group measure space construction* of Murray and von Neumann associates to every probability measure preserving (p.m.p.) action  $\Gamma \curvearrowright (X, \mu)$  of a countable group  $\Gamma$ , a finite von Neumann algebra  $M = L^\infty(X) \rtimes \Gamma$  [MvN36]. If the action is (essentially) free and ergodic, then  $M$  is a  $\text{II}_1$  factor and  $A = L^\infty(X)$  is a *Cartan subalgebra*, i.e. a maximal abelian von Neumann subalgebra whose normalizer,  $\mathcal{N}_M(A) = \{u \in \mathcal{U}(M) | uAu^* = A\}$ , generates  $M$ , as a von Neumann algebra.

During the last decade, S. Popa's *deformation/rigidity* theory has led to spectacular progress in the classification of group measure space  $\text{II}_1$  factors (see the surveys [Po07], [Va10a]). In particular, several large families of group measure space  $\text{II}_1$  factors  $L^\infty(X) \rtimes \Gamma$  have been shown to have a unique Cartan subalgebra [OP07], [OP08], [CS11], [CSU11], [PV11a], [PV11b] or a unique group measure space Cartan subalgebra [Pe09], [PV09], [Io10], [FV10], [IPV10], [CP10], [HPV10], [Va10b], [Io11]. Such "unique Cartan subalgebra" results are extremely useful because they enable one to reduce the classification of the factors  $L^\infty(X) \rtimes \Gamma$ , up to isomorphism, to the classification of the corresponding actions, up to *orbit equivalence*. Indeed, by [Si55], [FM77], two free, ergodic actions  $\Gamma \curvearrowright X$  and  $\Lambda \curvearrowright Y$  are orbit equivalent if and only if there exists an isomorphism of group measure space factors  $L^\infty(X) \rtimes \Gamma \cong L^\infty(Y) \rtimes \Lambda$  which identifies the Cartan subalgebras  $L^\infty(X)$  and  $L^\infty(Y)$ . Recall that the actions  $\Gamma \curvearrowright X$  and  $\Lambda \curvearrowright Y$  are orbit equivalent if there exists an isomorphism of probability spaces  $\theta : X \rightarrow Y$  such that  $\theta(\Gamma \cdot x) = \Lambda \cdot \theta(x)$ , almost everywhere; for recent progress on orbit equivalence, see the surveys [Fu09], [Ga10].

In their breakthrough work [OP07], S. Popa and N. Ozawa proved that  $\text{II}_1$  factors  $L^\infty(X) \rtimes \Gamma$  associated with free, ergodic, compact actions of free groups  $\Gamma = \mathbb{F}_n$  and their direct products  $\Gamma = \mathbb{F}_{n_1} \times \dots \times \mathbb{F}_{n_k}$  have a unique Cartan subalgebra, up to unitary conjugacy. Their proof makes crucial use of the fact that free groups satisfy a strong form of *weak amenability*, called the *complete metric approximation property* (*c.m.a.p.*). Subsequently, the results of [OP07] have been extended to other classes of groups  $\Gamma$ : lattices in products of  $\text{SO}(n, 1)$  and  $\text{SU}(n, 1)$  [OP08], groups having positive first  $\ell^2$ -Betti number and the c.m.a.p. [Da10], [Si10], hyperbolic groups [CS11] and direct products of hyperbolic groups [CSU11] (the last two results notably using [Oz07] and [Oz10]). Note, however, that in all of these cases one requires that both the action  $\Gamma \curvearrowright X$  is compact and, implicitly, that the group  $\Gamma$  is weakly amenable.

In a very recent breakthrough, S. Popa and S. Vaes obtained wide-ranging unique Cartan subalgebra results by removing the compactness assumption on the action. More precisely, they showed that if  $\Gamma$  is either a weakly amenable group with  $\beta_1^{(2)}(\Gamma) > 0$  [PV11a] or a hyperbolic group [PV11b] (or

a direct product of groups in one of these two classes), then  $\text{II}_1$  factors  $L^\infty(X) \rtimes \Gamma$  associated with *arbitrary* free, ergodic actions of  $\Gamma$  have a unique Cartan subalgebra, up to unitary conjugacy.

The main result of [OP07] shows in particular that any  $\text{II}_1$  factor arising from a free, ergodic, compact action of a free group  $\mathbb{F}_n$  has a unique group measure space Cartan subalgebra. The original motivation for our paper was to give a proof of this result which does not rely on the weak amenability of  $\mathbb{F}_n$ . By combining our recent work [Io11] with techniques from [OP07] and [CP10], we were able to not only prove this, but to show the following:

**Theorem 1.1.** *Let  $\Gamma \curvearrowright (X, \mu)$  be a free, ergodic, compact, p.m.p. action of a countable group  $\Gamma$  with  $\beta_1^{(2)}(\Gamma) > 0$ .*

*Then the  $\text{II}_1$  factor  $M = L^\infty(X) \rtimes \Gamma$  has a unique group measure space Cartan subalgebra, up to unitary conjugacy. More precisely, if  $M = L^\infty(Y) \rtimes \Lambda$ , for any free, ergodic, p.m.p. action  $\Lambda \curvearrowright (Y, \nu)$ , then we can find a unitary  $u \in M$  such that  $uL^\infty(X)u^* = L^\infty(Y)$ .*

Recall that a p.m.p. action  $\Gamma \curvearrowright (X, \mu)$  is called *compact* if we can decompose  $L^2(X) = \bigoplus_n \mathcal{H}_n$ , where  $\{\mathcal{H}_n\}_n$  is a sequence of  $\Gamma$ -invariant, finite dimensional subspaces of  $L^2(X)$ . For instance, if  $G$  is a compact group containing a dense copy of  $\Gamma$  and  $m_G$  denotes the Haar measure of  $G$ , then the left multiplication action  $\Gamma \curvearrowright (G, m_G)$  is a free, ergodic, p.m.p. compact action.

The condition  $\beta_1^{(2)}(\Gamma) > 0$  is equivalent to  $\Gamma$  being non-amenable and having an unbounded cocycle into its regular representation  $\lambda : \Gamma \rightarrow \mathcal{U}(\ell^2\Gamma)$  [PT07], and is satisfied by any free product group  $\Gamma = \Gamma_1 * \Gamma_2$  with  $|\Gamma_1| \geq 2$  and  $|\Gamma_2| \geq 3$ .

Note that in the case  $\Gamma$  does not have Haagerup's property, Theorem 1.1 is a consequence of the main result of [Pe09].

A far-reaching conjecture predicts that  $\text{II}_1$  factors  $M = L^\infty(X) \rtimes \Gamma$  arising from free, ergodic, p.m.p. actions of groups  $\Gamma$  with  $\beta_1^{(2)}(\Gamma) > 0$  have a unique Cartan subalgebra. Recently, there has been some progress on the ‘‘group measure space’’ version of the conjecture. Firstly, it has been shown in [PV09] that if  $\Gamma = \Gamma_1 * \Gamma_2$ , where  $\Gamma_1$  is an infinite property (T) group and  $\Gamma_2$  is a non-trivial group, then  $M$  has a unique group measure space Cartan subalgebra. More generally, the same has been proven in [CP10] (see also [Va10b]) under the assumption that  $\Gamma$  admits a non-amenable subgroup with the relative property (T). Lastly, we proved in [Io11] that  $M$  also has a unique group measure space Cartan subalgebra if the action  $\Gamma \curvearrowright (X, \mu)$  is rigid.

As mentioned above, this conjecture has been very recently established in full generality for arbitrary actions of weakly amenable groups  $\Gamma$  with  $\beta_1^{(2)}(\Gamma) > 0$  [PV11a].

Theorem 1.1 provides further positive evidence towards this conjecture. It implies that any residually finite group  $\Gamma$  satisfying  $\beta_1^{(2)}(\Gamma) > 0$  admits at least one action whose  $\text{II}_1$  factor has a unique group measure space decomposition. Indeed, if  $\{\Gamma_n\}_n$  is a descending chain of normal, finite index subgroups of  $\Gamma$  with trivial intersection, then the left multiplication action  $\Gamma \curvearrowright (G, m_G)$  on the *profinite completion*  $G = \varprojlim \Gamma/\Gamma_n$  is a free, ergodic, compact p.m.p. action.

To outline the proof of Theorem 1.1, denote  $A = L^\infty(X)$  and  $M = A \rtimes \Gamma$ . Consider another group measure space decomposition  $M = B \rtimes \Lambda$  and let  $\Delta : M \rightarrow M \overline{\otimes} M$  be the associated *comultiplication* [PV09]. In the first part of the proof, we show that a corner of  $\Delta(A)$  embeds into  $M \overline{\otimes} A$ , in the sense of [Po03]. To achieve this, we combine the fact that the action  $\Delta(\Gamma) \curvearrowright \Delta(A)$  is compact (hence, a fortiori, weakly compact) with a result from [OP07] relating weak compactness of actions to uniform convergence of deformations, and results of [CP10] on malleable deformations coming from cocycles into the regular representation (see Section 4). In the second part of the proof, we use techniques from [Io11] to conclude that if  $\Gamma$  has positive first  $\ell^2$ -Betti number, then the condition  $\Delta(A) \prec M \overline{\otimes} A$  automatically implies that  $A$  and  $B$  are conjugate (see Section 3).

## 2. PRELIMINARIES

Throughout the paper we work with *tracial von Neumann algebras*  $(M, \tau)$ , i.e. von Neumann algebras  $M$  endowed with a faithful normal tracial state  $\tau$ , and assume that all von Neumann algebras have separable predual.

**2.1. Intertwining-by-bimodules.** We first recall from [Po03, Theorem 2.1 and Corollary 2.3] S. Popa's powerful *intertwining-by-bimodules* technique.

**Theorem 2.1.** [Po03] *Let  $(M, \tau)$  be a tracial von Neumann algebra and  $P, Q \subset M$  be von Neumann subalgebras. Then the following are equivalent:*

- *There exist non-zero projections  $p \in P, q \in Q$ , a  $*$ -homomorphism  $\phi : pPp \rightarrow qQq$  and a non-zero partial isometry  $v \in qMp$  such that  $\phi(x)v = vx$ , for all  $x \in pPp$ .*
- *There is no sequence  $u_n \in \mathcal{U}(P)$  satisfying  $\|E_Q(xu_ny)\|_2 \rightarrow 0$ , for all  $x, y \in M$ .*

*If one of these conditions holds true, then we say that a corner of  $P$  embeds into  $Q$  inside  $M$  and write  $P \prec_M Q$ .*

**2.2. Relative amenability.** Let  $(M, \tau)$  be a tracial von Neumann algebra. Recall that  $M$  is said to be *amenable* if we can find a sequence  $\xi_n \in L^2(M) \overline{\otimes} L^2(M)$  such that  $\langle x\xi_n, \xi_n \rangle \rightarrow \tau(x)$  and  $\|[x, \xi_n]\|_2 \rightarrow 0$ , for every  $x \in M$ . By A. Connes' celebrated theorem [Co76] this is equivalent to  $M$  being *approximately finite dimensional*.

Now, let  $Q \subset M$  be a von Neumann subalgebra. *Jones' basic construction*  $\langle M, e_Q \rangle$  is defined as the von Neumann subalgebra of  $\mathbb{B}(L^2(M))$  generated by  $M$  and the orthogonal projection  $e_Q$  from  $L^2(M)$  onto  $L^2(Q)$ . Recall that  $\langle M, e_Q \rangle$  is equipped with a faithful semi-finite trace given by  $Tr(xe_Qy) = \tau(xy)$  for all  $x, y \in M$ . We denote by  $L^2(\langle M, e_Q \rangle)$  the associated Hilbert space and endow it with the natural  $M$ -bimodule structure.

Following [OP07, Definition 2.2] we say that a (not necessarily unital) von Neumann subalgebra  $P \subset M$  is *amenable relative to  $Q$  inside  $M$*  if there exists a sequence  $\xi_n \in L^2(\langle M, e_Q \rangle)$  such that  $\langle x\xi_n, \xi_n \rangle \rightarrow \tau(x)$ , for every  $x \in M$ , and  $\|[y, \xi_n]\|_2 \rightarrow 0$ , for every  $y \in P$ .

**Lemma 2.2.** [IPV10] *Let  $(B, \tau)$  be a tracial von Neumann algebra and  $\Lambda \curvearrowright (B, \tau)$  be a trace preserving action. Denote  $M = B \rtimes \Lambda$ . Define the  $*$ -homomorphism  $\Delta : M \rightarrow M \overline{\otimes} M$  by letting  $\Delta(b) = b \otimes 1$ , for all  $b \in B$ , and  $\Delta(v_s) = v_s \otimes v_s$ , for every  $s \in \Lambda$ . Let  $P, Q \subset M$  be von Neumann subalgebras such that  $P$  has no amenable direct summand and  $Q$  is amenable.*

*Then there is no non-zero projection  $p \in \Delta(P)' \cap M \overline{\otimes} M$  such that the von Neumann algebra  $\Delta(P)p$  is amenable relative to  $M \overline{\otimes} Q$ .*

*Proof.* Since  $B$  and  $Q$  are amenable, the  $M$ - $M \overline{\otimes} M$ -bimodule  ${}_{\Delta(M)}L^2(\langle M \overline{\otimes} M, e_{M \overline{\otimes} Q} \rangle)_{M \overline{\otimes} M}$  is weakly contained in the coarse  $M$ - $M \overline{\otimes} M$ -bimodule,  $L^2(M) \overline{\otimes} L^2(M \overline{\otimes} M)$ . Repeating the proof of [IPV10, Proposition 7.2.4] now gives the conclusion. ■

**2.3. Weakly compact actions.** We next recall N. Ozawa and S. Popa's notion of weakly compact actions. Let  $(P, \tau)$  be a tracial von Neumann algebra and denote by  $\bar{P} = \{\bar{x} | x \in P\}$  the complex conjugate von Neumann algebra.

**Definition 2.3.** [OP07, Definition 3.1]. A trace preserving action  $\Gamma \curvearrowright^\sigma (P, \tau)$  is *weakly compact* if we can find a sequence of unit vectors  $\eta_n \in L^2(P \overline{\otimes} \bar{P})_+$  such that

- $\|\eta_n - (v \otimes \bar{v})\eta_n\|_2 \rightarrow 0$ , for every  $v \in \mathcal{U}(P)$ .
- $\|\eta_n - (\sigma_g \otimes \bar{\sigma}_g)(\eta_n)\|_2 \rightarrow 0$ , for every  $g \in \Gamma$ .

- $\langle (x \otimes 1)\eta_n, \eta_n \rangle = \tau(x) = \langle \eta_n, (1 \otimes \bar{x})\eta_n \rangle$ , for every  $x \in P$  and every  $n$ .

Note that these conditions force  $P$  to be amenable. On the other hand, by [OP07, Proposition 3.2], if  $P$  is amenable and  $\Gamma \curvearrowright^\sigma P$  is a *compact* action (i.e. if  $L^2(P)$  is the direct sum of finite dimensional  $\sigma(\Gamma)$ -invariant subspaces) then  $\sigma$  is weakly compact.

N. Ozawa and S. Popa showed in [OP07, Theorem 4.9] that weakly compact actions can be used to deduce relative amenability. An obvious modification of their proof (in the case  $p = 1$ ) gives the following:

**Theorem 2.4.** [OP07, Theorem 4.9] *Let  $(M, \tau)$  be a tracial von Neumann algebra and  $Q_1, \dots, Q_k \subset M$  be von Neumann subalgebras. Assume that there are a tracial von Neumann algebra  $\tilde{M} \supset M$  and automorphisms  $\{\alpha_t\}_{t \in \mathbb{R}}$  of  $\tilde{M}$  such that*

- $\lim_{t \rightarrow 0} \|\alpha_t(x) - x\|_2 = 0$ , for all  $x \in M$ , and
- $L^2(\tilde{M}) \ominus L^2(M)$  is isomorphic as an  $M$ -bimodule to a submodule of a multiple of  $\bigoplus_{j=1}^k L^2(\langle M, e_{Q_j} \rangle)$ .

Let  $P \subset M$  be a von Neumann subalgebra and define  $c_t := \inf_{v \in \mathcal{U}(P)} \|E_M(\alpha_t(v))\|_2$ , for  $t \in \mathbb{R}$ . Let  $\mathcal{G} \subset \mathcal{N}_M(P)$  be a subgroup which acts weakly compactly on  $P$  by conjugation and denote  $N = \mathcal{G}''$ .

Then either  $c_t \rightarrow 1$ , as  $t \rightarrow 0$ , or there exist  $j \in \{1, \dots, k\}$  and a non-zero projection  $p_j \in N' \cap M$  such that  $Np_j$  is amenable relative to  $Q_j$ .

**2.4. Deformations coming from group cocycles.** Let  $\Gamma \curvearrowright (A, \tau)$  be a trace preserving action and set  $M = A \rtimes \Gamma$ . Let  $\pi : \Gamma \rightarrow \mathcal{O}(H_{\mathbb{R}})$  be an orthogonal representation, where  $H_{\mathbb{R}}$  is a separable real Hilbert space. Also, let  $b : \Gamma \rightarrow H_{\mathbb{R}}$  be a *cocycle*, i.e. a map satisfying the identity  $b(gh) = b(g) + \pi(g)b(h)$ , for all  $g, h \in \Gamma$ .

From this data, T. Sinclair constructed a *malleable deformation* in the sense of S. Popa, i.e. a tracial von Neumann algebra  $\tilde{M} \supset M$  and a 1-parameter group of automorphisms  $\{\alpha_t\}_{t \in \mathbb{R}}$  of  $\tilde{M}$  such that  $\|\alpha_t(x) - x\|_2 \rightarrow 0$ , for all  $x \in \tilde{M}$  (see [Si10, Section 3] and [Va10b, Section 3.1]).

To recall this construction, let  $(D, \tau_0)$  be the unique tracial von Neumann algebra generated by unitary elements  $\omega(\xi)$ ,  $\xi \in H_{\mathbb{R}}$ , subject to the relations  $\omega(\xi + \eta) = \omega(\xi)\omega(\eta)$ ,  $\omega(\xi)^* = \omega(-\xi)$  and  $\tau_0(\omega(\xi)) = \exp(-\|\xi\|^2)$ , for all  $\xi, \eta \in H_{\mathbb{R}}$ . Consider the Gaussian action  $\Gamma \curvearrowright^\sigma D$  which on the generating functions  $\omega(\xi)$  is given by  $\sigma_g(\omega(\xi)) = \omega(\pi(g)(\xi))$ . Finally, let  $\Gamma \curvearrowright D \overline{\otimes} A$  be the diagonal action and define  $\tilde{M} = (D \overline{\otimes} A) \rtimes \Gamma$ .

Then the formula

$$\alpha_t(u_g) = (\omega(tb(g)) \otimes 1)u_g, \text{ for all } g \in \Gamma, \text{ and } \alpha_t(x) = x, \text{ for all } x \in D \overline{\otimes} A$$

gives the desired 1-parameter group of automorphisms  $\{\alpha_t\}_{t \in \mathbb{R}}$  of  $\tilde{M}$ .

Moreover, the formula  $\beta(\omega(\xi)) = \omega(-\xi)$ , for all  $\xi \in H_{\mathbb{R}}$ , and  $\beta(x) = x$ , for all  $x \in M$ , defines an automorphism of  $\tilde{M}$  which satisfies  $\beta^2 = id$  and  $\beta \circ \alpha_t = \alpha_{-t} \circ \beta$ , for all  $t \in \mathbb{R}$ . As shown in [Po06a, Lemma 2.1] the existence of such a  $\beta$  implies a “transversality” property for  $\alpha_t$ :

**Lemma 2.5.** [Po06a] *For every  $x \in M$  and  $t \in \mathbb{R}$  we have that*

$$\|\alpha_{2t}(x) - x\|_2 \leq 2\|\alpha_t(x) - E_M(\alpha_t(x))\|_2.$$

Next we record a direct consequence of S. Popa’s *spectral gap principle* [Po06b, Lemma 1.5].

**Lemma 2.6.** [Po06b] *Assume that  $\pi$  is weakly contained in the regular representation of  $\Gamma$ . Let  $B \subset M$  be a von Neumann subalgebra with no amenable direct summand.*

*Then  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(B' \cap M)_1$  (the unit ball of the commutant of  $B$ ).*

For a proof, see e.g. [Io11, Lemma 2.2].

Now, let  $B \subset M$  be a von Neumann subalgebra. J. Peterson [Pe06, Theorem 4.5] and I. Chifan and J. Peterson [CP10, Theorem 2.5] proved that if  $\alpha_t \rightarrow id$  uniformly on  $(B)_1$  and  $B \not\prec_M A$  then  $\alpha_t \rightarrow id$  uniformly on  $\mathcal{N}_M(B)$ .

**Theorem 2.7.** [Pe06], [CP10] *Assume that  $\pi$  is mixing. Let  $B \subset M$  be a von Neumann subalgebra. Suppose that  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(B)_1$  and that  $B \not\prec_M A$ . Denote  $P = \mathcal{N}_M(B)''$ .*

*Then  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(P)_1$ .*

Conversely, I. Chifan and J. Peterson proved in [CP10, Theorem 3.2] that if  $B$  is abelian and  $\alpha_t \rightarrow id$  uniformly on a “large” sequence  $u_n \in \mathcal{N}_M(B)$ , then  $\alpha_t \rightarrow id$  uniformly on  $(B)_1$ .

**Theorem 2.8.** [CP10] *Assume that  $\pi$  is mixing. Let  $B \subset M$  be an abelian von Neumann subalgebra. Assume that we can find a sequence  $u_n \in \mathcal{N}_M(B)$  such that*

- $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $\{u_n\}_{n \geq 1}$  and
- $\|E_A(xu_ny)\|_2 \rightarrow 0$ , for all  $x, y \in M$ .

*Then  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(B)_1$ .*

Theorems 2.7 and 2.8 have been proven in [Pe06] and [CP10] using J. Peterson’s technique of unbounded derivations [Pe06]. For proofs using the automorphisms  $\{\alpha_t\}_{t \in \mathbb{R}}$ , see S. Vaes’s paper [Va10b, Theorems 3.9 and 4.1].

### 3. CONJUGACY OF GROUP MEASURE SPACE CARTAN SUBALGEBRAS

In this section we assemble together the main technical results of [Io11] to prove a new conjugacy criterion for group measure space Cartan subalgebras of  $\text{II}_1$  factors  $L^\infty(X) \rtimes \Gamma$  arising from actions of groups with positive first  $\ell^2$ -Betti number.

**Theorem 3.1.** *Let  $\Gamma$  be a countable group satisfying  $\beta_1^{(2)}(\Gamma) > 0$ . Let  $\Gamma \curvearrowright (X, \mu)$  be a free ergodic p.m.p. action and set  $M = L^\infty(X) \rtimes \Gamma$ . Assume that  $M = L^\infty(Y) \rtimes \Lambda$  for a free ergodic p.m.p. action  $\Lambda \curvearrowright (Y, \nu)$ . Let  $\Delta : M \rightarrow M \overline{\otimes} M$  be the  $*$ -homomorphism given by  $\Delta(b) = b \otimes 1$  for all  $b \in L^\infty(Y)$  and  $\Delta(v_g) = v_g \otimes v_g$  for all  $g \in \Lambda$ , where  $\{v_g\}_{g \in \Lambda} \subset M$  denote the canonical unitaries. If  $\Delta(L^\infty(X)) \prec_{M \overline{\otimes} M} M \overline{\otimes} L^\infty(X)$ , then  $L^\infty(X)$  and  $L^\infty(Y)$  are unitarily conjugate.*

*Proof.* Denote  $A = L^\infty(X)$  and  $B = L^\infty(Y)$ . Since  $\beta_1^{(2)}(\Gamma) > 0$ , by [Io11, Theorem 4.2], in order to get the conclusion, it suffices to find an amenable subgroup  $\Lambda_0 < \Lambda$  such that  $A \prec_M B \rtimes \Lambda_0$ . Let us assume by contradiction that this is false.

Since  $\Delta(A) \prec M \overline{\otimes} A$ , the second condition in Theorem 2.1 holds true. Thus, we can find a finite set  $F \subset (M \overline{\otimes} M)_1$  and  $\delta > 0$  such that  $\sum_{x \in F} \|E_{M \overline{\otimes} A}(\Delta(a)x)\|_2^2 \geq 2\delta$ , for every  $a \in \mathcal{U}(A)$ . By Kaplansky’s theorem we may assume that  $F \subset 1 \otimes (M)_1$ , i.e. we have

$$(3.1) \quad \sum_{x \in F} \|E_{M \overline{\otimes} A}(\Delta(a)(1 \otimes x))\|_2^2 \geq 2\delta, \quad \forall a \in \mathcal{U}(A)$$

For  $g \in \Lambda$ , we define  $f(g) = \sum_{x \in F} \|E_A(v_g x)\|_2^2$ . Then we have the following:

**Claim.** Let  $g_1, \dots, g_m, h_1, \dots, h_m \in \Lambda$  and  $\Lambda_1, \dots, \Lambda_m < \Lambda$  be amenable subgroups. Put  $S = \cup_{i=1}^m g_i \Lambda_i h_i$ . Then there exists  $g \in \Lambda \setminus S$  such that  $f(g) \geq \delta$ .

*Proof of the claim.* Since  $\Lambda_i$  is amenable by our assumption we have that  $A \not\prec_M B \rtimes \Lambda_i$ , for all  $i \in \{1, \dots, m\}$ . It follows from S. Popa's intertwining-by-bimodules technique (see Theorem 2.1 and e.g. [Io11, Remark 1.2]) that we can find  $a \in \mathcal{U}(A)$  such that

$$(3.2) \quad \sum_{i=1}^m \|E_{B \rtimes \Lambda_i}(v_{g_i}^* a v_{h_i}^*)\|_2^2 \leq \frac{\delta}{|F|}$$

If we write  $a = \sum_{g \in \Lambda} b_g v_g$ , where  $b_g \in B$ , then 3.2 implies that  $\sum_{g \in S} \|b_g\|_2^2 \leq \frac{\delta}{|F|}$ . On the other hand, since  $E_{M \overline{\otimes} A}(\Delta(a)(1 \otimes x)) = \sum_{g \in \Lambda} b_g v_g \otimes E_A(v_g x)$ , for every  $x \in M$ , 3.1 can be rewritten as  $\sum_{g \in \Lambda} f(g) \|b_g\|_2^2 \geq 2\delta$ . Since  $f(g) \leq |F|$ , for all  $g \in \Lambda$ , combining the last two inequalities yields that  $\sum_{g \in \Lambda \setminus S} f(g) \|b_g\|_2^2 \geq \delta$ . Finally, since  $\sum_{g \in \Lambda \setminus S} \|b_g\|_2^2 \leq \|a\|_2^2 = 1$ , the claim follows.  $\square$

The claim shows that the conclusion of [Io11, Lemma 3.2] holds true. The second part of the proof of [Io11, Theorem 3.1], which only uses the conclusion of [Io11, Lemma 3.2], provides a decreasing sequence  $\{\Lambda_k\}_{k \geq 1}$  of non-amenable subgroups of  $\Lambda$  such that  $A \prec_M B \rtimes (\cup_{k \geq 1} C(\Lambda_k))$ , where  $C(\Lambda_k)$  denotes the centralizer of  $\Lambda_k$  in  $\Lambda$ .

Since  $\beta_1^{(2)}(\Gamma) > 0$ , there is an unbounded cocycle  $b : \Gamma \rightarrow \ell_{\mathbb{R}}^2 \Gamma$  for the regular representation [PT07]. Let  $\tilde{M} \supset M$  and  $\{\alpha_t\}_{t \in \mathbb{R}} \subset \text{Aut}(\tilde{M})$  be the malleable deformation constructed in Section 2.4.

Now, the group  $\cup_{k \geq 1} C(\Lambda_k)$  must be non-amenable by our assumption. Thus,  $C(\Lambda_k)$  is non-amenable for some  $k \geq 1$  and therefore its von Neumann algebra,  $L(C(\Lambda_k))$ , has no amenable direct summand. Lemma 2.6 implies that  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(L(\Lambda_k))_1$ .

Since  $\Lambda_k$  is non-amenable and  $A$  is abelian, we have that  $L(\Lambda_k) \not\prec_M A$ . By [Po03, Corollary 2.3] can find a sequence  $g_n \in \Lambda_k$  such that  $\|E_A(x v_{g_n} y)\|_2 \rightarrow 0$ , for all  $x, y \in M$ . Because  $v_{g_n} \in \mathcal{N}_M(B)$  and  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $\{v_{g_n}\}_{n \geq 1}$ , applying Theorem 2.8 gives that  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(B)_1$ .

Finally, if  $B \prec_M A$ , then by [Po01, Theorem A.1] we get that  $A$  and  $B$  are conjugate. Otherwise, if  $B \not\prec_M A$ , then Theorem 2.7 implies that  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(M)_1$ . In particular, we can find  $t > 0$  such that  $\Re \tau(\alpha_t(u_g) u_g^*) \geq \exp(-\frac{1}{4})$ , for all  $g \in \Gamma$ . Since  $\tau(\alpha_t(u_g) u_g) = \exp(-t^2 \|b(g)\|^2)$  we derive that  $\|b(g)\| \leq \frac{1}{2t}$  for all  $g \in \Gamma$ . This shows that  $b$  is bounded, a contradiction.  $\blacksquare$

#### 4. PROOF OF THEOREM 1.1

Let  $\Gamma \curvearrowright (X, \mu)$  be a free ergodic p.m.p. compact action. Denote  $A = L^\infty(X)$  and  $M = A \rtimes \Gamma$ . Assume that  $\Gamma$  has positive first  $\ell^2$ -Betti number and let  $b : \Gamma \rightarrow \ell_{\mathbb{R}}^2 \Gamma$  be an unbounded cocycle into the regular representation of  $\Gamma$  [PT07].

Let  $\Lambda \curvearrowright (Y, \nu)$  be a free ergodic p.m.p. action such that  $M = B \rtimes \Lambda$ , where  $B = L^\infty(Y)$ . Let  $\{v_s\}_{s \in \Lambda} \subset M$  be the canonical unitaries. Let  $\Delta : M \rightarrow M \overline{\otimes} M$  be the  $*$ -homomorphism given by  $\Delta(b) = b \otimes 1$  for  $b \in B$  and  $\Delta(v_s) = v_s \otimes v_s$  for  $s \in \Lambda$ .

Our goal is to show that  $A$  and  $B$  are unitarily conjugate. If we denote  $\mathcal{A} = M \overline{\otimes} A$  and  $\mathcal{M} = M \overline{\otimes} M$  then by Theorem 3.1 it suffices to prove that  $\Delta(A) \prec_{\mathcal{M}} \mathcal{A}$ . Assume this to be false.

Towards a contradiction, let  $\tilde{M} \supset M$  and  $\{\alpha_t\}_{t \in \mathbb{R}} \subset \text{Aut}(\tilde{M})$  be the malleable deformation defined in Section 2.4. We let  $\tilde{\mathcal{M}} = M \overline{\otimes} \tilde{M}$  and denote by  $\tilde{\alpha}_t$  the automorphism  $id \otimes \alpha_t$  of  $\tilde{\mathcal{M}}$  for every  $t \in \mathbb{R}$ . Note that  $\tilde{\mathcal{M}} \supset \mathcal{M}$  and  $\{\tilde{\alpha}_t\}_{t \in \mathbb{R}} \subset \text{Aut}(\tilde{\mathcal{M}})$  is precisely the malleable deformation associated to the cocycle  $b$  and the obvious crossed product decomposition  $\mathcal{M} = \mathcal{A} \rtimes \Gamma$ .

Since  $b$  is a cocycle for the regular representation, it is easy to see that the  $\mathcal{M}$ -bimodule  $L^2(\tilde{\mathcal{M}}) \ominus L^2(\mathcal{M})$  is isomorphic to  $\bigoplus_{i=1}^{\infty} L^2(\langle \mathcal{M}, e_A \rangle)$  (see [Va10b, Lemma 3.5]). Since the action  $\Gamma \curvearrowright A$  is compact, the action  $\Delta(\Gamma) \curvearrowright \Delta(A)$  is compact and thus weakly compact. By applying Theorem 2.4 to  $P = \Delta(A)$  and  $N = \Delta(M)$  we deduce that either (1)  $\Delta(M)p$  is amenable relative to  $\mathcal{A}$  for some non-zero projection  $p \in \Delta(M)' \cap \mathcal{M}$  or (2)  $\inf_{v \in \mathcal{U}(\Delta(A))} \|E_{\mathcal{M}}(\tilde{\alpha}_t(v))\|_2 \rightarrow 1$  as  $t \rightarrow 0$ .

Since  $M$  has no amenable direct summand, the first case is ruled out by Lemma 2.2.

In the second case, we get that  $\sup_{v \in \mathcal{U}(\Delta(A))} \|\tilde{\alpha}_t(v) - E_{\mathcal{M}}(\tilde{\alpha}_t(v))\|_2 \rightarrow 0$  as  $t \rightarrow 0$ . By Lemma 2.5 we deduce that  $\tilde{\alpha}_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(\Delta(A))_1$ . Since  $\Delta(A) \not\curvearrowright_{\mathcal{M}} \mathcal{A}$  by assumption and the von Neumann algebra generated by the normalizer of  $\Delta(A)$  in  $\mathcal{M}$  contains  $\Delta(M)$ , Theorem 2.7 gives that  $\tilde{\alpha}_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $(\Delta(M))_1$ .

Since  $\tilde{\alpha}_t(\Delta(v_s)) = v_s \otimes \alpha_t(v_s)$  for  $s \in \Lambda$  we derive that  $\alpha_t \rightarrow id$  in  $\|\cdot\|_2$  uniformly on  $\{v_s\}_{s \in \Lambda}$ . Since  $\Lambda$  is non-amenable and  $A$  is abelian we can find a sequence  $\{s_n\}_{n \geq 1}$  satisfying  $\|E_A(xv_{s_n}y)\|_2 \rightarrow 0$  for all  $x, y \in M$ . Continuing exactly as in the end of the proof of Theorem 3.1 provides the desired contradiction.  $\blacksquare$

**Remark 4.1.** It is easy to adapt the proof of the main result to show that if  $t > 0$  and  $M^t = L^\infty(Y) \rtimes \Lambda$ , for some free ergodic p.m.p. action  $\Lambda \curvearrowright (Y, \nu)$ , then  $L^\infty(Y)$  is unitarily conjugate to  $L^\infty(X)^t$ . It follows from D. Gaboriau's work on  $\ell^2$ -Betti numbers for equivalence relations [Ga01] that  $M$  has trivial fundamental group,  $\mathcal{F}(M) = \{1\}$ .

**Remark 4.2.** Let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integers, for some prime  $p$ . Consider the profinite action  $\Gamma := \mathbb{Z}^2 \rtimes \mathrm{SL}_2(\mathbb{Z}) \curvearrowright \varprojlim \mathbb{Z}_p^2$  and denote by  $M$  the associated  $\mathrm{II}_1$  factor. By [OP08, Corollary D] and [PV09, Section 5.5],  $M$  has two non-conjugate group measure space Cartan subalgebras,  $L^\infty(\varprojlim \mathbb{Z}_p^2)$  and  $L(\mathbb{Z}^2)$ . This shows that the assumption that  $\beta_1^{(2)}(\Gamma) > 0$  from the hypothesis of our main result cannot be replaced by the assumption that  $\Gamma$  is merely non-amenable.

## REFERENCES

- [Co76] A. Connes: *Classification of injective factors*, Ann. of Math. (2) **104** (1976), 73–115.
- [CP10] I. Chifan, J. Peterson: *Some unique group-measure space decomposition results*, preprint arXiv:1010.5194.
- [CS11] I. Chifan, T. Sinclair: *On the structural theory of  $\mathrm{II}_1$  factors of negatively curved groups*, preprint arXiv:1103.4299.
- [CSU11] I. Chifan, T. Sinclair, B. Udrea: *On the structural theory of  $\mathrm{II}_1$  factors of negatively curved groups, II. Actions by product groups*, preprint arXiv:1108.4200.
- [Da10] Y. Dabrowski: *A non-commutative Path Space approach to stationary free Stochastic Differential Equations*, preprint arXiv:1006.4351.
- [FM77] J. Feldman, C.C. Moore: *Ergodic equivalence relations, cohomology, and von Neumann algebras, II*, Trans. Amer. Math. Soc. **234** (1977), 325–359.
- [FV10] P. Fima, S. Vaes: *HNN extensions and unique group measure space decomposition of  $\mathrm{II}_1$  factors*, preprint arXiv:1005.5002, to appear in Trans. Amer. Math. Soc.
- [Fu09] A. Furman: *A survey of Measured Group Theory*, Geometry, Rigidity, and Group Actions, 296–374, The University of Chicago Press, Chicago and London, 2011.
- [Ga01] D. Gaboriau: *Invariants  $\ell^2$  de relations d'équivalence et de groupes*, Publ. Math. Inst. Hautes Études Sci., **95** (2002), 93–150.
- [Ga10] D. Gaboriau: *Orbit Equivalence and Measured Group Theory*, Proceedings of the ICM (Hyderabad, India, 2010), Vol. III, Hindustan Book Agency (2010), 1501–1527.
- [HPV10] C. Houdayer, S. Popa, S. Vaes: *A class of groups for which every action is  $W^*$ -superrigid*, preprint arXiv:1010.5077, to appear in Groups Geom. Dyn.
- [Io10] A. Ioana:  *$W^*$ -superrigidity for Bernoulli actions of property (T) groups*, J. Amer. Math. Soc. **24** (2011), 1175–1226.
- [Io11] A. Ioana: *Uniqueness of the group measure space decomposition for Popa's  $\mathcal{HT}$  factors*, preprint arXiv:1104.2913.

- [IPV10] A. Ioana, S. Popa, S. Vaes: *A class of superrigid group von Neumann algebras*, preprint arXiv:1007.1412.
- [MvN36] F. Murray, J. von Neumann: *On rings of operators*, Ann. of Math. **37** (1936), 116–229.
- [Oz07] N. Ozawa: *Weak amenability of hyperbolic groups*, Groups Geom. Dyn., **2** (2008), 271–280.
- [Oz10] N. Ozawa: *Examples of groups which are not weakly amenable*, preprint arXiv:1012.0613, to appear in Kyoto J. Math.
- [OP07] N. Ozawa, S. Popa: *On a class of  $II_1$  factors with at most one Cartan subalgebra*, Ann. of Math. (2), **172** (2010), 713–749.
- [OP08] N. Ozawa, S. Popa: *On a class of  $II_1$  factors with at most one Cartan subalgebra, II*, Amer. J. Math., **132** (2010), 841–866.
- [Pe06] J. Peterson:  *$L^2$ -rigidity in von Neumann algebras*, Invent. Math. **175** (2009), no. 2, 417–433.
- [Pe09] J. Peterson: *Examples of group actions which are virtually  $W^*$ -superrigid*, preprint arXiv:1002.1745.
- [Po01] S. Popa: *On a class of type  $II_1$  factors with Betti numbers invariants*, Ann. of Math. **163** (2006), 809–889.
- [Po03] S. Popa: *Strong Rigidity of  $II_1$  Factors Arising from Malleable Actions of  $w$ -Rigid Groups. I.*, Invent. Math. **165** (2006), 369–408.
- [Po06a] S. Popa: *On the superrigidity of malleable actions with spectral gap*, J. Amer. Math. Soc. **21** (2008), 981–1000.
- [Po06b] S. Popa: *On Ozawa’s Property for Free Group Factors*, Int. Math. Res. Notices (2007) Vol. 2007, article ID rnm036.
- [Po07] S. Popa: *Deformation and rigidity for group actions and von Neumann algebras*, International Congress of Mathematicians. Vol. I, 445–477, Eur. Math. Soc., Zürich, 2007.
- [PV09] S. Popa, S. Vaes: *Group measure space decomposition of  $II_1$  factors and  $W^*$ -superrigidity*, Invent. Math. **182** (2010), no. 2, 371–417.
- [PV11a] S. Popa, S. Vaes: *Unique Cartan decomposition for  $II_1$  factors arising from arbitrary actions of free groups*, preprint arXiv:1111.6951.
- [PV11b] S. Popa, S. Vaes: *Unique Cartan decomposition for  $II_1$  factors arising from arbitrary actions of hyperbolic groups*, in preparation.
- [PT07] J. Peterson, A. Thom: *Group cocycles and the ring of affiliated operators*, Invent. Math., **185** (2011) 561–592.
- [Si55] I.M. Singer: *Automorphisms of finite factors*, Amer. J. Math. **77** (1955), 117–133.
- [Si10] T. Sinclair: *Strong solidity of group factors from lattices in  $SO(n,1)$  and  $SU(n,1)$* , J. Funct. Anal. **260** (2011), 3209–3221.
- [Va10a] S. Vaes: *Rigidity for von Neumann algebras and their invariants*, Proceedings of the ICM (Hyderabad, India, 2010), Vol. III, Hindustan Book Agency (2010), 1624–1650.
- [Va10b] S. Vaes: *One-cohomology and the uniqueness of the group measure space decomposition of a  $II_1$  factor*, preprint arXiv:1012.5377.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, LA JOLLA, CA 92093-0112, USA

*E-mail address:* aioana@ucsd.edu