# A MURRAY-VON NEUMANN TYPE CLASSIFICATION OF $C^*$ -ALGEBRAS

#### CHI-KEUNG NG AND NGAI-CHING WONG

ABSTRACT. We define type  $\mathfrak{A}$ , type  $\mathfrak{C}$  as well as  $C^*$ -semi-finite  $C^*$ -algebras.

It is shown that a von Neumann algebra is a type  $\mathfrak{A}$ , type  $\mathfrak{B}$ , type  $\mathfrak{C}$  or  $C^*$ -semi-finite  $C^*$ -algebra if and only if it is, respectively, a type I, type II, type III or semi-finite von Neumann algebra. Any type I  $C^*$ -algebra is of type  $\mathfrak{A}$  (actually, type  $\mathfrak{A}$  coincides with the discreteness as defined by Peligrad and Zsidó), and any type II  $C^*$ -algebra (as defined by Cuntz and Pedersen) is of type  $\mathfrak{B}$ . Moreover, any type  $\mathfrak{C}$   $C^*$ -algebra is of type III (in the sense of Cuntz and Pedersen). Conversely, any separable purely infinite  $C^*$ -algebra (in the sense of Kirchberg and Rørdam) with either real rank zero or stable rank one is of type  $\mathfrak{C}$ .

We also prove that type  $\mathfrak{A}$ , type  $\mathfrak{C}$  and  $C^*$ -semi-finiteness are stable under taking hereditary  $C^*$ -subalgebras, multiplier algebras and strong Morita equivalence. Furthermore, any  $C^*$ -algebra A contains a largest type  $\mathfrak{A}$  closed ideal  $J_{\mathfrak{A}}$ , a largest type  $\mathfrak{B}$  closed ideal  $J_{\mathfrak{B}}$ , a largest type  $\mathfrak{C}$  closed ideal  $J_{\mathfrak{C}}$  as well as a largest  $C^*$ -semi-finite closed ideal  $J_{\mathfrak{S}\mathfrak{f}}$ . Among them, we have  $J_{\mathfrak{A}} + J_{\mathfrak{B}}$  being an essential ideal of  $J_{\mathfrak{S}\mathfrak{f}}$ , and  $J_{\mathfrak{A}} + J_{\mathfrak{B}} + J_{\mathfrak{C}}$  being an essential ideal of A. On the other hand,  $A/J_{\mathfrak{C}}$  is always  $C^*$ -semi-finite, and if A is  $C^*$ -semi-finite, then  $A/J_{\mathfrak{B}}$  is of type  $\mathfrak{A}$ .

This paper is dedicated to Charles Batty on the occasion of his 60th birthday.

#### 1. Introduction

In their seminal works ([27], see also [26]), Murray and von Neumann defined three types of von Neumann algebras (namely, type I, type II and type III) according to the properties of their projections. They showed that any von Neumann algebra is a sum of a type I, a type II, and a type III von Neumann subalgebras. This classification was shown to be very important and becomes the basic theory for the study of von Neumann algebras (see, e.g., [20]). Since a  $C^*$ -algebra needs not have any projection, a similar classification for  $C^*$ -algebras seems impossible. There is, however, an interesting classification scheme for  $C^*$ -algebras proposed by

Date: October 26, 2018.

<sup>2000</sup> Mathematics Subject Classification. 46L05, 46L35.

Key words and phrases. C\*-algebra; open projections; Murray-von Neumann type classification.

The authors are supported by National Natural Science Foundation of China (11071126), and Taiwan NSC grant (99-2115-M-110-007-MY3).

Cuntz and Pedersen in [14], which captures some features of the classification of Murray and von Neumann.

The classification theme of  $C^*$ -algebras took a drastic turn after an exciting work of Elliott on the classification of AF-algebras through the ordered K-theory, in the sense that two AF-algebras are isomorphic if and only if they have the same ordered K-theory ([16]). Elliott then proposed an invariant consisting of the tracial state space and some K-theory datum of the underlying  $C^*$ -algebra (called the *Elliott invariant*) which could be a suitable candidate for a complete invariant for simple separable nuclear  $C^*$ -algebras. Although it is known recently that it is not the case (see [38]), this Elliott invariant still works for a very large class of such  $C^*$ -algebras (namely, those satisfying certain regularity conditions as described in [18]). Many people are still making progress in this direction in trying to find the biggest class of  $C^*$ -algebras that can be classified through the Elliott invariant (see, e.g., [17, 36]). Notice that this classification is very different from the classification in the sense of Murray and von Neumann.

In this article, we reconsider the classification of  $C^*$ -algebras through the idea of Murray and von Neumann. Instead of considering projections in a  $C^*$ -algebra A, we consider open projections and we twist the definition of the finiteness of projections slightly to obtain our classification scheme.

The notion of open projections was introduced by Akemann (in [1]). A projection p in the universal enveloping von Neumann algebra (i.e. the biduals)  $A^{**}$ of a  $C^*$ -algebra A (see, e.g., [37, §III.2]) is an open projection of A if there is an increasing net  $\{a_i\}_{i\in\mathfrak{I}}$  of positive elements in  $A_+$  with  $\lim_i a_i = p$  in the  $\sigma(A^{**}, A^*)$ topology. In the case when A is commutative, open projections of A are exactly characteristic functions of open subsets of the spectrum of A. In general, there is a bijective correspondence between open projections of A and hereditary  $C^*$ subalgebras of A (where a hereditary  $C^*$ -subalgebra B corresponds to an open projection p such that  $B = pA^{**}p \cap A$ ; see, e.g., [31, 3.11.10]). Characterisations and further developments of open projections can be found in, e.g., [2, 3, 4, 9, 15, 30, 33]. Since every element in a  $C^*$ -algebra is in the closed linear span of its open projections, it is reasonable to believe that the study of open projections will provide fruitful information about the underlying  $C^*$ -algebra. Moreover, because of the correspondence between open projections (respectively, central open projections) and hereditary  $C^*$ -subalgebras (respectively, closed ideals), the notion of strong Morita equivalence as defined by Rieffel (see [34] and also [11, 35]) is found to be very useful in this scheme.

One might wonder why we do not consider the classification of the universal enveloping von Neumann algebras of  $C^*$ -algebras to obtain a classification of  $C^*$ -algebras. A reason is that for a  $C^*$ -algebra A, its bidual  $A^{**}$  always contains many minimum projections (see, e.g., [1, II.17]), and hence a reasonable theory of type classification cannot be obtained without serious modifications. Furthermore,  $A^{**}$  are usually very far away from A, and information of A might not always be

respected very well in  $A^{**}$ ; for example, c and  $c_0$  have isomorphic biduals, but the structure of their open projections can be used to distinguish them (see, e.g., Example 2.1 and also Proposition 2.3(b)).

As in the case of von Neumann algebras, in order to give a classification of  $C^*$ -algebras, one needs, first of all, to consider a good equivalence relation among open projections. After some thoughts and considerations, we end up with the "spatial equivalence" as defined in Section 2, which is weaker than the one defined by Peligrad and Zsidó in [32] and stronger than the ordinary Murray-von Neumann equivalence. One reason for making this choice is that it is precisely the "hereditarily stable version of Murray-von Neumann equivalence" that one might want (see Proposition 2.7(a)(5)), and it also coincides with the "spatial isomorphism" of the hereditary  $C^*$ -subalgebras (see Proposition 2.7(a)(2)).

Using the spatial equivalence relation, we introduce in Section 3, the notion of  $C^*$ -finite  $C^*$ -algebras. It is shown that the sum of all  $C^*$ -finite hereditary  $C^*$ -subalgebra is a (not necessarily closed) ideal of the given  $C^*$ -algebra. In the case when the  $C^*$ -algebra is  $\mathfrak{B}(H)$  or  $\mathfrak{K}(H)$ , this ideal is the ideal of all finite rank operators on H. Moreover, through  $C^*$ -finiteness, we define type  $\mathfrak{A}$ , type  $\mathfrak{B}$ , type  $\mathfrak{C}$  as well as  $C^*$ -semi-finite  $C^*$ -algebras, and we study some properties of them. In particular, we will show that these properties are stable under taking hereditary  $C^*$ -subalgebras, multiplier algebras, unitalization (if the algebra is not unital) as well as strong Morita equivalence. We will also show that the notion of type  $\mathfrak{A}$  coincides precisely with the discreteness as defined in [32].

In Section 4, we will compare these notions with some results in the literature and give some examples. In particular, we show that any type I  $C^*$ -algebra (see, e.g., [31]) is of type  $\mathfrak{A}$ ; any type II  $C^*$ -algebra (as defined by Cuntz and Pedersen) is of type  $\mathfrak{B}$ ; any semi-finite  $C^*$ -algebras (in the sense of Cuntz and Pedersen) is  $C^*$ -semi-finite; any purely infinite  $C^*$ -algebra (in the sense of Kirchberg and Rørdam) with real rank zero and any separable purely infinite  $C^*$ -algebra with stable rank one are of type  $\mathfrak{C}$ ; and any type  $\mathfrak{C}$   $C^*$ -algebra is of type III (as introduced by Cuntz and Pedersen). Using our arguments for these results, we also show that any purely infinite  $C^*$ -algebra is of type III. Moreover, a von Neumann algebra M is a type  $\mathfrak{A}$ , a type  $\mathfrak{B}$ , a type  $\mathfrak{C}$  or a  $C^*$ -semi-finite  $C^*$ -algebra if and only if M is, respectively, a type I, a type II, a type III, or a semi-finite von Neumann algebra.

In Section 5, we show that any  $C^*$ -algebra A contains a largest type  $\mathfrak A$  closed ideal  $J_{\mathfrak A}^A$ , a largest type  $\mathfrak B$  closed ideal  $J_{\mathfrak B}^A$ , a largest type  $\mathfrak C$  closed ideal  $J_{\mathfrak C}^A$  as well as a largest  $C^*$ -semi-finite closed ideal  $J_{\mathfrak s\mathfrak f}^A$ . It is further shown that  $J_{\mathfrak A}^A + J_{\mathfrak B}^A$  is an essential ideal of  $J_{\mathfrak s\mathfrak f}^A$ , and  $J_{\mathfrak A}^A + J_{\mathfrak B}^A + J_{\mathfrak C}^A$  is an essential ideal of  $J_{\mathfrak B}^A$ . On the other hand,  $J_{\mathfrak C}^A$  is always a  $J_{\mathfrak S}^A$ -semi-finite  $J_{\mathfrak A}^A$ -algebra, while  $J_{\mathfrak B}^A$  is always of type  $J_{\mathfrak A}^A$  if one sets  $J_{\mathfrak S}^A$ . We also compare  $J_{\mathfrak A}^{M(A)}$ ,  $J_{\mathfrak B}^{M(A)}$ ,  $J_{\mathfrak C}^{M(A)}$  and  $J_{\mathfrak s\mathfrak f}^{M(A)}$  with  $J_{\mathfrak A}^A$ ,  $J_{\mathfrak C}^A$  and  $J_{\mathfrak s\mathfrak f}^A$ , respectively.

Notation 1.1. Throughout this paper, A is a non-zero  $C^*$ -algebra, M(A) is the multiplier algebra of A, Z(A) is the center of A, and  $A^{**}$  is the bidual of A. Furthermore, Proj(A) is the set of all projections in A, while  $OP(A) \subseteq Proj(A^{**})$  is the set of all open projections of A. All ideals in this paper are two-sided ideals (not assumed to be closed unless specified).

If  $x,y\in A^{**}$  and E is a subspace of  $A^{**}$ , we set  $xEy:=\{xzy:z\in E\}$ , and denote by  $\overline{E}$  the norm closure of E. For any  $x\in A^{**}$ , we set  $\operatorname{her}_A(x)$  to be the hereditary  $C^*$ -subalgebra  $\overline{x^*A^{**}x}\cap A$  of A (note that if  $u\in A^{**}$  is a partial isometry, then  $\operatorname{her}_A(u)=u^*A^{**}u\cap A=\{x\in A:x=u^*uxu^*u\}=\operatorname{her}_A(u^*u)$ ). When A is understood, we will use the notation  $\operatorname{her}(x)$  instead. Moreover,  $p_x$  is the right support projection of a norm one element  $x\in A$ , i.e.  $p_x$  is the  $\sigma(A^{**},A^*)$ -limit of  $\{(x^*x)^{1/n}\}_{n\in\mathbb{N}}$  and is the smallest open projection in  $A^{**}$  with  $xp_x=x$ .

Acknowledgement: The authors would like to thank L. Brown, E. Effros and G. Elliott for giving some comments.

#### 2. Spatial equivalence of open projections

In this section, we will consider a suitable equivalence relation on the set of open projections of a  $C^*$ -algebra. Let us start with the following example, which shows that the structure of open projections is rich enough to distinguish c and  $c_0$ , while they have isomorphic biduals (see Proposition 2.3(b) below for a more general result).

Example 2.1. The sets of open projections of  $c_0$  and c can be regarded as the collections  $\mathcal{X}$  and  $\mathcal{Y}$ , of open subsets of  $\mathbb{N}$  and of open subsets of the one point compactification of  $\mathbb{N}$ , respectively. As ordered sets,  $\mathcal{X}$  and  $\mathcal{Y}$  are not isomorphic. In fact, suppose on the contrary that there is an order isomorphism  $\Psi: \mathcal{Y} \to \mathcal{X}$ . Then  $\Psi(\mathbb{N})$  is a proper open subset of  $\mathbb{N}$ . Let  $k \notin \Psi(\mathbb{N})$  and  $U \in \mathcal{Y}$  with  $\Psi(U) = \{k\}$ . As U is a minimal element, it is a singleton set. Thus,  $U \subseteq \mathbb{N}$ , which gives the contradiction that  $\{k\} \subseteq \Psi(\mathbb{N})$ .

Secondly, we give the following well-known remarks which says that open projections and the hereditary  $C^*$ -subalgebras they define, are "hereditarily invariant". These will clarify some discussions later on.

Remark 2.2. Let  $B \subseteq A$  be a hereditary  $C^*$ -subalgebra and  $e \in OP(A)$  be the open projection with  $her_A(e) = B$ .

- (a) For any  $p \in \text{Proj}(B^{**})$ , one has  $\text{her}_B(p) = \text{her}_A(p)$ .
- (b)  $\operatorname{OP}(B) = \operatorname{OP}(A) \cap B^{**}$ . In fact, if  $p \in \operatorname{OP}(A) \cap B^{**}$  and  $\{a_i\}_{i \in \mathfrak{I}}$  is an approximate unit in  $\operatorname{her}_A(p) = \operatorname{her}_B(p)$ , then  $\{a_i\}_{i \in \mathfrak{I}}$  will  $\sigma(B^{**}, B^*)$ -converge to p and  $p \in \operatorname{OP}(B)$ .

- (c) If  $z \in A$  satisfying  $zz^*, z^*z \in B$ , then  $z \in B$ . In fact, as  $z^*z \in \text{her}_A(e) = eA^{**}e \cap A$ , by considering the polar decomposition of z, we see that ze = z. Similarly, we have ez = z.
- (d) If  $f \in \mathrm{OP}(A)$ , the open projections corresponding to  $\ker(e) \cap \ker(f)$  and the hereditary  $C^*$ -subalgebra generated by  $\ker(e) + \ker(f)$  are  $e \wedge f$  and  $e \vee f$  respectively.

Let  $j_A: M(A) \to A^{**}$  be the canonical \*-monomorphism, i.e.  $j_A(x)(f) = \tilde{f}(x)$   $(x \in M(A), f \in A^*)$ , where  $\tilde{f} \in M(A)^*$  is the unique strictly continuous extension of f. The proposition below can be regarded as a motivation behind the study of  $C^*$ -algebras through their open projections. It could be a known result (especially, part (a)). However, since we need it for the equivalence of (1) and (5) in Proposition 2.7(a), we give a proof here for completeness.

**Proposition 2.3.** Suppose that A and B are  $C^*$ -algebras, and  $\Phi: A^{**} \to B^{**}$  is a \*-isomorphism.

- (a) If  $\Phi(j_A(M(A))) = j_B(M(B))$ , then  $\Phi(A) = B$ .
- (b) If  $\Phi(OP(A)) = OP(B)$ , then  $\Phi(A) = B$ .

**Proof:** (a) Let  $p_A \in \mathrm{OP}(M(A))$  such that  $\ker_{M(A)}(p_A) = A$ . It is not hard to verify that  $p_A$  is the support of  $\tilde{j}_A$ , where  $\tilde{j}_A : M(A)^{**} \to A^{**}$  is the \*-epimorphism induced by  $j_A$ . Consider  $\Psi := j_B^{-1} \circ \Phi_{|j_A(M(A))} \circ j_A : M(A) \to M(B)$  (which is well-defined by the hypothesis). Since  $j_B \circ \Psi = \Phi_{|j_A(M(A))} \circ j_A$ , we see that  $\tilde{j}_B \circ \Psi^{**} = \Phi \circ \tilde{j}_A$  (as  $\Phi$  is automatically weak-\*-continuous). Thus,  $\tilde{j}_B(\Psi^{**}(p_A)) = 1_{B^{**}}$  which implies  $\Psi^{**}(p_A) \geq p_B$ . Similarly,

$$(\Psi^{**})^{-1}(p_B) = (j_A^{-1} \circ \Phi^{-1}_{|j_B(M(B))} \circ j_B)^{**}(p_B) \ge p_A$$

and we have  $\Psi^{**}(p_A) = p_B$ . Consequently,  $\Psi(\operatorname{her}_{M(A)}(p_A)) = \operatorname{her}_{M(B)}(p_B)$  as required.

(b) If  $a \in M(A)_{sa}$  and U is an open subset of  $\sigma(a) = \sigma(\Phi(j_A(a)))$ , then  $\chi_U(\Phi(j_A(a))) = \Phi(\chi_U(j_A(a)))$  is an element of OP(B) (by [5, Theorem 2.2] and the hypothesis). Thus, by [5, Theorem 2.2] again, we have  $\Phi(j_A(a)) \in j_B(M(B))$ . A similar argument shows that  $\Phi^{-1}(j_B(M(B))) \subseteq j_A(M(A))$ . Now, we can apply part (a) to obtain the required conclusion.

Remark 2.4. Note that if A and B are separable and  $\Psi: M(A) \to M(B)$  is a \*-isomorphism, then  $\Psi(A) = B$ , by a result of Brown in [10]. However, the same result is not true if one of them is not separable (e.g. take A = M(B) and  $\Psi = \mathrm{id}$ , where B is non-unital). Proposition 2.3(a) shows that one has  $\Psi(A) = B$  if (and only if)  $\Psi$  extends to a \*-isomorphism from  $A^{**}$  to  $B^{**}$ .

We now consider a suitable equivalence relation on OP(A). A naive choice is to use the original "Murray-von Neumann equivalence"  $\sim_{Mv}$ . However, this choice

is not good because [23] tells us that two open projections that are Murray-von Neumann equivalent might define non-isomorphic hereditary  $C^*$ -subalgebras. On the other hand, one might define  $p \sim_{\text{her}} q$   $(p, q \in \text{OP}(A))$  whenever  $\text{her}(p) \cong \text{her}(q)$  as  $C^*$ -algebras. The problem of this choice is that two distinct open projections of C([0,1]) can be equivalent (if they correspond to homeomorphic open subsets of [0,1]), which means that the resulting classification, even if possible, will be very different from the Murray-von Neumann classification.

After some thoughts, we end up with an equivalence relation  $\sim_{\text{sp}}$  on OP(A):  $p \sim_{\text{sp}} q$  if there is a partial isometry  $v \in A^{**}$  satisfying

$$v^* \operatorname{her}_A(p) v = \operatorname{her}_A(q)$$
 and  $v \operatorname{her}_A(q) v^* = \operatorname{her}_A(p)$ .

Note that this relation is precisely the "hereditarily stable version" of the Murray-von Neumann equivalence (see Proposition 2.7(a)(5) below and the discussion following it).

In [32, Definition 1.1], Peligrad and Zsidó introduced another equivalence relation on  $\operatorname{Proj}(A^{**})$ :  $p \sim_{\operatorname{PZ}} q$  if there is a partial isometry  $v \in A^{**}$  such that

$$(2.1) p = vv^*, q = v^*v, v^* \operatorname{her}_A(p) \subseteq A \text{and} v \operatorname{her}_A(q) \subseteq A.$$

It is not difficult to see that  $\sim_{\rm PZ}$  is stronger than  $\sim_{\rm sp}$ , and a natural description of  $\sim_{\rm PZ}$  on the set of range projections of positive elements of A is given in [29, Proposition 4.3]. Moreover, we also gave in [28, Proposition 3.1] an equivalent description of  $\sim_{\rm PZ}$  that is similar to  $\sim_{\rm sp}$  but use right ideals instead of hereditary  $C^*$ -subalgebras. However, it is now known that  $\sim_{\rm PZ}$  and  $\sim_{\rm sp}$  are actually different even for very simple kind of  $C^*$ -algebras (see [28, Theorem 5.3]). We decide to use  $\sim_{\rm sp}$  as it seems to be more natural in the way of using open projections (see Proposition 2.7(a) below).

Let us start with an extension of  $\sim_{\rm sp}$  to the whole of  $\operatorname{Proj}(A^{**})$ .

**Definition 2.5.** We say that  $p, q \in \text{Proj}(A^{**})$  are spatially equivalent with respect to A, denoted by  $p \sim_{\text{sp}} q$ , if there exists a partial isometry  $v \in A^{**}$  satisfying

(2.2) 
$$p = vv^*$$
,  $q = v^*v$ ,  $v^* \operatorname{her}_A(p)v = \operatorname{her}_A(q)$  and  $v \operatorname{her}_A(q)v^* = \operatorname{her}_A(p)$ .

In this case, we also say that the hereditary  $C^*$ -subalgebras  $\operatorname{her}_A(p)$  and  $\operatorname{her}_A(q)$  are spatially isomorphic.

It might happen that her(p) = 0 but  $p \neq 0$  and this is why we need to consider the first two conditions in (2.2). We will see in Proposition 2.7(a) that the first two conditions are redundant if p and q are both open projections.

Obviously,  $\sim_{\rm sp}$  is stronger than  $\sim_{\rm Mv}$  (for elements in  $\operatorname{Proj}(A^{**})$ ). Moreover, if  $p \sim_{\rm sp} q$ , then  $x \mapsto v^*xv$  is a \*-isomorphism from  $\operatorname{her}(p)$  to  $\operatorname{her}(q)$ , which means that  $\sim_{\rm sp}$  is stronger than  $\sim_{\rm her}$  in the context of open projections.

A good point of the spatial equivalence is that open projections are stable under  $\sim_{\rm sp}$ , as can be seen in part (b) of the following lemma.

**Lemma 2.6.** (a)  $\sim_{\text{sp}}$  is an equivalence relation in  $\text{Proj}(A^{**})$ .

- (b) Let  $p, q \in \operatorname{Proj}(A^{**})$  and  $u \in A^{**}$  be a partial isometry. If p is open,  $u^*pu = q$ ,  $\operatorname{her}_A(p) \subseteq u \operatorname{her}_A(q)u^*$  and  $\operatorname{her}_A(q) \subseteq u^* \operatorname{her}_A(p)u$ , then q is open and  $p \sim_{\operatorname{sp}} q$ . Consequently, if  $p \sim_{\operatorname{sp}} q$  and p is open, then q is open.
- (c) If  $B \subseteq A$  is a hereditary  $C^*$ -subalgebra and  $p, q \in \text{Proj}(B^{**})$ , then p and q are spatially equivalent with respect to B if and only if they are spatially equivalent with respect to A.

**Proof:** (a) It suffices to verify the transitivity. Suppose that p, q and v are as in Definition 2.5. If  $w \in A^{**}$  and  $r \in \text{Proj}(A^{**})$  satisfy that

$$p = w^*w$$
,  $r = ww^*$ ,  $w \operatorname{her}_A(p)w^* = \operatorname{her}_A(r)$  and  $w^* \operatorname{her}_A(r)w = \operatorname{her}_A(p)$ ,

then the partial isometry wv gives the equivalence  $r \sim_{sp} q$ .

- (b) As p is open and  $\operatorname{her}_A(p)$  is contained in the weak-\*-closed subspace  $uA^{**}u^*$ , one has  $p \leq uu^*$ . Let v := pu. Then  $vv^* = p$  and  $v^*v = u^*pu = q$ . Moreover, it is clear that  $\operatorname{her}_A(p) \subseteq v \operatorname{her}_A(q)v^*$  and  $\operatorname{her}_A(q) \subseteq v^* \operatorname{her}_A(p)v$ . Now, it is easy to see that the relations in (2.2) are satisfied. Furthermore, if  $\{a_i\}_{i\in\mathfrak{I}}$  is an approximate unit in  $\operatorname{her}_A(p)$ , then  $\{v^*a_iv\}$  is an increasing net in  $\operatorname{her}_A(q)$  that weak-\*-converges to  $v^*pv = q$ , and so q is open. The second statement follows directly from the first one.
- (c) Suppose that p and q are spatially equivalent with respect to A and  $v \in A^{**}$  satisfies the relations in (2.2). As  $vv^*, v^*v \in B^{**}$ , Remark 2.2(c) tells us that  $v \in B^{**}$ . Now the equivalence follows from Remark 2.2(a).

**Proposition 2.7.** (a) If  $p, q \in OP(A)$ , the following statements are equivalent.

- (1)  $p \sim_{\rm sp} q$ .
- (2)  $her(q) = u^* her(p)u$  and  $her(p) = u her(q)u^*$  for a partial isometry  $u \in A^{**}$ .
- (3)  $her(q) \subseteq u^* her(p)u$  and  $her(p) \subseteq u her(q)u^*$  for a partial isometry  $u \in A^{**}$ .
- (4)  $q \le v^*v$  and  $v \operatorname{her}(q)v^* = \operatorname{her}(p)$  for a partial isometry  $v \in A^{**}$ .
- (5) There is a partial isometry  $w \in A^{**}$  such that  $p = ww^*$  and

$$\{w^*rw : r \in OP(A); r \le p\} = \{s \in OP(A) : s \le q\}.$$

(b) If M is a von Neumann algebra and  $p, q \in \text{Proj}(M)$ , then  $p \sim_{\text{sp}} q$  if and only if  $p \sim_{\text{Mv}} q$  as elements in Proj(M).

**Proof:** (a) The implications  $(1) \Rightarrow (2) \Rightarrow (3)$  and  $(1) \Rightarrow (4)$  are clear.

- (3)  $\Rightarrow$  (1). Since q is open, one has  $q \leq u^*u$ . Thus,  $(uq)^*uq = q$  and Statement (3) also holds when u is replaced by uq. As p is also open, a similar argument shows that  $p \leq uqu^*$  and Statement (3) holds if we replace u by v := puq and that  $p = vv^*$ . Furthermore, since  $vqv^* = vv^* = p$ , Lemma 2.6(b) tells us that  $p \sim_{sp} q$ .
- $(4) \Rightarrow (2)$ . This follows from  $v^* \operatorname{her}(p)v = v^*v \operatorname{her}(q)v^*v = \operatorname{her}(q)$ .
- (1)  $\Rightarrow$  (5). Notice that  $OP(her(p)) = \{r \in OP(A) : r \leq p\}$  (see Remark 2.2(b)). Suppose that  $v \in A^{**}$  satisfies (2.2) and  $r \in OP(her(p))$ . If  $\{a_i\}_{i \in \mathfrak{I}}$  is an increasing

net in her(p) that  $\sigma(A^{**}, A^*)$ -converge to r, then  $\{v^*a_iv\}_{i\in\mathfrak{I}}$  is an increasing net in her(q) that  $\sigma(A^{**}, A^*)$ -converge to  $v^*rv$  and hence  $v^*rv \in OP(her(q))$ . The argument for the other inclusion is similar.

- (5)  $\Rightarrow$  (1). By Statement (5), we have  $q = w^*pw$ , and the map  $\Phi : x \mapsto w^*xw$  is a \*-isomorphism from  $her(p)^{**}$  to  $her(q)^{**}$ . By Proposition 2.3(b), we see that  $\Phi(her(p)) = her(q)$  and Statement (4) holds.
- (b) If  $p \sim_{\text{sp}} q$ , then  $p \sim_{\text{Mv}} q$  as elements in  $\text{Proj}(M^{**})$ , which implies that  $p \sim_{\text{Mv}} q$  as elements in Proj(M) (by considering the canonical \*-homomorphism  $\Lambda_M : M^{**} \to M$ ). Conversely, if  $v \in M$  satisfying  $p = vv^*$  and  $q = v^*v$ , then clearly  $v^* \text{her}(p)v = \text{her}(q)$ .

One can reformulate Statement (5) of Proposition 2.7(a) in the following way. There is a partial isometry  $w \in A^{**}$  that induces Murray-von Neumann equivalences between open subprojections of p (including p) and open subprojections of q (including q).

Therefore, one may regard  $\sim_{\rm sp}$  as the "hereditarily stable version" of the Murray-von Neumann equivalence. Moreover, if  $v \in A^{**}$  satisfies the relations in (2.2), then by Lemma 2.6(b),  $r \sim_{\rm sp} v^*rv$  for all  $r \in {\rm OP}({\rm her}(p))$ , which means that spatial equivalence is automatically "hereditarily stable".

Remark 2.8. (a) Let  $p, q \in \operatorname{Proj}(A^{**})$ . We call the unique  $p_{\operatorname{int}} \in \operatorname{OP}(A)$  with  $\operatorname{her}(p) = \operatorname{her}(p_{\operatorname{int}})$  the interior of p. By the bijective correspondence between hereditary  $C^*$ -subalgebras and open projections,  $p_{\operatorname{int}}$  is the largest open projection dominated by p. As a direct consequence of Proposition 2.7(a), we know that  $p_{\operatorname{int}} \sim_{\operatorname{sp}} q_{\operatorname{int}}$  if and only if

 $\operatorname{her}(q) \subseteq u^* \operatorname{her}(p)u$  and  $\operatorname{her}(p) \subseteq u \operatorname{her}(q)u^*$  for a partial isometry  $u \in A^{**}$ .

(b) Suppose that  $p,q \in \mathrm{OP}(A)$ . One might attempt to define  $p \lesssim_{\mathrm{sp}} q$  if there is  $q_1 \in \mathrm{OP}(A)$  with  $p \sim_{\mathrm{sp}} q_1 \leq q$ . However, unlike the Murray-von Neumann equivalence situation,  $p \lesssim_{\mathrm{sp}} q$  and  $q \lesssim_{\mathrm{sp}} p$  does not imply that  $p \sim_{\mathrm{sp}} q$ . This can be shown by using a result of Lin. More precisely, it was shown in [23, Theorem 9] that there exist a separable unital simple  $C^*$ -algebra A as well as  $p \in \mathrm{Proj}(A)$  and  $u \in A$  such that  $uu^* = 1$  and  $p_1 = u^*u \leq p$ , but  $\mathrm{her}(p)$  and A are not \*-isomorphic. In particular,  $p \nsim_{\mathrm{sp}} 1$ . Now, we clearly have  $p \lesssim_{\mathrm{sp}} 1$ . On the other hand, as  $u \in A$ , we have

$$u^*Au = her(p_1)$$
 and  $u her(p_1)u^* = A$ ,

which implies that  $1 \lesssim_{sp} p$ .

This example also shows that the same problematic situation appears even if we replace  $\sim_{\rm sp}$  with the stronger equivalence relation  $\sim_{\rm PZ}$  as defined in (2.1) (because  $u \in A$ ). Nevertheless, it was shown in [32, Theorem 1.13] that a weaker conclusion holds if one adds an extra assumption on either p or q, but we will not recall the details here.

Let us end this section with the following well-known example. We give an explicit argument here for future reference. Note that parts (a) and (b) of it mean that if  $a, b \in A_+$  are equivalent in the sense of Blackadar (i.e., there exists  $x \in A$  with  $a = x^*x$  and  $b = xx^*$ ; see, e.g., [29, Definition 2.1]), then their support projections are spatially equivalence (which is also a corollary of [29, Proposition 4.3], since  $\sim_{PZ}$  is stronger than  $\sim_{sp}$ ).

Example 2.9. Suppose that  $x \in A$  with ||x|| = 1. Set  $a = x^*x$  and  $b = xx^*$ . Let  $x = ua^{1/2}$  be the polar decomposition.

- (a) It is easy to see that  $\overline{aAa} = u^*(\overline{xAx^*})u$  and  $\overline{xAx^*} = u(\overline{aAa})u^*$ , i.e.,  $\overline{xAx^*}$  is spatially isomorphic to  $\overline{aAa}$  (by Proposition 2.7(a)).
- (b) Notice that  $u(\overline{aAa})u^* = \overline{xAx^*} \supseteq \overline{xx^*Axx^*} \supseteq \overline{xx^*xAx^*xx^*} \supseteq u\overline{a^{3/2}Aa^{3/2}}u^* = u(\overline{aAa})u^*$ , and we have  $\overline{xAx^*} = \overline{bAb}$ . Similarly,  $\overline{x^*Ax} = \overline{aAa}$  and  $\overline{x^*A^{**}x} = \overline{aA^{**}a}$ , which implies that  $\operatorname{her}(x) = \operatorname{her}(a)$ . On the other hand, as  $\overline{aAa}$  is a hereditary  $C^*$ -subalgebra of  $\operatorname{her}(a)$  and  $\{a^{1/k}\}_{k\in\mathbb{N}}$  is a sequence in  $\overline{aAa}$  which is an approximate unit for  $\operatorname{her}(a)$ , one has  $\overline{aAa} = \operatorname{her}(a)$ . Consequently,  $\operatorname{her}(x) = \overline{x^*Ax}$ .
- (c) Suppose that  $B \subseteq A$  is a hereditary  $C^*$ -subalgebra and  $x \in B$ . Since  $\overline{aAa} = \overline{a^2Aa^2}$ , we see that  $\overline{aBa} = \overline{aAa}$ . Therefore,  $\operatorname{her}_B(x) = \operatorname{her}_A(x)$  by part (b).

## 3. $C^*$ -semi-finiteness and three types of $C^*$ -algebras

As in the case of von Neumann algebras ([27]), in order to define different "types" of  $C^*$ -algebras, we need to define "abelian" and "finite" open projections. "Abelian" open projections are defined in the same way as that of von Neumann algebras. However, in order to define "finite" open projections, we need to use our "hereditarily stable version" of Murray-von Neumann equivalence in Section 2. Note that one cannot go very far with the original Murray-von Neumann equivalence, because there exist  $p, q \in \mathrm{OP}(A)$  with  $p \sim_{\mathrm{Mv}} q$  but  $\mathrm{her}(p)$  and  $\mathrm{her}(q)$  are not isomorphic (see [23]). Moreover, one cannot use a direct verbatim translation of the Murray-von Neumann finiteness.

**Definition 3.1.** (a) Let  $q \in OP(A)$  and  $p \in Proj(qA^{**}q)$ . The closure of p in q, denoted by  $\bar{p}^q$ , is the smallest closed projection of her(q) that dominates p.

- (b) Let  $p, q \in OP(A)$  with  $p \leq q$ . The projection p is said to be
  - i. dense in q if  $\bar{p}^q = q$ ;
- ii. abelian if her(p) is a commutative  $C^*$ -algebra;
- iii.  $C^*$ -finite if for any  $r, s \in OP(her(p))$  with  $r \leq s$  and  $r \sim_{sp} s$ , one has  $\bar{r}^s = s$ .

If p is dense in q, we say that her(p) is essential in her(q). We denote by  $OP_{\mathcal{C}}(A)$  and  $OP_{\mathcal{F}}(A)$  the set of all abelian open projections and the set of all  $C^*$ -finite open projections of A, respectively.

The terminology "p is dense in q" is used in many places (e.g. [32]), while the terminology "essential" comes from [39].

Some people might wonder why we do not use the finiteness as defined in [14]. The reason is that we want to give a classification scheme for  $C^*$ -algebras using open projections (and the definition of finiteness in [14] seems not related to open projections).

# Remark 3.2. Let $p \in OP(A)$ .

- (a) Suppose that p is abelian. If  $r, s \in OP(her(p))$  satisfying  $r \leq s$  and  $r \sim_{sp} s$ , then r = s. Thus, p is  $C^*$ -finite.
- (b) If her(p) is finite dimensional, then p is  $C^*$ -finite.
- (c) One might ask why we do not define  $C^*$ -finiteness of p in the following way: for any  $r \in \mathrm{OP}(\mathrm{her}(p))$  with  $r \sim_{\mathrm{sp}} p$ , one has  $\bar{r}^p = p$ . The reason is that the stronger condition in Definition 3.1(b) can ensure every open subprojection of a  $C^*$ -finite projection being  $C^*$ -finite. Such a phenomena is automatic for von Neumann algebras.
- (d) A hereditary  $C^*$ -subalgebra  $B \subseteq A$  is essential in A if and only if for any non-zero hereditary  $C^*$ -subalgebra  $C \subseteq A$ , one has  $B \cdot C \neq \{0\}$ . Thus, a closed ideal  $I \subseteq A$  is essential in the sense of Definition 3.1 if and only it is essential in the usual sense (i.e., any non-zero closed ideal of A intersects I non-trivially).

# **Definition 3.3.** A $C^*$ -algebra A is said to be:

- i.  $C^*$ -finite if  $1 \in \mathrm{OP}_{\mathfrak{F}}(A)$ ;
- ii.  $C^*$ -semi-finite if every element in  $OP(A)\setminus\{0\}$  dominates an element in  $OP_{\mathcal{F}}(A)\setminus\{0\}$ :
- iii. of Type  $\mathfrak{A}$  if every element in  $OP(A) \cap Z(A^{**}) \setminus \{0\}$  dominates an element in  $OP_{\mathfrak{C}}(A) \setminus \{0\}$ ;
- iv. of Type  $\mathfrak{B}$  if  $OP_{\mathfrak{C}}(A) = \{0\}$  but each element in  $OP(A) \cap Z(A^{**}) \setminus \{0\}$  dominates an element in  $OP_{\mathfrak{F}}(A) \setminus \{0\}$ ;
- v. of Type  $\mathfrak{C}$  if  $OP_{\mathfrak{F}}(A) = \{0\}$ .

Let us give an equivalent form of the above abstract definition through the relation between open projections (respectively, central open projections) and hereditary  $C^*$ -subalgebras (respectively, ideals). A  $C^*$ -algebra A is

- $C^*$ -finite if and only if for each hereditary  $C^*$ -subalgebra  $B \subseteq A$ , every hereditary  $C^*$ -subalgebra of B that is spatially isomorphic to B is essential in B;
- $C^*$ -semi-finite if and only if every non-zero hereditary  $C^*$ -subalgebra of A contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra;
- of type  $\mathfrak{A}$  if and only if every non-zero closed ideal of A contains a non-zero abelian hereditary  $C^*$ -subalgebra;

- of type  $\mathfrak{B}$  if and only if A does not contain any non-zero abelian hereditary  $C^*$ -subalgebra and every non-zero closed ideal of A contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra;
- of type  $\mathfrak C$  if and only if A does not contain any non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra.

Remark 3.4. Suppose that A is simple.

- (a) A is either of type  $\mathfrak{A}$ , type  $\mathfrak{B}$  or type  $\mathfrak{C}$ .
- (b) We will see in Corollary 4.5 that A is of type  $\mathfrak{A}$  if and only if A is of type I (see, e.g., [31, 6.1.1] for its definition). Moreover, if A is of type II (in the sense of [14]), then A is of type  $\mathfrak{B}$  (by Proposition 4.7 below), while if A is purely infinite (in the sense of [13]), then A is of type  $\mathfrak{C}$  (by Proposition 4.11(a) below and [40, Theorem 1.2(ii)]). However, we do not know if the converse of the last two statements hold.

A positive element  $a \in A_+$  is said to be  $C^*$ -finite if her(a) (i.e.,  $\overline{aAa}$ ) is  $C^*$ -finite.

**Proposition 3.5.** (a) The sum,  $\mathcal{C}(A)$ , of all abelian hereditary  $C^*$ -subalgebras of A is a (not necessarily closed) ideal of A. If  $\mathcal{C}(A)_+ := \mathcal{C}(A) \cap A_+$ , then  $\mathcal{C}(A)$  coincides with the vector space span  $\mathcal{C}(A)_+$  generated by  $\mathcal{C}(A)_+$ .

- (b) The sum,  $\mathfrak{F}(A)$ , of all  $C^*$ -finite hereditary  $C^*$ -subalgebras of A is a (not necessarily closed) ideal of A. If  $\mathfrak{F}(A)_+ := \mathfrak{F}(A) \cap A_+$ , then  $\mathfrak{F}(A) = \operatorname{span} \mathfrak{F}(A)_+$ .
- (c) If  $B \subseteq A$  is a hereditary  $C^*$ -subalgebra, then  $\mathfrak{C}(B)_+ = \mathfrak{C}(A) \cap B_+$  and  $\mathfrak{F}(B)_+ = \mathfrak{F}(A) \cap B_+$ .

**Proof:** Since parts (a) and (b) follow from the arguments of [31, Proposition 6.1.7], we will only give the proof for part (c). Moreover, we will only establish the second equality as the argument for the first one is similar. As  $K_A$  is a hereditary cone, the argument of part (b) tells us that  $\mathcal{F}(A)_+ = K_A$ . It is clear that  $\mathcal{F}(B) \subseteq \mathcal{F}(A) \cap B$ . Conversely, if  $w \in K_A \cap B$  and  $w_1, ..., w_n \in F_A$  such that  $w = \sum_{i=1}^n w_i$ , then  $w_i \leq w \in B_+$ , which implies that  $w_i \in F_A \cap B = F_B$  (see Example 2.9(c)). Consequently,  $w \in K_B$  as required.

Clearly,  $\mathcal{C}(A) \subseteq \mathcal{F}(A)$ . We will see in Theorem 5.2(d) below that the closed ideal  $\overline{\mathcal{C}(A)}$  is of type  $\mathfrak{A}$ , while  $\overline{\mathcal{F}(A)}$  is  $C^*$ -semi-finite.

Example 3.6. (a) If A is commutative, then A is of type  $\mathfrak{A}$  and is  $C^*$ -finite. Moreover,  $\mathfrak{C}(A) = \mathfrak{F}(A) = A$ .

(b) Let  $p \in \mathrm{OP}(\mathfrak{B}(\ell^2)) \subseteq \mathfrak{B}(\ell^2)^{**}$  such that  $\ker(p) = \mathfrak{K}(\ell^2)$  (the  $C^*$ -algebra of all compact operators). Then  $p \neq 1$  but  $\ker(1-p) = (0)$ . In fact, if  $T \in \ker(1-p)$ , we have pT = 0 and ST = SpT = 0 for any  $S \in \mathfrak{K}(\ell^2)$ , which gives T = 0. Moreover, p is dense in 1 because  $\mathfrak{K}(\ell^2)$  is an essential closed ideal of  $\mathfrak{B}(\ell^2)$  (see Remark 3.2(d)).

(c) If H is an infinite dimensional Hilbert space, then  $\mathcal{K}(H)$  is a  $C^*$ -algebra of type  $\mathfrak{A}$ , which is not  $C^*$ -finite but is  $C^*$ -semi-finite. In fact, as  $\mathcal{K}(H)$  is simple and contains many rank-one projections, it is of type  $\mathfrak{A}$ . On the other hand, suppose that  $e \in \operatorname{Proj}(\mathcal{K}(H))$  is a rank-one projection. Then  $1 - e \in \operatorname{OP}(\mathcal{K}(H)) \subseteq \mathcal{B}(H)$  and there is an isometry  $v \in \mathcal{B}(H)$  with  $vv^* = 1 - e$ . Thus,

$$v^* \operatorname{her}(1-e)v = \mathcal{K}(H)$$
 and  $1-e \sim_{\operatorname{sp}} 1$ .

Moreover, as  $e \in \operatorname{Proj}(\mathcal{K}(H))$ , we see that 1-e is also a closed projection and hence it is not dense in 1. Finally, as all hereditary  $C^*$ -subalgebras of  $\mathcal{K}(H)$  are given by projections in  $\mathcal{B}(H)$ , they are of the form  $\mathcal{K}(K)$  for some subspaces  $K \subseteq H$ . Hence,  $\mathcal{K}(H)$  is  $C^*$ -semi-finite (see Remark 3.2(b)).

(d) Let H be a Hilbert space. Clearly,  $\operatorname{Proj}(\mathfrak{K}(H)) \subseteq \operatorname{OP}_{\mathfrak{F}}(\mathfrak{B}(H))$ . Hence, if  $\mathfrak{F}(H)$  is the set of all finite rank operators, then  $\mathfrak{F}(H) \subseteq \mathfrak{F}(\mathfrak{B}(H))$ . Suppose that  $B \subseteq \mathfrak{B}(H)$  is a  $C^*$ -finite hereditary  $C^*$ -subalgebra and  $p \in \operatorname{Proj}(B)$ . As p is  $C^*$ -finite and  $pBp = p\mathfrak{B}(H)p \cong \mathfrak{B}(K)$  for a subspace  $K \subseteq H$ , we see that K is finite dimensional (see part (c)) and so  $p \in \mathcal{K}(H)$ . Since  $B \subseteq \mathfrak{B}(H)$  is a hereditary  $C^*$ -subalgebra, B is generated by its projections. Thus, B is a hereditary  $C^*$ -subalgebra of  $\mathcal{K}(H)$ , and  $B \cong \mathcal{K}(H')$  for a subspace  $H' \subseteq H$ . The  $C^*$ -finiteness of B again implies that  $\dim H' < \infty$ , and  $B \subseteq \mathfrak{F}(H)$ . Consequently,

$$\mathfrak{F}(\mathfrak{B}(H)) = \mathfrak{F}(H).$$

On the other hand, since any finite rank projection is a sum of rank-one projections and any rank-one projection belongs to  $\mathcal{C}(\mathcal{B}(H))$ , we see that  $\mathfrak{F}(H) = \mathcal{C}(\mathcal{B}(H)) = \mathcal{F}(\mathcal{B}(H))$ . Furthermore, by Proposition 3.5(c), we also have  $\mathcal{F}(\mathcal{K}(H)) = \mathcal{C}(\mathcal{K}(H)) = \mathfrak{F}(H)$ .

Remark 3.7. Let  $e \in OP(A)$  and z(e) be the central support of e in  $A^{**}$ .

- (a)  $z(e) = \sup_{u \in U_{M(A)}} ueu^*$  (see, e.g., [31, Lemma 2.6.3]), and z(e) is an open projection (see Remark 2.2(d)) with her(z(e)) being the smallest closed ideal containing her(e).
- (b) Recall that  $B := her(e) \subseteq A$  is said to be *full* if her(z(e)) = A. In this case, B is strongly Morita equivalent to A (see, e.g., [35]). Consequently, her(e) is always strongly Morita equivalent to her(z(e)).

The following provides an important tool to us in this paper. An essential ingredient of its proof (in particular, part (b)) is a result of Peligrad and Zsidó in [32].

**Proposition 3.8.** Let A and B be two strongly Morita equivalent  $C^*$ -algebras.

- (a) A contains a non-zero abelian hereditary  $C^*$ -subalgebra if and only if B does.
- (b) A contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra if and only if B does.

**Proof:** There exist a  $C^*$ -algebra D and  $e \in \text{Proj}(M(D))$  such that both A and B are full hereditary  $C^*$ -subalgebras of D and we have

$$A \cong eDe$$
 and  $B \cong (1-e)D(1-e)$ 

(see, e.g., [8, Theorem II.7.6.9]). Thus, z(e) = 1 = z(1 - e).

(a) It suffices to show that A contains a non-zero abelian hereditary  $C^*$ -subalgebra whenever D does. Let  $p \in \operatorname{OP}_{\mathfrak{C}}(D) \setminus \{0\}$ . As  $pz(e) = p \neq 0$ , we see that  $pueu^* \neq 0$  for some  $u \in U_{M(D)}$ . By replacing p with  $u^*pu$ , we may assume that  $pe \neq 0$ , and hence  $e \operatorname{her}_D(p)e \neq (0)$ . If  $x, y \in \operatorname{her}_D(p)$  and  $\{b_j\}_{j \in \mathfrak{I}}$  is an approximate unit of  $\operatorname{her}_D(p)$ , then  $b_i e b_j \in \operatorname{her}_D(p)$  which implies that

$$xey = \lim xb_ieb_iy = \lim yb_ieb_ix = yex.$$

Consequently,  $e \operatorname{her}_D(p)e$  is an abelian hereditary  $C^*$ -subalgebra of A.

(b) It suffices to show that if D contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra, then so does A. Suppose that  $p \in \mathrm{OP}_{\mathcal{F}}(D) \setminus \{0\}$ . By [32, Theorem 1.9], there exist  $e_0, e_1 \in \mathrm{OP}(\mathrm{her}_D(e))$  and  $p_0, p_1 \in \mathrm{OP}(\mathrm{her}_D(p))$  satisfying

$$\overline{e_0 + e_1}^e = e$$
,  $\overline{p_0 + p_1}^p = p$ ,  $z(e_0)z(p_0) = 0$  and  $e_1 \sim_{PZ} p_1$ .

Suppose that  $p_1 = 0$ . Then  $e_1 = 0$  and  $z(e_0)$  is dense in z(e) = 1 (by [32, Lemma 1.8]). This implies that  $z(p_0) = 0$ , and we have a contradiction that  $p_0 = 0$  is dense in the non-zero open projection p. Therefore,  $p_1 \neq 0$  and is  $C^*$ -finite. Since  $\operatorname{her}_D(e_1) \cong \operatorname{her}_D(p_1)$  (note that  $\sim_{\operatorname{PZ}}$  is stronger than  $\sim_{\operatorname{sp}}$ ), we see that  $\operatorname{her}_D(e_1)$  is a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra of  $A = \operatorname{her}_D(e)$ .

One may also use the argument of part (b) to obtain part (a), but we keep the alternative argument since it is also interesting.

Suppose that E is a full Hilbert A-module implementing the strong Morita equivalence between A and B, i.e.,  $B \cong \mathcal{K}_A(E)$  (see, e.g., [22]). If I is a closed ideal of A, then EI is a full Hilbert I-module and  $\mathcal{K}_I(EI)$  is a closed ideal of B.

We recall from [32, Definition 2.1] that A is said to be *discrete* if any non-zero open projection of A dominates a non-zero abelian open projection.

**Theorem 3.9.** (a) Let A and B be two strongly Morita equivalent  $C^*$ -algebras. Then A is of type  $\mathfrak{A}$  (respectively, type  $\mathfrak{B}$  or type  $\mathfrak{C}$ ) if and only if B is of the same type.

(b) A  $C^*$ -algebra A is of type  $\mathfrak{A}$  if and only if it is discrete.

**Proof:** (a) Suppose that A is of type  $\mathfrak{B}$ . If  $\mathrm{OP}_{\mathfrak{C}}(B) \neq \{0\}$ , then  $\mathrm{OP}_{\mathfrak{C}}(A) \neq \{0\}$  (because of Proposition 3.8(a)), which is a contradiction. Let J be a non-zero closed ideal of B. As in the paragraph above, the strong Morita equivalence of A and B gives a closed ideal  $J_0$  of A that is strongly Morita equivalent to J. As  $J_0$  contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra, so is J (by Proposition 3.8(b)). This shows that B is of type  $\mathfrak{B}$ . The argument for the other two types are similar and easier.

(b) It suffices to show that if A is of type  $\mathfrak{A}$ , then it is discrete. Let  $B \subseteq A$  be a non-zero hereditary  $C^*$ -subalgebra and  $J \subseteq A$  be the closed ideal generated by B (which is strongly Morita equivalent to B; see Remark 3.7(b)). As J contains a non-zero abelian hereditary  $C^*$ -subalgebra, so does B (by Proposition 3.8(a)).  $\square$ 

The following result follows from Proposition 3.8(b) and the argument of Theorem 3.9.

Corollary 3.10. (a) A is  $C^*$ -semi-finite if and only if any non-zero closed ideal of A contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra.

- (b) If A is strongly Morita equivalent to a  $C^*$ -semi-finite  $C^*$ -algebra, then A is also  $C^*$ -semi-finite.
- (c) A is of type  $\mathfrak{B}$  if and only if it is  $C^*$ -semi-finite and anti-liminary (i.e., it does not contain any non-zero commutative hereditary  $C^*$ -subalgebra).
- Remark 3.11. (a) As in the case of von Neumann algebra, strong Morita equivalence does not preserve  $C^*$ -finiteness. In fact, for any  $C^*$ -algebra A, the algebra  $A \otimes \mathcal{K}(\ell^2)$  is not  $C^*$ -finite (using the same argument as Example 3.6(c); note that  $1 \otimes (1-e)$  is both an open and a closed projection of  $A \otimes \mathcal{K}(\ell^2)$ ). Consequently, any stable  $C^*$ -algebra is not  $C^*$ -finite.
- (b) By Remark 3.7(b), Theorem 3.9(a) and Corollary 3.10(b), any type  $\mathfrak{A}$ , type  $\mathfrak{B}$ , type  $\mathfrak{C}$  or  $C^*$ -semi-finite hereditary  $C^*$ -subalgebra is contained in a closed ideal of the same type.

Recall that a  $C^*$ -algebra A has real rank zero in the sense of Brown and Pedersen if the set of elements in  $A_{sa}$  with finite spectrum is norm dense in  $A_{sa}$  (see, e.g., [12, Corollary 2.6]). The following result follows from Theorem 3.9(b), Corollary 3.10(c) as well as the fact that any hereditary  $C^*$ -subalgebra of a real rank zero  $C^*$ -algebra is again of real rank zero (see, e.g., [12, Corollary 2.8]).

# Corollary 3.12. Let A be a $C^*$ -algebra with real rank zero.

- (a) A is of type  $\mathfrak{A}$  if and only if every projection in  $\operatorname{Proj}(A) \setminus \{0\}$  dominates an abelian projection in  $\operatorname{Proj}(A) \setminus \{0\}$ .
- (b) A is of type  $\mathfrak{B}$  if and only if every projection in  $Proj(A) \setminus \{0\}$  is non-abelian but dominates a  $C^*$ -finite projection in  $Proj(A) \setminus \{0\}$ .
- (c) A is of type  $\mathfrak{C}$  if and only if A does not contain any non-zero  $C^*$ -finite projection.
- (d) A is  $C^*$ -semi-finite if and only if every projection in  $Proj(A) \setminus \{0\}$  dominates a  $C^*$ -finite projection in  $Proj(A) \setminus \{0\}$ .

Remark 3.13. Suppose that A is a  $C^*$ -finite  $C^*$ -algebra with real rank zero. If  $r, p \in \operatorname{Proj}(A)$  such that  $r \leq p$  and there exists  $u \in A$  with  $uu^* = r$  and  $u^*u = p$ , then  $r \sim_{\operatorname{sp}} p$  and so,  $r = \overline{r}^p = p$ .

**Corollary 3.14.** If A is of real rank zero, then the closures of the ideals C(A) and F(A) (see Proposition 3.5) are the closed linear spans of abelian projections and of  $C^*$ -finite projections in P(A), respectively.

**Proof:** If  $B \subseteq A$  is a  $C^*$ -finite hereditary  $C^*$ -subalgebra, then B is the closed linear span of  $\operatorname{Proj}(B) \cap \operatorname{OP}_{\mathfrak{F}}(B)$ . Thus,  $\mathfrak{F}(A)$  lies inside the closed linear span of  $\operatorname{Proj}(A) \cap \operatorname{OP}_{\mathfrak{F}}(A)$ . Conversely, it is clear that  $\operatorname{Proj}(A) \cap \operatorname{OP}_{\mathfrak{F}}(A) \subseteq \mathfrak{F}(A)$ . The argument for the statement concerning  $\mathfrak{C}(A)$  is similar.

Corollary 3.15. Let A be of type  $\mathfrak{A}$  (respectively, of type  $\mathfrak{B}$ , of type  $\mathfrak{C}$  or  $C^*$ -semifinite).

- (a) If B is a hereditary  $C^*$ -subalgebra of A, then B is of type  $\mathfrak{A}$  (respectively, of type  $\mathfrak{B}$ , of type  $\mathfrak{C}$  or  $C^*$ -semi-finite).
- (b) If A is a hereditary  $C^*$ -subalgebra of  $A_0$  that generates an essential ideal  $I \subseteq A_0$ , then  $A_0$  is of type  $\mathfrak{A}$  (respectively, of type  $\mathfrak{B}$ , of type  $\mathfrak{C}$  or  $C^*$ -semi-finite).
- **Proof:** (a) As any hereditary  $C^*$ -subalgebra of B is a hereditary  $C^*$ -subalgebra of A, this result follows directly from the definitions, Theorem 3.9(b) and Corollary 3.10(c).
- (b) Note that A is strongly Morita equivalent to I and any hereditary  $C^*$ -subalgebra of  $A_0$  intersects I non-trivially. Thus, this part follows from the definitions, Theorem 3.9 and Corollary 3.10.

Consequently, we have the following result.

Corollary 3.16. Suppose that A is non-unital, and  $\tilde{A}$  is the unitalization of A. Then A is of type  $\mathfrak{A}$  (respectively, of type  $\mathfrak{B}$ , of type  $\mathfrak{C}$  or  $C^*$ -semi-finite) if and only if  $\tilde{A}$  is of type  $\mathfrak{A}$  (respectively, of type  $\mathfrak{B}$ , of type  $\mathfrak{C}$  or  $C^*$ -semi-finite). The same is true when  $\tilde{A}$  is replaced by M(A).

Our next lemma is probably well-known, but we give a simple argument here for completeness.

**Lemma 3.17.** Let  $e, f \in OP(A)$  and  $p, q \in OP(A) \cap Z(A^{**})$ .

- (a)  $ep \in OP(A)$  and  $her(ep) = her(e) \cap her(p)$ .
- (b) If  $e \neq 0$  and  $her(e) \subseteq her(p) + her(q)$ , then  $her(e) \cap her(p) \neq (0)$  or  $her(e) \cap her(q) \neq (0)$ .
- (c) If z(e)z(f) = 0, then her(e) + her(f) = her(e+f).

**Proof:** Parts (a) and (c) are obvious (see Remark 2.2(d)). To show part (b), note that as  $her(p) + her(q) \subseteq her(p+q-pq)$ , we have  $e \le p+q-pq$ . If ep = 0 = eq, one obtains a contradiction that e = e(p+q-pq) = 0. Thus, the conclusion follows from part (a).

**Lemma 3.18.** If  $\{p_i\}_{i\in\Im}$  is a family in  $\mathrm{OP}_{\mathfrak{F}}(A)$  with  $z(p_i)z(p_j)=0$  for  $i\neq j$ , then  $p:=\sum_{i\in\Im}p_i\in\mathrm{OP}_{\mathfrak{F}}(A)$ .

**Proof:** It is clear that p is an open projection and  $z(p) = \sum_{i \in \mathfrak{I}} z(p_i)$ . Suppose that  $r, q \in \mathrm{OP}(\mathrm{her}(p))$  with  $r \leq q$  and  $r \sim_{\mathrm{sp}} q$ . Let  $u \in A^{**}$  with  $q = u^*u$  and  $u \cdot \mathrm{her}(q)u^* = \mathrm{her}(r)$ . For any  $i \in \mathfrak{I}$ , we set  $q_i := z(p_i)q, r_i := z(p_i)r \in \mathrm{OP}(A)$  and  $u_i := z(p_i)u$ . It is easy to see that  $q = \sum_{i \in \mathfrak{I}} q_i, r = \sum_{i \in \mathfrak{I}} r_i, q_i = u_i^*u_i$  and  $r_i \leq q_i \leq z(p_i)p = p_i$ . By Lemma 3.17(c), we see that

$$z(p_i) \operatorname{her}(q) \ = \ z(p_i) \big( \operatorname{her}(q_i) + \operatorname{her} \big( \sum\nolimits_{j \in \Im \backslash \{i\}} q_j \big) \big) \ = \ \operatorname{her}(q_i).$$

Similarly,  $z(p_i) \operatorname{her}(r) = \operatorname{her}(r_i)$  and we have  $u_i \operatorname{her}(q_i)u_i^* = \operatorname{her}(r_i)$ . By Proposition 2.7(a), we know that  $r_i \sim_{\operatorname{sp}} q_i$  and the  $C^*$ -finiteness of  $p_i$  tells us that  $r_i$  is dense in  $q_i$ . If  $e \in \operatorname{OP}(\operatorname{her}(q))$  with re = 0, then  $e_i := z(p_i)e \in \operatorname{OP}(\operatorname{her}(q_i))$  with  $r_ie_i = 0$ , which means that  $e_i = 0$  (because  $\overline{r_i}^{q_i} = q_i$ ). Consequently,  $e = \sum_{i \in \mathfrak{I}} e_i = 0$  and r is dense in q as required.

Part (a) of the following result is the equivalence of statements (i) and (iii) in [32, Theorem 2.3], while part (b) follows from the proof of [32, Theorem 2.3], Lemma 3.18, Theorem 3.9(a) and Corollary 3.15(b).

**Proposition 3.19.** (a) A  $C^*$ -algebra A is of type  $\mathfrak{A}$  if and only if there is an abelian hereditary  $C^*$ -subalgebra of A that generates an essential closed ideal of A. (b) A  $C^*$ -algebra A is  $C^*$ -semi-finite if and only if there is a  $C^*$ -finite hereditary  $C^*$ -subalgebra of A that generates an essential closed ideal of A.

#### 4. Comparison with existing theories

In this section, we compare our "Murray-von Neumann type classification" with existing results in the literature. Through these comparisons, we obtain many new examples of  $C^*$ -algebras of different types. Moreover, we will show that a von Neumann algebra is a type  $\mathfrak{A}$ , type  $\mathfrak{B}$ , type  $\mathfrak{C}$  or  $C^*$ -semi-finite  $C^*$ -algebra if and only if it is, respectively, a type I, type II, type III or semi-finiteness von Neumann algebra.

# 4.1. Comparison with type I algebras.

Recall that a  $C^*$ -algebra A is said to be of  $type\ I$  if for any irreducible representation  $(\pi, H)$  of A, one has  $\mathcal{K}(H) \subseteq \pi(A)$ . We have already seen in Theorem 3.9(b) that type  $\mathfrak{A}$  is the same as discreteness. Thus, the following result is a direct consequence of [32, Theorem 2.3]. Note that one can also obtain it using Theorem 3.9(a) and [6, Theorems 1.8 and 2.2].

# Corollary 4.1. Any type I $C^*$ -algebra is of type $\mathfrak{A}$ .

The converse of the above is not true even for real rank zero  $C^*$ -algebras, as can be seen in the following example.

Example 4.2. Example 3.6(c) and Corollary 3.15(b) tell us that  $\mathcal{B}(\ell^2)$  is of type  $\mathfrak{A}$ . However,  $\mathcal{B}(\ell^2)$  is not a type I  $C^*$ -algebra (see, e.g., [31, 6.1.2]).

**Proposition 4.3.** (a) A is of type I if and only if every primitive quotient of A is of type  $\mathfrak{A}$ .

- (b) If A is of type  $\mathfrak{A}$  and contains no essential primitive ideal, then A is of type I.
- **Proof:** (a) Because of Corollary 4.1 and the fact that quotients of type I  $C^*$ -algebras are also of type I, we only need to show the "if" part. Let  $\pi: A \to \mathcal{B}(H)$  be an irreducible representation and B be a non-zero abelian hereditary  $C^*$ -subalgebra of  $A/\ker \pi$ . If  $\tilde{\pi}: A/\ker \pi \to \mathcal{B}(H)$  is the induced representation, the restriction  $\tilde{\pi}_B: B \to \mathcal{B}(\tilde{\pi}(B)H)$  is non-zero and irreducible. Thus, dim  $\tilde{\pi}(B)H = 1$  and  $\tilde{\pi}(b)$  is a rank-one operator (and hence is compact) for any  $b \in B \setminus \{0\}$ . This shows that  $\tilde{\pi}(A/\ker \pi) \cap \mathcal{K}(H) \neq (0)$ , and  $\pi(A) \supseteq \mathcal{K}(H)$ .
- (b) Suppose that  $\pi: A \to \mathcal{B}(H)$  is an irreducible representation and J is a non-zero closed ideal of A with  $J \cap \ker \pi = (0)$ . If  $B \subseteq J$  is a non-zero abelian hereditary  $C^*$ -subalgebra, the restriction  $\pi_B: B \to \mathcal{B}(\pi(B)H)$  is non-zero and irreducible. The same argument as in part (a) tells us that  $\pi(A) \supseteq \mathcal{K}(H)$ .
- Remark 4.4. (a) Proposition 4.3(a) actually shows that A is of type I if and only if any primitive quotient contains a non-zero abelian hereditary  $C^*$ -subalgebra, which is likely to be a known fact.
- (b) If every quotient of  $\mathcal{B}(\ell^2)$  were of type  $\mathfrak{A}$ , then Proposition 4.3(a) told us that  $\mathcal{B}(\ell^2)$  were a type I  $C^*$ -algebra, which contradicted [31, 6.1.2]. Consequently, not every quotient of a type  $\mathfrak{A}$   $C^*$ -algebra is of type  $\mathfrak{A}$ .

If A is simple and of type  $\mathfrak{A}$ , then by Proposition 4.3(b), it is of type I. This, together with Example 3.6(c), gives the following.

Corollary 4.5. If A is a simple  $C^*$ -algebra of type  $\mathfrak{A}$ , then  $A = \mathcal{K}(H)$  for some Hilbert space H. If, in addition, A is  $C^*$ -finite, then  $A = M_n$  for some positive integer n.

# 4.2. Comparison with type II and (semi-)finite $C^*$ -algebras.

The following is a direct consequence of Remark 3.4(a) and Corollary 4.5.

Corollary 4.6. Any infinite dimensional  $C^*$ -finite simple  $C^*$ -algebra is of type  $\mathfrak{B}$ .

In the following, we compare type  $\mathfrak{B}$  and type  $\mathfrak{C}$  with the notions of type II and type III as introduced by Cuntz and Pedersen in [14]. Let us recall from [14, p. 140] that  $x \in A_+$  is said to be *finite* if for any sequence  $\{z_k\}_{k \in \mathbb{N}}$  in A with  $x = \sum_{k=1}^{\infty} z_k^* z_k$ , the condition  $\sum_{k=1}^{\infty} z_k z_k^* \leq x$  will imply  $x = \sum_{k=1}^{\infty} z_k z_k^*$ . We also recall that A is said to be *finite* (respectively, semi-finite) if every  $x \in A_+ \setminus \{0\}$  is finite (respectively, x dominates a non-zero finite element). Furthermore, A is said to be of type II if it is anti-liminary and finite, while A is said to be of type III if it has no non-zero finite elements (see [14, p. 149]).

Let  $T_s(A)$  be the set of all tracial states on A. It follows from [14, Theorem 3.4] that  $T_s(A)$  separates points of  $A_+$  if A is finite.

**Proposition 4.7.** If  $T_s(A)$  separates points of  $A_+$ , then A is  $C^*$ -finite. Consequently, if A is finite, then A is  $C^*$ -finite.

**Proof:** Suppose on the contrary that there exist  $r, q \in \mathrm{OP}(A)$  with  $r \leq q$ ,  $r \sim_{\mathrm{sp}} q$  but  $\bar{r}^q \leq q$ . For any  $\tau \in T_s(A)$ , if  $\tilde{\tau}$  is the normal tracial state on  $A^{**}$  extending  $\tau$ , then  $\tilde{\tau}(r) = \tilde{\tau}(q)$  (because  $r = vv^*$  and  $q = v^*v$  for some  $v \in A^{**}$ ). Moreover, if  $\{a_i\}_{i\in \mathfrak{I}}$  is an approximate unit in  $\mathrm{her}(r)$ , one has  $\tilde{\tau}(r) = \lim \tau(a_i)$ . Since  $\bar{r}^q \leq q$ , there exists  $s \in \mathrm{OP}(\mathrm{her}(q)) \setminus \{0\}$  with rs = 0. If  $x \in \mathrm{her}(s)_+$  with ||x|| = 1, one can find  $\tau_0 \in T_s(A)$  with  $\tau_0(x) > 0$ . Thus, we have  $\tau_0(a_i) + \tau_0(x) \leq \tilde{\tau}_0(q)$  (as  $a_i + x \leq q$  because  $a_i x = 0$ ), which gives the contradiction that  $\tilde{\tau}_0(r) + \tau_0(x) \leq \tilde{\tau}_0(q)$ .

As in [14], we denote by  $\mathscr{F}^A$  the set of all finite elements in  $A_+$ . If  $B \subseteq A$  is a hereditary  $C^*$ -subalgebra, then

$$\mathscr{F}^B = \mathscr{F}^A \cap B$$
.

In fact, it is obvious that  $\mathscr{F}^A \cap B \subseteq \mathscr{F}^B$ . Conversely, suppose that  $x \in \mathscr{F}^B$ . Consider  $y \in A_+$  and a sequence  $\{z_k\}_{k \in \mathbb{N}}$  in A satisfying  $y \leq x$ ,  $y = \sum_{k=1}^{\infty} z_k z_k^*$  and  $x = \sum_{k=1}^{\infty} z_k^* z_k$ . Since  $B_+$  is a hereditary cone of  $A_+$ , we have  $y \in B_+$  and  $z_k^* z_k, z_k z_k^* \in B_+$   $(k \in \mathbb{N})$ . By Remark 2.2(c), we know that  $z_k \in B$  and so, y = x as required.

Corollary 4.8. (a) A is semi-finite if and only if every non-zero hereditary  $C^*$ -subalgebra of A contains a non-zero finite hereditary  $C^*$ -subalgebra.

(b) If A is semi-finite (respectively, of type II), then A is  $C^*$ -semi-finite (respectively, of type  $\mathfrak{B}$ ).

**Proof:** (a) For the necessity, let  $B \subseteq A$  be a non-zero hereditary  $C^*$ -subalgebra. If  $y \in B_+ \setminus \{0\}$ , there is  $x \in \mathscr{F}^A \setminus \{0\}$  with  $x \leq y$ . By [14, Lemma 4.1] and [14, Theorem 4.8] as well as their arguments, one can find a non-zero finite hereditary  $C^*$ -subalgebra of her(x). More precisely, let  $f \in C(\sigma(x))_+$  such that f vanishes in a neighborhood of 0 and  $f(t) \leq t \leq f(t) + \frac{\|x\|}{2}$   $(t \in \sigma(x))$ . There exists  $g \in C(\sigma(x))_+$  and  $\lambda > 0$  such that f = fg and  $g(t) < \lambda t$   $(t \in \sigma(x))$ . Then  $g(x) \in \mathscr{F}^A$  and f(x) = f(x)g(x), i.e.,

$$f(x) \in \mathscr{F}_0 := \{a \in A_+ : a = ay \text{ for some } y \in \mathscr{F}^A\} \subseteq \mathscr{F}^A.$$

For any  $z \in \operatorname{her}(f(x))_+$ , we have zg(x) = z and  $z \in \mathscr{F}_0 \cap \operatorname{her}(f(x)) \subseteq \mathscr{F}^A \cap \operatorname{her}(f(x)) = \mathscr{F}^{\operatorname{her}(f(x))}$ . Thus,  $\operatorname{her}(f(x))$  is a non-zero finite hereditary  $C^*$ -subalgebra of  $\operatorname{her}(x)$ .

For the sufficiency, let  $y \in A_+ \setminus \{0\}$  and C be a non-zero finite hereditary  $C^*$ -subalgebra of her(y). Observe that  $C_+ = \mathscr{F}^C = \mathscr{F}^A \cap C$ . Take any  $x \in C_+$  with ||x|| = 1. Since  $x^{1/2}yx^{1/2} \leq ||y||x \in \mathscr{F}^A$ , we know, from [14, Lemma 4.1], that

$$y^{1/2}xy^{1/2} = y^{1/2}x^{1/2}(y^{1/2}x^{1/2})^* \in \mathscr{F}^A.$$

Moreover, as  $y^{1/2}xy^{1/2} \leq y$ , we see that A is semi-finite.

(b) This follows from part (a), Proposition 4.7 and Corollary 3.10(c).

Example 4.9. (a) If A is an infinite dimensional simple  $C^*$ -algebra with a faithful tracial state, then A is of type  $\mathfrak{B}$  (by Corollary 4.6 and Proposition 4.7). In particular, if  $\Gamma$  is an infinite discrete group such that  $C_r^*(\Gamma)$  is simple (see, e.g., [7] for some examples of such groups), then  $C_r^*(\Gamma)$  is of type  $\mathfrak{B}$ .

(b) Every simple AF algebra which is not of the form  $\mathcal{K}(H)$  is of type  $\mathfrak{B}$  (because of [14, Proposition 4.11] as well as Corollaries 4.5 and 4.8(b)).

# 4.3. Comparison with type III and purely infinite $C^*$ -algebras.

If a  $C^*$ -algebra A contains a non-zero (positive) finite element x, the argument of the necessity of Corollary 4.8(a) tells us that there is a non-zero finite hereditary  $C^*$ -subalgebra of A, and hence A is not of type  $\mathfrak{C}$ , because of Proposition 4.7. This gives the following corollary.

Corollary 4.10. If A is of type  $\mathfrak{C}$ , then it is of type III.

In the following, we will also compare type  $\mathfrak{C}$  with the notion of pure infinity as defined by Cuntz (in the case of simple  $C^*$ -algebras) and by Kirchberg and Rørdam (in the general case). Suppose that  $a \in M_n(A)$  and  $b \in M_m(A)$   $(m, n \in \mathbb{N})$ . As in [21, Definition 2.1], we say that  $a \preceq b$  relative to  $M_{m,n}(A)$  if there is a sequence  $\{x_k\}_{k\in\mathbb{N}}$  in  $M_{m,n}(A)$  such that  $\|x_k^*bx_k - a\| \to 0$ . An element  $a \in A$  is said to be properly infinite if  $a \oplus a \preceq a$  relative to  $M_{1,2}(A)$ . Moreover, A is said to be purely infinite if every element in  $A_+$  is properly infinite (see [21, Theorem 4.16]). Note that if A is simple, this notion coincides with the one in [13], namely, every hereditary  $C^*$ -subalgebra of A contains a non-zero infinite projection (see, e.g., the work of Lin and Zhang in [24]).

**Proposition 4.11.** (a) If A has real rank zero and is purely infinite, then it is of type  $\mathfrak{C}$ .

(b) If A is a separable purely infinite  $C^*$ -algebra with stable rank one, then A is of type  $\mathfrak{C}$ .

**Proof:** (a) By [21, Theorem 4.16], any element  $p \in \operatorname{Proj}(A) \setminus \{0\}$  is properly infinite and hence is infinite, in the sense that there exist  $q \in \operatorname{Proj}(A)$  and  $v \in A$  such that  $q \leq p$ ,  $v^*v = p$  and  $q = vv^*$  (see, e.g., [21, Lemma 3.1]). Thus,  $p \sim_{\operatorname{sp}} q$  (as  $v \in A$ ) but q is not dense in p (because  $p - q \in \operatorname{Proj}(A) \setminus \{0\}$ ). Consequently, any non-zero projection in A is not  $C^*$ -finite, and Corollary 3.12(c) shows that A is of type  $\mathfrak{C}$ .

(b) Suppose on contrary that A contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra B and we take any  $z \in B_+$  with ||z|| = 1. By [21, Theorem 4.16], one has  $z \oplus z \lesssim z \oplus 0$  relative to  $M_2(A)$ , and so,  $z \oplus z \lesssim z \oplus 0$  relative to  $M_2(\text{her}(z))$  (by [21, Lemma 2.2(iii)]). Thus, [29, Proposition 4.13] implies

$$p_z \oplus p_z = p_{z \oplus z} \lesssim_{\text{Cu}} p_{z \oplus 0} = p_z \oplus 0$$

(see [29, §3] for the meaning of  $\preceq_{\text{Cu}}$ ). Moreover, one obviously has  $p_{z\oplus 0} \preceq_{\text{Cu}} p_{z\oplus z}$ . Since A has stable rank one, we conclude that  $p_z \oplus p_z \sim_{\text{PZ}} p_z \oplus 0$  (by [29, 6.2(1)'&(2)']) and hence  $p_z \oplus p_z \sim_{\text{sp}} p_z \oplus 0$ . This means that  $M_2(\text{her}(z))$  is spatially isomorphic (and hence \*-isomorphic) to its hereditary  $C^*$ -subalgebra her $(z) \oplus (0)$ , which is not essential in  $M_2(\text{her}(z))$  (because  $(0) \oplus \text{her}(z)$  is a non-zero hereditary  $C^*$ -subalgebra and we can apply Remark 3.2(d)). As her(z) is \*-isomorphic to  $\text{her}(z) \oplus (0)$  and hence to  $M_2(\text{her}(z))$ , we know that her(z) is also spatially isomorphic to an inessential hereditary  $C^*$ -subalgebra. Consequently, her(z) is not  $C^*$ -finite, which contradicts the fact that B is  $C^*$ -finite.

One may regard parts (a) and (b) of the above as two extremes, because any real rank zero  $C^*$ -algebras has plenty of projections, while a purely infinite  $C^*$ -algebra with stable rank one is stably projectionless. Let us make the following conjecture.

Conjecture 4.12. Every purely infinite  $C^*$ -algebra is of type  $\mathfrak{C}$ .

On the other hand, by Proposition 4.11 and Corollary 4.10, we know that any separable purely infinite  $C^*$ -algebra A having real rank zero or stable rank one is of type III. This implication actually holds without these extra assumptions, as can be seen in the following proposition, which gives another evidence for Conjecture 4.12. Note that this proposition also implies [21, Proposition 4.4]. To show this result, let us recall the following notation from [29, p. 3476]. For any  $\epsilon > 0$ , let  $f_{\epsilon} : \mathbb{R}_+ \to \mathbb{R}_+$  be the function

$$f_{\epsilon}(t) = \begin{cases} t/\epsilon & \text{if } t \in [0, \epsilon) \\ 1 & \text{if } t \in [\epsilon, \infty). \end{cases}$$

If  $\mu \in T_s(A)$  and  $a \in A_+$ , we define

$$d_{\mu}(a) := \sup_{\epsilon > 0} \mu(f_{\epsilon}(a))$$

(note that the definition in [29] is for tracial weights but we only need tracial states here).

**Proposition 4.13.** Any purely infinite  $C^*$ -algebra A is of type III.

**Proof:** Suppose on the contrary that  $\mathscr{F}^A \neq \{0\}$ . By the argument of the necessity of Corollary 4.8(a), there is  $z \in A_+$  with ||z|| = 1 and  $\operatorname{her}(z)$  being a finite  $C^*$ -algebra. By the argument of Proposition 4.11(b), one has  $z \oplus z \preceq z \oplus 0$  relative to  $M_2(\operatorname{her}(z))$ . By [29, Remark 2.5], we see that  $d_{\mu}(z \oplus z) \leq d_{\mu}(z \oplus 0)$  for each  $\mu \in T_s(M_2(\operatorname{her}(z)))$ . Now, if  $\tau \in T_s(\operatorname{her}(z))$ , then  $\tau \otimes \operatorname{Tr}_2 \in T_s(M_2(\operatorname{her}(z)))$  (where  $\operatorname{Tr}_2$  is the canonical tracial state on  $M_2$ ), and the above tells us that

$$\sup_{\epsilon>0} \tau(f_{\epsilon}(z)) = \sup_{\epsilon>0} (\tau \otimes \operatorname{Tr}_2)(f_{\epsilon}(z) \oplus f_{\epsilon}(z)) \leq \sup_{\epsilon>0} (\tau \otimes \operatorname{Tr}_2)(f_{\epsilon}(z) \oplus 0) = \sup_{\epsilon>0} \frac{\tau(f_{\epsilon}(z))}{2},$$
which gives  $d_{\tau}(z) = 0$  and hence  $\tau(z) = 0$ . This contradicts [14, Theorem 3.4].  $\square$ 

If one can show that her(a) is not  $C^*$ -finite, for every properly infinite positive element a in any  $C^*$ -algebra, then the above conjecture is verified. Let us recall from [21, Proposition 3.3(iv)] that  $a \in A_+$  is properly infinite if and only if there are sequences  $\{x_n\}_{n\in\mathbb{N}}$  and  $\{y_n\}_{n\in\mathbb{N}}$  in her(a) such that  $x_n^*x_n \to a$ ,  $y_ny_n^* \to a$  and  $x_n^*y_n \to 0$ . The following remark tells us that if  $a \in A_+$  satisfies a stronger condition than the above, then her(a) is indeed non- $C^*$ -finite.

Remark 4.14. Let  $a \in A_+$  such that there exist  $x, y \in \text{her}(a)$  with  $x^*x = a = y^*y$  as well as  $x^*y = 0$ . By Example 2.9(a)&(b), we see that her(a) is spatially isomorphic to its hereditary  $C^*$ -subalgebra  $\text{her}(x^*)$ . As  $\text{her}(x^*) \text{her}(y^*) = (0)$ , we see that  $\text{her}(x^*)$  is not essential in her(a). Thus, her(a) is not  $C^*$ -finite.

Example 4.15. For any AF-algebra B, the  $C^*$ -algebra  $\mathcal{O}_2 \otimes B$  is purely infinite (by [21, Proposition 4.5]) and is of real rank zero (by [12, Theorem 3.2]), which means that  $\mathcal{O}_2 \otimes B$  is of type  $\mathfrak{C}$  (by Proposition 4.11(a)). Note that one may replace  $\mathcal{O}_2$ 

with any unital, simple, separable, purely infinite, nuclear  $C^*$ -algebra (which has real rank zero because of [40, Theorem 1.2(ii)]).

# 4.4. The case of von Neumann algebras.

In this subsection, we consider the case of von Neumann algebras. Let us start with the following lemma. Note that the necessity of part (a) of this result follows directly from Proposition 4.7, but we give an alternative proof here as this argument is also interesting (see Remark 4.17 below).

**Lemma 4.16.** (a) Let M be a von Neumann algebra. Then  $p \in \text{Proj}(M)$  is finite as a projection in M if and only if it is  $C^*$ -finite.

(b) The ideal  $\mathfrak{F}(M)$  in Proposition 3.5 is a dense subalgebra of the ideal J(M) generated by finite projections (as defined in [19]).

**Proof:** (a) Assume that p is finite. Let  $\Lambda_M: M^{**} \to M$  be the canonical \*-epimorphism. If  $q \in \mathrm{OP}(pMp)$ , then  $\mathrm{her}_M(q) \subseteq \mathrm{her}_M(\Lambda_M(q))$  and  $\Lambda_M(q) \leq p$ , which imply that  $\Lambda_M(q) = \bar{q}^p$  (notice that  $\bar{q}^p \in pMp$  because of [2, Theorem II.1]).

Suppose that  $r, q \in \mathrm{OP}(pMp)$  such that  $r \leq q$  and  $r \sim_{\mathrm{sp}} q$ . Consider  $w \in M^{**}$  satisfying

$$q = ww^*$$
,  $r = w^*w$ ,  $w^* \operatorname{her}(q)w = \operatorname{her}(r)$  and  $w \operatorname{her}(r)w^* = \operatorname{her}(q)$ .

Define  $v := \Lambda_M(w)$ . Then  $\Lambda_M(q) = vv^*$  and  $\Lambda_M(r) = v^*v$ . Since  $\Lambda_M(r) \le \Lambda_M(q) \le p$ , the finiteness of p tells us that  $\bar{r}^p = \Lambda_M(r) = \Lambda_M(q) = \bar{q}^p$ . If  $\bar{r}^q \le q$ , there is  $e \in \mathrm{OP}(\mathrm{her}(q)) \setminus \{0\}$  with re = 0. Since  $e \in \mathrm{OP}(\mathrm{her}(p))$ , we obtain a contradiction that  $\bar{r}^p \ne \bar{q}^p$  (as  $r \le p - e$  but  $q \not\le p - e$ ). This shows that p is  $C^*$ -finite.

Conversely, if p is  $C^*$ -finite, then Remark 3.13 implies that p is finite.

(b) This follows from part (a) and Corollary 3.14.

Remark 4.17. (a) Let  $p \in M$  be a finite projection. If  $r \in \text{Proj}(pMp)$  with  $r \sim_{\text{sp}} p$ , then Lemma 4.16(a) and Remark 3.13 tell us that r = p. The same is true if we relax the assumption to  $r \in \text{OP}(pMp)$ . In fact, we first notice that the  $C^*$ -finiteness of p gives  $\bar{r}^p = p$ . Moreover, suppose that  $w \in M^{**}$  and  $v \in M$  are as in the proof of Lemma 4.16 for the case when q = p. Then  $vv^* = p = \bar{r}^p = v^*v$ . This means that v is a unitary in pMp. As  $v \ker(r)v^* = \Lambda_M(w \ker(r)w^*) = pMp$ , we have  $\ker(r) = pMp$  and hence r = p.

(b) If A is a  $C^*$ -algebra and  $p \in \mathrm{OP}(A)$  satisfying  $\bar{r}^p = \bar{q}^p$  for any  $r, q \in \mathrm{OP}(\mathrm{her}(p))$  with  $r \leq q$  and  $r \sim_{\mathrm{sp}} q$ , then by the argument of Lemma 4.16, we see that p is  $C^*$ -finite.

The following is a direct consequence of Lemma 4.16 and Corollary 3.12.

**Theorem 4.18.** Let M be a von Neumann algebra.

- (a) M is of type  $\mathfrak{A}$  if and only if M is a type I von Neumann algebra.
- (b) M is of type  $\mathfrak{B}$  if and only if M is a type II von Neumann algebra.
- (c) M is of type  $\mathfrak{C}$  if and only if M is a type III von Neumann algebra.
- (d) M is C\*-semi-finite if and only if M is a semi-finite von Neumann algebra.

#### 5. Factorisations

In this section, we give two factorization type results for general  $C^*$ -algebras. Let us first state the following easy lemma. Notice that if A contains a non-zero abelian hereditary  $C^*$ -subalgebra B, the closed ideal generated by B is of type  $\mathfrak{A}$  (by Corollary 3.15(b) and Remark 3.7(b)), and the same is true for  $C^*$ -finite hereditary  $C^*$ -subalgebra.

**Lemma 5.1.** If A is not of type  $\mathfrak{C}$ , then A contains a non-zero closed ideal of either type  $\mathfrak{A}$  or type  $\mathfrak{B}$ .

The following is our first factorization type result, which mimics the corresponding situation for von Neumann algebras.

**Theorem 5.2.** Let A be a  $C^*$ -algebra.

- (a) There is a largest type  $\mathfrak{A}$  (respectively, type  $\mathfrak{B}$ , type  $\mathfrak{C}$  and  $C^*$ -semi-finite) hereditary  $C^*$ -subalgebra  $J_{\mathfrak{A}}$  (respectively,  $J_{\mathfrak{B}}$ ,  $J_{\mathfrak{C}}$  and  $J_{\mathfrak{sf}}$ ) of A, which is also an ideal of A.
- (b)  $J_{\mathfrak{A}}$ ,  $J_{\mathfrak{B}}$  and  $J_{\mathfrak{C}}$  are mutually disjoint such that  $J_{\mathfrak{A}} + J_{\mathfrak{B}} + J_{\mathfrak{C}}$  is an essential closed ideal of A. If  $e_{\mathfrak{A}}, e_{\mathfrak{B}}, e_{\mathfrak{C}} \in \mathrm{OP}(A) \cap Z(A^{**})$  with  $J_{\mathfrak{A}} = \mathrm{her}(e_{\mathfrak{A}}), J_{\mathfrak{B}} = \mathrm{her}(e_{\mathfrak{B}})$  and  $J_{\mathfrak{C}} = \mathrm{her}(e_{\mathfrak{C}})$ , then

$$1 = \overline{e_{\mathfrak{A}} + e_{\mathfrak{B}}}^{1} + e_{\mathfrak{C}}.$$

(c)  $J_{\mathfrak{A}} + J_{\mathfrak{B}}$  is an essential closed ideal of  $J_{\mathfrak{sf}}$ . If  $e_{\mathfrak{sf}} \in \mathrm{OP}(A)$  with  $J_{\mathfrak{sf}} = \mathrm{her}(e_{\mathfrak{sf}})$ , then

$$e_{\mathfrak{sf}} = \overline{e_{\mathfrak{A}}}^{e_{\mathfrak{sf}}} + e_{\mathfrak{B}}.$$

- (d) The closure of C(A) and F(A) (in Proposition 3.5) are essential closed ideals of  $J_{\mathfrak{A}}$  and  $J_{\mathfrak{sf}}$ , respectively.
- **Proof:** (a) We first consider the situation of type  $\mathfrak{B}$  hereditary  $C^*$ -subalgebra. Let  $\mathcal{J}_{\mathfrak{B}}$  be the set of all type  $\mathfrak{B}$  closed ideals of A. If  $\mathcal{J}_{\mathfrak{B}} = \{(0)\}$ , then  $J_{\mathfrak{B}} := (0)$  is the largest type  $\mathfrak{B}$  hereditary  $C^*$ -subalgebra of A (see Remark 3.11(b)). Suppose that there exist distinct elements  $J_1$  and  $J_2$  in  $\mathcal{J}_{\mathfrak{B}}$ . If  $J_1 + J_2$  contains a non-zero abelian hereditary  $C^*$ -algebra B, then by Lemma 3.17(b), one of the two abelian hereditary  $C^*$ -subalgebras  $B \cap J_1$  and  $B \cap J_2$  is non-zero, which contradicts  $J_1, J_2 \in \mathcal{J}_{\mathfrak{B}}$ . On the other hand, consider a non-zero closed ideal I of  $J_1 + J_2$ . Again, by Lemma 3.17(b), we may assume that the closed ideal  $I \cap J_1$  is non-zero. Thus,  $I \cap J_1$  contains a

non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra B. This shows that  $J_1 + J_2 \in \mathcal{J}_{\mathfrak{B}}$  and  $\mathcal{J}_{\mathfrak{B}}$  is a directed set.

For any ideal J of A, we consider  $e_J \in \mathrm{OP}(A) \cap \mathrm{Z}(A^{**})$  with  $J = \mathrm{her}(e_J)$ . Set

$$J_{\mathfrak{B}} := \overline{\sum_{J \in \mathcal{J}_{\mathfrak{B}}} J}.$$

Then  $e_{J_{\mathfrak{B}}} = w^*$ - $\lim_{J \in \mathcal{J}_{\mathfrak{B}}} e_J$ . If there is  $p \in \mathrm{OP}_{\mathfrak{C}}(A) \setminus \{0\}$  such that  $\mathrm{her}(p) \subseteq J_{\mathfrak{B}}$ , then

$$p = pe_{J_{\mathfrak{B}}} = pe_{J_{\mathfrak{B}}}p = w^* - \lim_{J \in \mathcal{J}_{\mathfrak{B}}} pe_J p,$$

and one can find  $J \in \mathcal{J}_{\mathfrak{B}}$  with the abelian algebra  $\ker(p) \cap J$  being non-zero (because of Lemma 3.17(a)), which is absurd. On the other hand, suppose that I is a non-zero closed ideal of  $J_{\mathfrak{B}}$ . The argument above tells us that  $I \cap J \neq (0)$  for some  $J \in \mathcal{J}_{\mathfrak{B}}$ , and hence it contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra. Consequently,  $J_{\mathfrak{B}} \in \mathcal{J}_{\mathfrak{B}}$ . Finally, if  $B \subseteq A$  is a hereditary  $C^*$ -subalgebra of type  $\mathfrak{B}$ , then, by Remark 3.11(b), one has  $B \subseteq J_{\mathfrak{B}}$ .

The arguments for the statements concerning  $J_{\mathfrak{A}}$ ,  $J_{\mathfrak{C}}$  and  $J_{\mathfrak{sf}}$  are similar and easier.

- (b) The first statement follows directly from Lemma 5.1 (any non-type  $\mathfrak{C}$  ideal interests either  $J_{\mathfrak{A}}$  or  $J_{\mathfrak{B}}$ ). For the second statement, one obviously has  $e_{\mathfrak{A}} + e_{\mathfrak{B}} \leq 1 e_{\mathfrak{C}}$ . Suppose that  $p \in \mathrm{OP}(A)$  with  $e_{\mathfrak{A}} + e_{\mathfrak{B}} \leq 1 p$ . We have  $p(e_{\mathfrak{A}} + e_{\mathfrak{B}}) = 0$ . If  $p \nleq e_{\mathfrak{C}}$ , then her(p) will contain a hereditary  $C^*$ -subalgebra of either type  $\mathfrak{A}$  or type  $\mathfrak{B}$  (by Lemma 5.1) and Lemma 3.17(a) will give a contradiction that either  $pe_{\mathfrak{A}} \neq 0$  or  $pe_{\mathfrak{B}} \neq 0$ . Thus,  $1 e_{\mathfrak{C}}$  is the smallest closed projection dominating  $e_{\mathfrak{A}} + e_{\mathfrak{B}}$ .
- (c) This follows from a similar (but easier) argument as part (b).
- (d) Clearly,  $\mathcal{F}(A) \subseteq J_{\mathfrak{sf}}$  and  $\mathcal{C}(A) \subseteq J_{\mathfrak{A}}$  (see Remark 3.11(b)). Their closure are both essential because of Proposition 3.19.

By Proposition 3.19, there is an abelian (respectively, a  $C^*$ -finite) hereditary  $C^*$ -subalgebra that generates an essential ideal of  $J_{\mathfrak{A}}$  (respectively, of  $J_{\mathfrak{B}}$ ). Moreover, by [32, Theorem 2.3(vi)], the largest type I closed ideal  $A_{postlim}$  of A is an essential ideal of  $J_{\mathfrak{A}}$ .

Remark 5.3. For any closed ideal J of A, we write  $J^{\perp}$  for the closed ideal  $\{a \in A : aJ = (0)\}$ . It is easy to see that if  $J_0$  is an essential ideal of J, then  $J_0^{\perp} = J^{\perp}$ .

- (a)  $J_{\mathfrak{A}}^{\perp} = A_{postlim}^{\perp}$  is the largest anti-liminary hereditary  $C^*$ -subalgebra of A (note that  $\overline{aJ_{\mathfrak{A}}a}$  is a hereditary  $C^*$ -subalgebra of  $J_{\mathfrak{A}}$  for every  $a \in A_+$ ). Furthermore,  $J_{\mathfrak{B}} + J_{\mathfrak{C}}$  is an essential ideal of  $J_{\mathfrak{A}}^{\perp}$  (by Lemma 5.1).
- (b)  $J_{\mathfrak{sf}}^{\perp} = (J_{\mathfrak{A}} + J_{\mathfrak{B}})^{\perp} = J_{\mathfrak{C}}.$
- (c)  $J_{\mathfrak{A}}^{\perp} \cap J_{\mathfrak{sf}} = J_{\mathfrak{B}}$  (compare with Corollary 3.10(c)).

From now on, we denote by  $J_{\mathfrak{A}}^{A}$ ,  $J_{\mathfrak{B}}^{A}$ ,  $J_{\mathfrak{C}}^{A}$  and  $J_{\mathfrak{sf}}^{A}$ , respectively, the largest type  $\mathfrak{A}$ , the largest type  $\mathfrak{C}$  and the largest  $C^*$ -semi-finite closed ideals of a  $C^*$ -algebra A.

The following is a direct application of Theorem 4.18.

**Corollary 5.4.** Let M be a von Neumann algebra. If  $M_I$ ,  $M_{II}$  and  $M_{III}$  are respectively the type I summand, the type II summand and the type III summand of M, then  $J_{\mathfrak{A}}^M = M_I$ ,  $J_{\mathfrak{B}}^M = M_{II}$  and  $J_{\mathfrak{C}}^M = M_{III}$ .

Our next theorem is the second factorization type result, which seems to be more interesting for  $C^*$ -algebra (c.f. [14, Proposition 4.13]).

**Theorem 5.5.** Let A be a  $C^*$ -algebra.

- (a)  $A/J_{\sigma}^{A}$  is  $C^{*}$ -semi-finite and  $A/(J_{\mathfrak{A}}^{A})^{\perp}$  is of type  $\mathfrak{A}$ .
- (b) If A is  $C^*$ -semi-finite, then  $A/J_{\mathfrak{B}}^A$  is of type  $\mathfrak{A}$ .

**Proof:** (a) Assume, without loss of generality, that  $A/J_{\mathfrak{C}}^A \neq (0)$  and consider  $Q: A \to A/J_{\mathfrak{C}}^A$  to be the canonical map. Let I be a non-zero closed ideal of  $A/J_{\mathfrak{C}}^A$  and  $J:=Q^{-1}(I)$ . Since  $J \supsetneq J_{\mathfrak{C}}^A$ , one knows that J contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra B. Since  $B \cap J_{\mathfrak{C}}^A = (0)$ , the \*-homomorphism Q restricts to an injection on B. Thus,  $Q(B) \subseteq I$  is also a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra, and  $A/J_{\mathfrak{C}}^A$  is  $C^*$ -semi-finite (by Corollary 3.10(a)). The proof of the second statement is similar.

(b) This follows from part (a) and Remark 5.3(c).
---

Remark 5.6. Let S be a statement concerning  $C^*$ -algebras that is stable under extensions of  $C^*$ -algebras (i.e. if I is a closed ideal of a  $C^*$ -algebra A such that S is true for both I and A/I, then S is true for A).

- (a) If S is true for all type  $\mathfrak{A}$  and all type  $\mathfrak{B}$   $C^*$ -algebras, S is true for all  $C^*$ -semifinite  $C^*$ -algebras. If, in addition, S is true for all type  $\mathfrak{C}$   $C^*$ -algebras, it is true for all  $C^*$ -algebras.
- (b) If S is true for all discrete  $C^*$ -algebras and all anti-liminary  $C^*$ -algebras, then S is true for all  $C^*$ -algebras.

The following results follows from Theorem 3.9(a).

Corollary 5.7. If A and B are strongly Morita equivalent, then the closed ideal of B that corresponds to  $J_{\mathfrak{A}}^{A}$  (respectively,  $J_{\mathfrak{B}}^{A}$ ,  $J_{\mathfrak{C}}^{A}$  and  $J_{\mathfrak{sf}}^{A}$ ) under the strong Morita equivalence (see the paragraph preceding Theorem 3.9) is precisely  $J_{\mathfrak{A}}^{B}$  (respectively,  $J_{\mathfrak{B}}^{B}$ ,  $J_{\mathfrak{C}}^{B}$  and  $J_{\mathfrak{sf}}^{B}$ ).

Remark 5.8. It is natural to ask if the closure  $\overline{\mathbb{C}(\cdot)}$  of  $\mathbb{C}(\cdot)$  (see Proposition 3.5) is also stable under strong Morita equivalence. Unfortunately, it is not the case. Suppose that A is any type I  $C^*$ -algebra. Then by [6, Theorems 1.8 and 2.2], there is a commutative  $C^*$ -algebra B that is strongly Morita equivalent to A. Notice that  $\mathbb{C}(B) = B$  and  $\overline{\mathbb{C}(A)}$  is of type I<sub>0</sub> (by [31, Proposition 6.1.7]). Thus, if  $\overline{\mathbb{C}(\cdot)}$  is stable under strong Morita equivalence, then any type I  $C^*$ -algebra A will coincide with  $\overline{\mathbb{C}(A)}$  and hence is liminary (see, e.g., [31, Corollary 6.1.6]), which is absurd.

To end this section, we compare  $J_*^A$  with  $J_*^{M(A)}$ .

**Proposition 5.9.** (a) If  $B \subseteq A$  is a hereditary  $C^*$ -subalgebra, then  $J_{\mathfrak{A}}^B = J_{\mathfrak{A}}^A \cap B$ ,  $J_{\mathfrak{B}}^B = J_{\mathfrak{C}}^A \cap B$  and  $J_{\mathfrak{sf}}^B = J_{\mathfrak{sf}}^A \cap B$ .

- (b)  $J_{\mathfrak{A}}^{M(A)} = \{x \in M(A) : xA \subseteq J_{\mathfrak{A}}^A\}$ . Similar statements hold for  $J_{\mathfrak{B}}$ ,  $J_{\mathfrak{C}}$  and  $J_{\mathfrak{sf}}$ .
- (c)  $J_{\mathfrak{B}}^{M(A)} = \{x \in M(A) : xJ_{\mathfrak{A}}^A = (0) \text{ and } xA \subseteq J_{\mathfrak{sf}}^A\}$
- $(d)\ J^{M(A)}_{\mathfrak{C}} = \{x \in M(A) : xJ^{A}_{\mathfrak{sf}} = (0)\} = \{x \in M(A) : xJ^{A}_{\mathfrak{A}} = (0)\ and\ xJ^{A}_{\mathfrak{B}} = (0)\}.$

**Proof:** (a) Clearly,  $J_{\mathfrak{A}}^{B} \subseteq B \cap J_{\mathfrak{A}}^{A}$ . Conversely, since  $B \cap J_{\mathfrak{A}}^{A}$  is a type  $\mathfrak{A}$  closed ideal of B (by Corollary 3.15(a)), we have  $B \cap J_{\mathfrak{A}}^{A} \subseteq J_{\mathfrak{A}}^{B}$ . The other cases follow from similar arguments.

(b) We will only consider the case of  $J_{\mathfrak{B}}$  (since the other cases follow from similar and easier arguments). Notice that  $J_{\mathfrak{B}}^{M(A)} \cdot A = J_{\mathfrak{B}}^{M(A)} \cap A = J_{\mathfrak{B}}^{A}$  (by part (a)) and

$$J_{\mathfrak{B}}^{M(A)} \subseteq J_0 := \{ x \in M(A) : xA \subseteq J_{\mathfrak{B}}^A \}.$$

Suppose that the closed ideal  $J_0 \subseteq M(A)$  contains a non-zero abelian hereditary  $C^*$ -subalgebra B. The abelian hereditary  $C^*$ -subalgebra  $B \cap A = B \cdot A \cdot B$  is contained in  $J_{\mathfrak{B}}^A$  and so,  $B \cdot A = (0)$ , which contradicts the fact that A is essential in M(A) (see Remark 3.2(d)). Furthermore, let I be a non-zero closed ideal of  $J_0$ . Then  $I \cdot A = I \cap A \neq (0)$  and is a closed ideal of  $J_{\mathfrak{B}}^A$ . Thus,  $I \cap A$  contains a non-zero  $C^*$ -finite hereditary  $C^*$ -subalgebra. Consequently,  $J_0$  is of type  $\mathfrak{B}$  and is a subset of  $J_{\mathfrak{B}}^{M(A)}$ .

- (c) Obviously,  $xJ_{\mathfrak{A}}^{A}=(0)$  if and only if  $xAJ_{\mathfrak{A}}^{A}=(0)$ . Thus, this part follows from part (b) and Remark 5.3(c).
- (d) This part follows from a similar argument as part (c) as well as Remark 5.3(b).  $\Box$

## References

- [1] C. A. Akemann, The General Stone-Weierstrass problem, J. Funct. Anal., 4 (1969), 277-294.
- [2] C. A. Akemann, Left ideal structure of  $C^*$ -algebras, J. Funct. Anal., 6 (1970), 305-317.
- [3] C. A. Akemann and S. Eilers, Regularity of projections revisited, J. Oper. Theory, 48 (2002), 515-534.

- [4] C. A. Akemann and G. K. Pedersen, Complications of semicontinuity in C\*-algebra theory, Duke Math. J., 40 (1973), 785-795.
- [5] C. A. Akemann, G. K. Pedersen and J. Tomiyama, Multipliers of  $C^*$ -algebras, J. Funct. Anal., 13 (1973), 277-301.
- [6] W. Beer, On Morita equivalence of nuclear C\*-algebras, J. Pure Appl. Alg., 26 (1982), 249-267.
- [7] M. Bekka, M. Cowling and P. de la Harpe, Some groups whose reduced  $C^*$ -aglebra is simple, Publ. Math. I.H.E.S., **80** (1994), 117-134.
- [8] B. Blackadar, Operator algebras theory of C\*-algebras and von Neumann algebras, in Operator Algebras and Non-commutative Geometry III, Encyc. Math. Sci., 122, Springer-Verlag, Berlin (2006).
- [9] L. G. Brown, Semicontinuity and multipliers of  $C^*$ -algebras, Canad. J. Math., **XL**, **4** (1988), 865-988.
- [10] L. G. Brown, Determination of A from M(A) and related matters, C. R. Math. Rep. Acad. Sci. Canada, **10** (1988), 273-278.
- [11] L. G. Brown and M. A. Rieffel, Stable isomorphism and strong Morita equivalence of C\*-algebras, Pacific J. Math., **71** (1977), 349-363.
- [12] L. G. Brown and G. K. Pedersen,  $C^*$ -algebras of real rank zero, J. Funct. Anal., **99** (1991), 131-149.
- [13] J. Cuntz, K-theory for certain C\*-algebras, Ann. of Math., **113** (1981), 181-197.
- [14] J. Cuntz and G. K. Pedersen, Equivalence and traces on  $C^*$ -algebras, J. Funct. Anal., **33** (1979), 135-164.
- [15] E. G. Effros, Order ideals in  $C^*$ -algebras and its dual, Duke Math. J., **30** (1963), 391-412.
- [16] G. A. Elliott, On the classification of inductive limits of sequences of semisimple finite-dimensional algebras, J. Alg. **38** (1976), 29-44.
- [17] G. A. Elliott, On the classification of  $C^*$ -algebras of real rank zero, J. Reine Angew. Math., 443 (1993), 179-219.
- [18] G. A. Elliott and A. S. Toms, Regularity properties in the classification program for separable amenable  $C^*$ -algebras, Bull. Amer. Math. Soc., **45** (2008), 229 245.
- [19] H. Halpern, V. Kaftal, P. W. Ng and S. Zhang, Finite sums of projections in von Neumann algebras, preprint (arXiv:1007.4679v1).
- [20] R. V. Kadison and J. R. Ringrose, Fundamentals of the theory of operator algebras Vol. II: Advanced theory, Pure and Applied Mathematics 100, Academic Press (1986).
- [21] E. Kirchberg and M. Rørdam, Non-simple purely infinite  $C^*$ -algebras, Amer. J. Math., **122** (2000), 637-666.
- [22] E. C. Lance, *Hilbert C\*-modules A toolkit for operator algebraists*, Lond. Math. Soc. Lect. Note Ser. **210**, Camb. Univ. Press (1995).
- [23] H. Lin, Equivalent open projections and corresponding hereditary  $C^*$ -subalgebras, J. Lond. Math. Soc., **41** (1990), 295-301.
- [24] H. Lin and S. Zhang, On infinite simple  $C^*$ -algebras, J. Funct. Anal., 100 (1991), 221-231.
- [25] G.J. Murphy, C\*-algebras and operator theory, Academic Press (1990).
- [26] F. J. Murray, The rings of operators papers, in *The legacy of John von Neumann* (Hempstead, NY, 1988), 57-60, Proc. Sympos. Pure Math. **50**, Amer. Math. Soc., Providence, R.I. (1990).
- [27] F. J. Murray and J. Von Neumann, On rings of operators, Ann. of Math. (2), 37 (1936), 116-229.
- [28] C.K. Ng and N.C. Wong, Comparisons of equivalence relations on open projections, preprint.
- [29] E. Ortega, M. Rørdam and H. Thiel, The Cuntz semigroup and comparison of open projections, J. Funct. Anal., 260 (2011), 3474-3493.
- [30] G. K. Pedersen, Applications of weak-\*-semicontinuity in  $C^*$ -algebra theory, Duke Math. J., **39** (1972), 431-450.

- [31] G. K. Pedersen, C\*-algebras and their automorphism groups, Academic Press (1979).
- [32] C. Peligrad and L. Zsidó, Open projections of C\*-algebras: Comparison and Regularity, Operator Theoretical Methods, 17th Int. Conf. on Operator Theory, Timisoara (Romania), June 23-26, 1998, Theta Found. Bucharest (2000), 285-300
- [33] R. T. Prosser, On the ideal structure of operator algebras, Memoirs Amer. Math. Soc., **45** (1963).
- [34] M. A. Rieffel, Morita equivalence for  $C^*$ -algebras and  $W^*$ -algebras, J. Pure Appl. Alg., 5 (1974), 51-96.
- [35] M. A. Rieffel, Morita equivalence for operator algebras, in *Operator algebras and applications*, Part I (Kingston, Ont., 1980), Proc. Sympos. Pure Math. 38, Amer. Math. Soc., Providence, R.I. (1982), 285 298.
- [36] M. Rørdam, Classification of nuclear, simple C\*-algebras, in Classification of nuclear C\*-algebras, Entropy in operator algebras, Encyclopaedia Math. Sci. 126, Springer, Berlin (2002), 1-145.
- [37] M. Takesaki, Theory of Operator algebras I, Springer-Verlag New York (1979).
- [38] A. S. Toms, On the classification problem for nuclear  $C^*$ -algebras, Ann. of Math. (2), 167 (2008), 1029-1044.
- [39] S. Zhang, Stable isomorphism of hereditary  $C^*$ -subalgebras and stable equivalence of open projections, Proc. Amer. Math. Soc., **105** (1989), 677-682.
- [40] S. Zhang, Certain  $C^*$ -algebras with real rank zero and their corona and multiplier algebras part I, Pac. J. Math. **155** (1992), 169 197.

(Chi-Keung Ng) Chern Institute of Mathematics and LPMC, Nankai University, Tianjin 300071, China.

E-mail address: ckng@nankai.edu.cn

(Ngai-Ching Wong) Department of Applied Mathematics, National Sun Yat-sen University, Kaohsiung, 80424, Taiwan.

E-mail address: wong@math.nsysu.edu.tw