

# The Skitovich-Darmois theorem for discrete and compact totally disconnected Abelian groups

I.P. Mazur

November 9, 2018

## 1 Introduction

The classical Skitovich-Darmois theorem states ([14],[1], see also [11, Ch. 3]): Let  $\xi_i, i = 1, 2, \dots, n, n \geq 2$ , be independent random variables, and  $\alpha_i, \beta_i$  be nonzero numbers. Suppose that the linear forms  $L_1 = \alpha_1\xi_1 + \dots + \alpha_n\xi_n$   $L_2 = \beta_1\xi_1 + \dots + \beta_n\xi_n$  are independent. Then all random variables  $\xi_i$  are Gaussian.

The Skitovich-Darmois theorem was generalized into various algebraic structures, in particular, into locally compact Abelian groups([3]-[9],[10],[12]). In these researches the random variables take values in a locally compact Abelian group  $X$ , coefficients of the linear forms are topological automorphisms of  $X$ , and the number of the linear forms are two. In the article we continue these researches and study the Skitovich-Darmois theorem in the case when random variables take values in different classes of locally compact Abelian groups but the number of linear forms more than 2.

Throughout the article  $X$  is a second countable locally compact Abelian group. Let  $Aut(X)$  be the group of topological automorphisms of  $X$ ,  $\mathbb{Z}(k) = \{0, 1, 2, \dots, k - 1\}$  be the group of residue modulo  $k$ . Let  $x \in X$ . Denote by  $E_x$  the degenerate distribution, concentrated at the point  $x$ . If  $K$  is a compact subgroup of  $X$ , denote by  $m_K$  the Haar distribution on  $K$ . Denote by  $I(X)$  the set of shifts of such distributions, i.e. the distributions of the form  $m_K * E_x$ , where  $K$  is a compact subgroup of  $X$ ,  $x \in X$ .

The first step to generalize the Skitovich-Darmois theorem on locally compact Abelian groups is to consider the case of finite Abelian groups. This problem was studied first in [5]. Note that the idempotent distributions on a finite Abelian group can be regarded as analogues of the Gaussian distributions on real line. Let  $\xi_i, i = 1, 2, \dots, n, n \geq 2$ , be independent random variables with values in a finite Abelian group  $X$  and distributions  $\mu_i$ . Let  $\alpha_j, \beta_j$  be automorphisms of  $X$ . Consider the linear forms  $L_1 = \alpha_1 \xi_1 + \dots + \alpha_n \xi_n$   $L_2 = \beta_1 \xi_1 + \dots + \beta_n \xi_n$ . G.M. Feldman proved that the class of groups, on which the independence of  $L_1$   $L_2$  implies that all  $\mu_i \in I(X)$ , are poor and consists of the groups of the form ([5])

$$\mathbb{Z}(2^{m_1}) \times \dots \times \mathbb{Z}(2^{m_l}), 0 \leq m_1 < \dots < m_l. \quad (1)$$

On the other hand if we consider  $n$  linear forms of  $n$  independent random variables, then the Skitovich-Darmois theorem is valid for an arbitrary finite Abelian group. Namely, the following theorem holds ([12]):

**Theorem 1.1** *Let  $\xi_i, i = 1, 2, \dots, n, n \geq 2$ , be independent random variables with values in a finite Abelian group  $X$  and distributions  $\mu_i$ . If the linear forms  $L_j = \sum_{i=1}^n \alpha_{ij} \xi_i$ , where  $\alpha_{ij} \in \text{Aut}(X), i, j = 1, 2, \dots, n$ , are independent, then  $\mu_i \in I(X), i = 1, 2, \dots, n$ .*

Our first aim is to generalize Theorem 1.1 into discrete groups. Using this result we extend Theorem 1.1 into groups of the form  $\mathbb{R}^m \times D \times K$ , where  $D$  is a discrete,  $K$  is a compact totally disconnected group of the special form,  $m \geq 0$ . Finally we describe conditions on compact totally disconnected group, which are necessary and sufficient for the validity of the Theorem 1.1.

## 2 Definitions and designations

Let  $X$  be a second countable locally compact Abelian group. Let  $\{X_\lambda : \lambda \in \Lambda\}$  be a non empty family of groups. Denote by  $\mathbf{P}_{\lambda \in \Lambda} X_\lambda$  the direct product of the groups  $X_\lambda$ . If  $X = X_\lambda$  for all  $\lambda \in \Lambda$ , then the direct product of the groups  $X_\lambda$  we denote by  $X^n$ , where  $n$  is a cardinal number of  $\Lambda$ .

Let  $p$  be a prime number. Denote by  $\mathfrak{P}$  the set of all prime number. Denote by  $\Delta_p$  the group of  $p$ -adic integers. We recall that an Abelian group is called a  $p$ -primary if every nonzero element of this group has order  $p^m$  for some natural  $m$ .

Let  $\mu$  be a distribution on  $X$ . The characteristic function of the distribution  $\mu$  is defined by the formula ([2])

$$\hat{\mu}(y) = \int_X (x, y) d\mu(x), y \in Y.$$

Recall ([13]), that a distribution  $\mu$  on  $X$  is called Gaussian, if its characteristic function can be represented in the form

$$\hat{\mu}(y) = (x, y) \exp\{-\phi(y)\}, \quad y \in Y,$$

where the function  $\phi(y)$  satisfy the equation

$$\phi(u + v) + \phi(u - v) = 2\phi(u) + 2\phi(v), \quad u, v \in Y.$$

Denote by  $\Gamma(X)$  the set of all Gaussian distributions on  $X$ .

### 3 Main results

The main result of the article is Theorem 3.2. The proof of Theorem 3.2 uses the following theorem, which is interesting in itself.

**Theorem 3.1** *Let  $X$  be a discrete Abelian group. Let  $\xi_i, i = 1, 2, \dots, n$ , be independent random variables with values in  $X$  and distributions  $\mu_i$ . Consider the linear forms  $L_j = \sum_{i=1}^n \alpha_{ij} \xi_i$ , where  $\alpha_{ij} \in \text{Aut}(X), i, j = 1, 2, \dots, n$ . Then the independence of the linear forms  $L_j$  implies that  $\mu_i = E_{x_i} * m_{G_i}$ , where  $G_i$  are finite subgroups of  $X, x_i \in X, i = 1, 2, \dots, n$ .*

Note that Theorem 3.1 was proved earlier in ([3, 13.17]) in the case  $n = 2$ .

**Theorem 3.2** *Let  $X$  be a Abelian group of the form*

$$X = \mathbb{R}^m \times K \times D, \tag{2}$$

where,  $m \geq 0$   $K$  — compact totally disconnected group, which is topologically isomorphic to a group of the form

$$\mathbf{P}_{p \in \mathfrak{P}}(\Delta_p^{n_p} \times G_p), \tag{3}$$

where  $n_p$  are some non negative integers,  $G_p$  are finite  $p$ -primary groups,  $D$  is a discrete countable group. Let  $\xi_i, i = 1, 2, \dots, n$ , be independent random variables with values in  $X$  and distributions  $\mu_i$ . Then the independence of the linear forms  $L_j = \sum_{i=1}^n \alpha_{ij} \xi_i$ , where  $\alpha_{ij} \in \text{Aut}(X), i, j = 1, 2, \dots, n$ , implies that  $\mu_i \in \Gamma(X) * I(X), i = 1, 2, \dots, n$ .

From the conditions of theorem 3.2 it follows that condition (3) is sufficient for validity of the Skitovich-Darmois theorem on compact totally disconnected Abelian groups in the case of  $n$  linear forms of  $n$  random variables. It turns out, that this condition is necessary.

**Theorem 3.3** *Let  $X$  be a compact totally disconnected Abelian group,  $\xi_i, i = 1, 2, \dots, n$ , be independent random variables with values in  $X$  and distributions  $\mu_i, \alpha_{ij} \in \text{Aut}(X)$ . From the independence of the linear forms  $L_j = \sum_{i=1}^n \alpha_{ij} \xi_i, j = 1, 2, \dots, n$ , follow that all  $\mu_i \in \Gamma(X) * I(X)$ , if and only if  $X$  is topologically isomorphic to the group of the form (3).*

Note that the theorem above was proved earlier in ([3, 13.25]) in the case  $n = 2$ .

**Remark.** There are locally compact Abelian groups which are non isomorphic to the group of the form (2), and on which The Skitovich-Darmois theorem is valid in the case of  $n$  linear forms of  $n$  random variables. The needed example was constructed for the case  $n = 2$  in [4, 3]. It easily can be checked, that this example can be generalized in the case of the arbitrary  $n$ .

## References

- [1] G. Darmois, Analyse generale des liasions stochastiques. Etude particuliere de l'analyse factorielle lineaire. *Rev. Inst. Internat. Statistique.* **21** (1953), 2-8.
- [2] G.M. Fel'dman, *Arithmetic of Probability Distributions and Characterization Problems on Abelian Groups* AMS translation of mathematical monographs **116**, (American Math. Society, Providence, RI, 1993).
- [3] G. Feldman, *Functional equations and Characterizations problems on locally compact Abelian Groups.* (EMS, 2008).
- [4] G. M. Feldman and P. Graczyk, The Skitovich-Darmois theorem for locally compact Abelian groups, *J. of the Austral. Math. Soc.* **88** (2010), 339-352.
- [5] G. M. Feldman, On the Skitovich-Darmois theorem for finite Abelian groups. *Theory Probabl. Appl.* **37** (1992), 621-631.

- [6] G. M. Feldman, On the Skitovich-Darmois theorem on compact groups. *Theory Probab. Appl.* **41** (1996), 768-773.
- [7] G. M. Feldman, The Skitovich-Darmois theorem for discrete periodic Abelian groups. *Theory Probab. Appl.* **42** (1997), 611-617.
- [8] G. M. Feldman and P. Graczyk, On the Skitovich-Darmois theorem on compact Abelian groups. *J. of Theoretical Probability*, **13** (2000), 859-869.
- [9] G. M. Feldman and P. Graczyk, On the Skitovich-Darmois theorem for discrete Abelian groups. *Theory Probab. Appl.* **49** (2005), 527-531.
- [10] P. Graczyk and G. M. Feldman, Independent linear statistics on finite Abelian groups. *Ukrainian Math. J.* **53** (2001), 499-506.
- [11] A. M. Kagan, Yu. V. Linnik and C. R. Rao, *Characterization Problems in Mathematical Statistics*. Wiley Series in Probability and Mathematical Statistics. (John Wiley & Sons, New York-London-Sydney, 1973).
- [12] I. Mazur, The Skitovich-Darmois theorem for finite Abelian groups, arXiv:1106.4134.
- [13] K. R. Parthasarathy , R. Ranga Rao, and S. R. S. Varadhan, Probability distributions on locally compact Abelian groups, *Illinois. J. Math.* **7**, 337-369 (1963).
- [14] V. P. Skitovich, On a property of the normal distribution. *Dokl. Akad. Nauk SSSR (N.S.)* **89** (1953), 217-219.

Mathematical Division  
 B. Verkin Institute for Low  
 Temperature Physics and Engineering  
 of the National Academy  
 of Sciences of Ukraine  
 47, Lenin Ave, Kharkov  
 61103, Ukraine