# Classical simulation of dissipative fermionic linear optics

Sergey Bravyi and Robert König

IBM T.J. Watson Research Center, Yorktown Heights, NY 10598, USA

October 13, 2018

#### Abstract

Fermionic linear optics is a limited form of quantum computation which is known to be efficiently simulable on a classical computer. We revisit and extend this result by enlarging the set of available computational gates: in addition to unitaries and measurements, we allow dissipative evolution governed by a Markovian master equation with linear Lindblad operators. We show that this more general form of fermionic computation is also simulable efficiently by classical means. Given a system of  $N$  fermionic modes, our algorithm simulates any such gate in time  $O(N^3)$  while a single-mode measurement is simulated in time  $O(N^2)$ . The steady state of the Lindblad equation can be computed in time  $O(N^3)$ .

### 1 Introduction

Quantum dynamical processes have inspired numerous models of computation: adiabatic quantum computation [\[8\]](#page-16-0), dissipative quantum computation [\[29,](#page-18-0) [19\]](#page-17-0) or computations based on modular functors [\[11\]](#page-16-1) are all examples of computational models associated with certain timeevolving quantum systems. In trying to characterize their computational power, arguably the most practically relevant question is how they compare to universal classical respectively quantum computation. Indeed, if computations in a model are efficiently simulable on a classical computer, the corresponding physical system may be accessible to numerical studies, but is unlikely to be a suitable substrate for building a quantum computer. In contrast, simulability by quantum circuits means that the underlying physics could be studied using a quantum computer, the prime application of such machines originally envisioned by Feynman [\[9\]](#page-16-2). Finally, in cases where the computational resources provided by a model are sufficient to implement universal quantum computation, the corresponding physical system is a candidate for the realization of a quantum computer.

A number of physically motivated models can be understood as the result of restricting the available set of initial states, gates and measurements in the standard quantum circuit model. For topological [\[18,](#page-17-1) [22,](#page-17-2) [10\]](#page-16-3) or permutational [\[13\]](#page-17-3) quantum computing, there is a preferred initial (vacuum) state, the available gates represent braid group generators or transpositions and there is a set of allowed (charge) measurements. In bosonic linear quantum optics [\[21\]](#page-17-4), the available repertoire includes preparation of the vacuum initial state, single photon sources, beam splitters, phase shifters and photo-detectors. In fermionic quantum optics [\[26,](#page-17-5) [20\]](#page-17-6), we permit

preparation of the vacuum state, free unitary evolution and occupation number measurements. The computational power of topological computing depends on the representation: for example, it is universal in the case of the Fibonacci model but classically simulable for Ising anyons. Bosonic linear optics was shown to be universal [\[21\]](#page-17-4) for quantum computation. Fermionic linear optics can be simulated efficiently on a classical computer [\[26,](#page-17-5) [20\]](#page-17-6) (see below).

The model of dissipative quantum computing is of a conceptually different origin: it is the result of considering more general quantum dynamics beyond unitary evolutions. Here the standard unitary gate set is augmented or completely replaced by 'dissipative gates'. Mathematically, the latter are completely positive trace-preserving maps corresponding to the timeevolution under a Markovian master equation. Such Markovian evolution is usually associated with noise in implementing a quantum computer, and one seeks to reduce its detrimental effect by the use of error-correction. Recently, however, it was realized that purely dissipative processes can actually be useful for quantum computation (see e.g., [\[12,](#page-17-7) [29,](#page-18-0) [30\]](#page-18-1)). In [\[29\]](#page-18-0), it was shown that dissipative quantum computation is universal for quantum computation. Conversely, Markovian dynamics with local Lindblad operators can be simulated efficiently (i.e., with polynomial overhead) on a quantum computer [\[19\]](#page-17-0).

Here we consider a model of computation that extends fermionic linear optics with dissipative processes. We will show that this extended model is still efficiently simulable classically. Since our simulation algorithm does not rely on Trotter-type expansions (as e.g., [\[19\]](#page-17-0)), dissipative processes can be simulated exactly for any evolution time without affecting complexity or accuracy. The algorithm reproduces the statistics of measurement outcomes and also provides a complete description of the state at any instant in the computation. Additionally, the Liouvillian may be non-local, but we restrict to Lindblad operators which are linear in fermionic annihilation and creation operators.

The physics underlying fermionic linear optics – non-interacting fermions – encompasses a number of systems of interest in condensed matter physics, including Kitaev's Majorana chain [\[16\]](#page-17-8) or honeycomb model [\[17\]](#page-17-9). Such systems exhibit topological order and could be used as fault-tolerant quantum memories or topological quantum computers. The classical simulation algorithm discussed here may be particularly suited to assess their potential to serve such information-processing purposes. Typically, this involves studying not only the dynamics of the system, but also the effect of simple manipulations (such as syndrome measurements or error correction). As an example, the known simulation technique in the non-dissipative case has been used to study the beneficial effect of disorder on the performance of the Majorana chain as a quantum memory [\[6\]](#page-16-4). The extended toolkit provided here may be applied to evaluate the robustness of certain proposals, e.g., for state transfer [\[32,](#page-18-2) [33\]](#page-18-3), in the presence of dissipation.

### <span id="page-1-1"></span>2 Fermionic linear optics

Fermionic linear optics is defined in terms of N creation and annihilation operators  $a_i^{\dagger}$  $j$  and  $a_j$ satisfying Fermi-Dirac canonical commutation relations  $\{a_j, a_k\} = 0$  and  $\{a_j, a_k\}$  $\{^\mathsf{T}_k\} = \delta_{j,k} I.$  We will refer to such a family of operators  $\{a_j, a_j^{\dagger}\}$  $j_{j}^{T}$ <sub>j=1</sub> as N Dirac fermions or modes to distinguish them from Majorana fermions introduced below (the latter will be denoted by the letter  $c$ instead of a). The corresponding Hilbert space is spanned by the number states

<span id="page-1-0"></span>
$$
|n_1, \ldots, n_N\rangle = (a_1^\dagger)^{n_1} \cdots (a_N^\dagger)^{n_N} |0\rangle \qquad \text{where } n_j \in \{0, 1\} \;, \tag{1}
$$

and  $|0\rangle$  is the fermionic vacuum state satisfying  $a_j|0\rangle = 0$  for all j. A state of the form [\(1\)](#page-1-0) is an eigenstate of the occupation number operator  $N_j = a_j^{\dagger} a_j$  with eigenvalue  $n_j$ .

<span id="page-2-1"></span>The allowed operations defining fermionic linear optics are:

- <span id="page-2-2"></span>(i) preparation of the fermionic vacuum  $|0\rangle$
- <span id="page-2-0"></span>(ii) measurement of the occupation numbers  $N_j$ ,  $j \in S$  for any subset S of modes,
- (iii) evolution under a quadratic fermion Hamiltonian  $H$  for a time  $t$ , that is, according to the equation of motion

<span id="page-2-5"></span>
$$
\frac{d}{dt}\rho = -i[H,\rho] \tag{2}
$$

Fermionic parity preservation implies that  $H$  is a linear combination of terms of the form  $\epsilon_j a_j^{\dagger} a_j$  giving energy  $\epsilon_j \in \mathbb{R}$  to mode j, 'hopping terms' of the form  $t_{j,k} a_j^{\dagger} a_k + t_{j,k}^* a_k^{\dagger}$  $_{k}^{\intercal}a_{j}$ where  $t_{j,k} \in \mathbb{C}$  for  $j \neq k$  and 'pair creation/annihilation terms' of the form  $s_{j,k}a_j^{\dagger}$  $j^{\dagger}a_{k}^{\dagger} +$  $s_{j,k}^* a_k a_j, s_{j,k} \in \mathbb{C}.$ 

These operations can be performed in an arbitrary order, and, in particular, may depend on measurement results obtained in the course of the computation. The computation concludes with a final measurement whose outcome is the classical result produced by the computation.

Simulating such a fermionic linear optics computation on a classical computer amounts to sampling from the distribution of measurement outcomes at the end of the computation. A classical polynomial-time (in N) algorithm for doing so was found by Terhal and DiVincenzo  $[26]$ and Knill [\[20\]](#page-17-6). A stronger form of simulation outputs a description of the state at any instant during the computation. An efficient algorithm for this problem was provided by Bravyi [\[4\]](#page-16-5). We review these techniques in Section [5.](#page-7-0)

Fermionic linear optics was motivated by the quantum universality of bosonic linear optics [\[21\]](#page-17-4), but is closely related to another computational model: the unitaries [\(iii\)](#page-2-0) are so-called matchgates. The corresponding computational model – matchgate computation – was earlier shown to be classically simulable by Valiant [\[27\]](#page-17-10). Josza and Miyake [\[15\]](#page-17-11) extended the simulation results by showing that more general initial (product) states can be allowed without losing classical simulability. They also showed that a slight modification of Valiant's gate set provides quantum universality (see also [\[2\]](#page-16-6) for a physically motivated universal gate set extending fermionic linear optics). More recently, new and elegant characterizations of the power of fermionic linear optics have been provided: Jozsa and coauthors [\[14\]](#page-17-12) showed that the model is equivalent to space-bounded quantum computation (see also [\[5\]](#page-16-7)), and van den Nest [\[28\]](#page-18-4) gave a characterization in terms of linear threshold gates.

### 3 Dissipative fermionic linear optics

<span id="page-2-3"></span>Here we generalize these results as follows: in addition to the unitary set of gates defined by [\(iii\)](#page-2-0), we allow dissipative processes. More specifically, we replace [\(iii\)](#page-2-0) with

(iii') evolution under the Lindblad master equation

<span id="page-2-4"></span>
$$
\frac{d}{dt}\rho = \mathcal{L}(\rho) := -i[H, \rho] + \sum_{\mu} \left( 2L_{\mu}\rho L_{\mu}^{\dagger} - \{ L_{\mu}^{\dagger}L_{\mu}, \rho \} \right) , \qquad (3)
$$

for a time t, where H is a quadratic fermion Hamiltonian and each Lindblad operator  $L_{\mu}$ is a linear combination  $L_{\mu} = \sum_{j} \alpha_{\mu,j} a_j^{\dagger} + \beta_{\mu,j} a_j$  with  $\alpha_{\mu,j}, \beta_{\mu,j} \in \mathbb{C}$ .

We show that a fermionic computation composed of  $(i)$ , $(ii)$  and  $(iii)$  can be efficiently simulated on a classical computer.

#### Outline

The remainder of this paper is structured as follows. In Section [4,](#page-3-0) we discuss fermionic Hamiltonians and Liouvillians and we show that dissipative dynamics with linear Lindblad operators preserves the set of Gaussian fermionic states. In Section [5,](#page-7-0) we review the classical simulation of fermionic linear optics. In Section [6](#page-10-0) we explain how to simulate dissipative fermionic linear optics.

### <span id="page-3-0"></span>4 Background

In this section, we give some background on quadratic open fermion systems with linear Lindblad operators. It will be convenient to work with the Hermitian operators

$$
c_{2j-1} = a_j + a_j^{\dagger}
$$
 and  $c_{2j} = i(a_j - a_j^{\dagger})$  for  $j = 1, ..., N$ .

The 2N operators  $c_1, \ldots, c_{2N}$  satisfy the canonical anticommutation relations

<span id="page-3-1"></span>
$$
\{c_j, c_k\} = 2\delta_{j,k}I \quad \text{for all} \quad j, k = 1, \dots, 2N,
$$

and will be referred to as Majorana fermions or modes. The Hamiltonian and Lindblad operators [\(3\)](#page-2-4) then take the form

$$
H(\mathbf{H}) = \frac{i}{4} \sum_{j,k=1}^{2N} \mathbf{H}_{jk} c_j c_k = \frac{i}{4} \underline{c} \cdot \mathbf{H} \underline{c} \qquad L_{\mu} = \sum_{j=1}^{2N} \ell_{\mu,j} c_j = \underline{\ell}_{\mu} \cdot \underline{c}
$$
(4)

where  $c = (c_1, \ldots, c_{2N})$ . Here  $\mathbf{H} = -\mathbf{H}^T \in \mathfrak{so}(2N, \mathbb{R})$  is an antisymmetric matrix with real entries. In contrast, the vectors  $\underline{\ell}_{\mu} = (\ell_{\mu,1}, \ldots, \ell_{\mu,2N}) \in \mathbb{C}^{2N}$  are generally complex-valued.

#### <span id="page-3-2"></span>4.1 Third quantization description of Liouvillians

We will present two methods for simulating dissipative dynamics of the form  $(3)$ . One of these methods is based on the language of 'third quantization'. This refers to the fact that the Liouvillian  $\mathcal L$  is quadratic in a set of fermionic superoperators as discussed by Prosen [\[23\]](#page-17-13) (see also [\[25\]](#page-17-14) and Dzhioev and Kosov [\[7\]](#page-16-8)). Here we follow his presentation.

Let K be the 4<sup>N</sup>-dimensional vector space spanned by the monomials  $c^{\alpha} = c_1^{\alpha_1}$  $i_1^{\alpha_1}\cdots i_{2N}^{\alpha_{2N}}$ where  $\underline{\alpha} = (\alpha_1, \ldots, \alpha_{2N}) \in \{0, 1\}^{2N}$ . To emphasize the vector space structure, we will write an operator  $\rho = \sum_{\underline{\alpha}} \lambda_{\underline{\alpha}} c^{\underline{\alpha}}$  (with  $\lambda_{\underline{\alpha}} \in \mathbb{R}$ ) as  $|\rho\rangle = \sum_{\underline{\alpha}} \lambda_{\underline{\alpha}} |c^{\underline{\alpha}}\rangle$  when it is considered as an element of K. The space K is equipped with the Hilbert-Schmidt inner product  $\langle \varphi | \sigma \rangle = 2^{-N} \text{Tr} \rho^{\dagger} \sigma$ , and with respect to the latter, the monomials  $|c^{\alpha}\rangle$  are an orthonormal basis. The space K decomposes into a direct sum  $K = \mathcal{K}^+ \oplus \mathcal{K}^-$  of spaces spanned by monomials  $|c^{\alpha}\rangle$  with even  $(\mathcal{K}^+)$ 

and odd  $(\mathcal{K}^-)$  parity  $\sum_{j=1}^{2N} \alpha_j$  mod 2. In the following, we will often restrict our attention to the subspace  $K^+$ , although identical arguments apply to  $K^-$ . By fermionic parity superselection, this is sufficient to cover physical states.

We define 2N pairs  $(\hat{a}_j, \hat{a}_j^{\dagger})$  $j$ ) of mutually adjoint linear operators acting on  $\mathcal{K}$  by

$$
\hat{a}_j|c^{\underline{\alpha}}\rangle = \delta_{\alpha_j,1}|c_j c^{\underline{\alpha}}\rangle \quad \text{and} \quad \hat{a}_j^{\dagger}|c^{\underline{\alpha}}\rangle = \delta_{\alpha_j,0}|c_j c^{\underline{\alpha}}\rangle \quad \text{for } j = 1,\dots,2N. \quad (5)
$$

These satisfy canonical Dirac anticommutation relations

<span id="page-4-1"></span><span id="page-4-0"></span>
$$
\{\hat{a}_j, \hat{a}_k\} = 0 \qquad \{\hat{a}_j, \hat{a}_k^{\dagger}\} = \delta_{j,k} \qquad \text{for } j, k = 1, ..., 2N \ ,
$$

which has motivated the expression 'third quantization'. Observe that the state  $|c^{\mathcal{Q}}\rangle = |I\rangle$  corresponding to the identity operator  $I$  is the vacuum state associated with this set of operators, i.e.,  $\hat{a}_j |I\rangle = 0$  for all j.

It is again often more convenient to work with Hermitian Majorana fermions

$$
\hat{c}_{2j-1} = \hat{a}_j + \hat{a}_j^{\dagger}
$$
 and  $\hat{c}_{2j} = i(\hat{a}_j - \hat{a}_j^{\dagger})$  for  $j = 1, ..., 2N$ . (6)

The restriction  $\mathcal{L}_+ = \mathcal{L}|_{\mathcal{K}^+}$  of the Liouvillian  $\mathcal L$  to operators supported on  $\mathcal{K}^+$  will be called the even part of  $\mathcal{L}$ . Remarkably, this superoperator can be expressed as a quadratic form of the operators [\(5\)](#page-4-0) (or equivalently [\(6\)](#page-4-1)). More precisely, we have

$$
|\mathcal{L}(\rho)\rangle = \widehat{\mathcal{L}_+}|\rho\rangle \qquad \text{for all } \rho \text{ with } supp(\rho) \subset \mathcal{K}^+ \text{ , where } \qquad \widehat{\mathcal{L}_+} = \frac{1}{4}\hat{\underline{c}} \cdot \mathbf{L}\hat{\underline{c}} \text{ , } \qquad (7)
$$

and where  $\hat{\mathbf{c}} = (\hat{c}_1, \dots, \hat{c}_{4N})$  for an antisymmetric complex-valued matrix  $\mathbf{L} = -\mathbf{L}^T \in \mathfrak{so}(4N, \mathbb{C})$ . The matrix **L** depends linearly on **H** and quadratically on the entries of the vectors  $\underline{\ell}_{\mu}$  specifying the Lindblad operators. Explicitly, it is given by [\[23\]](#page-17-13)

$$
\begin{array}{rcl}\n\mathbf{L}_{2j-1,2k-1} & = & \mathbf{H}_{j,k} - 2\mathbf{M}_{j,k} + 2\mathbf{M}_{k,j} \\
\mathbf{L}_{2j,2k} & = & \mathbf{H}_{j,k} + 2\mathbf{M}_{j,k} - 2\mathbf{M}_{k,j} \\
\end{array} \quad\n\begin{array}{rcl}\n\mathbf{L}_{2j-1,2k} & = & 4i\mathbf{M}_{k,j} \\
\mathbf{L}_{2j,2k-1} & = & -4i\mathbf{M}_{j,k}\n\end{array}\n\tag{8}
$$

with  $\mathbf{M}_{j,k} = \sum_{\mu} \ell_{\mu,j} \ell_{\mu,k}^*$ . In [\[23,](#page-17-13) [25\]](#page-17-14), it is assumed that the Liouvillian is generic in the sense that L is diagonalizable. Here we do not require such an assumption. Note also that in [\[24\]](#page-17-15), a normal form for such matrices **L** is derived.

Observe that for any integer n and any state  $\rho$  supported on  $K^+$ , we have

<span id="page-4-4"></span>
$$
|\mathcal{L}_+^n(\rho)\rangle = |\underbrace{\mathcal{L}_+ \circ \cdots \circ \mathcal{L}_+}_{n}(\rho)\rangle| = \widehat{\mathcal{L}_+}^n|\rho\rangle,
$$

which implies that the superoperator  $\exp(t\mathcal{L}_+)$  corresponding to time evolution for a time t under [\(3\)](#page-2-4) is given by

<span id="page-4-3"></span><span id="page-4-2"></span>
$$
\widehat{\exp(t\mathcal{L}_+)} = \exp(t\widehat{\mathcal{L}_+})
$$
\n(9)

when restricted to operators supported on  $K^+$ . Because det  $\exp(A) = \exp(\text{Tr}(A))$ , the opera-tor [\(9\)](#page-4-2) is invertible with inverse  $\exp(-t\widehat{\mathcal{L}_+})$ .

Finally, consider the adjoint Liouvillian defined by

$$
\mathcal{L}^{\dagger}(O) = i[H, O] + \sum_{\mu} \left( 2L_{\mu}^{\dagger} O L_{\mu} - \{L_{\mu}^{\dagger} L_{\mu}, O\} \right) \tag{10}
$$

Evolution for a time t under  $\mathcal{L}^{\dagger}$  generates the adjoint superoperator of  $\exp(t\mathcal{L})$ , that is,  $\text{Tr}(\exp(t\mathcal{L})(\rho)O) = \text{Tr}(\rho \exp(t\mathcal{L}^{\dagger})(O))$  for all operators  $\rho$ , O. This follows by comparing the derivative

$$
\frac{d}{dt}\operatorname{Tr}(Oe^{t\mathcal{L}}(\rho)) = \operatorname{Tr}(O\mathcal{L}(\rho)) = \operatorname{Tr}(\mathcal{L}^{\dagger}(O)\rho) = \frac{d}{dt}\operatorname{Tr}(e^{t\mathcal{L}^{\dagger}}(O)\rho)
$$

and observing that the two expressions agree for  $t = 0$ . Since  $\exp(t\mathcal{L})$  is a completely positive trace-preserving map, the map  $\exp(t\mathcal{L}^{\dagger})$  is unital, that is,  $\exp(t\mathcal{L}^{\dagger})(I) = I$ . The even part  $\mathcal{L}^{\dagger}_{+}$ + of the adjoint Liouvillian takes the form

<span id="page-5-1"></span><span id="page-5-0"></span>
$$
\widehat{\mathcal{L}}_{+}^{\dagger} = \frac{1}{4}\hat{\underline{c}} \cdot \mathbf{L}^{\dagger}\hat{\underline{c}} \tag{11}
$$

where the matrix  $\mathbf{L}^{\dagger} = -\mathbf{L}^* \in \mathfrak{so}(4N, \mathbb{C})$  is the Hermitian conjugate of  $\mathbf{L}$ .

#### 4.2 Gaussian states

Here we collect a few facts about Gaussian states of 2N Majorana fermions  $c_1, \ldots, c_{2N}$  (see [\[4\]](#page-16-5) for details and proofs). A Gaussian state  $\rho$  is completely determined by its covariance matrix

$$
M_{j,k} = \frac{i}{2} \text{Tr}(\rho[c_j, c_k]) \qquad \text{for } j, k = 1, ..., 2N \tag{12}
$$

and Wick's formula (see e.g., [\[4,](#page-16-5) Eq. 17])

$$
i^{p} \operatorname{Tr}(\rho c_{j_{1}} c_{j_{2}} \cdots c_{j_{2p}}) = \operatorname{Pf}(M[j_{1}, \ldots, j_{2p}]) \qquad \text{for all } 1 \leq j_{1} < j_{2} < \cdots < j_{2p} \leq 2N \ . \tag{13}
$$

Here  $M[j_1,\ldots,j_{2p}]$  is the submatrix of M of size  $2p \times 2p$  obtained by removing all columns and rows except those indexed by  $j_1, \ldots, j_{2p}$ , and Pf denotes the Pfaffian. The antisymmetric matrix  $M \in \mathfrak{so}(2N, \mathbb{R})$  can be brought into block-diagonal form by a special orthogonal matrix

$$
M = R \bigoplus_{j=1}^{N} \begin{pmatrix} 0 & \lambda_j \\ -\lambda_j & 0 \end{pmatrix} R^T \qquad R \in SO(2N), \lambda_j \in \mathbb{R}.
$$

This is called the Williamson normal form of M. The transformation  $\underline{c} \to \underline{c}' = R\underline{c}$ , that is,

$$
c'_{a} = \sum_{b=1}^{2N} R_{a,b} c_b \qquad \text{for } a = 1, ..., 2N
$$

corresponds to the adjoint action of a unitary as explained below. In particular,  $c'_1, \ldots, c'_{2N}$ again satisfy canonical commutation relations. We call these the eigenmodes of  $\rho$ . Expressed in terms of these operators,  $\rho$  takes the simple form

$$
\rho = \frac{1}{2^N} \prod_{j=1}^N (I + i\lambda_j c'_{2j-1} c'_{2j}) \tag{14}
$$

#### 4.3 Time evolution of Gaussian states

<span id="page-6-2"></span>Evolution of Gaussian states under the dissipative dynamics described by Eq. [\(3\)](#page-2-4) is particularly simple due to the following fact.

Lemma 1. *Let* L *be a Liouvillian for* N *fermions whose unitary part is given by a quadratic Hamiltonian and whose Lindblad operators are linear in the creation- and annihilation operators* (*cf.* [\(3\)](#page-2-4)). Let  $\rho$  be a Gaussian state. Then the time-evolved state  $e^{t\mathcal{L}}(\rho)$  is Gaussian for all  $t \geq 0$ .

*Proof.* We will construct a family of trace preserving completely positive maps  $\{\Phi_{\epsilon}\}\$  depending smoothly on a parameter  $\epsilon \geq 0$  such that each map  $\Phi_{\epsilon}$  preserves the set of Gaussian states and

<span id="page-6-1"></span>
$$
\left. \frac{d}{d\epsilon} \Phi_{\epsilon}(\rho) \right|_{\epsilon=0} = \mathcal{L}(\rho) \tag{15}
$$

for any state  $\rho$ . This implies that

$$
\rho(t)=\lim_{n\to\infty}\Phi^n_{t/n}(\rho)
$$

is a limiting point of a sequence of Gaussian states. Since the set of Gaussian states is compact,  $\rho(t)$  must be a Gaussian state itself.

Let us first consider a Liouvillian with a single Lindblad operator,

$$
\mathcal{L}(\rho) = 2L\rho L^{\dagger} - \{L^{\dagger}L, \rho\},\
$$

where L is a linear combination of the Majorana operators  $c_1, \ldots, c_{2N}$  with complex coefficients. Decompose  $L = K + iM$ , where K, M are real linear combinations of the Majorana operators. In particular, both  $K$  and  $M$  are Hermitian. Introduce one ancillary pair of Majorana operators  $c_{2N+1} \equiv b_1$  and  $c_{2N+2} \equiv b_2$  representing an environment and consider a unitary operator

$$
U_{\epsilon} = \exp \left[ -\sqrt{2\epsilon} (Kb_1 + Mb_2) \right] .
$$

Simple algebra shows that

<span id="page-6-0"></span>
$$
U_{\epsilon}\eta U_{\epsilon}^{\dagger} = \eta - \sqrt{2\epsilon}[Kb_1 + Mb_2, \eta] - 2\epsilon(Kb_1 + Mb_2)\eta(Kb_1 + Mb_2) + \epsilon\{(Kb_1 + Mb_2)^2, \eta\} + O(\epsilon^{3/2})
$$
(16)

for any state  $\eta$ . Define a map

$$
\Phi_{\epsilon}(\rho) = \text{Tr}_{E} U_{\epsilon} \,\rho \rho_{E} U_{\epsilon}^{\dagger} \,. \tag{17}
$$

Here  $\text{Tr}_E$  represents the partial trace over the environment, and  $\rho_E$  is the initial state of the environment which we choose as the vacuum state, that is,

$$
\rho_E = \frac{1}{2}(I - ib_1b_2) \; .
$$

Note that the terms in Eq. [\(16\)](#page-6-0) that contain half-integer powers of  $\epsilon$  do not contribute to  $\Phi_{\epsilon}(\rho)$ since  $\text{Tr}_E b_1 = \text{Tr}_E b_2 = 0$ . Using the identities

$$
\text{Tr}_E b_1 b_2 \rho_E = i \quad \text{and} \quad \text{Tr}_E b_1 \rho_E b_2 = -i
$$

it is easy to check that

<span id="page-7-1"></span>
$$
\Phi_{\epsilon}(\rho) = \rho + \epsilon \mathcal{L}(\rho) + O(\epsilon^2) \tag{18}
$$

Given a general Liouvillian with multiple Lindblad operators and the unitary evolution term as in Eq. [\(3\)](#page-2-4), the desired map  $\Phi_{\epsilon}$  can be constructed by taking a composition

$$
\Phi_{\epsilon}(\rho) = e^{-iH\epsilon} \Phi_{\epsilon}^{(1)} \circ \ldots \circ \Phi_{\epsilon}^{(m)}(\rho) e^{iH\epsilon},
$$

where  $\Phi_{\epsilon}^{(\mu)}$  is the map defined above with  $L = L_{\mu}$ . By construction,  $\Phi_{\epsilon}$  is a finite composition of unitary evolutions under quadratic Hamiltonians, addition of ancillary two-mode vacuum states  $\rho_E$ , and partial traces over some pairs of modes. It is well-known that all these operations preserve the set of Gaussian states, see e.g. [\[4\]](#page-16-5). Finally, Eq. [\(15\)](#page-6-1) follows from Eq. [\(18\)](#page-7-1) and the product rule for derivatives.  $\Box$ 

### <span id="page-7-0"></span>5 Classical simulation of fermionic linear optics

In this section, we review the known simulation techniques for a computation composed of the operations  $(i)$ – $(iii)$  introduced in Section [2.](#page-1-1) The basis of the simulation algorithm is the fact that the vacuum state  $|0\rangle$  prepared by (i) is a Gaussian state and all subsequent operations preserve the Gaussian nature of the state.

The covariance matrix  $M(0)$  of the vacuum state  $|0\rangle$  at the beginning of the computation is given by the non-zero entries

$$
M(0)_{2j-1,2j} = 1 \qquad \text{for } j = 1, \dots, 2N \tag{19}
$$

above the main diagonal. The remaining task is to find update rules for the covariance matrix. These updates can be done efficiently as discussed below: measuring (time) complexity in terms of the number of additions, multiplications, and divisions on complex numbers that are required, measurements and unitary gates can be simulated in time  $O(N^3)$ .

#### 5.1 Simulating measurements

Here we describe the method from [\[6\]](#page-16-4) which is a more efficient version of Terhal and DiVin-cenzo's algorithm [\[26\]](#page-17-5) (the latter has time complexity  $O(N^4)$  when measuring N modes).

Consider a (non-destructive) measurement of the occupation number  $N_j = a_j^{\dagger} a_j = \frac{1}{2}$  $rac{1}{2}(I$  $ic_{2j-1}c_{2j}$  in a Gaussian state  $\rho$  with covariance matrix M. By definition [\(12\)](#page-5-0), the probability of obtaining outcome 1 when measuring  $\rho$  is

$$
P_j(1) = \text{Tr}(N_j \rho) = \frac{1}{2} (1 - M_{2j - 1, 2j}) \tag{20}
$$

Let  $\Pi_j(n_j) = (a_j^{\dagger} a_j)^{n_j} (a_j a_j^{\dagger})$  $_{j}^{\dagger}$ )<sup>1-n<sub>j</sub></sup> be the projection corresponding to the measurement outcome  $n_i \in \{0, 1\}$ . The post-measurement state

<span id="page-7-3"></span><span id="page-7-2"></span>
$$
\rho(n_j) = \frac{\Pi_j(n_j)\rho \Pi_j(n_j)}{P_j(n_j)}\tag{21}
$$

is Gaussian, as shown in [\[4\]](#page-16-5). Its covariance matrix  $M^{(j)}(n_j)$  can be computed from [\(20\)](#page-7-2) and [\(21\)](#page-7-3) using Wick's theorem  $(13)$ , giving  $(cf. [6, Eq. (7.3)])$  $(cf. [6, Eq. (7.3)])$  $(cf. [6, Eq. (7.3)])$ 

<span id="page-8-0"></span>
$$
M^{(j)}(n_j)_{p,q} = M_{p,q} - \frac{(-1)^{n_j}}{2P_j(n_j)} M_{2j-1,p} M_{2j,p} + \frac{(-1)^{n_j}}{2P_j(n_j)} M_{2j-1,q} M_{2j,p} .
$$
 (22)

The following algorithm then simulates a measurement of  $N_j$ : first, compute [\(20\)](#page-7-2) and sample a bit  $n_j \in \{0,1\}$  according to the probability distribution  $Pr[n_j = 1] = P_j(1)$ . Then update the covariance matrix according to  $(22)$ . This requires  $O(N^2)$  computational steps and one (non-uniform) bit of randomness.

A measurement of a subset  $S = \{j_1, \ldots, j_{|S|}\}\$  of modes can be simulated by iterative use of this procedure, using the recursion relation

$$
P_{j_{\ell}}^{j_{1}...j_{\ell-1}}(n_{j_{\ell}}|n_{j_{1}}\cdots n_{j_{\ell-1}}) = \text{Tr}(\Pi_{j_{\ell}}(n_{j_{\ell}})\rho(n_{j_{1}}\cdots n_{j_{\ell-1}}))
$$

$$
\rho(n_{j_{1}},\ldots,n_{j_{\ell}}) = \frac{\Pi_{j_{\ell}}(n_{j_{\ell}})\rho(n_{j_{1}},\ldots,n_{j_{\ell-1}})\Pi_{j_{\ell}}(n_{j_{\ell}})}{P_{j_{\ell}}^{j_{1}...j_{\ell-1}}(n_{j_{\ell}}|n_{j_{1}}\cdots n_{j_{\ell-1}})}
$$

for the probability and post-measurement state after the measurement of the  $j_{\ell}$ -th mode, given that the measurements of  $N_{j_1}, \ldots, N_{j_{\ell-1}}$  resulted in the sequence  $n_{j_1}, \ldots, n_{j_{\ell-1}} \in \{0, 1\}$ . This can be done in time  $O(|S| \cdot N^2)$  using  $|S|$  random bits.

#### 5.2 Simulating unitary evolution

The classical simulation of unitary dynamics (see [\[20,](#page-17-6) [26,](#page-17-5) [15\]](#page-17-11)) is particularly instructive for our generalization to the dissipative case. Consider time evolution under a quadratic Hamiltonian  $H = H(H)$  (cf. [\(4\)](#page-3-1)) for some time t, starting from a Gaussian initial state  $\rho$  with covariance matrix  $M(0)$ . According to [\(2\)](#page-2-5), the covariance matrix  $M(t)$  of the time-evolved state  $\rho(t)$ satisfies

$$
\frac{d}{dt}M_{j,k}(t) = i \operatorname{Tr} (c_j c_k(-i)[H,\rho]) = -\operatorname{Tr} (\rho[H,c_j c_k]) \qquad \text{for any } j < k \tag{23}
$$

where we used the cyclicity of the trace. From this expression, it follows that the covariance matrix  $M(t)$  satisfies the equation

<span id="page-8-2"></span><span id="page-8-1"></span>
$$
\frac{d}{dt}M(t) = [M(t), \mathbf{H}].
$$
\n(24)

Eq. [\(24\)](#page-8-1) has the solution

$$
M(t) = R(t)M(0)R(t)^{T} \qquad \text{where} \qquad R(t) = e^{-\mathbf{H}t} \in SO(2N, \mathbb{R}). \tag{25}
$$

The matrix  $R(t)$  can be computed in time  $O(N^2)$  from the Williamson normal form of H. The latter can be computed in time  $O(N^3)$  (by diagonalizing  $\mathbf{H}^T\mathbf{H}$ ), hence it follows that the evolution  $M \mapsto M(t)$  can be simulated in time  $O(N^3)$ . Eq. [\(25\)](#page-8-2) is sufficient for the purpose of simulating unitaries for fermionic linear optics computation starting from a Gaussian state.

Let us discuss an alternative derivation of Eq. [\(25\)](#page-8-2), which additionally provides a method for simulating the evolution of higher moments of a (possibly non-Gaussian) initial state  $\rho$ . It also motivates the approach to dissipative dynamics discussed in Section [6.2.](#page-12-0) For this purpose, consider the adjoint action

$$
c_j(t) = e^{iHt} c_j e^{-iHt} = \sum_{k=1}^{2N} R(t)_{j,k} c_k
$$
\n(26)

of the unitary  $e^{-iHt}$  on Majorana operators. Eq. [\(26\)](#page-9-0) can be shown as follows (cf. [\[15,](#page-17-11) Theorem 3]). Consider the derivative

<span id="page-9-2"></span><span id="page-9-1"></span><span id="page-9-0"></span>
$$
\frac{dc_j(t)}{dt} = [(-i)H, c_j(t)].
$$
\n(27)

Because  $[c_j c_k, c_\ell] = 0$  unless  $\ell \in \{j, k\}$  and  $[c_j c_k, c_j] = -2c_k$ , we have

$$
[(-i)H(\mathbf{H}),c_{\ell}] = \sum_{n} \frac{1}{4} \mathbf{H}_{m,n}[c_m c_n, c_{\ell}] = -\sum_{n} \mathbf{H}_{\ell,n} c_n .
$$
 (28)

Observe that if  $c_j(t)$  is a linear combination of the operators  $\{c_\ell\}$ , then so is  $\frac{d}{dt}c_j(t)$  because of [\(27\)](#page-9-1) and [\(28\)](#page-9-2). Since this applies to  $t = 0$ , we conclude that  $c_j(t)$  is of the form specified on the lhs. of [\(26\)](#page-9-0) for all t, that is, a linear combination of the operators  ${c_k}$  with some coefficients  $R_{j,k}(t)$ . It remains to find the matrix  $R(t)$ . Rewriting [\(27\)](#page-9-1) in terms of  $R(t)$  and using [\(28\)](#page-9-2), we obtain  $\frac{dR(t)}{dt} = -R(t)H$  by taking the anticommutator  $\frac{1}{2}\{c_k, \cdot\}$  on both sides, for  $k = 1, \ldots, 2N$ . This shows that  $-\mathbf{H} \in \mathfrak{so}(2N, \mathbb{R})$  indeed generates  $R(t)$ , proving the claim [\(26\)](#page-9-0).

Since the covariance matrix  $M(t)$  of the time-evolved state can be computed in the Heisenberg picture as

$$
M_{j,k}(t) = \text{Tr}(e^{-iHt}\rho e^{iHt}c_jc_k) = \text{Tr}(\rho c_j(t)c_k(t)),
$$

the claim [\(25\)](#page-8-2) immediately follows from [\(26\)](#page-9-0). For later reference, we point out that in this argument, we made use of [\(26\)](#page-9-0) only to compute the product of two time-evolved operators, that is, in the form

$$
c_j(t)c_k(t) = \sum_{\ell,m=1}^{2N} R(t)_{j,\ell} R(t)_{k,m} c_\ell c_m \qquad \text{for } j \neq k .
$$
 (29)

It is also clear how this generalizes to higher moments: for example, if

<span id="page-9-4"></span><span id="page-9-3"></span>
$$
M_{j,k,\ell}(0) = \text{Tr}(\rho c_j c_k c_\ell) \tag{30}
$$

are the moments of a possibly non-Gaussian state  $\rho(0)$ , the moments of the time-evolved state  $\rho(t)$  are given by

$$
M_{j,k,\ell}(t) = \sum_{j',k',\ell'} R_{j,j'}(t) R_{k,k'}(t) R_{\ell,\ell'}(t) M_{j',k',\ell'}.
$$

# <span id="page-10-0"></span>6 Simulating dissipative evolution

In this section, we show how to simulate dissipative dynamics of the form [\(3\)](#page-2-4). In more detail, since [\(3\)](#page-2-4) preserves the Gaussian nature of a state  $\rho(0)$  according to Lemma [1,](#page-6-2) it suffices to compute the covariance matrix  $M(t)$  of the time-evolved state  $\rho(t) = e^{t\mathcal{L}}(\rho)$ . Our main result can be stated as follows:

<span id="page-10-4"></span>Theorem 1. *Consider* N *fermionic modes and let* L *be a Liouvillian of the form* [\(3\)](#page-2-4)*. Let* ρ *be an even Gaussian state with covariance matrix*  $M(0) \in \mathfrak{so}(2N, \mathbb{R})$ *. The covariance matrix*  $M(t)$  *of the time-evolved Gaussian state*  $\exp(t\mathcal{L})(\rho)$  *can be computed in time*  $O(N^3)$ *.* 

We give two different proofs of this statement: in Section [6.1,](#page-10-1) we sketch how to solve the corresponding differential equation directly. This approach may be most numerically stable and relies on the Bartels-Stewart algorithm for finding the covariance matrix of a fixed point. Our second method, discussed in Section [6.2](#page-12-0) is based on computing the Heisenberg-evolved Majorana operators using the formalism of third quantization. This method can be adapted to simulate the evolution of higher moments starting from (possibly non-Gaussian) initial states.

#### <span id="page-10-1"></span>6.1 Simulation method based on evolution equation

The definition of the covariance matrix  $M(t)$  implies that

$$
\frac{d}{dt}M_{j,k}(t) = i \operatorname{Tr} c_j c_k \mathcal{L}(\rho(t)) = i \operatorname{Tr} \mathcal{L}^\dagger(c_j c_k) \rho(t) \quad \text{for any } j < k. \tag{31}
$$

The action of the adjoint Liouvillian  $\mathcal{L}^{\dagger}$  (cf. [\(10\)](#page-4-3)) on observables can be written as

<span id="page-10-2"></span>
$$
\mathcal{L}^{\dagger}(O) = i[H, O] + \sum_{\mu} L^{\dagger}_{\mu}[O, L_{\mu}] + [L^{\dagger}_{\mu}, O]L_{\mu}.
$$

Choosing  $O \sim c_i c_k$  one can easily check that the commutators [H, O] and [O,  $L_\mu$ ] are quadratic and linear functions of the Majorana fermion operators respectively. It follows that  $\mathcal{L}^{\dagger}$  preserves the subspace of operators spanned by the identity and  $c_j c_k$  with  $1 \leq j \leq k \leq 2N$ . Therefore Eq. [\(31\)](#page-10-2) provides a closed linear differential equation that governs the time evolution of  $M(t)$ . Simple algebra shows that

<span id="page-10-3"></span>
$$
\frac{d}{dt}M(t) = \mathbf{X}M(t) + M(t)\mathbf{X}^T + \mathbf{Y},\tag{32}
$$

where

$$
\mathbf{X} = -\mathbf{H} - 2(\mathbf{M} + \mathbf{M}^*) \quad \text{and} \quad \mathbf{Y} = 4i(\mathbf{M}^* - \mathbf{M}) .
$$

This generalizes [\(24\)](#page-8-1). Recall that **H** is a real anti-symmetric matrix of size  $2N \times 2N$  parametrizing the quadratic Hamiltonian H, while M is a complex Hermitian matrix of size  $2N \times 2N$ parameterizing the Lindblad operators  $L_{\mu}$ , see Section [4.1.](#page-3-2) It follows that **X** and **Y** are both real-valued  $2N \times 2N$ -matrices, and  $\mathbf{Y} = -\mathbf{Y}^T$  is antisymmetric.

Since any Lindblad equation has at least one steady state<sup>[1](#page-11-0)</sup>, there must exist at least one covariance matrix  $M_0$  which is a fixed point of Eq. [\(32\)](#page-10-3), that is,

<span id="page-11-1"></span>
$$
\mathbf{X}M_0 + M_0 \mathbf{X}^T + \mathbf{Y} = 0. \tag{33}
$$

<span id="page-11-2"></span>Given any covariance matrix  $M_0$  satisfying Eq. [\(33\)](#page-11-1), the solution  $M(t)$  of Eq. [\(32\)](#page-10-3) can be written as

$$
M(t) = M_0 + e^{\mathbf{X}t} (M(0) - M_0) e^{\mathbf{X}^T t}, \quad t \ge 0.
$$
 (34)

Consider as an example a unitary evolution, that is,  $L_{\mu} = 0$  for all  $\mu$ . In this case  $\mathbf{M} = 0$ ,  $X = -H$ , and  $Y = 0$ . The fixed point covariance matrix can be chosen as  $M_0 = 0$  and we recover [\(25\)](#page-8-2).

Finding the fixed point covariance matrix  $M_0$  in the general case requires solving the system of linear equations Eq. [\(33\)](#page-11-1), where  $M_0$  is considered as an unknown vector of size  $N(2N-1)$ . A naïve approach based on the Gaussian elimination would take time  $O(N^6)$  to find  $M_0$ . However, this approach ignores the special structure of the problem. To the best of our knowledge, the most efficient method of solving the matrix equation Eq. [\(33\)](#page-11-1) is the Bartels-Stewart algo-rithm [\[1\]](#page-16-9) which has running time  $O(N^3)$ . The key step of the algorithm is to perform a real Schur decomposition of  $X$ , that is, an orthogonal change of basis making  $X$  block upper triangular with blocks of size 1 and 2 on the main diagonal. In the new basis the resulting system of equations on matrix elements of  $M_0$  turns out to be quasi-triangular and can be solved in time  $O(N^3)$ . For completeness, we sketch the Bartels-Stewart algorithm in Appendix [A.](#page-15-0)

It is worth emphasizing that all matrices involved in Eq. [\(34\)](#page-11-2) have bounded norm indepen-dent of t. This makes Eq. [\(34\)](#page-11-2) suitable for numerical calculation of  $M(t)$ . Indeed, taking into account that  $X + X^T = -4Re(M) \leq 0$ , one gets

$$
||e^{\mathbf{X}t}|| \le ||e^{(\mathbf{X} + \mathbf{X}^T)t/2}|| \le 1.
$$

Here the first inequality follows from Theorem IX.3.1 of [\[3\]](#page-16-10).

Finally, let us remark that Eq. [\(32\)](#page-10-3) can be solved directly even without knowing a fixed point  $M_0$  by transforming it into a homogeneous linear system. This method takes time only  $O(N^3)$ , but unfortunately, it is computationally unstable for large evolution time  $t$ . Indeed, introduce an auxiliary  $2N \times 2N$  matrix  $K(t)$  whose time evolution is trivial,  $K(t) = K(0) = I$ . Then one can rewrite [\(32\)](#page-10-3) as

<span id="page-11-3"></span>
$$
\dot{M}(t) = \mathbf{X}M(t) + M(t)\mathbf{X}^T + \mathbf{Y}K(t),
$$
\n(35)

$$
\dot{K}(t) = -\mathbf{X}^T K(t) + K(t) \mathbf{X}^T,
$$
\n(36)

with initial condition  $K(0) = I$ . Note that  $K(t) = I$  is indeed the only solution of Eq. [\(36\)](#page-11-3). Define a  $4N \times 2N$  matrix  $\Omega$  such that

$$
\Omega(t) = \left[ \begin{array}{c} M(t) \\ K(t) \end{array} \right].
$$

<span id="page-11-0"></span><sup>&</sup>lt;sup>1</sup>Note that for any integer  $n \geq 0$  there exists at least one (mixed) state  $\rho_n$  such that  $e^{\mathcal{L}/n}(\rho_n) = \rho_n$ . It implies  $\mathcal{L}(\rho_n) = O(1/n)$  for large n. Since  $\rho_n$  belongs to a compact manifold, the sequence  $\{\rho_n\}$  has a convergent subsequence. Its limiting point  $\rho$  is a (mixed) state that obeys  $\mathcal{L}(\rho) = 0$ .

Then the system of Eqs. [\(35,36\)](#page-11-3) is equivalent to

<span id="page-12-1"></span>
$$
\dot{\Omega}(t) = \mathbf{Z}\Omega(t) + \Omega(t)\mathbf{X}^T, \quad \text{where} \quad \mathbf{Z} = \begin{bmatrix} \mathbf{X} & \mathbf{Y} \\ 0 & -\mathbf{X}^T \end{bmatrix}.
$$
 (37)

Its solution is

$$
\Omega(t) = \exp\left(\mathbf{Z}t\right)\Omega(0)\exp\left(\mathbf{X}^T t\right). \tag{38}
$$

This can be computed in time  $O(N^3)$ . Unfortunately, using Eq. [\(38\)](#page-12-1) for numerical calculations may be problematic due to the exponential growth of the factor  $\exp(Zt)$ .

### <span id="page-12-0"></span>6.2 Simulation method based on 'third quantization'

Without loss of generality, we assume that the even part  $\hat{\mathcal{L}}_+ = \hat{\mathcal{L}}(\mathbf{L})$  of the Liouvillian is specified by an antisymmetric matrix  $\mathbf{L} \in \mathfrak{so}(4N, \mathbb{C})$  as in [\(7\)](#page-4-4). We first derive the following analog of [\(26\)](#page-9-0) (or, more precisely, [\(29\)](#page-9-3)) which applies to dissipative evolution.

<span id="page-12-5"></span>**Lemma 2.** For  $j < k$ ,  $j, k \in \{1, ..., 2N\}$ , let  $\Lambda(j,k)$  be the  $4N \times 4N$  antisymmetric matrix *with non-zero entries*

<span id="page-12-7"></span><span id="page-12-6"></span>
$$
\Lambda(j,k)_{2j-1+x,2k-1+y} = i^{x+y} \qquad \text{for } x, y \in \{0,1\} \tag{39}
$$

*above the main diagonal and let*  $\Lambda(k, j) = -\Lambda(j, k)$ *. Furthermore, for*  $p = 2j - 1 + x$ *,*  $q =$  $2k-1+y$ , where  $x, y \in \{0,1\}$ , let  $\Gamma(p,q)$  be the antisymmetric  $2N \times 2N$ -matrix with non-zero *entries*

$$
\Gamma(2j - 1 + x, 2k - 1 + y)_{j,k} = (-i)^{x+y} \qquad \text{for } j < k \text{ and } x, y \in \{0, 1\} \tag{40}
$$

*above the main diagonal, and let*  $\Gamma(q, p) = -\Gamma(p, q)$ *. Then* 

$$
\exp(t\mathcal{L}_+^{\dagger})(c_jc_k) = \frac{1}{16} \sum_{r,s=1}^{4N} \sum_{\ell,m=1}^{2N} (R(t)^T \Lambda(j,k) R(t))_{r,s} \Gamma(r,s)_{\ell,m} c_\ell c_m \qquad \text{for all } j < k . \tag{41}
$$

*where*  $R(t) = \exp(\mathbf{L}^{\dagger}t)$ *.* 

*Proof.* We have by [\(5\)](#page-4-0) and [\(6\)](#page-4-1)

<span id="page-12-3"></span>
$$
|c_j c_k\rangle = \frac{1}{8} \sum_{p,q=1}^{4N} \Lambda(j,k)_{p,q} \hat{c}_p \hat{c}_q |I\rangle \qquad \text{for } j \neq k \in \{1, \dots, 2N\}
$$
 (42)

$$
\hat{c}_r \hat{c}_s |I\rangle = \frac{1}{2} \sum_{\ell,m=1}^{4N} \Gamma(r,s)_{\ell,m} |c_\ell c_m\rangle \qquad \text{for } r \neq s \in \{1, \dots, 4N\} \ . \tag{43}
$$

Using [\(9\)](#page-4-2) and unitality, we get

<span id="page-12-4"></span><span id="page-12-2"></span>
$$
\widehat{\exp(t\mathcal{L}_+^{\dagger})\hat{c}_p\hat{c}_q}|I\rangle\!\rangle = \hat{c}_p(t)\hat{c}_q(t)\exp(t\widehat{\mathcal{L}_+^{\dagger}})|I\rangle\!\rangle = \hat{c}_p(t)\hat{c}_q(t)|I\rangle\!\rangle \;, \tag{44}
$$

where

$$
\hat{c}_p(t) = \exp(t\widehat{\mathcal{L}_+^{\dagger}})\hat{c}_p \exp(-t\widehat{\mathcal{L}_+^{\dagger}}) = \sum_{r=1}^{4N} R(t)_{p,r}\hat{c}_r \text{ with } R(t) = e^{\mathbf{L}^{\dagger}t} \in SO(4N, \mathbb{C}) \tag{45}
$$

is the Heisenberg time-evolved super-Majorana operator  $\hat{c}_p$ . Equation [\(45\)](#page-13-0) follows in the same way as [\(26\)](#page-9-0) because we have

<span id="page-13-1"></span><span id="page-13-0"></span>
$$
[\mathcal{L}(\mathbf{L}), \hat{c}_{\ell}] = \sum_{k} \mathbf{L}_{\ell, k} \hat{c}_{k} .
$$
 (46)

in analogy with [\(28\)](#page-9-2). Reinserting [\(45\)](#page-13-0) into [\(44\)](#page-12-2) gives

$$
\widehat{\exp(t\mathcal{L}_+^{\dagger})\hat{c}_p\hat{c}_q}|I\rangle\!\rangle = \sum_{r \neq s}^{4N} R_{p,r}(t) R_{q,s}(t)\hat{c}_r\hat{c}_s|I\rangle\!\rangle \qquad \text{for } p \neq q . \tag{47}
$$

Here we can restrict the sum to  $r \neq s$  because  $R(t)$  is orthogonal and  $\hat{c}_r^2 = I$  for all r. Equation [\(47\)](#page-13-1) is formally analogous to [\(29\)](#page-9-3). The claim follows by computing  $\exp(t\mathcal{L}_+^{\dagger})|c_j c_k\rangle$ using expression [\(42\)](#page-12-3), linearity and [\(47\)](#page-13-1), and then translating the result back with [\(43\)](#page-12-4).

We can use Lemma [2](#page-12-5) to compute the time-evolved covariance matrix  $M(t)$  of a Gaussian state  $\rho$  with covariance matrix  $M(0)$ . Expressing the matrix elements using the Heisenberg picture gives

<span id="page-13-2"></span>
$$
M_{j,k}(t) = i \operatorname{Tr}(\rho \exp(t \mathcal{L}_+^\dagger)(c_j c_k)) \qquad \text{for } j < k \tag{48}
$$

since  $\exp(t\mathcal{L})(\rho) = \exp(t\mathcal{L}_+)(\rho)$ . Theorem [1](#page-10-4) essentially follows by combining this equation with Lemma [2.](#page-12-5) However, showing that the resulting expressions can be evaluated in time  $O(N^3)$ requires some care.

Let  $R = R(t) = \exp(\mathbf{L}^\dagger t)$  be as in Lemma [2](#page-12-5) and let  $\Lambda(j,k)$  and  $\Gamma(r,s)$  be defined by [\(39\)](#page-12-6) and [\(40\)](#page-12-7), respectively. With Lemma [2](#page-12-5) and [\(48\)](#page-13-2) we have

$$
M_{j,k}(t) = \frac{1}{16} \sum_{r,s=1}^{4N} \sum_{\ell,m=1}^{2N} (R^T \Lambda(j,k) R)_{r,s} \Gamma(r,s)_{\ell,m} M_{\ell,m}(0) = \frac{1}{16} \text{Tr} \left( (R^T \Lambda(j,k) R) \Omega(0)^T \right) , \tag{49}
$$

where  $\Omega(0)$  is the  $4N \times 4N$ -matrix defined by the entries

<span id="page-13-4"></span><span id="page-13-3"></span>
$$
\Omega_{r,s}(0) = \text{Tr}\left(\Gamma(r,s)M(0)^T\right) \tag{50}
$$

Using the general identity Tr  $((ABC)D^T)$  = Tr  $((A^TDC^T)B^T)$ , we can rewrite [\(49\)](#page-13-3) as  $M_{j,k}(t)$  =  $\frac{1}{16}$  Tr  $((R\Omega(0)R^T)\Lambda(j,k)^T)$  or

<span id="page-13-5"></span>
$$
M_{j,k}(t) = \frac{1}{16} \operatorname{Tr} \left( \Omega(t) \Lambda(j,k)^T \right) \tag{51}
$$

where

<span id="page-13-6"></span>
$$
\Omega(t) = \exp(\mathbf{L}^{\dagger}t)\Omega(0)\exp(\mathbf{L}^{\dagger}t)^{T}.
$$
\n(52)

Clearly,  $\Omega(0)$  can be obtained from  $M(0)$  in time  $O(N^2)$  since every entry  $\Omega_{r,s}(0)$  is a linear combination of a constant number of entries of M (cf.  $(50)$ ). Similarly, the matrix  $M(t)$  can be computed in time  $O(N^2)$  in an entrywise fashion according to [\(51\)](#page-13-5) given the matrix  $\Omega(t)$ . Therefore, the claim of Theorem [1](#page-10-4) follows since  $\Omega(t)$  can be computed from  $\Omega(0)$  in time  $O(N^3)$ using [\(52\)](#page-13-6). To avoid exponentially growing terms arising from the expression  $\exp(\mathbf{L}^{\dagger}t)$  in (52), we can decompose  $\Omega(t)$  as

<span id="page-14-0"></span>
$$
\Omega(t) = \Omega_b(t) + \Omega_u(t) \tag{53}
$$

using the Jordan decomposition of  $\mathbf{L}^{\dagger}$ , such that the matrix elements of  $\Omega_b(t)$  are bounded for all t, whereas the matrix elements of  $\Omega_u(t)$  are exponentially growing with t. Because  $M(t)$  is the covariance matrix of the time-evolved state  $\exp(t\mathcal{L})(\rho)$ , its entries are bounded for all t and we conclude from [\(51\)](#page-13-5) that the contribution of  $\Omega_u(t)$  must vanish, i.e.,

$$
M_{j,k}(t) = \frac{1}{16} \operatorname{Tr} \left( \Omega_b(t) \Lambda(j,k)^T \right) .
$$

To describe the decomposition [\(53\)](#page-14-0) in more detail, consider the Jordan normal form of  $\mathbf{L}^{\dagger}$ ,

<span id="page-14-1"></span>
$$
\mathbf{L}^{\dagger} = V \bigoplus_{\alpha} J_{\alpha}(\lambda_{\alpha}) V^{-1} \tag{54}
$$

where V is an invertible matrix. Here  $J_{\alpha}(\lambda_{\alpha})$  is a Jordan block

$$
J_{\alpha}(\lambda_{\alpha}) = \lambda_{\alpha} I_{m_{\alpha}} + N_{m_{\alpha}}
$$

where  $I_{m_{\alpha}}$  is the identity matrix of size  $m_{\alpha} \times m_{\alpha}$  and  $N_{m_{\alpha}}$  is the nilpotent matrix with ones on the first upper off-diagonal. With

$$
\exp(J_{\alpha}(\lambda_{\alpha})t) = e^{\lambda_{\alpha}t} \left( I_{m_{\alpha}} + \sum_{k=1}^{m_{\alpha}-1} \frac{t^k}{k!} N_{m_{\alpha}}^k \right) \equiv e^{\lambda_{\alpha}t} S_{m_{\alpha}}(t) ,
$$

and  $\exp(-J_{\alpha}(\lambda_{\alpha})t) = e^{-\lambda_{\alpha}t}S_{m_{\alpha}}(-t)$  we get

$$
\Omega(t) = V \left( \bigoplus_{\alpha,\beta} e^{(\lambda_{\alpha} - \lambda_{\beta})t} S_{m_{\alpha}}(t) V^{-1} \Omega(0) V S_{m_{\beta}}(-t) \right) V^{-1} . \tag{55}
$$

Clearly, all terms corresponding to pairs  $(\alpha, \beta)$  with  $Re(\lambda_{\alpha} - \lambda_{\beta}) > 0$  grow exponentially and must be absorbed in  $\Omega_u(t)$ . Furthermore, for pairs with  $Re(\lambda_\alpha - \lambda_\beta) = 0$ , the contribution of every nilpotent term  $N_m^k$  grows polynomially with t. The remaining terms remain bounded, and we conclude that  $\Omega_b(t)$  is given by

<span id="page-14-2"></span>
$$
\Omega_b(t) = V \left( \bigoplus_{\alpha,\beta:Re(\lambda_\alpha - \lambda_\beta) < 0} e^{(\lambda_\alpha - \lambda_\beta)t} S_{m_\alpha}(t) V^{-1} \Omega(0) V S_{m_\beta}(-t) \right) V^{-1}
$$
\n
$$
+ V \left( \bigoplus_{\alpha,\beta:Re(\lambda_\alpha - \lambda_\beta) = 0} e^{(\lambda_\alpha - \lambda_\beta)t} I_{m_\alpha} V^{-1} \Omega(0) V I_{m_\beta} \right) V^{-1} . \tag{56}
$$

Because computing the Jordan normal form [\(54\)](#page-14-1) takes time  $O(N^3)$ , the matrix  $\Omega_b(t)$  (and hence also  $M(t)$  can be computed in time  $O(N^3)$  using [\(56\)](#page-14-2).

Finally, let us sketch the modifications required to simulate the evolution of higher moments such as [\(30\)](#page-9-4) starting from a non-Gaussian initial state. Clearly, Lemma [2](#page-12-5) can be generalized to give an expansion of the Heisenberg evolved product  $\exp(t\mathcal{L}_+^{\dagger})(c_j c_k c_\ell)$  in terms of a linear combination of Majorana monomials  $c_{j'}c_{k'}c_{\ell'}$ . Inserting this into  $M_{j,k,\ell}(t) = \text{Tr}(\rho \exp(t\mathcal{L}_+^{\dagger})(c_jc_kc_{\ell}))$ immediately gives an explicit expression for the tensor of time-evolved moments  $M(t)$  in terms of the original moments  $M(0)$ .

## <span id="page-15-0"></span>A Computing the steady state

In this appendix we sketch the Bartels-Stewart algorithm [\[1\]](#page-16-9) for solving a matrix equation

<span id="page-15-2"></span>
$$
\mathbf{X}M + M\mathbf{X}^T + \mathbf{Y} = 0\tag{57}
$$

which determines the steady state covariance matrix  $M \equiv M_0$ , see Eq. [\(33\)](#page-11-1) in Section [6.1.](#page-10-1) Here **X** and Y are known real matrices such that  $Y^T = -Y$ , while M is an unknown matrix to be found. All matrices have size  $n \times n$ . As opposed to the third quantization method, where computing the steady state may involve manipulations with ill-conditioned invertible matrices, see Eq. [\(54\)](#page-14-1), the Bartels-Stewart method uses only transformations based on orthogonal (unitary) matrices. This is likely to offer a better numerical stability.

The first step of the algorithm involves the Schur decomposition<sup>[2](#page-15-1)</sup> of  $X$ . This defines a unitary matrix U such that  $X = U^{\dagger} X U$  is upper triangular. Introducing a new unknown matrix  $K = U^{\dagger} M U$  we can rewrite Eq. [\(57\)](#page-15-2) as

<span id="page-15-3"></span>
$$
XK + KX^{\dagger} = Y \tag{58}
$$

where  $Y = -U^{\dagger} Y U$ . Given any matrix Z, the j-th column of Z will be denoted  $Z_j$ . Taking the *n*-th column of Eq.  $(58)$  one arrives at

<span id="page-15-4"></span>
$$
(X + X_{n,n}^* I)K_n = Y_n.
$$
\n(59)

This is a triangular linear system of equations for the unknown vector  $K_n$  which can be solved in time  $O(n^2)$ . Taking the m-th column of Eq. [\(58\)](#page-15-3) one arrives at

$$
(X + X_{m,m}^* I)K_m = Y_m - \sum_{j=m+1}^n X_{m,j}^* K_j, \quad m = n - 1, \dots, 2, 1.
$$
 (60)

Suppose the columns  $K_{m+1}, \ldots, K_n$  are already known. Then the right-hand side of Eq. [\(60\)](#page-15-4) can be formed in time  $O(n^2)$ . We get a triangular system of equations for the unknown vector  $K_m$  which can be solved in time  $O(n^2)$ . Proceeding inductively from  $m = n - 1$  towards  $m = 1$ we can compute the entire matrix K in time  $O(n^3)$ . This gives a solution of Eq. [\(57\)](#page-15-2), namely  $M = UKU^{\dagger}$ . In general, this solution is neither real nor anti-symmetric. However, one can easily check that matrices  $(M + M^*)/2$  and  $(M - M^T)/2$  are solutions of Eq. [\(57\)](#page-15-2) for any (complex) solution  $M$ . Hence we can always transform  $M$  into a real anti-symmetric solution.

<span id="page-15-1"></span><sup>2</sup>To simplify notations, we slightly deviate from the original algorithm of [\[1\]](#page-16-9) which adopted a real Schur decomposition.

Computing the Schur decomposition of a matrix  $X$  is a standard subroutine available in many numerical linear algebra tools (such as MATLAB). Theoretically, it can be computed in time  $O(n^3)$  by finding generalized eigenvectors of **X** and applying the Gram-Schmidt orthog-onalization. A more practical method used in [\[1\]](#page-16-9) involves two steps. First, one transforms  $\bf{X}$ to the upper Hessenberg form by a sequence of  $n-2$  Householder reflections, see [\[31,](#page-18-5) p. 346], which requires  $O(n^3)$  elementary operations. Secondly, **X** is made upper triangular by the QRalgorithm. Each iteration of the QR-algorithm requires  $O(n^2)$  elementary operations, however the required number of iterations is generally unknown.

### Acknowledgments

The authors acknowledge partial support by the DARPA QuEST program under contract number HR0011-09-C-0047.

# <span id="page-16-9"></span>References

- <span id="page-16-6"></span>[1] R. Bartels and G. Stewart. Solution of the matrix equation AX + XB = C. *Comm. of ACM*, 1:820, 1972.
- <span id="page-16-10"></span>[2] C. W. J. Beenakker, D. P. DiVincenzo, C. Emary, and M. Kindermann. Charge detection enables free-electron quantum computation. *Phys. Rev. Lett.*, 93:020501, Jul 2004.
- <span id="page-16-5"></span>[3] R. Bhatia. *Matrix Analysis*. Springer, 1997.
- [4] S. Bravyi. Lagrangian representation for fermionic linear optics. *Quantum Inf. and Comp.*, 5(3):216–238, 2005.
- <span id="page-16-7"></span><span id="page-16-4"></span>[5] S. Bravyi and A. Kitaev. Fermionic quantum computation. *Annals of Physics*, 298(1):210  $-226, 2002.$
- <span id="page-16-8"></span>[6] S. Bravyi and R. K¨onig. Disorder-assisted error correction in Majorana chains, 2011, arXiv:1108.3845.
- [7] A. A. Dzhioev and D. D. Kosov. Super-fermion representation of quantum kinetic equations for the electron transport problem. *J. Chem. Phys.*, 134(044121), 2011.
- <span id="page-16-0"></span>[8] E. Farhi, J. Goldstone, S. Gutmann, J. Lapan, A. Lundgren, and D. Preda. A quantum adiabatic evolution algorithm applied to random instances of an np-complete problem. *Science*, 292(472), 2001.
- <span id="page-16-3"></span><span id="page-16-2"></span>[9] R. Feyman. Simulating physics with computers. *Int. J. Theory. Phys.*, pages 467–488, 1982.
- [10] M. H. Freedman, A. Kitaev, M. J. Larsen, and Z. Wang. Topological quantum computation. *Bull. Amer. Math. Soc.*, 40(1):31–38, 2003, arXiv:quant-ph/0101025.
- <span id="page-16-1"></span>[11] M. H. Freedman, M. Larsen, and Z. Wang. A modular functor which is universal for quantum computation. *Comm. Math. Phys*, pages 605–622, 2002.
- <span id="page-17-7"></span><span id="page-17-3"></span>[12] A. Griessner, A. J. Daley, D. Jaksch, and P. Zoller. Fault-tolerant dissipative preparation of atomic quantum registers with fermions. *Phys. Rev. A*, 72:032332, Sep 2005.
- <span id="page-17-12"></span>[13] S. P. Jordan. Permutational quantum computing. *Quant. Inf. Comput.*, 10:470–497, May 2010.
- [14] R. Jozsa, B. Kraus, A. Miyake, and J. Watrous. Matchgate and space-bounded quantum computations are equivalent. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Science*, 466(2115):809–830, 2009.
- <span id="page-17-11"></span>[15] R. Jozsa and A. Miyake. Matchgates and classical simulation of quantum circuits. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Science*, 464(2100):3089–3106, 2008.
- <span id="page-17-9"></span><span id="page-17-8"></span>[16] A. Kitaev. Unpaired Majorana fermions in quantum wires. In *Mesoscopic And Strongly Correlated Electron Systems conference*, Chernogolovka, Russia, 2000. cond-mat/0010440.
- <span id="page-17-1"></span>[17] A. Kitaev. Anyons in an exactly solved model and beyond. *Annals of Physics*, 321:2–111, 2006.
- <span id="page-17-0"></span>[18] A. Yu. Kitaev. Fault-tolerant quantum computation by anyons. *Annals of Physics*, 303(1):2  $-30, 2003$ .
- <span id="page-17-6"></span>[19] M. Kliesch, T. Barthel, C. Gogolin, M. Kastoryano, and J. Eisert. Dissipative quantum Church-Turing theorem. *Phys. Rev. Lett.*, 107:120501, Sep 2011.
- <span id="page-17-4"></span>[20] E. Knill. Fermionic linear optics and matchgates. Technical Report LAUR-01-4472, Los Alamos National Laboratory, 2001.
- <span id="page-17-2"></span>[21] E. Knill, R. Laflamme, and G. J. Milburn. A scheme for efficient quantum computation with linear optics. *Nature*, 409:46–52, 2001.
- <span id="page-17-13"></span>[22] J. Preskill. Topological quantum computation. Lecture Notes (Chapter 9), 2004.
- <span id="page-17-15"></span>[23] T. Prosen. Third quantization: a general method to solve master equations for quadratic open Fermi systems. *New Journal of Physics*, 10(4):043026, 2008.
- <span id="page-17-14"></span>[24] T. Prosen. Spectral theorem for the Lindblad equation for quadratic open fermionic systems. *Journal of Statistical Mechanics: Theory and Experiment*, 2010(07):P07020, 2010.
- [25] T. Prosen and B. Zunkovic. Exact solution of Markovian master equations for quadratic Fermi systems: thermal baths, open XY spin chains and non-equilibrium phase transition. *New Journal of Physics*, 12(2):025016, 2010.
- <span id="page-17-10"></span><span id="page-17-5"></span>[26] B. M. Terhal and D. P. DiVincenzo. Classical simulation of noninteracting-fermion quantum circuits. *Phys. Rev. A*, 65:032325, 2002.
- [27] L. G. Valiant. Quantum computers that can be simulated classically in polynomial time. *Proceedings of the thirtythird annual ACM symposium on Theory of computing*, pages 114–123, 2001.
- <span id="page-18-4"></span>[28] M. Van den Nest. Quantum matchgate computations and linear threshold gates. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Science*, 467(2127):821–840, 2011.
- <span id="page-18-1"></span><span id="page-18-0"></span>[29] F. Verstraete, M. Wolf, and I. J. Cirac. Quantum computation and quantum-state engineering driven by dissipation. *Nature Physics*, 5(9):633–636, September 2009.
- <span id="page-18-5"></span>[30] H. Weimer, M. Müller, I. Lesanovsky, P. Zoller, and H. P. Büchler. A Rydberg quantum simulator. *Nature Physics*, 6:382–388, 2010.
- <span id="page-18-2"></span>[31] J. H. Wilkinson. *The Algebraic Eigenvalue Problem*. Clarendon Press, Oxford, 1965.
- [32] N. Y. Yao, L. Jiang, A. V. Gorshkov, Z.-X. Gong, A. Zhai, L.-M. Duan, and M. D. Lukin. Robust quantum state transfer in random unpolarized spin chains. *Phys. Rev. Lett.*, 106:040505, Jan 2011.
- <span id="page-18-3"></span>[33] N. Y. Yao, C. R. Laumann, A. V. Gorshkov, H. Weimer, L. Jiang, J. I. Cirac, P. Zoller, and M. D. Lukin. Topologically protected quantum state transfer in a chiral spin liquid, 2011, arXiv:1110.3788.