

Riemann flow and Riemann wave via bialternate product Riemannian metric

Constantin Udriște*

May 3, 2019

Abstract

We illustrate the flow or wave character of the metrics and curvatures of evolution manifolds, introducing the Riemann flow and the Riemann wave via the bialternate product Riemannian metric. This kind of evolutions are new and very natural to understand certain flow or wave phenomena in the nature as well as the geometry of manifolds. It possesses many interesting properties from both mathematical and physical point of views.

Section 1 is a motivation for our point of view. Section 2 discusses the bialternate product Riemannian metric and formulates two basic problems regarding the Riemann flow and Riemann wave. Section 3 defines and studies the Riemann flow. The original results include the meaning of the Riemann flows on constant curvature manifolds and the Riemann solitons. Section 4 analyzes the infinitesimal deformations (linearizations) of Ricci flow and of Riemann flow PDEs. Section 5 shows the existence of Ricci or Riemann curvature blow-up at finite-time singularities of the flow. Also here it is introduced the gradient Riemann soliton. Section 6 defines and studies the Riemann waves and their meaning on constant curvature manifolds. Section 7 proves the existence of Ricci or Riemann curvature blow-up at finite-time singularities of the wave. Section 8 points out the general form of some essential PDE systems on Riemannian manifolds.

Key Words: Riemann flow, Riemann wave, blow-up, ultrahyperbolic PDE system, geometric evolution PDEs.

Mathematics Subject Classification (2010): 58J35; 58J45; 53C44.

1 Motivation

The papers of Richard Hamilton [3]-[7] and some lectures at 1990 Summer Research Institute, Differential Geometry, University of California, Los Angeles,

*Prof. Dr., University Politehnica of Bucharest, Faculty of Applied Sciences, Department of Mathematics-Informatics, Splaiul Independenței 313, Bucharest, 060042, Romania.
E-mail address: udriste@mathem.pub.ro, anet.udri@yahoo.com

July 8-28, stimulated me to think about the Riemann flow and to discuss with other mathematicians about the subject. I have introduced these ideas by an open problem in the book [13] and some explicit results in the paper [17]. Since the Hamilton Ricci flow, defined by

$$\frac{\partial g}{\partial t}(t, x) = -2 \operatorname{Ric}(g(t, x)),$$

was used recently by Grisha Perelman [8]-[10] to prove two very deep theorems in topology, namely the geometrization and Poincaré conjectures, I decided to bring into attention my original ideas, writing again about the Riemann flow and the Riemann wave via the bialternate product Riemannian metric. The Riemann wave is connected to the Ricci wave introduced recently by De-Xing Kong and Kefeng Liu [14], [15].

Within the last years, I was impressed by several researchers which have obtained original results starting from the theory of Hamilton and Perelman (see also [1], [2], [11], [12], [16], [18]). Also, the Research Blog [23] rises the problem of the curvature flow.

2 Bialternate product Riemannian metric

Let (M, g) be a *Riemannian manifold* of dimension n . The Riemannian metric $g = (g_{ij})$ determines the *Christoffel symbols*

$$\begin{aligned} \Gamma_{jk}^i &= \frac{1}{2} g^{il} \left(\frac{\partial g_{lj}}{\partial x^k} + \frac{\partial g_{lk}}{\partial x^j} - \frac{\partial g_{jk}}{\partial x^l} \right) \\ &= \frac{1}{2} g^{im} (\delta_m^r \delta_j^s \delta_k^t + \delta_m^r \delta_k^s \delta_j^t - \delta_m^t \delta_j^r \delta_k^s) \frac{\partial g_{rs}}{\partial x^t} \end{aligned}$$

and this connection determines the *curvature (Riemannian) tensor field* $\operatorname{Riem}(g)$ of components

$$\begin{aligned} R_{ijkl} &= \frac{1}{2} \left(\frac{\partial^2 g_{ik}}{\partial x^j \partial x^l} + \frac{\partial^2 g_{jl}}{\partial x^i \partial x^k} - \frac{\partial^2 g_{jk}}{\partial x^i \partial x^l} - \frac{\partial^2 g_{il}}{\partial x^j \partial x^k} \right) - g_{mn} (\Gamma_{jk}^m \Gamma_{il}^n - \Gamma_{jl}^m \Gamma_{ik}^n) \\ &= 2 \delta_{[i}^p \delta_{j]}^q \delta_{[k}^r \delta_{l]}^s \frac{\partial^2 g_{pr}}{\partial x^q \partial x^s} + 2 \delta_j^q \delta_i^p \delta_{[k}^r \delta_{l]}^s g_{mn} \Gamma_{qr}^m \Gamma_{ps}^n. \end{aligned}$$

It follows the *Ricci tensor field* $\operatorname{Ric}(g)$ of components $R_{jk} = g^{il} R_{ijkl}$ and the *curvature scalar* $R = g^{jk} R_{jk}$.

The Riemannian metric g induces the *bialternate product Riemannian metric*

$$G = g \odot g, \quad G_{ijkl} = g_{ik} g_{jl} - g_{il} g_{jk} = 2 \delta_j^q \delta_i^p \delta_{[k}^r \delta_{l]}^s g_{pr} g_{qs}$$

on 2-forms. Then $(\Lambda^2(M), G = g \odot g)$ is the Riemannian manifold of skew-symmetric 2-forms. If $n = 1$, then $G = 0$. If $n = 2$, then

$$G_{1212} = -G_{1221} = -G_{2112} = G_{2121} = \det(g_{ij}) = \mathcal{G}.$$

If $n \geq 3$, then the bialternate product Riemannian metric G determines the Riemannian metric g since $\text{rank } g = n \geq 3$ and

$$2g_{ij}(g_{ks}G_{mlnr} + g_{kn}G_{lmsr} + g_{kr}G_{mlsn}) = G_{mijn}G_{klrs} + G_{mijs}G_{klnr} + G_{mijr}G_{klsn} \\ - G_{lijn}G_{kmrs} - G_{lijr}G_{kmnr} - G_{lijr}G_{kmsn} - G_{mljs}G_{kirn} - G_{mljr}G_{kins} - G_{mljn}G_{kisir}$$

is an identity.

Problems. 1) Study the flow of the Riemannian tensor field $\text{Riem}(g(t, x))$, via bialternate product Riemannian metric $G(t, x) = g(t, x) \odot g(t, x)$, $t \in \mathbb{R}$, $x \in M$ [13], [18].

2) Study the wave of the Riemannian tensor field $\text{Riem}(g(t, x))$, via bialternate product Riemannian metric $G(t, x) = g(t, x) \odot g(t, x)$, $t \in \mathbb{R}$, $x \in M$.

3 Riemann flow via the bialternate product Riemannian metric

Let M be a smooth closed (compact and without boundary) manifold. Can we equippe the manifold M with a family of smooth Riemannian metrics $g(t, x)$ satisfying the *backward heat ultrahyperbolic PDEs system*

$$\frac{\partial G}{\partial t}(t, x) = -2 \text{Riem}(g(t, x)), \quad (1)$$

where $(\Lambda^2(M), G(t, x) = g(t, x) \odot g(t, x))$ is the associated Riemannian manifold of skew-symmetric 2-forms? The *Riemann flow* via the bialternate product Riemannian metric G is a mean of processing the Riemannian metric $g(t, x)$ by allowing it to evolve under the PDE (1). The classical (backward or forward) heat ultrahyperbolic PDEs with standard Goursat conditions, Cauchy conditions, Picard conditions, mixed conditions are well studied with the help of the energy inequality method and Riemann functions.

Let $\mathcal{G} = \det(g_{ij})$ be the determinant of the metric g . Then

$$\frac{\partial \mathcal{G}}{\partial t} = \frac{\partial \mathcal{G}}{\partial g_{ik}} \frac{\partial g_{ik}}{\partial t} = \mathcal{G} g^{ik} \frac{\partial g_{ik}}{\partial t}, \quad \frac{\partial \ln \mathcal{G}}{\partial t} = g^{ik} \frac{\partial g_{ik}}{\partial t}.$$

Theorem 3.1. *If $n \geq 3$, then the Riemann type flow (1) induces a Ricci flow of the form*

$$\frac{\partial g_{jk}}{\partial t} = -\frac{2}{2-n} R_{jk} + \frac{1}{2-n} \frac{\partial \mathcal{G}}{\partial t} g_{jk}, \quad (1-n) \frac{\partial \ln \mathcal{G}}{\partial t} = -R.$$

Theorem 3.2. *If $n \geq 3$, then the Riemann type flow*

$$\frac{\partial G}{\partial t}(t) = \alpha \text{Riem}(g(t)) + \beta \frac{\partial \mathcal{G}}{\partial t}(t) G(t),$$

with $\alpha = 2(n-2)$, $\beta = \frac{1}{n-1}$, determines a Ricci flow

$$\frac{\partial g_{jk}}{\partial t} = -2 R_{jk}.$$

In simple situations, the Ricci flow was used to deform the metric g into a metric distinguished by its curvature [1]-[13], [16].

Conversely, we have the following statements.

Theorem 3.3. 1) On 3-dimensional Riemannian manifolds, the Ricci flow

$$\frac{\partial g_{ij}}{\partial t} = \frac{R}{2} g_{ij} - 2 R_{ij}$$

determines the Riemann flow (1).

2) On conformally flat Riemannian manifolds with dimension $n \geq 4$, the Ricci flow

$$\frac{\partial g_{ij}}{\partial t} = \frac{R}{2(n-1)(n-2)} g_{ij} - \frac{2}{n-2} R_{ij}$$

determines the Riemann flow (1).

3) On a Riemannian manifold with dimension $n \geq 4$, the Ricci flow

$$\frac{\partial g_{ij}}{\partial t} = \frac{R}{2(n-1)(n-2)} g_{ij} - \frac{2}{n-2} R_{ij}$$

determines a Riemann type flow

$$\frac{\partial G_{ijkl}}{\partial t} = -2 (R_{ijkl} - C_{ijkl}),$$

where C_{ijkl} is the Weyl conformal tensor field.

Hint. The PDEs system

$$\frac{\partial g_{ij}}{\partial t} = \alpha g_{ij} + \beta R_{ij}$$

implies the PDEs system

$$\frac{\partial G_{ijkl}}{\partial t} = 2\alpha G_{ijkl} + \beta (R_{ik}g_{jl} + g_{ik}R_{jl} - R_{il}g_{jk} - g_{il}R_{jk}).$$

On the other hand,

1) in dimension three, the curvature tensor field has the local expression

$$R_{ijkl} = g_{ik}R_{jl} + g_{jl}R_{ik} - g_{il}R_{jk} - g_{jk}R_{il} - \frac{R}{2} G_{ijkl};$$

2) the curvature tensor field of a conformally flat Riemannian manifold, with dimension $n \geq 4$, is given by

$$R_{ijkl} = \frac{1}{n-2} (g_{ik}R_{jl} + g_{jl}R_{ik} - g_{il}R_{jk} - g_{jk}R_{il}) - \frac{R}{(n-1)(n-2)} G_{ijkl}.$$

3) The Weyl conformal tensor field is

$$C_{ijkl} = R_{ijkl} - \frac{1}{n-2} (g_{ik}R_{jl} + g_{jl}R_{ik} - g_{il}R_{jk} - g_{jk}R_{il}) + \frac{R}{(n-1)(n-2)} G_{ijkl}.$$

Theorem 3.4 (short time existence). *Given a smooth metric g_0 on a closed manifold M , there exist $\epsilon > 0$ and a smooth family of metrics $g(t)$ for $t \in [0, \epsilon]$ such that*

$$G(t) = g(t) \circ g(t), \quad \frac{\partial G}{\partial t} = -2 \text{Riem}(g(t)), \quad t \in [0, \epsilon], \quad g(0) = g_0.$$

Proof. Let \bar{g}_t be a solution to the Ricci flow $\frac{\partial g_{jk}}{\partial t} = -2 R_{jk}$. Let us find a time-dependent diffeomorphism φ_t from our manifold M to itself, with $\varphi_0 = id$, so that the metric $g_t = \varphi_t^* \bar{g}_t$ is a solution to the Riemann flow. Since

$$\frac{\partial g_t}{\partial t} = \varphi_t^* \frac{\partial \bar{g}_t}{\partial t} + \varphi_t^* \mathcal{L}_{\frac{\partial \varphi_t}{\partial t}} \bar{g}_t,$$

we obtain

$$\frac{\partial G_t}{\partial t} = \varphi_t^* \frac{\partial \bar{G}_t}{\partial t} + \varphi_t^* \mathcal{L}_{\frac{\partial \varphi_t}{\partial t}} \bar{G}_t.$$

On the other hand,

$$\varphi_t^* \frac{\partial \bar{G}_t}{\partial t} + \varphi_t^* \mathcal{L}_{\frac{\partial \varphi_t}{\partial t}} \bar{G}_t = -2 \varphi_t^* (\text{Riem}(\bar{g}_t) - C(\bar{g}_t)).$$

We select the diffeomorphism φ_t by the condition $\mathcal{L}_{\frac{\partial \varphi_t}{\partial t}} \bar{G}_t = 2 \varphi_t^* C(\bar{g}_t)$.

Theorem 3.5 (uniqueness of solutions). *Suppose $g_1(t)$ and $g_2(t)$ are two Riemann flows on a closed manifold M , for $t \in [0, \epsilon]$, $\epsilon > 0$. If $g_1(s) = g_2(s)$ for some $s \in [0, \epsilon]$, then $g_1(t) = g_2(t)$ for all $t \in [0, \epsilon]$.*

Combining these two Theorems, we can talk about the Riemann flow with initial metric g_0 , on a maximal time interval $[0, T)$. In this situation, maximal means that either $T = \infty$, or that $T < \infty$ but there do not exist $\epsilon > 0$ and a smooth Riemann flow $\hat{g}(t)$ for $t \in [0, T + \epsilon)$ such that $\hat{g}(t) = g(t)$ for $t \in [0, T)$.

If the initial metric $g(0, x)$ is Riemann flat, i.e., $\text{Riem}(g(0, x)) = 0$, then $g(t, x) = g(0, x)$ is obviously a solution of the evolution PDE (1). Consequently, each Riemann flat metric $g(x)$ is a steady solution of the heat ultrahyperbolic PDEs system.

4 Infinitesimal deformations

4.1 Linearizing Ricci flow PDE

Let $g(t, x)$, $t \in I \subset \mathbb{R}$, be a Ricci flow. Let $g(t, x; \epsilon)$ be a differentiable variation of $g(t, x)$, i.e.,

$$\frac{\partial g}{\partial t}(t, x; \epsilon) = -2 \text{Ric}(g(t, x; \epsilon)), \quad g(t, x; 0) = g(t, x).$$

We take the partial derivative with respect to ϵ and we denote $\frac{\partial g}{\partial \epsilon}|_{\epsilon=0} = h$. The relations $g_{ij}g^{jk} = \delta_i^k$ implies

$$\frac{\partial g^{jk}}{\partial \epsilon}|_{\epsilon=0} = -h_{ij}g^{jk}g^{il} = -h^{lk}.$$

We find the *single-time infinitesimal deformation PDE*

$$\frac{\partial h}{\partial t} = -2 \left(\frac{\partial \text{Ric}}{\partial g^{ij}}(-h^{ij}) + \frac{\partial \text{Ric}}{\partial \frac{\partial g_{kl}}{\partial x^i}} \frac{\partial h_{kl}}{\partial x^i} + \frac{\partial \text{Ric}}{\partial \frac{\partial^2 g_{kl}}{\partial x^i \partial x^j}} \frac{\partial^2 h_{kl}}{\partial x^i \partial x^j} \right)$$

around a solution $g(t, x)$ of the Ricci PDE. Of course, this new PDE is the linearization of the Ricci PDE along a solution $g(t, x)$.

4.2 Linearizing Riemann flow PDE

Let $g(t, x)$, $t \in I \subset R$, be a Riemann flow. Let $g(t, x; \epsilon)$ be a differentiable variation of $g(t, x)$, i.e.,

$$\frac{\partial G}{\partial t}(t, x; \epsilon) = -2 \text{Riem}(g(t, x; \epsilon)), \quad g(t, x; 0) = g(t, x).$$

We take the partial derivative with respect to ϵ and we denote $\frac{\partial g}{\partial \epsilon}|_{\epsilon=0} = h$. The relations $g_{ij}g^{jk} = \delta_i^k$ implies

$$\frac{\partial g^{jk}}{\partial \epsilon}|_{\epsilon=0} = -h_{ij}g^{jk}g^{il} = -h^{lk}.$$

We find the *single-time infinitesimal deformation PDE*

$$\frac{\partial}{\partial t}(h \odot g + g \odot h) = -2 \left(\frac{\partial \text{Riem}}{\partial g^{ij}}(-h^{ij}) + \frac{\partial \text{Riem}}{\partial \frac{\partial g_{kl}}{\partial x^i}} \frac{\partial h_{kl}}{\partial x^i} + \frac{\partial \text{Riem}}{\partial \frac{\partial^2 g_{kl}}{\partial x^i \partial x^j}} \frac{\partial^2 h_{kl}}{\partial x^i \partial x^j} \right)$$

around a solution $g(x, t)$ of the Riemann PDE. Of course, this new PDE is the linearization of the Riemann PDE along a solution $g(t, x)$.

5 Blow-ups at finite-time singularities for geometric flows

5.1 Ricci estimation and Ricci curvature blow-up at finite-time singularities

This subsection will be devoted to proving that, if the Ricci flow $g(t, x)$, defined by

$$\frac{\partial g}{\partial t}(t, x) = -2 \text{Ric}(g(t, x)),$$

becomes singular in finite time, i.e., the Ricci curvature must explode as we approach the singular time T .

Lemma 4.1 (Riemannian metric equivalence). *If $g(t)$ is a Ricci flow for $t \in [0, T)$ and $\|\text{Ric}\| < m$ on $M \times [0, T)$, then*

$$e^{-2mt}g(0) \leq g(t) \leq e^{2mt}g(0), \forall t \in [0, T),$$

$$\|g(t)\|^2 - \|g(0)\|^2 \leq C e^{4m^2t}, \forall t \in [0, T).$$

Remark. Given two symmetric $(0, 2)$ -tensors a and b , we write $a \geq b$ to mean that $a - b$ is positive semidefinite.

Proof. Since $\frac{\partial}{\partial t} g(X, X) = -2 \text{Ric}(X, X)$, we have

$$\left| \frac{\partial}{\partial t} g(X, X) \right| = 2 \|\text{Ric}\| g(X, X)$$

and hence

$$\left| \frac{\partial}{\partial t} \ln g(X, X) \right| \leq 2 \|\text{Ric}\|,$$

for any nonzero tangent vectors X, Y . In this way,

$$\left| \ln \frac{g(t)(X, X)}{g(0)(X, X)} \right| \leq 2mt.$$

To obtain the second formula, we remark that

$$\frac{1}{2} \frac{\partial}{\partial t} \|g\|^2 = -2 \langle \text{Ric}, g \rangle \leq 2 \|\text{Ric}\|^2 \|g\|^2$$

implies

$$\|g(t)\|^2 - \|g(0)\|^2 \leq C e^{4 \int_0^t \|\text{Ric}\|^2 ds}.$$

Theorem 4.2 *Let M be a closed manifold. If $g(t)$ is a Ricci flow on a maximal time interval $[0, T)$ and $T < \infty$, then*

$$\lim_{t \nearrow T} \left(\sup_{x \in M} \|\text{Ric}(t, x)\| \right) = \infty.$$

Proof. We will deal with the contrapositive of this theorem. Let us show that if $\|\text{Ric}(t, x)\| \leq m$, $t \in [0, T)$, then the Ricci flow $g(t)$ can be extended to a larger time interval $[0, T + \epsilon)$. Indeed, the relation

$$g(t) = g(0) - 2 \int_0^t \text{Ric}(s, x) ds, \quad t \in [0, T)$$

implies

$$\|g(t') - g(t'')\| \leq 2m|t' - t''|, \quad \forall t', t'' \in [0, T).$$

The Cauchy Criterion shows that $\lim_{t \nearrow T} g(t)$ exists, while $\lim_{t \nearrow T} \frac{\partial g}{\partial t}(t)$ exists since $\lim_{t \nearrow T} \text{Ric}(t, x)$ exists (see the definition of the Ricci flow). Consequently $g(t)$, $t \in [0, T]$ is a solution for the Ricci flow.

In this case, the Ricci flow $g(t)$ may be extended from being a smooth solution on $[0, T)$ to a smooth solution on $[0, T]$. Then we take $g(T)$ to be an 'initial' metric in our short-time existence theorem (Theorem 3.4) in order to extend the flow to a Ricci flow for $t \in [0, T + \epsilon)$, contradicting the assumption that $[0, T)$ is a maximal time interval.

5.2 Riemann estimation and curvature blow-up at finite-time singularities

This subsection will be devoted to proving that, if the Riemann flow becomes singular in finite time, i.e., the curvature must explode as we approach the singular time T .

Lemma 4.3 (bialternate product Riemannian metric equivalence). *If $g(t)$ is a Riemann flow for $t \in [0, T)$ and $\|\text{Riem}\| < m$ on $M \times [0, T)$, then*

$$e^{-2mt}G(0) \leq G(t) \leq e^{2mt}G(0), \forall t \in [0, T),$$

$$\|G(t)\|^2 - \|G(0)\|^2 \leq C e^{4m^2t}, \forall t \in [0, T).$$

Proof. Since $\frac{\partial}{\partial t} G(X, Y, X, Y) = -2 \text{Riem}(X, Y, X, Y)$, we have

$$\left| \frac{\partial}{\partial t} G(X, Y, X, Y) \right| = 2 \|\text{Riem}\| |G(X, Y, X, Y)|$$

and hence

$$\left| \frac{\partial}{\partial t} \ln G(X, Y, X, Y) \right| \leq 2 \|\text{Riem}\|,$$

for any nonzero tangent vectors X, Y . In this way,

$$\left| \ln \frac{G(t)(X, Y, X, Y)}{G(0)(X, Y, X, Y)} \right| \leq 2mt.$$

To obtain the second formula, we remark that

$$\frac{1}{2} \frac{\partial}{\partial t} \|G\|^2 = -2 \langle \text{Riem}, G \rangle \leq 2 \|\text{Riem}\|^2 \|G\|^2$$

implies

$$\|G(t)\|^2 - \|G(0)\|^2 \leq C e^{4 \int_0^t \|\text{Riem}\|^2 ds}.$$

Theorem 4.4 *Let M be a closed manifold with the dimension $n \geq 3$. If $g(t)$ is a Riemann flow on a maximal time interval $[0, T)$ and $T < \infty$, then*

$$\lim_{t \nearrow T} \left(\sup_{x \in M} \|\text{Riem}(t, x)\| \right) = \infty.$$

Proof. Let us show that if $\|\text{Riem}(t, x)\| \leq m$, $t \in [0, T)$, then the Riemann flow $g(t)$ can be extended to a larger time interval $[0, T + \epsilon)$. Indeed, the relation

$$G(t) = G(0) - 2 \int_0^t \text{Riem}(s, x) ds, \quad t \in [0, T)$$

implies

$$\|G(t') - G(t'')\| \leq 2m|t' - t''|, \quad \forall t', t'' \in [0, T).$$

The Cauchy Criterion shows that $\lim_{t \nearrow T} G(t)$ exists, while $\lim_{t \nearrow T} \frac{\partial G}{\partial t}(t)$ exists since $\lim_{t \nearrow T} \text{Riem}(t, x)$ exists (see the definition of the Riemann flow). Consequently $g(t)$, $t \in [0, T]$ is a solution for the Riemann flow.

In this case, the Riemann flow $g(t)$ may be extended from being a smooth solution on $[0, T)$ to a smooth solution on $[0, T]$. Then we take $g(T)$ to be an 'initial' metric in our short-time existence theorem (Theorem 3.4) in order to extend the flow to a Riemann flow for $t \in [0, T + \epsilon)$, contradicting the assumption that $[0, T)$ is a maximal time interval.

5.3 Riemann flows on constant curvature manifolds

If we take a metric g_0 such that $\text{Riem}(g_0) = \lambda G_0$ for some constant $\lambda \in R$ (these metrics are known as constant curvature metrics), then a solution $G(t)$ of PDE (1) with $G(0) = G_0$ is given by $G(t) = (1 - 2\lambda t) G_0$ and a solution $g(t)$ of PDE (1) with $g(0) = g_0$ is given by $g(t) = \sqrt{1 - 2\lambda t} g_0$. (It is worth pointing out here that the Riemann tensor is invariant under uniform scalings of the metric). In particular, for the round unit sphere (S^n, g_0) , $n \geq 2$, we have $\text{Riem}(g_0) = (n - 1)G_0$, so the evolution metric is $g(t) = \sqrt{1 - 2(n - 1)t} g_0$ and the sphere collapses to a point at finite time $T = \frac{1}{2(n - 1)}$. An alternative example of this type would be if g_0 were a hyperbolic metric, that is, of constant sectional curvature -1 . In this case, $n \geq 2$, $\text{Riem}(g_0) = -(n - 1)G_0$, the evolution metric is $g(t) = \sqrt{1 + 2(n - 1)t} g_0$ and the manifold expands homothetically for all time. Of course a metric is flat, i.e., the Riemannian manifold is locally isometric to the Euclidean space, if and only if the associated Riemann curvature tensor vanishes. These metrics can be regarded as the fixed points of the Riemann flow.

5.4 Gradient Riemann soliton

We can regard as "generalized fixed points" of the Riemann flow those manifolds which change only by a diffeomorphism and a rescaling under the Riemann flow. To simplify, instead of $g(t, x)$ we shall write $g(t)$. Let $(M, g(t))$ be a solution of the Riemann flow, and suppose $\varphi_t : M \rightarrow M$, $\varphi_0 = id$ is a time-dependent family of diffeomorphisms and $\sigma(t)$ is a time-dependent scale factor (with $\sigma(0) = 1$). If we have $g(t) = \sigma(t)\varphi_t^* g(0)$, then the solution $(M, g(t))$ or $(\Lambda^2(M), G(t) = g(t) \odot g(t), t \in R)$ is called *Riemann soliton*. Of course

$$G_{ijkl}(t) = g_{ik}(t)g_{jl}(t) - g_{il}(t)g_{jk}(t)$$

$$= \sigma^2(t) \varphi_t^* g_{ik}(0) \varphi_t^* g_{jl}(0) - \sigma^2(t) \varphi_t^* g_{il}(0) \varphi_t^* g_{jk}(0).$$

Computing the derivative $\frac{\partial G}{\partial t}(t)$ and evaluating it at $t = 0$, we find

$$\begin{aligned} \frac{\partial G}{\partial t}(t)|_{t=0} &= 2\sigma(t) \frac{d\sigma}{dt}(t) \varphi_t^* g(t) \circ \varphi_t^* g(t)|_{t=0} \\ &+ \sigma^2(t) \left(\frac{\partial}{\partial t} \varphi_t^* g(t) \circ \varphi_t^* g(t) + \varphi_t^* g(t) \circ \frac{\partial}{\partial t} \varphi_t^* g(t) \right) |_{t=0} \\ &= 2\sigma'(0) g(0) \circ g(0) + \mathcal{L}_V g(0) \circ g(0) + g(0) \circ \mathcal{L}_V g(0) = -2 \text{Riem}(g(0)), \end{aligned}$$

where $V = \frac{\partial \varphi_t}{\partial t}$, by the definition of *Lie derivative*. We replace $\sigma'(0) = \lambda$ and we write in coordinates

$$\begin{aligned} -2R_{ijkl} &= 2\lambda(g_{ik}(0)g_{jl}(0) - g_{il}(0)g_{jk}(0)) \\ &+ \mathcal{L}_V g_{ik}(0)g_{jl}(0) + g_{ik}(0)\mathcal{L}_V g_{jl}(0) - \mathcal{L}_V g_{il}(0)g_{jk}(0) - g_{il}(0)\mathcal{L}_V g_{jk}(0) \\ &= 2\lambda(g_{ik}(0)g_{jl}(0) - g_{il}(0)g_{jk}(0)) \\ &+ (\nabla_i V_k + \nabla_k V_i)g_{jl}(0) + g_{ik}(0)(\nabla_j V_l + \nabla_l V_j) \\ &- (\nabla_i V_l + \nabla_l V_i)g_{jk}(0) - g_{il}(0)(\nabla_j V_k + \nabla_k V_j). \end{aligned}$$

Suppose $V = \nabla f$. Then the foregoing relation becomes

$$\begin{aligned} R_{ijkl} + \lambda(g_{ik}(0)g_{jl}(0) - g_{il}(0)g_{jk}(0)) \\ + g_{jl}(0)\nabla_i \nabla_k f + g_{ik}(0)\nabla_j \nabla_l f - g_{jk}(0)\nabla_i \nabla_l f - g_{il}(0)\nabla_j \nabla_k f = 0. \end{aligned}$$

Such solutions we call *gradient Riemann solitons*. A gradient Riemann soliton is called *shrinking* if $\lambda < 0$, *static* if $\lambda = 0$, and *expanding* if $\lambda > 0$.

6 Riemann wave via the bialternate product Riemannian metric

Let M be a smooth closed (compact and without boundary) manifold. Can we equippe the manifold M with a family of smooth Riemannian metrics $g(t, x)$ satisfying the *wave ultrahyperbolic PDEs system*

$$\frac{\partial^2 G}{\partial t^2}(t, x) = -2 \text{Riem}(g(t, x)), \quad (2)$$

where $(\Lambda^2(M), G(t, x) = g(t, x) \odot g(t, x))$ is the associated Riemannian manifold of skew-symmetric 2-forms? The *Riemann wave* via the bialternate product Riemannian metric G is a mean of processing the Riemannian metric $g(t, x)$ by allowing it to evolve under the PDE (2).

Theorem 4.1 (uniqueness of solution). *Let $(M, g_0(x))$ be a compact Riemannian manifold and $k_1(x)$ be a $(0, 2)$ symmetric tensor field on M . Then there exists a constant $\epsilon > 0$ such that the initial value problem*

$$\frac{\partial^2 G}{\partial t^2}(t, x) = -2 \text{Riem}(g(t, x)), \quad g(0, x) = g_0(x), \quad \frac{\partial g}{\partial t}(0, x) = k_1(x)$$

has a unique smooth solution $g(t, x)$ on $M \times [0, \epsilon]$.

Theorem 4.2. *If $(M, g_{ij}(0, x) = u_0(x^1)\delta_{ij}, u_0(x^1) > 0)$ is a conformally flat Riemannian manifold, then the Cauchy problem*

$$\frac{\partial^2 G}{\partial t^2}(t, x) = -2 \text{Riem}(g(t, x)), \quad g(0, x) = g_0(x), \quad \frac{\partial g}{\partial t}(0, x) = k_1(x)$$

has a unique smooth solution for all time $t \in R$, and the solution metric has the form $g_{ij} = u(t, x^1)\delta_{ij}$, $u(t, x^1) > 0$.

Hint. Let (M, g) be a conformally flat Riemannian manifold. If we take a Riemannian metric with the components $g_{ii} = H_i^2$, $g_{ij} = 0$, $i \neq j$ (see Lamé coefficients), then

$$R_{hijk} = 0, \quad (h, i, j, k \neq)$$

$$R_{hiik} = -H_i \left(\frac{\partial^2 H_i}{\partial x^h \partial x^k} - \frac{\partial H_i}{\partial x^h} \frac{\partial \ln H_h}{\partial x^k} - \frac{\partial H_i}{\partial x^k} \frac{\partial \ln H_h}{\partial x^h} \right), \quad (h, i, k \neq)$$

$$R_{hiih} = -H_h H_i \left(\frac{\partial}{\partial x^h} \left(\frac{1}{H_h} \frac{\partial H_i}{\partial x^h} \right) + \frac{\partial}{\partial x^i} \left(\frac{1}{H_i} \frac{\partial H_h}{\partial x^i} \right) + \sum_l \frac{1}{H_l^2} \frac{\partial H_h}{\partial x^l} \frac{\partial H_i}{\partial x^l} \right),$$

where the index of summation l indicates the sum for the values $1, \dots, n$ excepting h and i . Replacing $g_{ij} = u(t, x^1)\delta_{ij}$, we find

$$\left(\frac{\partial u}{\partial t} \right)^2 + \left(\frac{\partial u}{\partial x^1} \right)^2 + u \left(\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^{1^2}} \right) = 0,$$

with the initial conditions $u(0, x^1) = u_0(x^1)$, $\frac{\partial u}{\partial t}(0, x^1) = u_1(x^1)$.

6.1 Riemann waves on constant curvature manifolds

If we take a metric g_0 such that $\text{Riem}(g_0) = \lambda G_0$ for some constant $\lambda \in R$ (these metrics are known as constant curvature metrics), then a solution $G(t)$ of PDE (2) with $G(0) = G_0$ is given by $G(t) = e^{-2\lambda t} G_0$ and a solution $g(t)$ of PDE (2) with $g(0) = g_0$ is given by $g(t) = e^{-\lambda t} g_0$. (It is worth pointing out here that the Riemann tensor is invariant under uniform scalings of the metric). In particular, for the round unit sphere (S^n, g_0) , $n \geq 2$, we have $\text{Riem}(g_0) = (n-1)G_0$, so the evolution metric is $g(t) = e^{-(n-1)t} g_0$ and the sphere collapses to a point when $t \rightarrow \infty$. An alternative example of this type would be if g_0 were a hyperbolic metric, that is, of constant sectional curvature -1 . In this case, $n \geq 2$, $\text{Riem}(g_0) = -(n-1)G_0$, the evolution metric is $g(t) = e^{(n-1)t} g_0$ and

the manifold expands homothetically for all time. If the initial metric $g(0, x)$ is Riemann flat, i.e., $\text{Riem}(g(0, x)) = 0$, then $g(t, x) = g(0, x)$ is obviously a solution of the evolution PDE (2). Consequently, each Riemann flat metric $g(x)$ is a steady solution of the wave ultrahyperbolic PDEs system.

Open problem. The case of multitime Ricci wave or multitime Riemann wave PDEs does not seem to have been investigated. The study of this case is caused not only by theoretical interest, but also by practical necessity of describing the deformations (see [18]-[22] for inspiration).

7 Blow-ups at finite-time singularities for geometric waves

Theorem 6.1 *Let M be a closed manifold. If $g(t)$ is a Ricci wave, i.e.,*

$$\frac{\partial^2 g}{\partial t^2}(t, x) = -2 \text{Ric}(g(t, x)),$$

on a maximal time interval $[0, T)$ and $T < \infty$, then

$$\lim_{t \nearrow T} \left(\sup_{x \in M} \|\text{Ric}(t, x)\| \right) = \infty.$$

Proof. Let us show that if $\|\text{Ric}(t, x)\| \leq m$, $t \in [0, T)$, then the Ricci wave $g(t)$ can be extended to a larger time interval $[0, T + \epsilon)$. Indeed, the relations

$$\frac{\partial g}{\partial t}(t) = \frac{\partial g}{\partial t}(0) - 2 \int_0^t \text{Ric}(s, x) ds, \quad t \in [0, T),$$

$$g(t) = g(0) + \frac{\partial g}{\partial t}(0) t - 2 \int_0^t \int_0^u \text{Ric}(s, x) ds du, \quad t \in [0, T)$$

imply

$$g(t') - g(t'') = \frac{\partial g}{\partial t}(0) (t' - t'') + 2 \int_{t'}^{t''} \int_0^u \text{Ric}(s, x) ds du, \quad t \in [0, T)$$

and hence

$$\|g(t') - g(t'')\| \leq \left(\left\| \frac{\partial g}{\partial t}(0) \right\| + 2mT \right) |t' - t''|, \quad \forall t', t'' \in [0, T).$$

The Cauchy Criterion shows that $\lim_{t \nearrow T} g(t)$ exists, while $\lim_{t \nearrow T} \frac{\partial g}{\partial t}(t)$ and $\lim_{t \nearrow T} \frac{\partial^2 g}{\partial t^2}(t)$ exist since $\lim_{t \nearrow T} \text{Ric}(t, x)$ exists (see the definition of the Ricci wave). Consequently $g(t)$, $t \in [0, T]$ is a Ricci wave.

In this case, the Ricci wave $g(t)$ may be extended from being a smooth solution on $[0, T)$ to a smooth solution on $[0, T]$. Then we take $g(T)$, $\frac{\partial g}{\partial t}(T)$ to

be an 'initial' metric in a short-time existence theorem in order to extend the wave to a Ricci wave for $t \in [0, T + \epsilon)$, contradicting the assumption that $[0, T)$ is a maximal time interval.

Theorem 6.2 *Let M be a closed manifold with the dimension $n \geq 3$. If $g(t)$ is a Riemann wave, i.e.,*

$$\frac{\partial^2 G}{\partial t^2}(t, x) = -2 \text{Riem}(g(t, x)),$$

on a maximal time interval $[0, T)$ and $T < \infty$, then

$$\lim_{t \nearrow T} \left(\sup_{x \in M} \|\text{Riem}(t, x)\| \right) = \infty.$$

Proof. Let us show that if $\|\text{Riem}(t, x)\| \leq m$, $t \in [0, T)$, then the Riemann wave $g(t)$ can be extended to a larger time interval $[0, T + \epsilon)$. Indeed, the relations

$$\frac{\partial G}{\partial t}(t) = \frac{\partial G}{\partial t}(0) - 2 \int_0^t \text{Riem}(s, x) ds, \quad t \in [0, T),$$

$$G(t) = G(0) + \frac{\partial G}{\partial t}(0)t - 2 \int_0^t \int_0^u \text{Riem}(s, x) ds du, \quad t \in [0, T)$$

imply

$$G(t') - G(t'') = \frac{\partial g}{\partial t}(0)(t' - t'') + 2 \int_{t'}^{t''} \int_0^u \text{Riem}(s, x) ds du, \quad t \in [0, T)$$

and hence

$$\|G(t') - G(t'')\| \leq \left(\left\| \frac{\partial G}{\partial t}(0) \right\| + 2mT \right) |t' - t''|, \quad \forall t', t'' \in [0, T).$$

The Cauchy Criterion shows that $\lim_{t \nearrow T} G(t)$ exists, while $\lim_{t \nearrow T} \frac{\partial G}{\partial t}(t)$ and $\lim_{t \nearrow T} \frac{\partial^2 G}{\partial t^2}(t)$ exist since $\lim_{t \nearrow T} \text{Riem}(t, x)$ exists (see the definition of the Riemann wave). Consequently $g(t)$, $t \in [0, T]$ is a Riemann wave.

In this case, the Riemann wave $g(t)$ may be extended from being a smooth solution on $[0, T)$ to a smooth solution on $[0, T]$. Then we take $g(T)$, $\frac{\partial g}{\partial t}(T)$ to be an 'initial' metric in a short-time existence theorem in order to extend the wave to a Riemann wave for $t \in [0, T + \epsilon)$, contradicting the assumption that $[0, T)$ is a maximal time interval.

8 Conclusions

On Riemannian manifolds we can discuss about the ultrahyperbolic PDEs system (constant curvature manifolds) $R_{ijkl} = \lambda G_{ijkl}$. On Riemannian manifolds

whose metric depends on the evolution parameter t , we can add: heat ultrahyperbolic PDEs system (Riemann flow), $\frac{\partial G_{ijkl}}{\partial t} = -2 R_{ijkl}$; wave ultrahyperbolic PDEs system (Riemann wave), $\frac{\partial^2 G_{ijkl}}{\partial t^2} = -2 R_{ijkl}$. In fact, the Riemann flow and the Riemann wave are PDE systems for evolving the Riemannian metric to make it "rounder", in the hope that one may draw topological conclusions from the existence of such "round" metrics. The general form of these PDE systems is

$$\alpha_{ijkl} \frac{\partial^2 G_{ijkl}}{\partial t^2} + \beta_{ijkl} \frac{\partial G_{ijkl}}{\partial t} + \gamma_{ijkl} G_{ijkl} + \delta_{ijkl} R_{ijkl} = 0,$$

where the coefficients are smooth functions depending on t, x . All these determine new classes of Riemannian metrics.

There are more ingenious ways to define such PDEs using spacetimes.

Acknowledgement

Partially supported by University Politehnica of Bucharest and by Academy of Romanian Scientists, Bucharest, Romania.

References

- [1] C. Hopper, B. Andrews, *The Ricci Flow in Riemannian Geometry*, Springer, 2009.
- [2] D. Máximo, *Non-negative Ricci curvature on closed manifolds under Ricci flow*, Proceedings of the American Mathematical Society, 139, 2 (2011), 675-685.
- [3] H. Richard, *Three-manifolds with positive Ricci curvature*, J. Diff. Geo., 17 (1982), 255–306.
- [4] H. Richard, *Four-manifolds with positive curvature operator*, J. Diff. Geo., 24 (1986), 153–179.
- [5] H. Richard, *A compactness property for solutions of the Ricci flow*, American Journal of Mathematics, 117 (1992), 545–572.
- [6] H. Richard, *The formation of singularities in the Ricci flow*, Surveys in Differential Geometry, 2 (1995), 1–136.
- [7] H. Richard, *Non-singular solutions of the Ricci flow on three-manifolds*, Comm. Anal. Geom., 7 (1999), 695–729.
- [8] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, <http://arXiv.org/math.DG/0211159v1>, 2002.

- [9] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, <http://arXiv.org/math.DG/0307245v1>, 2002.
- [10] G. Perelman, *Ricci flow with surgery on three-manifolds*, <http://arXiv.org/math.DG/0303109v1>, 2003.
- [11] N. Sheridan, *Hamilton's Ricci Flow*, PhD Thesis, The University of Melbourne, Department of Mathematics and Statistics, 2006.
- [12] J. Porti, *Hamilton-Ricci flow on three dimensional manifolds*, Fall Workshop in Geometry and Physics, Bilbao, 2005.
- [13] B. Chow, D. Knopf, *The Ricci Flow: An Introduction*, AMS Math. Surveys and Monographs, 2004.
- [14] D.-X. Kong, K. Liu, *Wave character of metrics and hyperbolic geometric flow*, <http://www.cms.zju.edu.cn/UploadFiles/AttachFiles/200682885946597.pdf>
- [15] D.-X. Kong, K. Liu, D.-L. Xu, *The hyperbolic geometric flow on Riemann surfaces*, arXiv:0709.1607v3 [math.DG] 9 Jan 2008.
- [16] P. Topping, *Lectures on the Ricci flow*, 2006.
- [17] C. Udriște, *Riemann flow and Riemann wave*, An. Univ. Vest, Timișoara, Ser. Mat.-Inf., 47, 2 (2009), 1-10.
- [18] C. Udriște, M. Ferrara, D. Opreș, *Economic Geometric Dynamics*, Monographs and Textbooks 6, Geometry Balkan Press, Bucharest, 2004,
- [19] C. Udriște, *Simplified multitime maximum principle*, Balkan J. Geom. Appl., 14, 1 (2009), 102-119.
- [20] C. Udriște, *Nonholonomic approach of multitime maximum principle*, Balkan J. Geom. Appl., 14, 2 (2009), 111-126.
- [21] C. Udriște, *Multitime maximum principle for curvilinear integral cost*, Balkan J. Geom. Appl., 16, 1 (2011), 128-149.
- [22] C. Udriște, A. Bejenaru, *Multitime optimal control with area integral costs on boundary*, Balkan J. Geom. Appl., 16, 2 (2011), 138-154.
- [23] Research Blog 2/28/03, www.math.uic.edu/agol/blog/030228.pdf