

Open circle maps: Small hole asymptotics

Carl Dettmann

School of Mathematics, University of Bristol, BS8 1TW, UK

May 14, 2022

Abstract

We consider escape from chaotic maps through a subset of phase space, the hole. Escape rates are known to be locally constant functions of the hole position and size. In spite of this, for the doubling map we can extend the current best result for small holes, a linear dependence on hole size h , to include a smooth $h^2 \ln h$ term and explicit fractal terms to h^2 and higher orders. The results are confirmed by numerical simulations, followed by a discussion of the occurrence of $h^2 \ln h$ in more general chaotic maps.

1 Introduction

Here we consider open dynamical systems, in which motion is considered only until the trajectory reaches a specified subset of phase space, the “hole(s)”. The initial conditions are distributed with respect to some measure, typically an invariant measure on the phase space of the corresponding closed system (the “escape” problem), or on such a measure restricted to hole(s) (“recurrence” or “transport” problems). There are also relevant questions where there are two systems connected by a hole (the “metastability” problem). There is a vast literature in mathematics and physics on open dynamical systems and applications, some of which is mentioned in [1, 7, 8].

For the escape problem, we expect that for strongly chaotic systems (eg with exponential decay of correlations), the survival probability $P(n) = \mu(M_n)$ decays exponentially with time n , so that an escape rate can be defined:

$$\gamma = \lim_{n \rightarrow \infty} -\frac{1}{n} \ln P(n) \quad (1)$$

Here μ is the measure of initial conditions (assumed to be invariant under the dynamics) and M_n the subset of the phase space M that survives for at least n iterations before reaching the hole $H \subset M$. In general the limit might not exist, and we need upper and lower escape rates, possibly infinite. Here we restrict to maps, considering only a discrete time n ; similar questions apply to flows.

It is of interest to consider how the escape rate depends on the hole, in particular its size $h = \mu(H)$. On one hand, there have been attempts to quantify the small hole asymptotics of the escape rate, that is, its behaviour for a sequence of holes contracting to a point $x \in M$. Ref. [13] numerically observed the effects of periodic orbits on the escape rate from a chaotic map with a small hole. Ref. [4] expresses escape rates in terms of correlation functions, finding good numerical agreement in the case of diamond billiard with continuous time escape rate computed to order h^2 ; the heuristic arguments proposed there suggest that for billiards with holes typically considered (holes that allow particles to escape that reach a small region of the boundary, but from any direction) the first order result is impervious to a short periodic orbit located in the hole, but that corrections would be needed for the second order formulas. Ref. [10] shows rigorously that

$$\gamma = h(1 - \Lambda_x^{-1}) + o(h) \quad (2)$$

for a wide class of hyperbolic maps, where Λ_x is the stability eigenvalue of the periodic orbit starting from $x \in M$, infinite if the orbit of x is aperiodic. This result (amongst others) was also shown independently in a study of the open doubling map [5].

On the other hand it is known that the largest invariant set that never reaches the hole, a subset of M_∞ , is a locally constant function of the hole. This is also mentioned in [10] but was noted much earlier [16]. It then implies the escape rate, and also properties such as the relevant Lyapunov exponents and entropy, are

locally constant. This can be seen (but was not remarked on) for the diamond billiard in Fig. 4 of Ref. [4]. While putting an apparent limit on small hole asymptotic results, this effect is also related to the very helpful observation that open dynamics restricted to the repeller is a normally a subshift of finite type; almost always in the case of the doubling map considered here. For very recent results on the existence, continuity and locally constant properties of the escape rate for more general hyperbolic systems, see Ref. [6].

We note that the locally constant property does not exclude the possibility that the escape rate is smoother as $h \rightarrow 0$ than the single derivative required by Eq. (2). This paper is an attempt to push the small hole asymptotics as far as possible in the simplest possible dynamical system, the doubling map, with the hope of elucidating similar effects in more general contexts.

2 Existing results

The doubling map $x \rightarrow 2x \pmod{1}$ has an invariant measure μ simply given by Lebesgue, so the hole size h is simply the length of the relevant interval(s). There is a correspondence between the binary representation of a point $x = 0.a_1a_2\dots$ and symbolic dynamics for the partition $\{[0, 1/2), [1/2, 1)\}$, modulo minor details to do with the dyadic rationals (ie fractions with denominator a power of two).

The case of a hole between 0 and h has received some attention in the literature, also including more general expanding circle maps [9, 16]. In brief, given a hole size h , find

$$h_- = \sup_{x \leq h} \{2^n x > x \pmod{1}, \quad \forall n \in \mathbb{N}\} \quad (3)$$

This is closely related to the concept of a “minimal prefix” [12]. The quantity h_- is almost always a dyadic rational $i/2^n$ for some integer i , and in this case h lies in the interval of constant escape rate $i/2^n \leq h \leq i/(2^n - 1)$, often abbreviated here as just “interval.” Note that each periodic orbit of the map appears exactly once as the upper limit of one of these intervals.

Within an interval, the dynamics is a subshift of finite type. If we write the (n term) binary expansion

$$1 - h_- = \sum_{k=1}^n a_k 2^{-k} \quad (4)$$

then the leading eigenvalue of the transition matrix is $\beta/2$ where β is the solution of

$$1 = \sum_{k=1}^n a_k \beta^{-k} \quad (5)$$

with $\beta \approx 2$. Thus we find the escape rate

$$\gamma = -\ln \frac{\beta}{2} \quad (6)$$

The set of irrational h_- has local dimension $\beta/2$, and the escape rate is computed as in the rational case (but taking $n \rightarrow \infty$) since it is continuous and monotonic. Thus the case of $H = [0, h]$ is of interest for β -expansions.

3 Small hole asymptotics

We now consider the question as to what asymptotic results are possible in this case. The intervals of constant escape rate give an important constraint on the possible form of such asymptotics: γ is constant, but h has varied by an amount proportional to h^2 (roughly equal to h^2 when $i = 1$). Thus γh^2 has jumps of unit size occurring over a short distance relative to h itself: the $O(h^2)$ term in the expansion, if it exists, must have jump discontinuities.

For any h contained in an interval, we have $\gamma(h) = \gamma(h_-)$ so it is sufficient to consider just values h_- on the left of an interval. From the monotonicity and continuity we may furthermore calculate γ as a limit of

rational or irrational h_- as convenient. For the following argument we use irrational $h = h_-$ and the binary expansion of h rather than the $1 - h$ above:

$$h = 2^{-J} K_0 = 2^{-J} \sum_{k=1}^{\infty} A_k 2^{-k} \quad (7)$$

Here, $J = (\ln K_0 - \ln h)/\ln 2$ is a non-negative integer so that $K_0 < 1$. There are at least two logical choices for J : We can set $1/2 \leq K_0 < 1$ so that K_0 (and each K_m below) is bounded above and below; this is most helpful in determining the magnitude of the relevant terms. The other choice is to set $J = 0$ which simplifies the computations. For now we need to keep J arbitrary. The other new quantities in the above expression are $A_k = 1 - a_{J+k}$, with all the non-positive A_k equal to zero. Thus the expansion for β becomes

$$1 = \sum_{k=1}^{\infty} a_k \beta^{-k} = (\beta - 1)^{-1} - \sum_{k=1}^{\infty} A_k \beta^{-J-k} \quad (8)$$

We know from [10] that to leading order $\gamma \approx h/2$, since the fixed point at $x = 0$ has $\Lambda = 2$. Thus we write

$$\beta = 2 - \sum_{l=1}^{\infty} \epsilon_l h^l \quad (9)$$

where $\epsilon_1 = 1$, and we allow the other ϵ_l to be functions $o(h^{-1})$ as $h \rightarrow 0$. Substituting for β , expanding both powers of β in a binomial expansion, substituting h/K_0 for 2^{-J} and equating coefficients with equal powers of h , we find that

$$\epsilon_2 = -1 + \frac{J}{2} + \frac{K_1}{2K_0} \quad (10)$$

$$\epsilon_3 = 1 + \frac{3J^2}{8} + \frac{J}{8} \left(\frac{6K_1}{K_0} - 11 \right) + \frac{1}{8} \left(\frac{K_2}{K_0} + \frac{2K_1^2}{K_0^2} - \frac{11K_1}{K_0} \right) \quad (11)$$

and so on, where

$$K_m = \sum_{k=1}^{\infty} A_k k^m 2^{-k} \quad (12)$$

consistent with the definition of K_0 above. The K_m for $m > 0$ are fractal functions bounded on $[1/2, 1)$ as shown in Fig. 1; they are discontinuous at dyadic rationals.

We also see that at second order the expansion involves Jh^2 which gives rise to a smooth $h^2 \ln h$ term. Thus we can extend Ref. [10] result to

$$\gamma = \frac{h}{2} - \frac{h^2 \ln h}{4 \ln 2} + \frac{h^2}{4} \left(-\frac{3}{2} + \frac{\ln K_0}{\ln 2} + \frac{K_1}{K_0} \right) + \dots \quad (13)$$

for h an irrational value in the fractal set not contained in any interval.

If h lies in an interval $[h_-, h_+] = [i/2^n, i/(2^n - 1)]$ we identify it with the discontinuity occurring at $K_0 = i/2^{(n-J)}$. The jump at this point is $K_1^- - K_1^+ = 2^{(1-(n-J))}$ independent of i , which leads to a cancellation to second order with the discrepancy in the first term, since $h_- - h_+ = i/2^{2n}$ to this order; all other terms are smooth. Thus, the jumps in γ are exactly those required by the locally constant intervals.

In what sense does this expression converge in the $h \rightarrow 0$ limit? Considering a fixed domain of K_0 , intervals containing $2^{-J} K_0$ decrease with increasing J . Thus we expect convergence for almost all fixed $K_0 \in (1/2, 1)$ as $J \rightarrow \infty$, where $h = 2^{-J} K_0$, with an error of smaller order than the last retained term. On the other hand, there is no guarantee that for fixed J the terms with higher powers of h continue to decrease.

Recall that we did not specify J exactly; we can see that the transformation $J \rightarrow J + 1$ has the effects $K_0 \rightarrow K_0/2$ and $K_1 \rightarrow (K_1 + K_0)/2$, leaving the above expression for γ unaffected. But this also means that the coefficient of h^2 is log-periodic, ie a periodic function of $\ln h$. We can thus write the expansion of γ as a series involving h , $h^2 \ln h$ and $h^{2+2\pi i q/\ln 2}$ for $q \in \mathbb{Z}$, however the coefficients (Fourier expansion of the K functions in the variable $\ln h$) do not appear to have a simple analytic form. They are also slow to converge due to the discontinuities. We can continue analogously with h^3 and higher, indicating that the Mellin transform of γ has a semi-infinite periodic forest of poles reminiscent of those for Julia sets [2].




Figure 1: The K_m functions defined in Eq. (12).

Using the arbitrariness in J , we now set $J = 0$ to yield the simplest form of these expressions: we can substitute h for K_0 and evaluate the other K functions at h , giving the main result:

$$\gamma = \gamma_1 + \gamma_2 + \gamma_3 + O(h^4 \ln^3 h) \quad (14)$$

$$\gamma_1 = \frac{h}{2} \quad (15)$$

$$\gamma_2 = \frac{2K_1 h - 3h^2}{8} \quad (16)$$

$$\gamma_3 = \frac{6K_1^2 h + 3K_2 h^2 - 27K_1 h^2 + 14h^3}{48} \quad (17)$$

where we note that $K_m(h) \sim h \ln^m h / \ln^m 2$ as $h \rightarrow 0$.

Fig. 2 shows a comparison between the escape rate calculated directly and with the above formulas. The direct numerical computation uses a Markov matrix of side length 2^{16} (2^{19} for $h < 0.01$) that does not need to be stored due to its regular structure. The leading eigenvalue corresponding to γ is obtained by repeated multiplication of the matrix on a vector with initially identical nonzero entries, using long double (19 digit) precision. The direct escape rate is plotted, then subtracting each of the γ_i in turn, leading to smaller and smaller remainders. We can see that the locally constant intervals (smooth curves in the direct computation) become smaller for smaller h , but remain as spikes at dyadic rationals. Elsewhere, the remainder decreases with the order of the approximation, showing good convergence despite the fractal character of the functions involved.

4 Discussion

We have argued that, despite its locally constant form, the escape rate of the doubling map for small holes is smoother than required by the existing linear result, Eq. (2), having a smooth $h^2 \ln h$ term, and non-smooth h^2 and higher terms which is however expressible in terms of explicit discontinuous fractal functions K_m , with the expansion formally proceeding to arbitrary order $h^n \ln^m h$ with $m < n$.




Figure 2: Numerical test of the small hole expansion for the escape rate, Eq. (14).

Various connections may be made between the K functions and self-affine functions and sets appearing in the literature. The graph of K_1 is easily seen to be a discontinuous self-affine set, see Ref. [14] for some relevant properties. It shares the relation

$$K_1(x/2) = (K_1(x) + x)/2 \quad (18)$$

with the Takagi function, used to study diffusion in extended doubling maps [11], but the similarity appears to end there, as the Takagi function is continuous. We can construct a generating function for all the K functions, (cf Eq. 5)

$$G_\beta(x) = \sum_{k=1}^{\infty} A_k \beta^{-k} \quad (19)$$

so that, at least formally, we have

$$K_n(x) = (-\beta \partial_\beta)^n G_\beta(x)|_{\beta=2} \quad (20)$$

The limits here are quite subtle, since $G_\beta(x)$ effectively computes the Bernoulli convolution corresponding to β , and it is known that the distribution of the latter is singular when β is Pisot, including a sequence accumulating at $\beta = 2$ [15]; it might be interesting to investigate these connections further.

A more challenging question concerns the applicability of the escape rate results to more general maps, for example considered in Ref. [6]. While the present calculation depends rather heavily on an explicit formula for a polynomial equation satisfied by γ , there are good reasons to believe the $h^2 \ln h$ term is much more widely applicable. In a general hyperbolic map, the linear dependence of the escape rate on the periodic orbit stability arises because that is the extent of the “shadow” [3] cast by the hole on itself under iteration around the periodic orbit. Beyond this, we need to study the shadow of the hole on itself along orbits that do not remain close to the periodic orbit (if there is one) passing through the hole. The shortest of these are typically of length $|\ln h|$, with stabilities of order h^{-1} . This will make a piece of the hole of size roughly h^2 inaccessible to the escaping trajectories. At the next length there will be more periodic orbits, balanced by greater instabilities, also yielding of order h^2 . This continues until double the original length, say $2|\ln h|$ beyond which the periodic orbits are effectively just shadowing combinations of shorter periodic orbits. Thus

we expect these to give a total contribution to the escape rate of order $h^2 \ln h$, and predict a term of this form irrespective of whether the hole includes a short periodic orbit, for a general class of hyperbolic maps.

Acknowledgements

The author gratefully acknowledges helpful discussions with Mark Demers during his visit to Bristol in June 2011, which was partly funded by a London Mathematical Society scheme 2 grant, and also discussions with Orestis Georgiou, Thomas Jordan, Rainer Klages and Georgie Knight.

References

- [1] Altmann E. & Tel, T. [2009], “Poincaré recurrences and transient chaos in systems with leaks,” *Phys. Rev. E* **79**, 016204.
- [2] Bessis, D., Geronimo, J. S. & Moussa, P. [1984], “Mellin transforms associated with Julia sets and physical applications,” *J. Stat. Phys.* **34**, 75-110.
- [3] Buljan H. & Paar, V. [2001], “Many-hole interactions and the average lifetimes of chaotic transients that precede controlled periodic motion,” *Phys. Rev. E* **63**, 066205.
- [4] Bunimovich, L. A. & Dettmann, C. P. [2007], “Peeping at chaos: Nondestructive monitoring of chaotic systems by measuring long-time escape rates,” *EPL*, **80** 40001.
- [5] Bunimovich, L. A. & Yurchenko, A. [2008,11], “Where to place a hole to achieve a maximal escape rate,” arXiv:0811.4438; *Israel. J. Math.*, **182**, 229-252.
- [6] Demers, M. F. & Wright, P. [2011], “Behavior of the escape rate function in hyperbolic dynamical systems”, arXiv:1112.4812.
- [7] Demers, M. F., & Young, L.-S. [2006], “Escape rates and conditionally invariant measures” *Nonlinearity* **19** 377-397.
- [8] Dettmann, C. P., “Recent advances in open billiards with some open problems” in *Frontiers in the study of chaotic dynamical systems with open problems*, (Ed. Z. Elhadj and J. C. Sprott, World Scientific, 2011)
- [9] Jensen, J. L. [2011], “On lower bounded orbits of the times- q map”, arXiv:1106.2932
- [10] Keller, G. & Liverani, C. [2009], “Rare events, escape rates and quasistationarity: Some exact formulae” *J. Stat. Phys.* **135**, 519-534.
- [11] Knight, G., Georgiou, O., Dettmann, C. P. & Klages, R. [2011], “Where to place a hole to achieve a maximal diffusion coefficient, ” arXiv:1112.3922.
- [12] Nilsson, J. [2009], “On numbers badly approximable by dyadic rationals,” *Israel J. Math.*, **171**, 93-110.
- [13] Paar, V. & Pavin, N. [1997], “Bursts in average lifetime of transients for chaotic logistic map with a hole”, *Phys. Rev E*, **55**, 4112-4115.
- [14] Protasov, V. Y., [2006], “Fractal curves and wavelets,” *Izv. Math.*, **70**, 975-1013.
- [15] Solomyak, B., [2004] “Notes on Bernoulli convolutions,” in *Fractal Geometry and Applications: A Jubilee of Benot Mandelbrot: Analysis, Number Theory, and Dynamical Systems* (Ed. M. L. Lapidus and M. van Frankenhuijsen, Amer. Math. Soc., 2004)
- [16] Urbanski, M. [1986], “On the Hausdorff dimension of invariant sets for expanding maps of a circle” *Ergod. Th. & Dynam. Sys.* **6**, 295-309.