

**ASYMPTOTIC STABILITY  
FOR A CLASS OF MARKOV SEMIGROUPS**

BEBE PRUNARU

ABSTRACT. Let  $U \subset K$  be an open and dense subset of a compact metric space and let  $\{\Phi_t\}_{t \geq 0}$  be a Markov semigroup on the space of bounded Borel measurable functions on  $U$  with the strong Feller property. Suppose that for each  $x \in \partial U$  there exists a barrier  $h \in C(K)$  at  $x$  such that  $\Phi_t(h) \geq h$  for all  $t \geq 0$ . Suppose also that every real-valued  $g \in C(K)$  with  $\Phi_t(g) \geq g$  for all  $t \geq 0$  and which attains its global maximum at a point inside  $U$  is constant. Then for each  $f \in C(K)$  there exists the uniform limit  $F = \lim_{t \rightarrow \infty} \Phi_t(f)$ . Moreover  $F$  is continuous on  $K$ , agrees with  $f$  on  $\partial U$  and  $\Phi_t(F) = F$  for all  $t \geq 0$ .

Let  $E$  be a locally compact Hausdorff space and let  $M_b(E)$  be the space of all complex-valued, bounded and Borel measurable functions on  $E$ . Let also  $C_b(E)$  be the set of all continuous functions in  $M_b(E)$ .

A linear map  $\Phi : M_b(E) \rightarrow M_b(E)$  is called a Markov operator if for each  $x \in E$  there exists a probability Borel measure  $\mu_x$  on  $E$  such that

$$\Phi(f)(x) = \int f d\mu_x \quad \forall f \in M_b(E).$$

A Markov operator  $\Phi$  is said to have the strong Feller property if  $\Phi(f) \in C_b(E)$  for every  $f \in M_b(E)$ . The main result of this paper is the following:

**Theorem 1.** *Let  $K$  be a compact metric space and let  $U \subset K$  be a dense open subset such that  $\partial U$  contains at least two points. Let  $C(K)$  be the space of all continuous complex-valued functions on  $K$  and let  $C(\partial U)$  be the corresponding space for  $\partial U$ .*

*Let  $\Phi : M_b(U) \rightarrow M_b(U)$  be a Markov operator with the strong Feller property. Suppose that  $\Phi$  satisfies the following conditions:*

- (A) *For each point  $x \in \partial U$  there exists  $h \in C(K)$  such that  $h(x) = 0$ ,  $h(y) < 0$  for all  $y \in K \setminus \{x\}$  and  $\Phi(h_U) \geq h_U$  on  $U$ , where  $h_U$  is the restriction of  $h$  to  $U$ ;*
- (B) *If  $g \in C(K)$  is a real valued function with  $\Phi(g_U) \geq g_U$  and if there exists  $z \in U$  such that  $g(z) = \max\{g(x) : x \in K\}$  then  $g$  is constant on  $K$ .*

*Then, for each  $f \in C(\partial U)$  there exists a unique function  $G \in C(K)$  such that  $\Phi(G_U) = G_U$  and  $G(x) = f(x)$  for all  $x \in \partial U$ . Moreover, if  $F \in C(K)$  is an arbitrary continuous extension of  $f$  to  $K$  then the sequence  $\{\Phi^n(F_U)\}_n$  converges uniformly on  $U$  to  $G_U$ .*

*Proof.* (1) First of all, it can be proved, exactly as in Proposition 1.3 from [1], that the boundary condition (A) implies the following. For each  $f \in C(K)$  and for each

---

2000 *Mathematics Subject Classification.* Primary: 47D07, Secondary: 60J05, 60J25, 37A30, 47A35.

*Key words and phrases.* Markov semigroup, boundary value problem, asymptotic stability.

$x \in \partial U$

$$\limsup_{y \rightarrow x, n \geq 1} |(\Phi^n(f_U))(y) - f(x)| = 0.$$

This enables us to define a map

$$\Psi : C(K) \rightarrow C(K)$$

by setting, for each  $f \in C(K)$ ,  $\Psi(f)(x) = \Phi(f_U)(x)$  for  $x \in U$  and  $\Psi(f)(x) = f(x)$  for  $x \in \partial U$ .

(2) We show that if  $f \in C(K)$  is a real-valued function such that  $\Phi(f_U) \geq f_U$  on  $U$ , then the sequence  $\{\Psi^n(f)\}$  converges uniformly on  $K$ . Indeed, this sequence is monotone increasing and if  $g$  is its pointwise limit then  $g$  is Borel measurable and  $\Phi(g_U) = g_U$ . Since  $\Phi$  has the strong Feller property, it follows that  $g_U$  is continuous on  $U$ . Moreover (1) implies that

$$\lim_{y \rightarrow x} g(y) = g(x)$$

for every  $x \in \partial U$ . This shows that  $g \in C(K)$  and Dini's theorem shows that the convergence is uniform on  $K$ .

(3) Let

$$\mathcal{T}(\Psi) = \{h \in C(K) : \Psi(h) = h\}$$

and let  $A$  be the norm closed subalgebra of  $C(K)$  generated by  $\mathcal{T}(\Psi)$ . Let  $C(\Psi)$  be the set of all  $f \in C(K)$  for which the sequence  $\{\Psi^n(f)\}$  is uniformly convergent on  $K$  and denote  $\pi(f)$  its limit. Let also

$$C(\Psi)_0 = \{f \in C(\Psi) : \pi(f) = 0\}.$$

We will show that  $A \subset C(\Psi)$ .

Let  $h \in \mathcal{T}(\Psi)$ . Then  $\Psi(|h|^2) \geq |h|^2$  therefore (2) shows that  $|h|^2 \in C(\Psi)$  and also that  $\pi(|h|^2) - |h|^2 \geq 0$ . This also shows that the norm closed ideal of  $C(K)$  generated by all the functions of the form  $\pi(|h|^2) - |h|^2$  with  $h \in \mathcal{T}(\Psi)$  is contained in  $C(\Psi)_0$ . Indeed it is easy to see that if  $f \in C(\Psi)_0$  is non-negative on  $K$  then  $fg \in C(\Psi)_0$  for every  $g \in C(K)$ .

We shall now prove that for any finite set of  $k$  functions from  $\mathcal{T}(\Psi)$  their product belongs to  $C(\Psi)$ . Let  $k = 2$ . If  $h_1, h_2 \in \mathcal{T}(\Psi)$  then

$$h_1 h_2 = (1/4) \sum_{m=0}^3 i^m |g_m|^2$$

where  $i = \sqrt{-1}$  and  $g_m = (h_1 + i^m h_2)$ . Since  $g_m \in \mathcal{T}(\Psi)$  we see that  $h_1 h_2 \in C(\Psi)$ .

Let  $k \geq 3$  and assume that every product of at most  $k - 1$  elements from  $\mathcal{T}(\Psi)$  belongs to  $C(\Psi)$ . Let  $h_1, \dots, h_k$  in  $\mathcal{T}(\Psi)$  and let  $g = h_1 \cdots h_k$ . Then

$$g = (h_1 h_2 - \pi(h_1 h_2)) \cdot h_3 \cdots h_k + \pi(h_1 h_2) \cdot h_3 \cdots h_k.$$

By what we have already proved the first summand belongs to  $C(\Psi)_0$  and the second belongs, by the induction hypothesis, to  $C(\Psi)$ . This shows that  $A \subset C(\Psi)$ .

(4) Consider the map

$$\rho : A \rightarrow C(\partial U)$$

that takes any  $f \in A$  into its restriction to  $\partial U$ . It turns out that  $\rho$  is onto. To see this, we first observe that the boundary condition (A) together with (2) implies that for each  $x \in \partial U$  there exists  $g \in C(K)$  such that  $g(x) = 0$ ,  $g(y) < 0$  for every  $y \in \partial U \setminus \{x\}$  and  $\Phi(g_U) = g_U$ . In particular  $g \in \mathcal{T}(\Psi)$ . This shows that the range

of  $\rho$  separates the points of  $\partial U$  therefore  $\rho$  is onto. This proves the existence part of the theorem. Uniqueness follows easily from (B).

(5) We shall denote

$$\theta : C(K) \rightarrow C(K)$$

the map which takes a function  $f \in C(K)$  into the uniquely determined function in  $\mathcal{T}(\Psi)$  which agrees with  $f$  on  $\partial U$ . It follows that for each  $f \in C(\Psi)$  we have  $\theta(f) = \pi(f)$ . In particular,  $\theta(f) = f$  for every  $f \in \mathcal{T}(\Psi)$ .

Let

$$L = \{g \in C(K) : g = \pi(|h|^2) - |h|^2 \text{ for some } h \in \mathcal{T}(\Psi)\}.$$

If  $g \in L$ , then it follows from (3) that  $g \geq 0$ . Moreover  $\Psi(g) \leq g$  hence  $\Phi(g_U) \leq g_U$ .

Suppose now that there exists a point  $z_0 \in U$  such that  $g(z_0) = 0$  for every  $g \in L$ . It then follows from (B) that  $g = 0$  on  $U$  hence on  $K$  for every  $g \in L$ . This easily implies that  $\mathcal{T}(\Psi)$  is closed under multiplication hence it equals  $A$ . Indeed if  $h_1$  and  $h_2$  are functions in  $\mathcal{T}(\Psi)$  then  $\pi(h_1 h_2) - h_1 h_2$  can be written as a linear combination of elements from  $L$  (see step 3). It then follows that  $\pi(h_1 h_2) = h_1 h_2$  therefore  $\mathcal{T}(\Psi) = A$ .

This shows that the map  $\theta : C(K) \rightarrow C(K)$  defined above is multiplicative on  $C(K)$  and its range equals  $A$ . Let  $M$  be the maximal ideal space of  $A$  and for each  $h \in A$  let  $\hat{h} \in C(M)$  be its Gelfand transform. Since  $A$  is self-adjoint, the Gelfand transform is an isometric isomorphism from  $A$  onto  $C(M)$  (the Banach algebra of all continuous, complex-valued functions on  $M$ ). It then follows that there exists a continuous one-to-one map  $\gamma : M \rightarrow K$  such that  $\widehat{\theta(f)} = f \circ \gamma$  for every  $f \in C(K)$ . Moreover, since  $A \subset C(K)$  there exists a continuous surjective map  $\lambda : K \rightarrow M$  such that  $h = \hat{h} \circ \lambda$  for every  $h \in A$ .

We now claim that  $\gamma(M) \subset \partial U$ . Suppose there exists  $\alpha \in M$  such that  $z = \gamma(\alpha) \in U$ . Then there exists a non-constant real valued function  $h \in A$  such that

$$\hat{h}(\alpha) = \sup\{h(x) : x \in K\}.$$

Every such  $h$  attains its maximum only on  $\partial U$ . However

$$\hat{h}(\alpha) = h(\gamma(\alpha)) = h(z)$$

This shows that  $\gamma(M) \subset \partial U$ .

Let  $z \in U$  and let  $x = \gamma \circ \lambda(z)$ . Then  $x \in \partial U$ . Let  $h \in \mathcal{T}(\Psi)$  which attains its maximum on  $K$  only at this point. Then

$$h(x) = (h \circ \gamma \circ \lambda)(z) = (\hat{h} \circ \lambda)(z) = h(z).$$

We get a contradiction. This means that there is no  $z_0 \in U$  such that  $g(z_0) = 0$  for all  $g \in L$ .

(6) It follows from (5) that

$$\partial U = \{x \in K : g(x) = 0 \text{ for every } g \in L\}.$$

It then follows that the closed ideal of  $C(K)$  generated by  $L$  is precisely the ideal of all  $f \in C(K)$  which vanish identically on all points of  $\partial U$ . Recall now that we already proved in part 3 that for any  $g$  in the closed ideal generated by  $L$  the sequence  $\{\Psi^n(g)\}$  converges uniformly to 0. In particular, this holds true for all functions  $g$  of the form  $g = f - \theta(f)$  with  $f \in C(K)$ . This completes the proof of this theorem. □

This proof works as well for Markov semigroups with continuous parameter. Examples of Markov operators on complex domains which satisfy all the conditions in Theorem 1 are given in [1]. As a matter of fact, the results and the methods used in [1] strongly motivated and inspired our research. We have also used some ideas appearing in [3]. Theorem 1 can be used to give an alternate proof for the main result in [2].

#### REFERENCES

- [1] J. Arazy and M. Englis, *Iterates and the boundary behavior of the Berezin transform*, Ann. Inst. Fourier, **51** (2001) 1101-1133.
- [2] C. Liu, *Iterates of a Berezin-type transform*, J. Math. Anal. Appl. **329** (2007) 822-829.
- [3] B. Prunaru, *Lifting fixed points of completely positive semigroups*, Integral Equations and Operator Theory, to appear.

INSTITUTE OF MATHEMATICS "SIMION STOILOW" OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA  
*E-mail address:* Bebe.Prunaru@imar.ro