Michiel Kosters - Mathematisch Instituut, Universiteit Leiden mkosters@math.leidenuniv.nl, November 5, 2021

The subset sum problem for finite abelian groups

1. Abstract

Let G be a finite abelian group. For $g \in G$ and i an integer we define $N(i,g) = \#\{S \subseteq G : \#S = i, \sum_{s \in S} s = g\}$, the number of subsets of G of size i which sum up to g. We will give a short proof, using character theory, of a formula for these N(i,g) due to Li and Wan. We also give a formula for $N(i,g)^* = \#\{S \subseteq G \setminus \{0\} : \#S = i, \sum_{s \in S} s = g\}$, which generalizes another result of Wan.

Keywords: subset sum, finite abelian group. Classification MSC2010: 11B75, 20K99.

2. Main theorem

In this article we fix a finite abelian group G of size n and we let $0 \le i \le n$.

Definition 2.1. For $g \in G$ we define $N(i,g) = \#\{S \subseteq G : \#S = i, \sum_{s \in S} s = g\}$.

Definition 2.2. For $g \in G$ we define $e(g) = \max\{d : d | \exp(G), g \in dG\}$.

Remark that $e(g) = \operatorname{lcm}\{d : d | \exp(G), g \in dG\}$. Indeed, if $g = d_1g_1 = d_2g_2$ with $g_1, g_2 \in G$ and $d_1, d_2 \in \mathbb{Z}$ with $\operatorname{gcd}(d_1, d_2) = 1$, then there are integers such that $1 = n_1d_1 + n_2d_2$. Hence we have

$$g = (n_1d_1 + n_2d_2)g = d_1d_2(n_1g_1 + n_2g_2).$$

Let μ be the Möbius function and for an integer d let $G[d] = \{h \in G : dh = 0\}$, the d-torsion of G.

Our main theorem is the following one.

Theorem 2.3. We have the following formula for $g \in G$:

$$N(i,g) = \frac{1}{n} \sum_{\substack{s \mid \gcd(\exp(G),i)}} (-1)^{i + \frac{i}{s}} \binom{n/s}{i/s} \sum_{\substack{d \mid \gcd(e(g),s)}} \mu(\frac{s}{d}) \# G[d].$$

Remark 2.4. The above theorem is a slight improvement of Theorem 1.1 in [1]. In [1] a rather long proof is given using some sieving techniques.

Let \hat{G} be the set of characters on G with identity $\chi_0 = 1$. Note that \hat{G} is a group isomorphic to G. We make the following observation:

$$\sum_{i=0}^{n} \sum_{g \in G} N(i,g)gX^{i} = \prod_{\sigma \in G} (1+\sigma X) \in \mathbf{C}[G][X].$$

We use the following easy fact. If $\alpha = \sum_{g \in G} \alpha_g g \in \mathbf{C}[G]$, then $\alpha_g = \frac{1}{n} \sum_{\chi \in \hat{G}} \overline{\chi}(g) \chi(\alpha)$ (here we extend χ to a **C**-linear map $\chi : \mathbf{C}[G] \to \mathbf{C}$). This gives the following formula, where we put Y = -X:

$$\sum_{i=0}^{n} N(i,g)(-Y)^{i} = \frac{1}{n} \sum_{\chi \in \hat{G}} \overline{\chi}(g) \prod_{\sigma \in G} \left(1 - \chi(\sigma)Y\right).$$

Lemma 2.5. Suppose that $\chi \in \hat{G}$ has order m. Then $\prod_{\sigma \in G} (1 - \chi(\sigma)Y) = (1 - Y^m)^{n/m}$.

Proof. Observe that $\prod_{i=0}^{m} (1 - \zeta_m^i Y) = 1 - Y^m$, and that we have n/m of such products.

Lemma 2.6. We have

$$\sum_{\chi \in \hat{G}: \chi^m = 1} \chi(g) = \begin{cases} \#G[m] & g \in mG \\ 0 & g \notin mG. \end{cases}$$

Proof. If $\chi^m = 1$, we know that the character factors through G/mG. A character on G/mG induces such a character on G. Hence we deduce the result from standard character theory.

Now define the following function:

$$f(s) = \sum_{d \mid s} \sum_{\chi \in \hat{G}: \operatorname{ord}(\chi) = d} \overline{\chi}(g).$$

From the lemma above we have $f(s) = \delta_{g \in sG} \# G[s]$. Using the Möbius inversion formula we find, for $s | \exp(G)$,

$$\sum_{\chi \in \hat{G}: \operatorname{ord}(\chi) = s} \overline{\chi}(g) = \sum_{d \mid s} \mu(s/d) \delta_{g \in dG} \# G[d]$$
$$= \sum_{d \mid (s, e(g))} \mu(s/d) \# G[d].$$

Hence we find

(1)
$$\sum_{i=0}^{n} N(i,g)(-Y)^{i} = \frac{1}{n} \sum_{s \mid \exp(G)} \sum_{d \mid (s,e(g))} \mu(s/d) \# G[d] (1-Y^{s})^{n/s}$$

We single out N(i,g) and obtain:

$$(-1)^{i}N(i,g) = \frac{1}{n} \sum_{s \mid (\exp(G),i)} \sum_{d \mid (s,e(g))} \mu(s/d) \# G[d](-1)^{i/s} \binom{n/s}{i/s}$$

Hence we find

$$N(i,g) = \frac{1}{n} \sum_{s \mid (\exp(G),i)} (-1)^{i+\frac{i}{s}} \binom{n/s}{i/s} \sum_{d \mid (s,e(g))} \mu(s/d) \# G[d].$$

This finishes the proof of Theorem 2.3.

3. Related problem

We will now generalize Theorem 1.2 from [2]. For $g \in G$ we define $N(i,g)^* = \{S \subseteq G \setminus \{0\} : \#S = i, \sum_{s \in S} s = g\}$. We have the following result.

Theorem 3.1. For $g \in G$ we have

$$N(i,g)^* = \frac{1}{n} \sum_{s \mid \exp(G)} (-1)^{i + \lfloor \frac{i}{s} \rfloor} \binom{n/s - 1}{\lfloor i/s \rfloor} \sum_{d \mid (s,e(g))} \mu(s/d) \# G[d].$$

Proof. We have the following identity:

$$\sum_{i=0}^n \sum_{g \in G} N(i,g)^* g X^i = \prod_{\sigma \in G, \sigma \neq 0} (1+\sigma X) \in \mathbf{C}[G][X].$$

As all characters have value one on 0, we deduce the following from Equation 1:

$$\sum_{i=0}^{n} N(i,g)^{*}(-Y)^{i} = \frac{1}{n} \sum_{s \mid \exp(G)} \sum_{d \mid (s,e(g))} \mu(s/d) \# G[d](1-Y^{s})^{n/s-1}(1+Y+\ldots+Y^{s-1})$$

Comparing the coefficients gives the result.

4. ACKNOWLEDGEMENTS

I would like to thank Hendrik Lenstra for helping me to find the proof of the main theorem.

References

- J. Li, D. Wan, Counting subset sums of finite abelian groups Journal of Combinatorial Theory, Series A, Volume 119, Issue 1, January 2012, Pages 170-182
- [2] D. Wan, On the subset sum problem over finite fields, arXiv:0708.2456v1 [math.NT]