

HIGHER-ORDER LAPLACE EQUATIONS AND HYPER-CAUCHY DISTRIBUTIONS

ENZO ORSINGHER AND MIRKO D'OVIDIO

ABSTRACT. In this paper we introduce new distributions which are solutions of higher-order Laplace equations. It is proved that their densities can be obtained by folding and symmetrizing Cauchy distributions. Another class of probability laws related to higher-order Laplace equations is obtained by composing pseudo-processes with positively-skewed stable distributions which produce asymmetric Cauchy densities in the odd-order case. A special attention is devoted to the third-order Laplace equation where the connection between the Cauchy distribution and the Airy functions is obtained and analyzed.

1. INTRODUCTION

The Cauchy density

$$p(x, t) = \frac{1}{\pi} \frac{t}{(x^2 + t^2)} \quad (1.1)$$

solves the Laplace equation (see Nane [9])

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial^2 u}{\partial x^2} = 0, \quad x \in \mathbb{R}, t > 0. \quad (1.2)$$

The n -dimensional counterpart of (1.1)

$$p(\mathbf{x}, t) = \frac{\Gamma(\frac{n}{2})}{\pi^{\frac{n}{2}}} \frac{t}{(t^2 + |\mathbf{x}|^2)^{\frac{n}{2}}}, \quad \mathbf{x} \in \mathbb{R}^{n-1}, t > 0 \quad (1.3)$$

with characteristic function

$$\int_{\mathbb{R}^{n-1}} e^{i(\boldsymbol{\alpha}, \mathbf{x})} p(\mathbf{x}, t) d\mathbf{x} = \exp(-t|\boldsymbol{\alpha}|) \quad (1.4)$$

solves the n -dimensional Laplace equation

$$\frac{\partial^2 p}{\partial t^2} + \sum_{j=1}^{n-1} \frac{\partial^2 p}{\partial x_j^2} = 0. \quad (1.5)$$

The inspiring idea of this paper is to investigate the class of distributions which satisfy the higher-order Laplace equations of the form

$$\frac{\partial^n u}{\partial t^n} + \frac{\partial^n u}{\partial x^n} = 0, \quad x \in \mathbb{R}, t > 0 \quad (1.6)$$

Date: October 25, 2018.

2010 Mathematics Subject Classification. 60G52, 35C05.

Key words and phrases. Pseudo-processes, stable processes, Cauchy processes, higher-order Laplace equations, Airy functions, modified Bessel functions.

In a previous paper of ours we have shown that the law

$$p_4(x, t) = \frac{t}{\pi\sqrt{2}} \frac{x^2 + t^2}{x^4 + t^4} \quad (1.7)$$

solves the fourth-order Laplace equation

$$\frac{\partial^4 u}{\partial t^4} + \frac{\partial^4 u}{\partial x^4} = 0, \quad x \in \mathbb{R}, t > 0. \quad (1.8)$$

In Section 2 we analyze distributions related to equations of the form

$$\frac{\partial^{2^n} u}{\partial t^{2^n}} + \frac{\partial^{2^n} u}{\partial x^{2^n}} = 0 \quad (1.9)$$

which can be expressed in many alternative forms. The decoupling of the 2^n -th order differential operator in (1.9)

$$\frac{\partial^{2^n}}{\partial t^{2^n}} + \frac{\partial^{2^n}}{\partial x^{2^n}} = \prod_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \left(\frac{\partial^2}{\partial t^2} + e^{i\frac{\pi k}{2^{n-1}}} \frac{\partial^2}{\partial x^2} \right)$$

suggests to represent distributions related to (1.9) as

$$p_{2^n}(x, t) = \frac{1}{\pi 2^{n-1}} \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \frac{t e^{i\frac{\pi k}{2^n}}}{x^2 + (t e^{i\frac{\pi k}{2^n}})^2}, \quad n \geq 2. \quad (1.10)$$

that is the superposition of Cauchy densities at imaginary times. Alternatively, we give a real-valued expression for (1.10) as

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{2^{n-2}\pi(x^{2^n} + t^{2^n})} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \cos \frac{k\pi}{2^n} \prod_{\substack{j=1, j \text{ odd} \\ j \neq k}}^{2^{n-1}-1} \left(x^4 + t^4 + 2x^2 t^2 \cos \frac{j\pi}{2^{n-1}} \right). \quad (1.11)$$

The density (1.11) can also be represented as

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{2^{n-2}\pi} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \frac{\cos \frac{k\pi}{2^n}}{x^4 + t^4 + 2x^2 t^2 \cos \frac{k\pi}{2^{n-1}}}, \quad n \geq 2. \quad (1.12)$$

Each component of the distribution (1.12) is produced by folding and symmetrizing the density of the r.v.

$$V(t) = C \left(t \cos \frac{k\pi}{2^n} \right) - t \sin \frac{k\pi}{2^n}, \quad t > 0, 1 \leq k \leq 2^{n-1} - 1, k \text{ odd}$$

where $C(t)$, $t > 0$ is the Cauchy symmetric process. The distributions (1.12) differ from the Cauchy laws since they have a bimodal structure for all $n \geq 2$ as figures below show. For $n = 2$, the distribution (1.11) reduces to (1.7) if we assume that the inner product appearing in formula (1.11) is equal to one. Of course, the density (1.12) coincides with (1.7) for $n = 2$. For $n = 3$ we get from (1.11) and (1.12) that

$$\begin{aligned} p_{2^3}(x, t) &= \frac{t(x^2 + t^2)}{2\pi(x^8 + t^8)} \left[(x^4 + t^4 - \sqrt{2}x^2 t^2) \cos \frac{\pi}{8} + (x^4 + t^4 + \sqrt{2}x^2 t^2) \sin \frac{\pi}{8} \right] \\ &= \frac{t(x^2 + t^2)}{2\pi} \left[\frac{\sin \frac{\pi}{8}}{x^4 + t^4 - \sqrt{2}x^2 t^2} + \frac{\cos \frac{\pi}{8}}{x^4 + t^4 + \sqrt{2}x^2 t^2} \right]. \end{aligned} \quad (1.13)$$

In Orsingher and D'Ovidio [12] we have shown that the density (1.7) is the probability distribution of

$$Q(t) = F(T_t), \quad t > 0$$

where F is the Fresnel pseudo-process and T_t , $t > 0$ is the first passage time of a Brownian motion independent from F . The pseudo-process F is constructed in [12] by means of the fundamental solution (representing the density of the pseudo-process F)

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \cos\left(\frac{x^2}{2t} - \frac{\pi}{4}\right), \quad x \in \mathbb{R}, t > 0$$

of the equation of vibrations of rods

$$\frac{\partial^2 u}{\partial t^2} = -\frac{1}{2^2} \frac{\partial^4 u}{\partial x^4}. \quad (1.14)$$

The Fourier transform $U(\beta, t)$ of $u(x, t)$ is

$$U(\beta, t) = \cos \frac{\beta^2 t}{2}.$$

It is a new non-Markovian pseudo-process which can be analysed by means of the decoupling of (1.14) into two Schrödinger equations. We note that

$$\mathcal{Q}(t) = F(|B(t)|), \quad t > 0$$

has density coinciding with the fundamental solution of the fourth-order heat equation

$$\frac{\partial u}{\partial t} = -\frac{\partial^4 u}{\partial x^4}.$$

We prove also that, for $k \in \mathbb{N}$, there are non-centered Cauchy distributions which solve the equations

$$\frac{\partial^{2k+1} u}{\partial t^{2k+1}} + \frac{\partial^{2k+1} u}{\partial x^{2k+1}} = 0. \quad (1.15)$$

If $X_{2k+1}(t)$, $t > 0$ is the pseudo-process whose density measure

$$\mu_{2k+1}(dx, t) = \mu\{X_{2k+1}(t) \in dx\}$$

solves the heat-type equations

$$\frac{\partial u}{\partial t} = -\frac{\partial^{2k+1} u}{\partial x^{2k+1}}, \quad k \in \mathbb{N} \quad (1.16)$$

and $S_{\frac{1}{2k+1}}(t)$, $t > 0$ is a positively skewed stable process of order $\frac{1}{2k+1}$ we have that

$$Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t)) \in dx\}/dx = \frac{t \cos \frac{\pi}{2(2k+1)}}{\pi \left[\left(x + (-1)^{k+1} t \sin \frac{\pi}{2(2k+1)} \right)^2 + t^2 \cos^2 \frac{\pi}{2(2k+1)} \right]}. \quad (1.17)$$

Pseudo-processes are constructed by attributing to cylinders

$$C = \{x : a_j \leq x(t_j) \leq b_j, j = 1, 2, \dots, n\}$$

of sample paths $x : t \mapsto x(t)$ the following signed measure

$$\mu(C) = \int_{a_1}^{b_1} \cdots \int_{a_n}^{b_n} \prod_{j=1}^n p_n(x_j - x_{j-1}; t_j - t_{j-1}) dx_j$$

and therefore has the following explicit form

$$\mu_{2k+1}(x, t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\xi x - t(-i\xi)^{2n+1}} d\xi.$$

The extension of the measure $\mu(C)$ to the field generated by the cylinders C is described in [4] for the fourth-order pseudo-process and can be adopted also in the odd-order case treated here (see also [5; 6; 7; 10]). A review of higher-order equations appearing in different areas of applied sciences can be found in [1].

We show below that the densities (1.17) solve also the following second-order p.d.e.

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial^2 u}{\partial x^2} = 2 \sin \frac{\pi}{2(2k+1)} \frac{\partial^2 u}{\partial t \partial x}.$$

We have investigated in detail the case of third-order Laplace equation

$$\frac{\partial^3 u}{\partial t^3} + \frac{\partial^3 u}{\partial x^3} = 0 \quad (1.18)$$

and have shown that

$$\begin{aligned} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} &= dx \int_0^\infty \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{x}{\sqrt[3]{3s}}\right) \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) ds \\ &= dx \frac{\sqrt{3}}{2} t \frac{x-t}{x^3-t^3} = dx \frac{\sqrt{3}}{2} \frac{t}{x^2+xt+t^2} \\ &= dx \frac{t \cos \frac{\pi}{6}}{(x+t \sin \frac{\pi}{6})^2 + t^2 \cos^2 \frac{\pi}{6}}. \end{aligned} \quad (1.19)$$

The pictures of the Cauchy distributions (1.17) show that the location parameter $t \sin \frac{\pi}{2(2k+1)}$ tends to zero as $k \rightarrow \infty$ while the scale parameter tends to one, $t \cos \frac{\pi}{2(2k+1)} \rightarrow t$. This means that the asymmetry of the Cauchy densities decreases as k increases and is maximal for $k = 1$. The decrease of parameters of (1.17) (with k increasing) is due to the growing symmetrization of the fundamental solutions of equations (1.16).

By suitably combining the distribution (1.17) for $k = 1$, we arrive at the density

$$p_6(x, t) = \frac{\sqrt{3}}{2\pi} t \frac{(x^2 + t^2) \cos \frac{\pi}{6} + xt}{(x^2 + t^2 + xt \cos \frac{\pi}{6})^2 - 3x^2 t^2 \sin^2 \frac{\pi}{6}} \quad (1.20)$$

which solves the equation

$$\frac{\partial^6 u}{\partial t^6} + \frac{\partial^6 u}{\partial x^6} = 0. \quad (1.21)$$

The probability density (1.20) displays the unimodal structure of the Cauchy distribution.

2. HYPER-CAUCHY DISTRIBUTIONS

In this section we analyze the distribution related to Laplace-type equations of the form

$$\left(\frac{\partial^{2n}}{\partial t^{2n}} + \frac{\partial^{2n}}{\partial x^{2n}} \right) u = 0, \quad n > 1. \quad (2.1)$$

For $n \geq 2$ we obtain a new class of distributions having the form

$$p_{2n}(x, t) = \frac{t(x^2 + t^2)}{2^{n-2} \pi (x^{2n} + t^{2n})} g(x, t), \quad x \in \mathbb{R}, t > 0 \quad (2.2)$$

where $g(x, t)$ is a polynomial of order $2^n - 2^2$. For $n = 2$, formula (2.2) yields the distribution

$$p_4(x, t) = \frac{t(x^2 + t^2)}{\sqrt{2\pi}(x^4 + t^4)}, \quad x \in \mathbb{R}, t > 0 \quad (2.3)$$

emerging in the analysis of Fresnel pseudo-processes (see Orsingher and D'Ovidio [12]).

The main result of this section is given in the next theorem.

Theorem 2.1. *The hyper-Cauchy density*

$$p_{2^n}(x, t) = \frac{1}{\pi 2^{2^n-1}} \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \frac{te^{i\frac{\pi k}{2^n}}}{x^2 + (te^{i\frac{\pi k}{2^n}})^2} \quad (2.4)$$

solves the equation

$$\left(\frac{\partial^{2^n}}{\partial t^{2^n}} + \frac{\partial^{2^n}}{\partial x^{2^n}} \right) u = 0, \quad x \in \mathbb{R}, t > 0, \quad n > 1. \quad (2.5)$$

A real-valued expression of (2.4) reads

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{\pi 2^{2^n-2}(x^{2^n} + t^{2^n})} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \cos \frac{k\pi}{2^n} \prod_{\substack{k \neq j=1 \\ j \text{ odd}}}^{2^{n-1}-1} \left(x^4 + t^4 + 2x^2t^2 \cos \frac{j\pi}{2^{n-1}} \right) \quad (2.6)$$

or equivalently

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{2^{n-2}\pi} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \frac{\cos \frac{k\pi}{2^n}}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}}, \quad \text{for } n > 1. \quad (2.7)$$

Proof. In order to check that (2.4) satisfies equation (2.5) we resort to Fourier transforms

$$U(\beta, t) = \int_{-\infty}^{+\infty} e^{i\beta x} u(x, t) dx.$$

Equation (2.5) becomes

$$\frac{\partial^{2^n} U}{\partial t^{2^n}} + (-i\beta)^{2^n} U = \frac{\partial^{2^n} U}{\partial t^{2^n}} + \beta^{2^n} U = 0. \quad (2.8)$$

The solutions of the algebraic equation associated to (2.8) have the form

$$r_j = |\beta| e^{i\pi \frac{2j+1}{2^n}}, \quad 0 \leq j \leq 2^n - 1. \quad (2.9)$$

In order to construct bounded solutions to (2.8) we restrict ourselves to

$$U(\beta, t) = \frac{1}{2^{n-1}} \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} e^{-t|\beta|e^{i\frac{k\pi}{2^n}}} \quad (2.10)$$

where the normalizing constant in (2.10) is chosen equal to $1/2^{n-1}$ so that $U(\beta, 0) = 1$. The inverse of (2.10) is (2.4). We check directly that each term of (2.4) has

Fourier transform solving equation (2.8). For all odd values of k , we have that

$$\begin{aligned}
& \int_{-\infty}^{+\infty} e^{i\beta x} \left(\frac{\partial^{2^n}}{\partial t^{2^n}} + \frac{\partial^{2^n}}{\partial x^{2^n}} \right) \frac{1}{\pi} \left(\frac{te^{i\frac{k\pi}{2^n}}}{x^2 + (te^{i\frac{k\pi}{2^n}})^2} \right) dx \\
&= \frac{\partial^{2^n}}{\partial t^{2^n}} e^{-t|\beta|e^{i\frac{k\pi}{2^n}}} + (-i\beta)^{2^n} e^{-t|\beta|e^{i\frac{k\pi}{2^n}}} \\
&= \left(\beta^{2^n} e^{ik\pi} + i^{2^n} \beta^{2^n} \right) e^{-t|\beta|e^{i\frac{k\pi}{2^n}}} \\
&= \left((-1)^k \beta^{2^n} + \beta^{2^n} \right) e^{-t|\beta|e^{i\frac{k\pi}{2^n}}} = 0
\end{aligned}$$

because k is odd. In order to obtain (2.6) we observe that, in view of (2.4) we can write

$$p_{2^n}(x, t) = \frac{1}{\pi} \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \frac{c_k t^{|2k-1|} x^{2^n - |2k-1| - 1}}{\prod_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} (x^2 + (te^{i\frac{k\pi}{2^n}})^2)}$$

where

$$\prod_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} (x^2 + (te^{i\frac{k\pi}{2^n}})^2) = x^{2^n} + t^{2^n} \quad (2.11)$$

and c_k are constants evaluated below. Result (2.11) can be obtained directly by solving the equation $x^{2^n} + t^{2^n} = 0$ or by successively regrouping the terms of the right-hand side of (2.11). We have at first that

$$\begin{aligned}
\prod_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} (x^2 + (te^{i\frac{k\pi}{2^n}})^2) &= \prod_{k=1, k \text{ odd}}^{2^{n-1}-1} \left(x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}} \right) \\
&= \prod_{k=1, k \text{ odd}}^{2^{n-2}-1} \left(x^8 + t^8 + 2x^4t^4 \cos \frac{k\pi}{2^{n-2}} \right) \\
&= \dots \\
&= \left(x^{2^n} + t^{2^n} + 2x^2t^2 \cos \frac{\pi}{2} \right) \\
&= x^{2^n} + t^{2^n}.
\end{aligned}$$

In view of (2.11) we can rewrite (2.4) as

$$p_{2^n}(x, t) = \frac{t}{\pi 2^{n-1} (x^{2^n} + t^{2^n})} \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \prod_{\substack{j=-(2^{n-1}-1) \\ j \text{ odd}, j \neq k}}^{2^{n-1}-1} (x^2 + (te^{i\frac{\pi j}{2^n}})^2) e^{i\frac{\pi k}{2^n}}$$

where

$$\sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \prod_{\substack{j=-(2^{n-1}-1) \\ j \text{ odd}, j \neq k}}^{2^{n-1}-1} (x^2 + (te^{i\frac{\pi j}{2^n}})^2) e^{i\frac{\pi k}{2^n}}$$

$$\begin{aligned}
 &= \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \prod_{\substack{j=1 \\ j \text{ odd}, j \neq k}}^{2^{n-1}-1} (x^4 + t^4 + 2x^2t^2 \cos \frac{\pi j}{2^{n-1}}) (x^2 + (te^{-i\frac{\pi k}{2^n}})^2) e^{i\frac{\pi k}{2^n}} \\
 &= \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \prod_{\substack{j=1 \\ j \text{ odd}, j \neq k}}^{2^{n-1}-1} (x^4 + t^4 + 2x^2t^2 \cos \frac{\pi j}{2^{n-1}}) (x^2 e^{i\frac{k\pi}{2^n}} + t^2 e^{-i\frac{k\pi}{2^n}} + x^2 e^{-i\frac{k\pi}{2^n}} + t^2 e^{i\frac{k\pi}{2^n}}) \\
 &= 2(x^2 + t^2) \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \cos \frac{k\pi}{2^n} \prod_{\substack{j=1 \\ j \text{ odd}, j \neq k}}^{2^{n-1}-1} (x^4 + t^4 + 2x^2t^2 \cos \frac{\pi j}{2^{n-1}})
 \end{aligned}$$

and thus

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{\pi 2^{n-2}(x^{2^n} + t^{2^n})} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \cos \frac{k\pi}{2^n} \prod_{\substack{j=1 \\ j \text{ odd}, j \neq k}}^{2^{n-1}-1} (x^4 + t^4 + 2x^2t^2 \cos \frac{\pi j}{2^{n-1}}).$$

Furthermore, from the fact that

$$x^{2^n} + t^{2^n} = \prod_{k=1, k \text{ odd}}^{2^{n-2}-1} \left(x^8 + t^8 + 2x^4t^4 \cos \frac{k\pi}{2^{n-2}} \right)$$

we obtain that

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{2^{n-2}\pi} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \frac{\cos \frac{k\pi}{2^n}}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}}.$$

□

Remark 2.2. In order to prove that the density (2.6) integrates to unity we present the following calculation

$$\begin{aligned}
 \int_{-\infty}^{+\infty} \frac{x^2 + t^2}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}} dx &= 2 \int_0^{+\infty} \frac{x^2 + t^2}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}} dx \\
 &= \frac{2}{t} \int_0^{+\infty} \frac{y^2 + 1}{y^4 + 1 + 2y \cos \frac{k\pi}{2^{n-1}}} dy \\
 &= \frac{2}{t} \int_0^{\frac{\pi}{2}} \frac{1}{\tan^4 \theta + 1 + 2 \tan^2 \theta \cos \frac{k\pi}{2^{n-1}}} \frac{d\theta}{\cos^4 \theta} \\
 &= \frac{2}{t} \int_0^{\frac{\pi}{2}} \frac{d\theta}{\sin^4 \theta + \cos^4 \theta + 2 \sin^2 \theta \cos^2 \theta \cos \frac{k\pi}{2^{n-1}}} \\
 &= \frac{2}{t} \int_0^{\frac{\pi}{2}} \frac{d\theta}{1 - \frac{1 - \cos \frac{k\pi}{2^{n-1}}}{2} \sin^2 2\theta} \\
 &= \frac{2}{t} \int_0^{\frac{\pi}{2}} \frac{d\theta}{1 - \frac{1}{2} (1 - \cos \frac{k\pi}{2^{n-1}}) \left(\frac{1 - \cos 4\theta}{2} \right)} \\
 &= \frac{1}{2t} \int_0^{2\pi} \frac{d\phi}{1 - \frac{1 - \cos \frac{k\pi}{2^{n-1}}}{4} + \frac{1}{4} (1 - \cos \frac{k\pi}{2^{n-1}}) \cos \phi}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{2}{t} \int_0^{2\pi} \frac{d\phi}{\left(3 + \cos \frac{k\pi}{2^{n-1}}\right) + \left(1 - \cos \frac{k\pi}{2^{n-1}}\right) \cos \phi} \\
&= \frac{2}{t} \frac{2\pi}{\sqrt{\left(3 + \cos \frac{k\pi}{2^{n-1}}\right)^2 - \left(1 - \cos \frac{k\pi}{2^{n-1}}\right)^2}} \\
&= \frac{\pi\sqrt{2}}{t} \frac{1}{\sqrt{1 + \cos \frac{k\pi}{2^{n-1}}}} \\
&= \frac{\pi}{t} \frac{1}{\cos \frac{k\pi}{2^n}}. \tag{2.12}
\end{aligned}$$

From (2.7), in view of (2.12), we can conclude that

$$\int_{-\infty}^{+\infty} p_{2^n}(x, t) dx = 1$$

Remark 2.3. From (2.4), for $n = 2$ we obtain that

$$p_4(x, t) = \frac{1}{2\pi} \left[\frac{te^{i\frac{\pi}{4}}}{x^2 + (te^{i\frac{\pi}{4}})^2} + \frac{te^{-i\frac{\pi}{4}}}{x^2 + (te^{-i\frac{\pi}{4}})^2} \right]$$

with Fourier transform

$$\int_{-\infty}^{+\infty} e^{i\beta x} p_4(x, t) dx = e^{-\frac{t}{\sqrt{2}}|\beta|} \cos \frac{\beta t}{\sqrt{2}}.$$

From (2.6) and (2.7) we have that

$$p_4(x, t) = \frac{t}{\sqrt{2}\pi} \frac{x^2 + t^2}{x^4 + t^4}. \tag{2.13}$$

The law (2.13) has two maxima as Figure 1 shows.

Remark 2.4. For $n = 3$, from (2.6), we have that

$$p_8(x, t) = \frac{t(x^2 + t^2)}{2\pi(x^8 + t^8)} \left[\left(x^4 + t^4 + 2x^2t^2 \cos \frac{\pi}{4}\right) \cos \frac{3\pi}{8} + \left(x^4 + t^4 + 2x^2t^2 \cos \frac{3\pi}{4}\right) \cos \frac{\pi}{8} \right].$$

From the fact that

$$\cos \frac{3\pi}{4} = -\cos \frac{\pi}{4} \quad \text{and} \quad \cos \frac{3\pi}{8} = \sin \frac{\pi}{8}$$

we write

$$p_8(x, t) = \frac{t(x^2 + t^2)}{2\pi(x^8 + t^8)} \left[\left(x^4 + t^4 + \sqrt{2}x^2t^2\right) \sin \frac{\pi}{8} + \left(x^4 + t^4 - \sqrt{2}x^2t^2\right) \cos \frac{\pi}{8} \right].$$

From (2.7) we have also that


$$p_8(x, t) = \frac{t}{2\pi} \left[\frac{x^2 + t^2}{x^4 + t^4 - \sqrt{2}x^2t^2} \sin \frac{\pi}{8} + \frac{x^2 + t^2}{x^4 + t^4 + \sqrt{2}x^2t^2} \cos \frac{\pi}{8} \right]. \tag{2.14}$$

From (2.4) we obtain the characteristic function

$$\int_{\mathbb{R}} e^{i\beta x} p_8(x, t) dx = \frac{1}{2^2} \left[e^{-t|\beta| \cos \frac{\pi}{8}} \cos \left(t\beta \sin \frac{\pi}{8}\right) + e^{-t|\beta| \sin \frac{\pi}{8}} \cos \left(t\beta \cos \frac{\pi}{8}\right) \right].$$

The density $p_8(x, t)$ is a bimodal curve as well as $p_4(x, t)$. The maxima of $p_8(x, t)$ are higher than those of $p_4(x, t)$ as Figure 1 shows. Also $p_{2^n}(x, t)$ displays a bimodal

FIGURE 1. The profile of the functions p_4 (dotted line), formula (2.13) and p_8 , formula (2.14).



structure with the height peaks increasing as n increases. The form of $p_{2^n}(x, t)$ reminds the structure of densities of fractional diffusions governed by equations

$$\frac{\partial^\nu u}{\partial t^\nu} = \lambda^2 \frac{\partial^2 u}{\partial x^2}$$

for $1 < \nu < 2$ (see [11]).

Remark 2.5. The result (2.7) can conveniently be rewritten as

$$p_{2^n}(x, t) = \frac{t}{\pi(x^2 + t^2)} \left[\frac{1}{2^{n-1}} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \frac{x^4 + t^4 + 2x^2t^2}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}} \cos \frac{k\pi}{2^n} \right]. \quad (2.15)$$

The factor in square parenthesis measures, in some sense, the disturbance of p_{2^n} on the classical Cauchy. For $n = 2$, we have in particular that


$$p_{2^2}(x, t) = \frac{t}{\pi(x^2 + t^2)} \frac{1}{\sqrt{2}} \left[1 + \frac{2x^2t^2}{x^4 + t^4} \right] = \frac{t}{\sqrt{2}\pi} \frac{x^2 + t^2}{x^4 + t^4}. \quad (2.16)$$

The density (2.16) has two symmetric maxima at $x = \pm t\sqrt{\sqrt{2} - 1}$ and a minimum at $x = 0$ (see Fig. 6 of Orsingher and D’Ovidio [12]). The terms

$$g_k(x, t) = \frac{x^4 + t^4 + 2x^2t^2}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}} \quad (2.17)$$

display two maxima at $x = \pm t$ with height depending on k and whose profile is depicted in Figure 2.

FIGURE 2. The profile of the function g_k for $n = 3$ and $k = 1$ (dotted line), $k = 3$.



Remark 2.6. The density $p_{2^n}(x, t)$ can be written as

$$p_{2^n}(x, t) = \frac{t(x^2 + t^2)}{2^{n-2}\pi(x^{2^n} + t^{2^n})}Q(x, t) \quad (2.18)$$

where $Q(x, t)$ is a polynomial of order $2^n - 2^2$. For $n = 2$ the function $Q(x, t)$ reduces to $\cos \frac{\pi}{4}$. For $n = 3$,

$$Q(x, t) = (x^4 + t^4 + \sqrt{2}x^2t^2) \sin \frac{\pi}{8} + (x^4 + t^4 - \sqrt{2}x^2t^2) \cos \frac{\pi}{8}.$$

The expression (2.18) shows that the probability law $p_{2^n}(x, t)$, $x \in \mathbb{R}$, $t > 0$ shares with the classical Cauchy density the property of non-existence of the mean value.


Remark 2.7. The density of the hyper-Cauchy can also be presented in an alternative form by regrouping the terms in the right-hand side of (2.7) as

$$\begin{aligned} & \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-2}-1} \left[\frac{\sin \frac{k\pi}{2^n}}{x^4 + t^4 - 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}} + \frac{\cos \frac{k\pi}{2^n}}{x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}} \right] \\ &= \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-2}-1} \frac{(x^4 + t^4 + 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}) \sin \frac{k\pi}{2^n} + (x^4 + t^4 - 2x^2t^2 \cos \frac{k\pi}{2^{n-1}}) \cos \frac{k\pi}{2^n}}{x^8 + t^8 - 2x^4t^4 \cos \frac{k\pi}{2^{n-2}}}. \end{aligned} \quad (2.19)$$

For $n = 3$, from (2.19), we get again that

$$p_8(x, t) = \frac{t(x^2 + t^2)}{2\pi(x^8 + t^8)} \left[(x^4 + t^4 + \sqrt{2}x^2t^2) \sin \frac{k\pi}{8} + (x^4 + t^4 - \sqrt{2}x^2t^2) \cos \frac{k\pi}{8} \right].$$

FIGURE 3. The profile of the functions p_{2^n} , formula (2.7), for $n = 5, 10, 15, 20$.



Remark 2.8. The r.v.

$$W(t) = \left| C \left(t \cos \frac{\pi k}{2^n} \right) - t \sin \frac{\pi k}{2^n} \right| \quad (2.20)$$

(where $C(t)$, $t > 0$ is the Cauchy process) has probability density

$$f_k(w, t) = \frac{2t(w^2 + t^2) \cos \frac{k\pi}{2^n}}{\pi(w^4 + t^4 + 2w^2t^2 \cos \frac{k\pi}{2^{n-1}})}, \quad w > 0. \quad (2.21)$$

Indeed, we have that

$$Pr \{W(t) < w\} = \int_{-w+t \sin \frac{k\pi}{2^n}}^{+w+t \sin \frac{k\pi}{2^n}} dy \frac{t \cos \frac{k\pi}{2^n}}{\pi(y^2 + t^2 \cos^2 \frac{k\pi}{2^n})} \quad (2.22)$$

and

$$\begin{aligned} f_k(w, t) &= \frac{d}{dw} Pr \left\{ \left| C \left(t \cos \frac{\pi k}{2^n} \right) - t \sin \frac{\pi k}{2^n} \right| < w \right\} \\ &= \frac{t \cos \frac{k\pi}{2^n}}{\pi \left((w + t \sin \frac{k\pi}{2^n})^2 + t^2 \cos^2 \frac{k\pi}{2^n} \right)} + \frac{t \cos \frac{k\pi}{2^n}}{\pi \left((-w + t \sin \frac{k\pi}{2^n})^2 + t^2 \cos^2 \frac{k\pi}{2^n} \right)} \end{aligned}$$

$$\begin{aligned}
&= \frac{t \cos \frac{k\pi}{2^n}}{\pi \left(w^2 + 2wt \sin \frac{k\pi}{2^n} + t^2 \right)} + \frac{t \cos \frac{k\pi}{2^n}}{\pi \left(w^2 - 2wt \sin \frac{k\pi}{2^n} + t^2 \right)} \\
&= \frac{2t(w^2 + t^2) \cos \frac{k\pi}{2^n}}{\pi(w^2 + t^2 + 2wt \sin \frac{k\pi}{2^n})(w^2 + t^2 - 2wt \sin \frac{k\pi}{2^n})} \\
&= \frac{2t(w^2 + t^2) \cos \frac{k\pi}{2^n}}{\pi(w^4 + t^4 + 2w^2t^2 \cos \frac{k\pi}{2^{n-1}})}
\end{aligned}$$

because

$$2 \sin^2 \frac{k\pi}{2^n} = 1 - \cos \frac{k\pi}{2^{n-1}}.$$

By symmetrizing (2.20) as follows

$$Z(t) = \frac{W_1(t) - W_2(t)}{2}$$

where $W_1(t), W_2(t)$ are independent copies of $W(t)$ we obtain a distribution of the form

$$h_k(w, t) = \frac{t(w^2 + t^2) \cos \frac{k\pi}{2^n}}{\pi(w^4 + t^4 + 2w^2t^2 \cos \frac{k\pi}{2^{n-1}})}, \quad w \in \mathbb{R} \quad (2.23)$$

which coincides with each term of (2.15). This construction explains the reason for which each term in (2.15) has two symmetric maxima at $w = \pm t \sqrt{2 \sin \frac{k\pi}{2^n} - 1}$ for $k : \sin \frac{\pi k}{2^n} > \frac{1}{2}$.

3. HIGHER-ORDER LAPLACE-TYPE EQUATION

Let us consider the pseudo-processes related to higher-order heat-type equations

$$\frac{\partial u}{\partial t} = c_n \frac{\partial^n u}{\partial x^n}, \quad x \in \mathbb{R}, \quad t > 0, \quad n > 2. \quad (3.1)$$

where $c_n = (-1)^{\frac{n}{2}+1}$ for n even and $c_n = \pm 1$ for n odd.

Pseudo-processes constructed by exploiting the sign-varying measures obtained as fundamental solutions to (3.1) have been examined in many papers since the beginning of the Sixties. A description of the procedure of construction of pseudo-processes can be found, for example in Krylov [5], Ladokhin [7], Hochberg [4], Orsingher [10], Lachal [6]. In the case where $n = 2k + 1$, $c_{2k+1} = -1$, the fundamental solution to (3.1) reads

$$u_{2k+1}(x, t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\beta x + i(-1)^k t \beta^{2k+1}} d\beta. \quad (3.2)$$

In particular, for $k = 1$


$$u_3(x, t) = \frac{1}{\pi} \int_0^\infty \cos(\beta x + \beta^3 t) d\beta = \frac{1}{\sqrt[3]{3t}} Ai\left(\frac{x}{\sqrt[3]{3t}}\right) \quad (3.3)$$

where

$$Ai(x) = \frac{\sqrt{x}}{3} \left[I_{-\frac{1}{3}}\left(\frac{2}{3}x^{3/2}\right) - I_{\frac{1}{3}}\left(\frac{2}{3}x^{3/2}\right) \right]$$

is the Airy function (see for example Lebedev [8]).

FIGURE 4. The figure shows how the distribution (2.23) can be constructed from the Cauchy density by folding and symmetrizing, in the cases $n = 3, k = 1$ (top figures) and $k = 3$ (bottom figures). The dotted line gives the density of the folded distribution (3.13).



In this section we study the composition of pseudo-processes with stable processes $S_\alpha(t)$, $t > 0$, $\alpha \in (0, 1)$ whose characteristic function reads

$$\mathbb{E}e^{i\beta S_\alpha(t)} = \exp\left(-t|\beta|^\alpha e^{-i\frac{\pi\gamma}{2}\frac{\beta}{|\beta|}}\right) = \exp\left(-\sigma t|\beta|^\alpha \left(1 - i\theta\frac{\beta}{|\beta|} \tan\frac{\pi\alpha}{2}\right)\right) \quad (3.4)$$

where $\sigma = \cos \pi\gamma/2 > 0$ and

$$\theta = \cot\left(\frac{\pi\alpha}{2}\right) \tan\left(\frac{\pi\gamma}{2}\right).$$

The parameter γ must be chosen in such a way that $\theta \in [-1, 1]$ for $\alpha \in (0, 1)$. The skewness parameter $\theta = 1$ (that is $\gamma = \alpha$) corresponds to positively skewed stable distributions. For the density

$$p_\alpha(x, \gamma, t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\beta x} \mathbb{E}e^{i\beta S_\alpha(t)} d\beta$$

we have the scaling property

$$p_\alpha(x, \gamma, t) = \frac{1}{t^{1/\alpha}} p_\alpha\left(\frac{x}{t^{1/\alpha}}, \gamma, 1\right). \quad (3.5)$$

For $\alpha \in (0, 1)$, we have the series representation of stable density (see [11, page 245])

$$p_\alpha(x; \gamma, 1) = \frac{\alpha}{\pi} \sum_{r=0}^{\infty} (-1)^r \frac{\Gamma(\alpha(r+1))}{r!} x^{-\alpha(r+1)-1} \sin\left(\frac{\pi}{2}(\gamma + \alpha)(r+1)\right). \quad (3.6)$$

Theorem 3.1. *The composition of the pseudo-process $X_{2k+1}(t)$, $t > 0$ with the stable process $S_{\frac{1}{2k+1}}(t)$, $t > 0$, $k \in \mathbb{N}$, has a Cauchy probability distribution which can be written as*

$$Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t) \in dx)/dx = \frac{t \cos \frac{\pi}{2(2k+1)}}{\pi \left[\left(x + (-1)^{k+1} t \sin \frac{\pi}{2(2k+1)} \right)^2 + t^2 \cos^2 \frac{\pi}{2(2k+1)} \right]} \quad (3.7)$$

with $x \in \mathbb{R}$, $t > 0$. The density function (3.7) is a solution to the higher-order Laplace equation

$$\frac{\partial^{2k+1} u}{\partial t^{2k+1}} + \frac{\partial^{2k+1} u}{\partial x^{2k+1}} = 0, \quad x \in \mathbb{R}, t > 0 \quad (3.8)$$

Proof. For $\theta = 1$, $\alpha = \gamma = 1/2k + 1$, in view of (3.4) we have that

$$\begin{aligned} U(\beta, t) &= \int_{-\infty}^{+\infty} e^{i\beta x} Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t)) \in dx\} \\ &= \int_0^{\infty} Pr\{S_{\frac{1}{2k+1}}(t) \in ds\} \int_{-\infty}^{+\infty} e^{i\beta x} u_{2k+1}(x, s) dx \\ &= \int_0^{\infty} e^{is(-1)^k \beta^{2k+1}} Pr\{S_{\frac{1}{2k+1}}(t) \in ds\} \\ &= \exp\left(-t \left| (-1)^k \beta^{2k+1} \right|^{\frac{1}{2k+1}} \cos \frac{\pi}{2(2k+1)} \left(1 - i \operatorname{sgn} \left((-1)^k \beta^{2k+1} \right) \tan \frac{\pi}{2(2k+1)} \right)\right) \\ &= \exp\left(-t|\beta| \left(\cos \frac{\pi}{2(2k+1)} - i(-1)^k \frac{\beta}{|\beta|} \sin \frac{\pi}{2(2k+1)} \right)\right) \\ &= \exp\left(-t|\beta| \cos \frac{\pi}{2(2k+1)} - i(-1)^k t \beta \sin \frac{\pi}{2(2k+1)}\right). \end{aligned} \quad (3.9)$$

This is the characteristic function of a Cauchy distribution with scale parameter $t \cos \frac{\pi}{2(2k+1)}$ and location parameter $t(-1)^{k+1} \sin \frac{\pi}{2(2k+1)}$. Formula (3.9) can also be rewritten as

$$\begin{aligned} U(\beta, t) &= \exp\left(-t|\beta| \left(\cos \frac{\pi}{2(2k+1)} - i(-1)^k \frac{\beta}{|\beta|} \sin \frac{\pi}{2(2k+1)} \right)\right) \\ &= \exp\left(-t|\beta| \left(\cos \left(\frac{\pi}{2(2k+1)} (-1)^k \frac{\beta}{|\beta|} \right) - i \sin \left(\frac{\pi}{2(2k+1)} (-1)^k \frac{\beta}{|\beta|} \right) \right)\right) \\ &= \exp\left(-t|\beta| e^{-i \frac{\pi}{2(2k+1)} (-1)^k \frac{\beta}{|\beta|}}\right). \end{aligned} \quad (3.10)$$

The Fourier transform of equation (3.8) becomes

$$\frac{\partial^{2k+1} U}{\partial t^{2k+1}} + (-i\beta)^{2k+1} U = 0. \quad (3.11)$$

The derivative of order $2k + 1$ of (3.10) is

$$\frac{\partial^{2k+1} U}{\partial t^{2k+1}}(\beta, t) = (-|\beta|)^{2k+1} \left(e^{-i \frac{\pi}{2(2k+1)} (-1)^k \frac{\beta}{|\beta|}} \right)^{2k+1} U(\beta, t) \quad (3.12)$$

and this shows that the Cauchy distribution (3.7) solves the higher-order Laplace equation (3.8). \square

Remark 3.2. We notice that

$$\begin{aligned} \int_0^\infty Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t)) \in dx\} &= \frac{1}{\pi} \int_{(-1)^{k+1} \tan \frac{\pi}{2(2k+1)}}^\infty \frac{dy}{1+y^2} \\ &= \frac{1}{2} \left(1 + \frac{(-1)^k}{2k+1} \right) \end{aligned} \quad (3.13)$$

which is somehow in accord with Lachal [6]. The results (3.7) and (3.13) show that the mode of the Cauchy law (3.7) approaches the origin as k increases.

Let us consider the process of the form $X_3(S_{\frac{1}{3}}(t))$, $t > 0$ where X_3 is a pseudo-process whose measure density is governed by the third-order heat equation

$$\frac{\partial u}{\partial t} = -\frac{\partial^3 u}{\partial x^3}, \quad x \in \mathbb{R}, t > 0 \quad (3.14)$$

and $S_{\frac{1}{3}}$ is the stable process of order $1/3$. The distribution of $X_3(S_{\frac{1}{3}}(t))$, $t > 0$ reads

$$Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} = dx \int_0^\infty \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{x}{\sqrt[3]{3s}}\right) \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) ds \quad (3.15)$$

where

$$Pr\{S_{\frac{1}{3}}(t) \in ds\} = ds \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right), \quad s \geq 0, t > 0 \quad (3.16)$$

for which

$$\begin{aligned} \int_0^\infty Pr\{S_{\frac{1}{3}}(t) \in ds\} &= \int_0^\infty ds \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) \\ &= (w = t/\sqrt[3]{3s}) = 3 \int_0^\infty Ai(w) dw = 1. \end{aligned}$$

Corollary 3.3. *The law (3.15) solves the higher-order Laplace equation*

$$\frac{\partial^3 u}{\partial t^3} + \frac{\partial^3 u}{\partial x^3} = 0, \quad x \in \mathbb{R}, t > 0 \quad (3.17)$$

and can be written as

$$\begin{aligned} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} &= \frac{dx}{\pi} \frac{\frac{\sqrt{3}}{2}t}{\left(x + \frac{t}{2}\right)^2 + \frac{3t^2}{4}} \\ &= \frac{dx}{\pi} \frac{3^{1/2}}{2} \frac{t}{x^2 + xt + t^2} \\ &= dx \frac{3^{1/2}t}{2\pi} \frac{x-t}{x^3 - t^3}. \end{aligned} \quad (3.18)$$

Proof. The Fourier transform of (3.15) becomes

$$\int_{-\infty}^\infty e^{i\beta x} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} = \int_0^\infty e^{-i\beta^3 s} Pr\{S_{\frac{1}{3}}(t) \in ds\}. \quad (3.19)$$

We show that (3.16) is a stable law of order $1/3$. In view of the representation of the the Airy function (4.10) of Orsingher and Beghin [11]

$$Ai(w) = \frac{3^{-2/3}}{\pi} \sum_{k=0}^{\infty} \frac{(3^{1/3}w)^k}{k!} \sin\left(\frac{2\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right) \quad (3.20)$$

we can write that

$$\frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) = \frac{t}{3\pi s \sqrt[3]{s}} \sum_{k=0}^{\infty} \left(\frac{t}{\sqrt[3]{s}}\right)^k \frac{1}{k!} \sin\left(\frac{2\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right)$$

We consider the series expansion (3.6) of the stable density (with $t = 1$) for which (3.4) holds true. For $\alpha = \gamma = 1/3$ (that is $\theta = +1$), $x = s/t^3$ in (3.6) we get that

$$\begin{aligned} p_{\frac{1}{3}}\left(\frac{s}{t^3}; \frac{1}{3}, 1\right) &= \frac{1}{3\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \left(\frac{s}{t^3}\right)^{-\frac{k+1}{3}-1} \sin\left(\frac{\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right) \\ &= (\text{by 4.5 of [11]}) \\ &= \frac{1}{3\pi} \frac{t^4}{s \sqrt[3]{s}} \sum_{k=0}^{\infty} \left(\frac{t}{\sqrt[3]{s}}\right)^k \frac{1}{k!} \sin\left(\frac{2\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right) \\ &= t^3 \left[\frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) \right] \end{aligned}$$

and thus, from (3.5), we have that

$$\frac{1}{t^3} p_{\frac{1}{3}}\left(\frac{s}{t^3}; \frac{1}{3}, 1\right) = p_{\frac{1}{3}}\left(s; \frac{1}{3}, t\right) = \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right), \quad s, t > 0.$$

We now evaluate the integral (3.19). We have that

$$\begin{aligned} &\int_0^{\infty} e^{-i\beta^3 s} Pr\{S_{\frac{1}{3}}(t) \in ds\} \\ &= \exp\left(-\cos\frac{\pi}{6}t - \beta^3 |t|^{\frac{1}{3}} \left(1 - i \operatorname{sgn}(-\beta^3) \tan\frac{\pi}{6}\right)\right) \\ &= \exp\left(-\frac{\sqrt{3}}{2}t|\beta| \left(1 + i \operatorname{sgn}(\beta) \frac{1}{\sqrt{3}}\right)\right) \\ &= \exp\left(-\frac{\sqrt{3}}{2}t|\beta| - i\frac{t}{2}\beta\right) \end{aligned} \quad (3.21)$$

since $\operatorname{sgn}(-\beta^3) = \operatorname{sgn}(-\beta) = -\operatorname{sgn}(\beta) = -\frac{\beta}{|\beta|}$. From (3.21) we infer that

$$\begin{aligned} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} &= \frac{dx}{2\pi} \int_{-\infty}^{+\infty} e^{-i\beta x} \exp\left(-\frac{\sqrt{3}}{2}t|\beta| - i\frac{t}{2}\beta\right) d\beta \\ &= \frac{dx}{\pi} \frac{\frac{\sqrt{3}}{2}t}{\left(x + \frac{t}{2}\right)^2 + \frac{3t^2}{4}} = \frac{dx}{\pi} \frac{3^{1/2}}{2} \frac{t}{x^2 + xt + t^2} \\ &= dx \frac{3^{1/2}t}{2\pi} \frac{x-t}{x^3 - t^3} \end{aligned} \quad (3.22)$$

□

Remark 3.4. We observe that the r.v. $X_3(S_{\frac{1}{3}}(t))$ possesses Cauchy distribution with scale parameter $\sqrt{3}t/2$ and location parameter $-t/2$. Furthermore, it solves the third-order Laplace-type equation

$$\frac{\partial^3 u}{\partial t^3} + \frac{\partial^3 u}{\partial x^3} = 0. \quad (3.23)$$

Remark 3.5. From the fact that

$$\frac{1}{\sqrt[3]{3t}} Ai\left(\frac{x}{\sqrt[3]{3t}}\right) = \frac{1}{3\pi} \sqrt{\frac{x}{t}} K_{1/3}\left(\frac{2}{3^{3/2}} \frac{x^{3/2}}{\sqrt{t}}\right), \quad x, t > 0 \quad (3.24)$$

we can write, for $x > 0$,

$$\begin{aligned} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\}/dx &= \int_0^\infty \frac{1}{3\pi} \sqrt{\frac{x}{s}} K_{1/3}\left(\frac{2}{3^{3/2}} \frac{x^{3/2}}{\sqrt{s}}\right) \frac{t}{s} \frac{1}{3\pi} \sqrt{\frac{t}{s}} K_{1/3}\left(\frac{2}{3^{3/2}} \frac{t^{3/2}}{\sqrt{s}}\right) ds \\ &= \frac{2\sqrt{xt^3}}{3^2\pi^2} \int_0^\infty s K_{1/3}\left(\frac{2x^{3/2}}{3^{3/2}s}\right) K_{1/3}\left(\frac{2t^{3/2}}{3^{3/2}s}\right) ds. \end{aligned} \quad (3.25)$$

In view of (see [3, formula 6.521])


$$\int_0^\infty s K_\nu(ys) K_\nu(zs) ds = \frac{\pi(yz)^{-\nu}(y^{2\nu} - z^{2\nu})}{2 \sin \pi\nu (y^2 - z^2)}, \quad \Re\{y + z\} > 0, |\Re\{\nu\}| < 1$$

we get that

$$Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} = dx \frac{3^{1/2} t}{2\pi} \frac{x - t}{x^3 - t^3}, \quad x, t > 0 \quad (3.27)$$

which coincides with (3.18).

FIGURE 5. The profile of the function (3.18).



The Cauchy densities pertaining to the composition $X_{\frac{1}{2k+1}}(S_{\frac{1}{2k+1}}(t))$, $t > 0$, solve also a second-order p.d.e. as we show in the next theorem.

Theorem 3.6. *The Cauchy densities*

$$f(x, t; m) = \frac{1}{\pi} \frac{t \cos \frac{\pi}{2m}}{(x + t \sin \frac{\pi}{2m})^2 + t^2 \cos^2 \frac{\pi}{2m}}, \quad m \in \mathbb{N} \quad (3.28)$$

satisfy the following second-order equation

$$\frac{\partial^2 f}{\partial t^2} + \frac{\partial^2 f}{\partial x^2} = 2 \sin \frac{\pi}{2m} \frac{\partial^2 f}{\partial x \partial t}, \quad x \in \mathbb{R}, t > 0. \quad (3.29)$$

Proof. It is convenient to write (3.28) as a composed function

$$f(u, v) = \frac{1}{\pi} \frac{u}{u^2 + v^2}$$

where

$$u = t \cos \frac{\pi}{2m}, \quad v = x + t \sin \frac{\pi}{2m}.$$

Since

$$\begin{aligned} \frac{\partial f}{\partial t} &= \cos \frac{\pi}{2m} \frac{\partial f}{\partial u} + \sin \frac{\pi}{2m} \frac{\partial f}{\partial v} \\ \frac{\partial^2 f}{\partial t^2} &= \cos^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial u^2} + 2 \cos \frac{\pi}{2m} \sin \frac{\pi}{2m} \frac{\partial^2 f}{\partial u \partial v} + \sin^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial v^2} \\ \frac{\partial f}{\partial x} &= \frac{\partial f}{\partial v} \quad \text{and} \quad \frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial v^2} \end{aligned}$$

and

$$\frac{\partial^2 f}{\partial u^2} + \frac{\partial^2 f}{\partial v^2} = 0$$

we have that

$$\begin{aligned} \frac{\partial^2 f}{\partial t^2} + \frac{\partial^2 f}{\partial x^2} &= \cos^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial u^2} + \frac{\partial^2 f}{\partial v^2} + 2 \sin \frac{\pi}{2m} \cos \frac{\pi}{2m} \frac{\partial^2 f}{\partial u \partial v} + \sin^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial v^2} \\ &= \frac{\partial^2 f}{\partial v^2} \left[1 - \cos^2 \frac{\pi}{2m} + \sin^2 \frac{\pi}{2m} \right] + 2 \sin \frac{\pi}{2m} \cos \frac{\pi}{2m} \frac{\partial^2 f}{\partial u \partial v} \\ &= 2 \sin \frac{\pi}{2m} \frac{\partial}{\partial v} \left[\sin \frac{\pi}{2m} \frac{\partial f}{\partial v} + \cos \frac{\pi}{2m} \frac{\partial f}{\partial u} \right] \\ &= 2 \sin \frac{\pi}{2m} \frac{\partial}{\partial x} \frac{\partial f}{\partial t} \end{aligned}$$

□

Remark 3.7. The characteristic function of (3.28) is

$$\int_{-\infty}^{+\infty} e^{i\beta x} f(x, t; m) dx = e^{-t|\beta| \cos \frac{\pi}{2m} - i\beta t \sin \frac{\pi}{2m}}$$

and can be obtained by considering the bounded solution to the Fourier transform of (3.29)

$$\frac{d^2 F}{dt^2} + 2i\beta \sin \frac{\pi}{2m} \frac{dF}{dt} - \beta^2 F = 0.$$

For the even-order Laplace equations we have the following result.

Theorem 3.8. *The solution to the higher-order Laplace-type equation*

$$\frac{\partial^{2n} u}{\partial t^{2n}} = (-1)^n \frac{\partial^{2n} u}{\partial x^{2n}}, \quad x \in \mathbb{R}, t > 0 \quad (3.30)$$

subject to the initial conditions

$$\begin{cases} u(x, 0) = \delta(x) \\ \left. \frac{\partial^k u}{\partial t^k}(x, t) \right|_{t=0^+} = \frac{(-1)^k k!}{\pi |x|^{k+1}} \cos \frac{\pi(k+1)}{2}, \quad 0 < k < 2n \end{cases} \quad (3.31)$$

is the classical Cauchy distribution given by

$$u(x, t) = Pr\{X_{2n}(S_{\frac{1}{2n}}(t)) \in dx\}/dx = \frac{t}{\pi(x^2 + t^2)}, \quad x \in \mathbb{R}, t > 0 \quad (3.32)$$

where $X_{2n}(t)$, $t > 0$ is a pseudo-process such that

$$\mathbb{E}e^{i\beta X_{2n}(t)} = e^{-t\beta^{2n}}.$$

Proof. The pseudo-process $X_{2n}(t)$, $t > 0$ related to the equation

$$\frac{\partial u}{\partial t} = (-1)^{n+1} \frac{\partial^{2n} u}{\partial t^{2n}}$$

has fundamental solution whose Fourier transform reads

$$\int_{-\infty}^{+\infty} e^{i\beta x} u(x, t) dx = e^{-t\beta^{2n}}.$$

If $S_{\frac{1}{2n}}(t)$, $t > 0$ is a stable subordinator with Laplace transform

$$\mathbb{E} \exp\left(-\lambda S_{\frac{1}{2n}}(t)\right) = \exp\left(-t\lambda^{\frac{1}{2n}}\right), \quad \lambda > 0, t > 0 \quad (3.33)$$

the characteristic function of $X_{2n}(S_{\frac{1}{2n}}(t))$, $t > 0$ becomes

$$\begin{aligned} \int_{-\infty}^{+\infty} e^{i\beta x} Pr\{X_{2n}(S_{\frac{1}{2n}}(t)) \in dx\} &= \int_0^{\infty} e^{-s\beta^{2n}} Pr\{S_{\frac{1}{2n}}(t) \in ds\} \\ &= \exp\left(-t|\beta|e^{i\frac{\pi r}{n}}\right), \quad r = 0, 1, \dots, 2n-1 \end{aligned} \quad (3.34)$$

For $r = 0$, we have the characteristic function of the Cauchy symmetric law. For $r \neq 0$ and $n \leq r \leq 2n-1$ we have a function which is not absolutely integrable and, for $0 < r < n-1$ is not a characteristic function (but can be regarded as a Cauchy r.v. at a complex time). The functions

$$F_r(\beta, t) = e^{-t|\beta|e^{i\frac{\pi r}{2k}}}$$

for all $0 \leq r \leq 2n-1$ are solutions to

$$\frac{\partial^{2n} F_r}{\partial t^{2n}} = (-1)^{n+1} F_r.$$

We now check that for $0 \leq k \leq 2n-1$ the initial conditions (3.31) are verified by the Cauchy distribution. Indeed,

$$\begin{aligned} \left. \frac{\partial^k u}{\partial t^k}(x, t) \right|_{t=0} &= \left. \frac{\partial^k}{\partial t^k} \left(\frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\beta x} e^{-t|\beta|} d\beta \right) \right|_{t=0} \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\beta x} (-1)^k |\beta|^k d\beta \\ &= \frac{(-1)^k k!}{\pi |x|^{k+1}} \cos\left(\frac{\pi(k+1)}{2}\right). \end{aligned}$$

□

Remark 3.9. We notice that for $n = 1$ the problem above becomes

$$\frac{\partial^2 u}{\partial t^2} = -\frac{\partial^2 u}{\partial x^2}, \quad x \in \mathbb{R}, t > 0$$

subject to the initial conditions

$$\begin{cases} u(x, 0) = \delta(x) \\ \left. \frac{\partial u}{\partial t}(x, t) \right|_{t=0^+} = \frac{-1}{\pi|x|^2} \cos \pi \end{cases}$$

which is in accord with

$$\left. \frac{\partial}{\partial t} \frac{t}{\pi(x^2 + t^2)} \right|_{t=0^+} = \frac{1}{\pi x^2}.$$

The connection between wave equations and the composition of two independent Cauchy processes $C^1(|C^2(t)|)$, $t > 0$ has been investigated in D'Ovidio and Orsingher [2] and more general results involving the Cauchy process have been presented in Nane [9].

Remark 3.10. We finally notice that the equation

$$\frac{\partial^6 u}{\partial t^6} + \frac{\partial^6 u}{\partial x^6} = 0 \quad (3.35)$$

can be decoupled as

$$\left(\frac{\partial^3}{\partial t^3} + i \frac{\partial^3}{\partial x^3} \right) \left(\frac{\partial^3}{\partial t^3} - i \frac{\partial^3}{\partial x^3} \right) u = 0. \quad (3.36)$$

Form the Corollary 3.3, the solution to (3.36) can be therefore written as

$$\begin{aligned} u(x, t) &= \frac{1}{2\pi} \left[\frac{\frac{\sqrt{3}}{2} t e^{i\frac{\pi}{6}}}{\left(x + \frac{t e^{i\frac{\pi}{6}}}{2}\right)^2 + \frac{3}{4} t^2 e^{i\frac{\pi}{3}}} + \frac{\frac{\sqrt{3}}{2} t e^{-i\frac{\pi}{6}}}{\left(x + \frac{t e^{-i\frac{\pi}{6}}}{2}\right)^2 + \frac{3}{4} t^2 e^{-i\frac{\pi}{3}}} \right] \\ &= \frac{\sqrt{3}}{2^2 \pi} t \left[\frac{e^{i\frac{\pi}{6}} \left(x^2 + \frac{t^2}{4} e^{-i\frac{\pi}{3}} + x t e^{-i\frac{\pi}{6}} + \frac{3}{4} t^2 e^{-i\frac{\pi}{3}}\right)}{\left(x^2 + \frac{t^2}{4} e^{-i\frac{\pi}{3}} + x t e^{-i\frac{\pi}{6}} + \frac{3}{4} t^2 e^{-i\frac{\pi}{3}}\right) \left(x^2 + \frac{t^2}{4} e^{i\frac{\pi}{3}} + x t e^{i\frac{\pi}{6}} + \frac{3}{4} t^2 e^{i\frac{\pi}{3}}\right)} \right. \\ &\quad \left. + \frac{e^{-i\frac{\pi}{6}} \left(x^2 + \frac{t^2}{4} e^{i\frac{\pi}{3}} + x t e^{i\frac{\pi}{6}} + \frac{3}{4} t^2 e^{i\frac{\pi}{3}}\right)}{\left(x^2 + \frac{t^2}{4} e^{-i\frac{\pi}{3}} + x t e^{-i\frac{\pi}{6}} + \frac{3}{4} t^2 e^{-i\frac{\pi}{3}}\right) \left(x^2 + \frac{t^2}{4} e^{i\frac{\pi}{3}} + x t e^{i\frac{\pi}{6}} + \frac{3}{4} t^2 e^{i\frac{\pi}{3}}\right)} \right] \\ &= \frac{\sqrt{3}}{2\pi} t \frac{(x^2 + t^2) \cos \frac{\pi}{6} + x t}{\left(x^2 + t^2 e^{-i\frac{\pi}{3}} + x t e^{-i\frac{\pi}{6}}\right) \left(x^2 + t^2 e^{i\frac{\pi}{3}} + x t e^{i\frac{\pi}{6}}\right)} \\ &= \frac{\sqrt{3}}{2\pi} t \frac{(x^2 + t^2) \cos \frac{\pi}{6} + x t}{\left(x^2 + t^2 + x t \cos \frac{\pi}{6}\right)^2 - 3x^2 t^2 \sin^2 \frac{\pi}{6}}. \quad (3.37) \end{aligned}$$

Equation (3.36) is satisfied by the Cauchy density and therefore by the probability law (3.37) which however is no longer a Cauchy distribution and possesses asymmetric structure.

REFERENCES

- [1] L. Beghin, Yu. Kozachenko, E. Orsingher and L. Sakhno. On the solutions of linear odd-order heat-type equations with random initial conditions. *J. Stat. Phys.* 127: 721739, 2007.
- [2] M. D'Ovidio and E. Orsingher. Composition of processes and related partial differential equations. *J. Theor. Probab.*, 24:342 – 375, 2011.

- [3] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series and products*. Academic Press, 2007. Seventh edition.
- [4] K. J. Hochberg. A signed measure on path space related to Wiener measure. *Ann. Probab.*, 6:433 – 458, 1978.
- [5] V. Yu. Krylov. Some properties of the distribution corresponding to the equation $\frac{\partial u}{\partial t} = (-1)^{q+1} \frac{\partial u^{2q}}{\partial x^{2q}}$. *Soviet Math. Dokl.*, 1:760 – 763, 1960.
- [6] A. Lachal. Distributions of sojourn time, maximum and minimum for pseudo-processes governed by higher-order heat-type equations. *Elect. J. Probab.*, 8 (20):1 – 53, 2003.
- [7] V.I. Ladokhin. On the measure on functional spaces corresponding to complex and diffusion coefficients. *Uchenye Zapiski Kazan Univ.*, 123:36 – 42, 1963. in Russian.
- [8] N. N. Lebedev. *Special functions and their applications*. Dover, New York, 1972.
- [9] E. Nane. Higher order pde's and iterated processes. *Trans. Amer. Math. Soc.*, 360(5):2681 – 2692, 2008.
- [10] E. Orsingher. Processes governed by signed measures connected with third-order "heat-type" equations. *Litovsk. Mat. Sb.*, 31(2):323 – 336, 1991. Translation in *Lithuanian Math. J.* 31 (1991), no. 2, 220 - 231 (1992).
- [11] E. Orsingher and L. Beghin. Fractional diffusion equations and processes with randomly varying time. *Ann. Probab.*, 37:206 – 249, 2009.
- [12] E. Orsingher and M. D'Ovidio. Vibrations and fractional vibrations of rods, plates and Fresnel pseudo-processes. *J. Stat. Phys.*, 145:143 – 174, 2011.

DIPARTIMENTO DI SCIENZE STATISTICHE, SAPIENZA UNIVERSITY OF ROME - P.LE A. MORO 5,
00185 ROMA, ITALY

E-mail address: enzo.orsingher@uniroma1.it

DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L'INGEGNERIA, SAPIENZA UNIVERSITY OF
ROME - VIA A. SCARPA 10, 00161 ROMA, ITALY

E-mail address: mirko.dovidio@uniroma1.it