

Tightened Exponential Bounds for Discrete Time, Conditionally Symmetric Martingales with Bounded Jumps

Igal Sason

Department of Electrical Engineering
Technion - Israel Institute of Technology
Haifa 32000, Israel
E-mail: sason@ee.technion.ac.il

Abstract

This paper derives some new exponential bounds for discrete time, real valued, conditionally symmetric martingales with bounded jumps. The new bounds are extended to conditionally symmetric sub/ supermartingales, and are compared to some existing bounds. The bounds are finally exemplified in the context of gambling.

Index Terms

Central limit theorems, concentration inequalities, conditional symmetry, discrete-time sub/ supermartingales, divergence (a.k.a. relative entropy), gambling, large deviations, moderate deviations principle, stopping times.

AMS 2000 subject classifications: 60F10, 60G40, 60G42, 60G48 (primary), 60F05, 91A60, 94A17 (secondary).

I. INTRODUCTION AND MAIN RESULTS

Classes of exponential bounds for discrete-time real-valued martingales were extensively studied in the literature (see, e.g., [1], [2], [4]–[6], [8]–[11], [14], and [17]–[22]). This paper further assumes conditional symmetry of these martingales, as it is defined in the following:

Definition 1: Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$, where $\mathbb{N}_0 \triangleq \mathbb{N} \cup \{0\}$, be a discrete-time and real-valued martingale, and let $\xi_k \triangleq X_k - X_{k-1}$ for every $k \in \mathbb{N}$ designate the jumps of the martingale. Then $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ is called a *conditionally symmetric martingale* if, conditioned on \mathcal{F}_{k-1} , the random variable ξ_k is symmetrically distributed around zero.

Our goal in this paper is to demonstrate how the assumption of the conditional symmetry improves existing exponential inequalities for discrete-time real-valued martingales with bounded increments. Earlier results, serving as motivation, appear in [9, Section 4] and [17, Section 6]. The new exponential bounds are also extended to conditionally symmetric sub or supermartingales, where the construction of these objects is exemplified later in this section. The relation of some of the exponential bounds derived in this work with some existing bounds is discussed later in this paper. Additional results addressing weak-type inequalities, maximal inequalities and ratio inequalities for conditionally symmetric martingales were derived in [15], [16] and [23].

A. Main Results of this Paper

Our main results for conditionally symmetric martingales with bounded jumps are introduced in Theorems 1, 3 and 4. Theorems 2 and 5 are existing bounds, for general martingales without the conditional symmetry assumption, that are introduced in connection to the new theorems. Corollaries 1 and 2 provide an extension of the new results to conditionally symmetric sub/ supermartingales with bounded jumps. Our first result is the following theorem:

Theorem 1: Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a discrete-time real-valued and conditionally symmetric martingale. Assume that, for some fixed numbers $d, \sigma > 0$, the following two requirements are satisfied a.s.

$$|X_k - X_{k-1}| \leq d, \quad \text{Var}(X_k | \mathcal{F}_{k-1}) = \mathbb{E}[(X_k - X_{k-1})^2 | \mathcal{F}_{k-1}] \leq \sigma^2 \quad (1)$$

for every $k \in \mathbb{N}$. Then, for every $\alpha \geq 0$ and $n \in \mathbb{N}$,

$$\mathbb{P}(|X_n - X_0| \geq \alpha n) \leq 2 \exp(-nE(\gamma, \delta)) \quad (2)$$

where

$$\gamma \triangleq \frac{\sigma^2}{d^2}, \quad \delta \triangleq \frac{\alpha}{d} \quad (3)$$

and for $\gamma \in (0, 1]$ and $\delta \in [0, 1]$

$$E(\gamma, \delta) \triangleq \delta x - \ln\left(1 + \gamma[\cosh(x) - 1]\right) \quad (4)$$

$$x \triangleq \ln\left(\frac{\delta(1 - \gamma) + \sqrt{\delta^2(1 - \gamma)^2 + \gamma^2(1 - \delta^2)}}{\gamma(1 - \delta)}\right). \quad (5)$$

If $\delta > 1$, then the probability on the left-hand side of (2) is zero (so $E(\gamma, \delta) \triangleq +\infty$), and $E(\gamma, 1) = \ln\left(\frac{2}{\gamma}\right)$.

Remark 1: From the above conditions, without any loss of generality, $\sigma^2 \leq d^2$ and therefore $\gamma \in (0, 1]$. This implies that Theorem 1 characterizes the exponent $E(\gamma, \delta)$ for all values of γ and δ .

This result should be compared to the statement in [13, Theorem 6.1] (see also [8, Corollary 2.4.7]), which does not require the conditional symmetry property. It gives the following result:

Theorem 2: Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a discrete-time real-valued martingale with bounded jumps. Assume that the two conditions in (1) are satisfied a.s. for every $k \in \mathbb{N}$. Then, for every $\alpha \geq 0$,

$$\mathbb{P}(|X_n - X_0| \geq \alpha n) \leq 2 \exp\left(-n D\left(\frac{\delta + \gamma}{1 + \gamma} \parallel \frac{\gamma}{1 + \gamma}\right)\right) \quad (6)$$

where γ and δ are introduced in (3), and

$$D(p||q) \triangleq p \ln\left(\frac{p}{q}\right) + (1 - p) \ln\left(\frac{1 - p}{1 - q}\right), \quad \forall p, q \in [0, 1] \quad (7)$$

is the divergence (a.k.a. relative entropy or Kullback-Leibler distance) to the natural base between the two probability distributions $(p, 1 - p)$ and $(q, 1 - q)$. If $\delta > 1$, then the probability on the left-hand side of (6) is zero.

Theorem 1 provides an improvement over the bound in Theorem 2 for conditionally symmetric martingales with bounded jumps. This bounds in Theorems 1 and 2 depend on the conditional variance of the martingale, but they do not take into consideration conditional moments of higher orders. In [21, Theorem 4], an alternative approach was used for the derivation of a bound that applies to discrete-time martingales with bounded jumps, where the latter bound may depend on conditional moments of higher orders (i.e., it depends on all the conditional moments of the martingale, up to the m -th order for an arbitrary even number $m \geq 2$). For conditionally symmetric martingales with bounded jumps, the result in [21, Theorem 4] is particularized to give the following theorem:

Theorem 3: Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a discrete-time real-valued and conditionally symmetric martingale. Let $m \in \mathbb{N}$ be an even number, and assume that the following conditions hold a.s. for every $k \in \mathbb{N}$

$$\begin{aligned} |X_k - X_{k-1}| &\leq d, \\ \mathbb{E}[(X_k - X_{k-1})^l | \mathcal{F}_{k-1}] &\leq \mu_l, \quad \forall l \in \{2, 4, \dots, m\} \end{aligned}$$

for some $d > 0$ and non-negative numbers $\{\mu_2, \mu_4, \dots, \mu_m\}$. Then, for every $\alpha \geq 0$,

$$\mathbb{P}(|X_n - X_0| \geq \alpha n) \leq 2 \left\{ \inf_{x \geq 0} e^{-\delta x} \left[1 + \sum_{l=1}^{\frac{m}{2}-1} \frac{\gamma_{2l} x^{2l}}{(2l)!} + \gamma_m \left(e^x - \sum_{l=0}^{m-1} \frac{x^l}{l!} \right) \right] \right\}^n \quad (8)$$

where

$$\delta \triangleq \frac{\alpha}{d}, \quad \gamma_l \triangleq \frac{\mu_l}{d^l}, \quad \forall l \in \{2, 4, \dots, m\}. \quad (9)$$

We consider in the following a different type of exponential inequalities for conditionally symmetric martingales with bounded jumps.

Theorem 4: Let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a discrete-time real-valued and conditionally symmetric martingale. Assume that there exists a fixed number $d > 0$ such that $\xi_k \triangleq X_k - X_{k-1} \leq d$ a.s. for every $k \in \mathbb{N}$. Let

$$Q_n \triangleq \sum_{k=1}^n \mathbb{E}[\xi_k^2 | \mathcal{F}_{k-1}] \quad (10)$$

with $Q_0 \triangleq 0$, be the predictable quadratic variation of the martingale up to time n . Then, for every $z, r > 0$,

$$\mathbb{P} \left(\max_{1 \leq k \leq n} (X_k - X_0) \geq z, Q_n \leq r \text{ for some } n \in \mathbb{N} \right) \leq \exp \left(-\frac{z^2}{2r} \cdot C \left(\frac{zd}{r} \right) \right) \quad (11)$$

where

$$C(u) \triangleq \frac{2[u \sinh^{-1}(u) - \sqrt{1+u^2} + 1]}{u^2}, \quad \forall u > 0. \quad (12)$$

This theorem will be compared to [10, Theorem 1.6] (see also [8, Exercise 2.4.21(b)]) that was stated without the requirement for the conditional symmetry of the martingale. It provides the following result:

Theorem 5: Let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a discrete-time real-valued martingale. Assume that there exists a fixed number $d > 0$ such that $\xi_k \triangleq X_k - X_{k-1} \leq d$ a.s. for every $k \in \mathbb{N}$. Then, for every $z, r > 0$,

$$\mathbb{P} \left(\max_{1 \leq k \leq n} (X_k - X_0) \geq z, Q_n \leq r \text{ for some } n \in \mathbb{N} \right) \leq \exp \left(-\frac{z^2}{2r} \cdot B \left(\frac{zd}{r} \right) \right) \quad (13)$$

where

$$B(u) \triangleq \frac{2[(1+u) \ln(1+u) - u]}{u^2}, \quad \forall u > 0. \quad (14)$$

The proof of [10, Theorem 1.6] is modified by using Bennett's inequality for the derivation of the original bound in Theorem 5 (without the conditional symmetry requirement). Furthermore, this modified proof serves to derive the improved bound in Theorem 4 under the conditional symmetry assumption.

Extension of The Inequalities to Discrete-Time Conditionally Symmetric Sub/ Supermartingales:

Definition 2: Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a discrete-time real-valued sub or supermartingale, and $\eta_k \triangleq X_k - \mathbb{E}[X_k | \mathcal{F}_{k-1}]$ for every $k \in \mathbb{N}$. Then the martingale $\{\eta_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ is called, respectively, a conditionally symmetric sub or supermartingale if, conditioned on \mathcal{F}_{k-1} , the random variable η_k is symmetrically distributed around zero.

Remark 2: For martingales, $\eta_k = \xi_k$ for every $k \in \mathbb{N}$, so we obtain consistency with Definition 1.

An extension of Theorem 1 to conditionally symmetric sub and supermartingales is introduced in the following:

Corollary 1: Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a discrete-time real-valued and conditionally symmetric supermartingale. Assume that, for some constants $d, \sigma > 0$, the following two requirements are satisfied a.s.

$$\eta_k \leq d, \quad \text{Var}(X_k | \mathcal{F}_{k-1}) \triangleq \mathbb{E}[\eta_k^2 | \mathcal{F}_{k-1}] \leq \sigma^2 \quad (15)$$

for every $k \in \mathbb{N}$. Then, for every $\alpha \geq 0$ and $n \in \mathbb{N}$,

$$\mathbb{P}(X_n - X_0 \geq \alpha n) \leq \exp(-n E(\gamma, \delta)) \quad (16)$$

where γ and δ are defined as in (3), and $E(\gamma, \delta)$ is introduced in (4). Alternatively, if $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ is a submartingale, the same bound holds for the probability $\mathbb{P}(X_n - X_0 \leq -\alpha n)$ provided that $\eta_k \geq -d$ and the second condition in (15) hold a.s. for every $k \in \mathbb{N}$. If $\delta > 1$, then these two probabilities are equal to zero.

The following statement extends Theorem 4 to conditionally symmetric supermartingales.

Corollary 2: Let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a discrete-time real-valued supermartingale. Assume that there exists a fixed number $d > 0$ such that $\eta_k \leq d$ a.s. for every $k \in \mathbb{N}$. Let $\{Q_n\}_{n \in \mathbb{N}_0}$ be the predictable quadratic variations of the supermartingale, i.e., $Q_n \triangleq \sum_{k=1}^n \mathbb{E}[\eta_k^2 | \mathcal{F}_{k-1}]$ for every $n \in \mathbb{N}$ with $Q_0 \triangleq 0$. Then, the result in (13) holds. Furthermore, if the above supermartingale is conditionally symmetric, then the improved bound in (11) holds.

B. Construction of Discrete-Time Conditionally Symmetric Sub/ Supermartingales

Before proving the tightened inequalities for discrete-time conditionally symmetric sub/ supermartingales, it is in place to exemplify the construction of these objects.

Example 1: Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $\{U_k\}_{k \in \mathbb{N}} \subseteq L^1(\Omega, \mathcal{F}, \mathbb{P})$ be a sequence of i.i.d. random variables with zero mean. Let $\{\mathcal{F}_k\}_{k \geq 0}$ be a filtration of sub σ -algebras of \mathcal{F} , where

$$\mathcal{F}_0 = \{\emptyset, \Omega\}, \quad \mathcal{F}_k = \sigma(U_1, \dots, U_k), \quad \forall k \in \mathbb{N}.$$

Furthermore, for $k \in \mathbb{N}$, let $A_k \in L^\infty(\Omega, \mathcal{F}_{k-1}, \mathbb{P})$ be an \mathcal{F}_{k-1} -measurable random variable with a finite essential supremum. Define a new sequence of random variables in $L^1(\Omega, \mathcal{F}, \mathbb{P})$ where

$$X_n = \sum_{k=1}^n A_k U_k, \quad \forall n \in \mathbb{N}$$

and $X_0 = 0$. Then, $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a martingale. Lets assume that the random variables $\{U_k\}_{k \in \mathbb{N}}$ are symmetrically distributed around zero. Note that $X_n = X_{n-1} + A_n U_n$ where A_n is \mathcal{F}_{n-1} -measurable and U_n is independent of the σ -algebra \mathcal{F}_{n-1} (due to the independence of the random variables U_1, \dots, U_n). It therefore follows that for every $n \in \mathbb{N}$, given \mathcal{F}_{n-1} , the random variable X_n is symmetrically distributed around its conditional expectation X_{n-1} . Hence, the martingale $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is conditionally symmetric.

Example 2: In continuation to Example 1, let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a martingale, and define $Y_0 = 0$ and

$$Y_n = \sum_{k=1}^n A_k (X_k - X_{k-1}), \quad \forall n \in \mathbb{N}.$$

The sequence $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a martingale. If $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a conditionally symmetric martingale then also the martingale $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is conditionally symmetric (since $Y_n = Y_{n-1} + A_n (X_n - X_{n-1})$, and by assumption A_n is \mathcal{F}_{n-1} -measurable).

Example 3: In continuation to Example 1, let $\{U_k\}_{k \in \mathbb{N}}$ be i.i.d. random variables with a symmetric distribution around their expected value, and assume that $\mathbb{E}(U_k) \leq 0$ for every $k \in \mathbb{N}$. Furthermore, let $A_k \in L^\infty(\Omega, \mathcal{F}_{k-1}, \mathbb{P})$, and assume that a.s. $A_k \geq 0$ for every $k \in \mathbb{N}$. Let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a martingale as defined in Example 1. Note that $X_n = X_{n-1} + A_n U_n$ where A_n is non-negative and \mathcal{F}_{n-1} -measurable, and U_n is independent of \mathcal{F}_{n-1} and symmetrically distributed around its average. This implies that $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a conditionally symmetric supermartingale.

Example 4: In continuation to Examples 2 and 3, let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a conditionally symmetric supermartingale. Define $\{Y_n\}_{n \in \mathbb{N}_0}$ as in Example 2 where A_k is non-negative a.s. and \mathcal{F}_{k-1} -measurable for every $k \in \mathbb{N}$. Then $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a conditionally symmetric supermartingale.

Example 5: Consider a standard Brownian motion $(W_t)_{t \geq 0}$. Define, for some $T > 0$, the discrete-time process

$$X_n = W_{nT}, \quad \mathcal{F}_n = \sigma(\{W_t\}_{0 \leq t \leq nT}), \quad \forall n \in \mathbb{N}_0.$$

The increments of $(W_t)_{t \geq 0}$ over time intervals $[t_{k-1}, t_k]$ are statistically independent if these intervals do not overlap (except of their endpoints), and they are Gaussian distributed with a zero mean and variance $t_k - t_{k-1}$. The random variable $\xi_n \triangleq X_n - X_{n-1}$ is therefore statistically independent of \mathcal{F}_{n-1} , and it is Gaussian distributed with a zero mean and variance T . The martingale $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is therefore conditionally symmetric.

C. Structure of the Paper

This paper is structured as follows: Introduction and the main results of this work are provided in this section. Proofs of these theorems are introduced in Section II. Discussion of the new results and some other related exponential inequalities in the literature are considered in Section III. An example of the possible use of the bounds is presented in Section IV in the context of gambling.

II. PROOFS

A. Combined Proof of Theorems 1 and 2

We rely here on the proof of the existing bound that is stated in Theorem 2, for discrete-time real-valued martingales with bounded jumps (see [13, Theorem 6.1] and [8, Corollary 2.4.7]), and then deviate from this proof at the point where the additional property of the conditional symmetry of the martingale is taken into consideration for the derivation of the improved exponential inequality in Theorem 1.

Write $X_n - X_0 = \sum_{k=1}^n \xi_k$ where $\xi_k = X_k - X_{k-1}$ for $k = 1, \dots, n$. By assumption $|\xi_k| \leq d$ a.s., and $\mathbb{E}[\xi_k | \mathcal{F}_{k-1}] = 0$ for a discrete-time martingale. Based on Chernoff's inequality, then for an arbitrary $\alpha \geq 0$,

$$\mathbb{P}(X_n - X_0 \geq \alpha n) \leq \exp(-\alpha n t) \mathbb{E} \left[\exp \left(t \sum_{k=1}^n \xi_k \right) \right], \quad \forall t \geq 0. \quad (17)$$

Furthermore,

$$\begin{aligned} & \mathbb{E} \left[\exp \left(t \sum_{k=1}^n \xi_k \right) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[\exp \left(t \sum_{k=1}^n \xi_k \right) \mid \mathcal{F}_{n-1} \right] \right] \\ &= \mathbb{E} \left[\exp \left(t \sum_{k=1}^{n-1} \xi_k \right) \mathbb{E} [\exp(t\xi_n) \mid \mathcal{F}_{n-1}] \right] \end{aligned} \quad (18)$$

where the last transition holds since $\exp(t \sum_{k=1}^{n-1} \xi_k)$ is \mathcal{F}_{n-1} -measurable. Theorem 2, for a discrete-time real-valued martingale with bounded jumps, follows by applying Bennett's inequality [3] (see, e.g., [8, Lemma 2.4.1]) for the conditional law of ξ_k given the σ -algebra \mathcal{F}_{k-1} . Since $\mathbb{E}[\xi_k | \mathcal{F}_{k-1}] = 0$, $\text{Var}[\xi_k | \mathcal{F}_{k-1}] \leq \sigma^2$ and $\xi_k \leq d$ a.s. for $k \in \mathbb{N}$, then a.s. (see (3))

$$\mathbb{E} [\exp(t\xi_k) \mid \mathcal{F}_{k-1}] \leq \frac{\gamma \exp(td) + \exp(-\gamma td)}{1 + \gamma}, \quad \forall t \geq 0. \quad (19)$$

By induction, from (18) and (19), it follows that for every $t \geq 0$

$$\mathbb{E} \left[\exp \left(t \sum_{k=1}^n \xi_k \right) \right] \leq \left(\frac{\gamma \exp(td) + \exp(-\gamma td)}{1 + \gamma} \right)^n. \quad (20)$$

Combining (17) and (20), and then optimizing the bound on $\mathbb{P}(X_n - X_0 \geq \alpha n)$ over the non-negative free parameter t gives after the inequality in Theorem 2 (note that the same bound holds also for the probability $\mathbb{P}(X_n - X_0 \leq -\alpha n)$, so the bound on $\mathbb{P}(|X_n - X_0| \geq \alpha n)$ follows by doubling the bound on $\mathbb{P}(X_n - X_0 \geq \alpha n)$).

In order to prove Theorem 1 for a real-valued conditionally symmetric martingale with bounded jumps, we deviate from the proof of Theorem 2 starting from a replacement of Bennett's inequality for the conditional expectation in (19) with a tightened bound under the conditional symmetry assumption. To this end, we need a lemma to proceed.

Lemma 1: Let X be a real-valued RV with a symmetric distribution around zero, a support $[-d, d]$, and assume that $\mathbb{E}[X^2] = \text{Var}(X) \leq \gamma d^2$ for some $d > 0$ and $\gamma \in [0, 1]$. Let h be a real-valued convex function, and assume that $h(d^2) \geq h(0)$. Then

$$\mathbb{E}[h(X^2)] \leq (1 - \gamma)h(0) + \gamma h(d^2) \quad (21)$$

where equality holds for the symmetric distribution

$$\mathbb{P}(X = d) = \mathbb{P}(X = -d) = \frac{\gamma}{2}, \quad \mathbb{P}(X = 0) = 1 - \gamma. \quad (22)$$

Proof: Since h is convex and the support of X is $[-d, d]$, then a.s. $h(X^2) \leq h(0) + \left(\frac{X}{d}\right)^2 (h(d^2) - h(0))$. Eq. (21) follows by taking expectation on both sides of the last inequality, and it holds with equality for the symmetric distribution in (22). ■

Corollary 3: If X is a random variable that satisfies the three requirements in Lemma 1 then, for every $\lambda \in \mathbb{R}$,

$$\mathbb{E}[\exp(\lambda X)] \leq 1 + \gamma[\cosh(\lambda d) - 1] \quad (23)$$

and (23) holds with equality for the symmetric distribution in Lemma 1, independently of the value of λ .

Proof: For every $\lambda \in \mathbb{R}$, due to the symmetric distribution of X , $\mathbb{E}[\exp(\lambda X)] = \mathbb{E}[\cosh(\lambda X)]$. The claim now follows from Lemma 1 since, for every $x \in \mathbb{R}$, $\cosh(\lambda x) = h(x^2)$ where $h(x) \triangleq \sum_{n=0}^{\infty} \frac{\lambda^{2n} |x|^{2n}}{(2n)!}$ is a convex function (h is convex since it is a linear combination, with non-negative coefficients, of convex functions), and $h(d^2) = \cosh(\lambda d) \geq 1 = h(0)$. ■

We continue with the proof of Theorem 1. Under the assumption of this theorem, for every $k \in \mathbb{N}$, the random variable $\xi_k \triangleq X_k - X_{k-1}$ satisfies a.s.

$$\mathbb{E}[\xi_k | \mathcal{F}_{k-1}] = 0, \quad \mathbb{E}[(\xi_k)^2 | \mathcal{F}_{k-1}] \leq \sigma^2.$$

Applying Corollary 3 for the conditional law of ξ_k given \mathcal{F}_{k-1} , it follows that for every $k \in \mathbb{N}$ and $t \in \mathbb{R}$ a.s.

$$\mathbb{E}[\exp(t\xi_k) | \mathcal{F}_{k-1}] \leq 1 + \gamma[\cosh(td) - 1] \quad (24)$$

and therefore (18) and (24), it follows that for every $t \in \mathbb{R}$

$$\mathbb{E} \left[\exp \left(t \sum_{k=1}^n \xi_k \right) \right] \leq \left(1 + \gamma[\cosh(td) - 1] \right)^n. \quad (25)$$

Therefore, from (17), for every $t \geq 0$,

$$\mathbb{P}(X_n - X_0 \geq \alpha n) \leq \exp(-\alpha n t) \left(1 + \gamma[\cosh(td) - 1] \right)^n.$$

From (3), the substitution $x = td$ gives that for an arbitrary $\alpha \geq 0$

$$\mathbb{P}(X_n - X_0 \geq \alpha n) \leq \inf_{x \geq 0} \left\{ \exp \left(-n \left[\delta x - \ln(1 + \gamma[\cosh(x) - 1]) \right] \right) \right\}. \quad (26)$$

An optimization over the non-negative parameter x gives the solution for the optimized parameter in (5).

Applying (26) for the martingale $\{-X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ gives the same upper bound on $\mathbb{P}(X_n - X_0 \leq -\alpha n)$ for an arbitrary $\alpha \geq 0$. This gives the factor 2 in the pre-exponent of the upper bound on $\mathbb{P}(|X_n - X_0| \geq \alpha n)$, which concludes the proof of Theorem 1.

B. Proof of Theorem 3

This inequality in Theorem 3 follows from [21, Theorem 4], which is satisfied for a general discrete-time real-valued martingale with bounded jumps. Due to the conditional symmetry of the martingale, then all the conditional moments of $\xi_k \triangleq X_k - X_{k-1}$ of odd orders are equal to zero, i.e., $\mathbb{E}[(\xi_k)^l | \mathcal{F}_{k-1}] = 0$ for every $l \in \{1, 3, 5, \dots\}$. Note that all the conditional moments are finite since $|\xi_k| \leq d$ a.s. for every $k \in \mathbb{N}$. Consequently, $\gamma_l = 0$ for all the odd values of $l \geq 1$. The substitution of the last equality in [21, Theorem 4] gives that, for an arbitrary $\alpha \geq 0$,

$$\begin{aligned} & \mathbb{P}(|X_n - X_0| \geq \alpha n) \\ & \leq 2 \left\{ \inf_{x \geq 0} e^{-\delta x} \left[1 + \sum_{l=2}^{m-1} \frac{(\gamma_l - \gamma_m) x^l}{l!} + \gamma_m (e^x - 1 - x) \right] \right\}^n \\ & = 2 \left\{ \inf_{x \geq 0} e^{-\delta x} \left[1 + \sum_{l=1}^{m-1} \frac{\gamma_l x^l}{l!} + \gamma_m \left(e^x - \sum_{l=0}^{m-1} \frac{x^l}{l!} \right) \right] \right\}^n \\ & = 2 \left\{ \inf_{x \geq 0} e^{-\delta x} \left[1 + \sum_{l=1}^{\frac{m}{2}-1} \frac{\gamma_{2l} x^{2l}}{(2l)!} + \gamma_m \left(e^x - \sum_{l=0}^{m-1} \frac{x^l}{l!} \right) \right] \right\}^n. \end{aligned}$$

C. A Combined Proof of Theorems 4 and 5

The proof of Theorem 4 relies on the proof of the known result in Theorem 5, where the latter dates back to Freedman's paper (see [10, Theorem 1.6], and also [8, Exercise 2.4.21(b)]). By revisiting the original proof of Theorem 5 (see [10, Section 3]), it is modified in a way which facilitates to realize, in light of Bennett's inequality and its improvement in (24) for conditionally symmetric distributions, how the bound can be improved for conditionally symmetric martingales with bounded jumps. Furthermore, the following revisited proof of Theorem 5 simplifies the derivation of the new and improved bound in Theorem 4 for the considered subclass of discrete-time, real-valued and conditionally symmetric martingales with bounded jumps.

Without any loss of generality, let's assume that $d = 1$ (otherwise, $\{X_k\}$ and z are divided by d , and $\{Q_k\}$ and r are divided by d^2 ; this normalization extends the bound to the case of an arbitrary $d > 0$). Let $S_n \triangleq X_n - X_0$ for every $n \in \mathbb{N}_0$, then $\{S_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a martingale, and $S_0 = 0$. The proof starts by introducing two lemmas.

Lemma 2: Under the assumptions of Theorem 5, let

$$U_n \triangleq \exp(\lambda S_n - \theta Q_n), \quad \forall n \in \{0, 1, \dots\} \quad (27)$$

where $\lambda \geq 0$ and $\theta \geq e^\lambda - \lambda - 1$ are arbitrary constants. Then, $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a supermartingale.

Proof: U_n in (27) is \mathcal{F}_n -measurable since Q_n in (10) is \mathcal{F}_{n-1} -measurable (where $\mathcal{F}_{n-1} \subseteq \mathcal{F}_n$) and S_n is \mathcal{F}_n -measurable. Since Q_n and U_n are non-negative, and $S_n = \sum_{k=1}^n \xi_k \leq n$ a.s. (note that $\xi_k \leq 1$ a.s. for every $k \in \mathbb{N}$, and $S_0 = 0$ a.s.) then for $\theta \geq 0$

$$\mathbb{E}[|U_n|] = \mathbb{E}[U_n] \leq e^{\lambda n} < \infty$$

so $U_n \in L^1(\Omega, \mathcal{F}_n, \mathbb{P})$. It is therefore required to show that $\mathbb{E}[U_n | \mathcal{F}_{n-1}] \leq U_{n-1}$ a.s. for every $n \in \mathbb{N}$ under the following assumptions on the fixed parameters λ and θ in (27). For every $n \in \mathbb{N}$, the following holds a.s.

$$\begin{aligned} & \mathbb{E}[U_n | \mathcal{F}_{n-1}] \\ & \stackrel{(a)}{=} \exp(-\theta Q_n) \exp(\lambda S_{n-1}) \mathbb{E}[\exp(\lambda \xi_n) | \mathcal{F}_{n-1}] \\ & \stackrel{(b)}{=} \exp(\lambda S_{n-1}) \exp(-\theta(Q_{n-1} + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}])) \mathbb{E}[\exp(\lambda \xi_n) | \mathcal{F}_{n-1}] \\ & \stackrel{(c)}{=} U_{n-1} \left(\frac{\mathbb{E}[\exp(\lambda \xi_n) | \mathcal{F}_{n-1}]}{\exp(\theta \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}])} \right) \end{aligned} \quad (28)$$

where (a) follows from (27) and because Q_n and S_{n-1} are \mathcal{F}_{n-1} -measurable and $S_n = S_{n-1} + \xi_n$, (b) follows from (10), and (c) follows from (27).

A modification to the original proof of Lemma 2 (see [10, Section 3]) is suggested in the following, which then enables to obtain an improvement of Theorem 5 for real-valued discrete-time conditionally symmetric martingales with bounded jumps. This will lead to the improved bound in Theorem 4 for the considered subclass of martingales.

Applying Bennett's inequality in (19) to the conditional expectation of $e^{\lambda \xi_n}$ given \mathcal{F}_{n-1} , since by assumption $\xi_n \leq 1$ and $\mathbb{E}[\xi_n | \mathcal{F}_{n-1}] = 0$ a.s., then (recall that $\lambda \geq 0$)

$$\mathbb{E}[\exp(\lambda \xi_n) | \mathcal{F}_{n-1}] \leq \frac{\exp(-\lambda \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}]) + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}] \exp(\lambda)}{1 + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}]} \quad (29)$$

Substituting the inequality in (29) into the right-hand of (28) gives that

$$\mathbb{E}[U_n | \mathcal{F}_{n-1}] \leq U_{n-1} \left(\frac{\exp(-(\lambda + \theta) \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}])}{1 + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}]} + \frac{\mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}] \exp(\lambda - \theta \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}])}{1 + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}]} \right). \quad (30)$$

In order to prove that $\mathbb{E}[U_n | \mathcal{F}_{n-1}] \leq U_{n-1}$ a.s., it is sufficient to prove that the second term on the right-hand side of (30) is a.s. less than or equal to 1. To this end let's, find the condition on $\lambda, \theta \geq 0$ such that for every $\alpha \geq 0$

$$\left(\frac{1}{1 + \alpha} \right) \exp(-\alpha(\lambda + \theta)) + \left(\frac{\alpha}{1 + \alpha} \right) \exp(\lambda - \alpha\theta) \leq 1 \quad (31)$$

which then assures that the second term on the right-hand side of (30) is less than or equal to 1 a.s. as required.

Lemma 3: If $\lambda \geq 0$ and $\theta \geq \exp(\lambda) - \lambda - 1$ then the condition in (31) is satisfied for every $\alpha \geq 0$.

Proof: This claim follows by calculus, showing that the function

$$g(\alpha) = (1 + \alpha) \exp(\alpha\theta) - \alpha \exp(\lambda) - \exp(-\alpha\lambda), \quad \forall \alpha \geq 0$$

is non-negative on \mathbb{R}_+ if $\lambda \geq 0$ and $\theta \geq \exp(\lambda) - \lambda - 1$. ■

From (30) and Lemma 3, it follows that $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a supermartingale if $\lambda \geq 0$ and $\theta \geq \exp(\lambda) - \lambda - 1$. This completes the proof of Lemma 2. ■

At this point we deviate from the (modified) proof of Theorem 5, to obtain the tightened bound in Theorem 4 when the martingale is conditionally symmetric. As earlier, it is assumed without any loss of generality that $d = 1$.

Lemma 4: Under the additional assumption of the conditional symmetry in Theorem 4, then $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ in (27) is a supermartingale if $\lambda \geq 0$ and $\theta \geq \cosh(\lambda) - 1$ are arbitrary constants.

Proof: By assumption $\xi_n = S_n - S_{n-1} \leq 1$ a.s., and ξ_n is conditionally symmetric around zero, given \mathcal{F}_{n-1} , for every $n \in \mathbb{N}$. By applying Corollary 3 to the conditional expectation of $\exp(\lambda\xi_n)$ given \mathcal{F}_{n-1} , for every $\lambda \geq 0$,

$$\mathbb{E}[\exp(\lambda\xi_n) | \mathcal{F}_{n-1}] \leq 1 + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}] (\cosh(\lambda) - 1). \quad (32)$$

Hence, combining (28) and (32) gives

$$\mathbb{E}[U_n | \mathcal{F}_{n-1}] \leq U_{n-1} \left(\frac{1 + \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}] (\cosh(\lambda) - 1)}{\exp(\theta \mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}])} \right). \quad (33)$$

Let $\lambda \geq 0$. Since $\mathbb{E}[\xi_n^2 | \mathcal{F}_{n-1}] \geq 0$ a.s. then in order to ensure that $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ forms a supermartingale, it is sufficient (based on (33)) that the following condition holds:

$$\frac{1 + \alpha (\cosh(\lambda) - 1)}{\exp(\theta\alpha)} \leq 1, \quad \forall \alpha \geq 0. \quad (34)$$

By calculus it follows that, for $\lambda \geq 0$, the condition in (33) is satisfied if and only if

$$\theta \geq \cosh(\lambda) - 1 \triangleq \theta_{\min}(\lambda). \quad (35)$$

From (33), $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a supermartingale if $\lambda \geq 0$ and $\theta \geq \theta_{\min}(\lambda)$. This proves Lemma 4. ■

Hence, due to the assumption of the conditional symmetry of the martingale in Theorem 4, the set of parameters for which $\{U_n, \mathcal{F}_n\}$ is a supermartingale was extended. This follows from a comparison of Lemma 2 and 4 where indeed $\exp(\lambda) - 1 - \lambda \geq \theta_{\min}(\lambda) \geq 0$ for every $\lambda \geq 0$.

Let $z, r > 0$, $\lambda \geq 0$ and either $\theta \geq \cosh(\lambda) - 1$ or $\theta \geq \exp(\lambda) - \lambda - 1$ with or without assuming the conditional symmetry property, respectively (see Lemma 2 and 4). In the following, we rely on Doob's sampling theorem. To this end, let $M \in \mathbb{N}$, and define two stopping times adapted to $\{\mathcal{F}_n\}$. The first stopping time is $\alpha = 0$, and the second stopping time β is defined to be the minimal value of $n \in \{0, \dots, M\}$ (if any) such that $S_n \geq z$ and $Q_n \leq r$ (note that S_n is \mathcal{F}_n -measurable, and Q_n is \mathcal{F}_{n-1} -measurable, and since $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \dots$ then the event $\{\beta \leq n\}$ is \mathcal{F}_n -measurable); if such a value of n does not exist, let $\beta \triangleq M$. Hence $\alpha \leq \beta$ are two bounded stopping times. From Lemma 2 or 4, $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a supermartingale for the corresponding set of parameters of λ and θ , and according to Doob's sampling theorem

$$\mathbb{E}[U_\beta] \leq \mathbb{E}[U_0] = 1 \quad (36)$$

($S_0 = Q_0 = 0$, so from (27), $U_0 = 1$ a.s.). Hence, this implies the following chain of inequalities:

$$\begin{aligned} & \mathbb{P}(\exists n \leq M : S_n \geq z, Q_n \leq r) \\ & \stackrel{(a)}{=} \mathbb{P}(S_\beta \geq z, Q_\beta \leq r) \\ & \stackrel{(b)}{\leq} \mathbb{P}(\lambda S_\beta - \theta Q_\beta \geq \lambda z - \theta r) \\ & \stackrel{(c)}{\leq} \frac{\mathbb{E}[\exp(\lambda S_\beta - \theta Q_\beta)]}{\exp(\lambda z - \theta r)} \\ & \stackrel{(d)}{=} \frac{\mathbb{E}[U_\beta]}{\exp(\lambda z - \theta r)} \\ & \stackrel{(e)}{\leq} \exp(-(\lambda z - \theta r)) \end{aligned} \quad (37)$$

where equality (a) follows from the definition of the stopping time $\beta \in \{0, \dots, M\}$, (b) holds since $\lambda, \theta \geq 0$, (c) follows from Chernoff's bound, (d) follows from the definition in (27), and finally (e) follows from (36). Since (37) holds for every $M \in \mathbb{N}$, then from the continuity theorem for non-decreasing events and (37)

$$\begin{aligned} & \mathbb{P}(\exists n \in \mathbb{N} : S_n \geq z, Q_n \leq r) \\ &= \lim_{M \rightarrow \infty} \mathbb{P}(\exists n \leq M : S_n \geq z, Q_n \leq r) \\ &\leq \exp(-(\lambda z - \theta r)). \end{aligned} \tag{38}$$

The choice of the non-negative parameter θ as the minimal value for which (38) is valid provides the tightest bound within this form. Hence, without assuming the conditional symmetry property for the martingale $\{X_n, \mathcal{F}_n\}$, let (see Lemma 2) $\theta = \exp(\lambda) - \lambda - 1$. This gives that for every $z, r > 0$,

$$\mathbb{P}(\exists n \in \mathbb{N} : S_n \geq z, Q_n \leq r) \leq \exp\left(-[\lambda z - (\exp(\lambda) - \lambda - 1)r]\right), \quad \forall \lambda \geq 0.$$

The minimization w.r.t. λ gives that $\lambda = \ln\left(1 + \frac{z}{r}\right)$, and its substitution in the bound yields that

$$\mathbb{P}(\exists n \in \mathbb{N} : S_n \geq z, Q_n \leq r) \leq \exp\left(-\frac{z^2}{2r} \cdot B\left(\frac{z}{r}\right)\right) \tag{39}$$

where the function B is introduced in (14).

Furthermore, under the assumption that the martingale $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is conditionally symmetric, let $\theta = \theta_{\min}(\lambda)$ (see Lemma 4) for obtaining the tightest bound in (38) for a fixed $\lambda \geq 0$. This gives the inequality

$$\mathbb{P}(\exists n \in \mathbb{N} : S_n \geq z, Q_n \leq r) \leq \exp\left(-[\lambda z - r \theta_{\min}(\lambda)]\right), \quad \forall \lambda \geq 0.$$

The optimized λ is equal to $\lambda = \sinh^{-1}\left(\frac{z}{r}\right)$. Its substitution in (35) gives that $\theta_{\min}(\lambda) = \sqrt{1 + \frac{z^2}{r^2}} - 1$, and

$$\mathbb{P}(\exists n \in \mathbb{N} : S_n \geq z, Q_n \leq r) \leq \exp\left(-\frac{z^2}{2r} C\left(\frac{z}{r}\right)\right) \tag{40}$$

where the function C is introduced in (12).

Finally, the proof of Theorems 4 and 5 is completed by showing that the following equality holds:

$$\begin{aligned} A &\triangleq \{\exists n \in \mathbb{N} : S_n \geq z, Q_n \leq r\} \\ &= \{\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} S_k \geq z, Q_n \leq r\} \triangleq B. \end{aligned} \tag{41}$$

Clearly $A \subseteq B$, so one needs to show that $B \subseteq A$. To this end, assume that event B is satisfied. Then, there exists some $n \in \mathbb{N}$ and $k \in \{1, \dots, n\}$ such that $S_k \geq z$ and $Q_n \leq r$. Since the quadratic variation process $\{Q_n\}_{n \in \mathbb{N}_0}$ in (10) is monotonic non-decreasing, then it implies that $S_k \geq z$ and $Q_k \leq r$; therefore, event A is also satisfied and $B \subseteq A$. The combination of (40) and (41) completes the proof of Theorem 4, and respectively the combination of (39) and (41) completes the proof of Theorem 5.

We prove in the following Corollaries 1 and 2 that extend, respectively, Theorems 1 and 4 to real-valued discrete-time conditionally symmetric sub/ supermartingales.

D. Proof of Corollary 1

The proof of Corollary 1 is similar to the proof of Theorem 1. The only difference is that for a supermartingale, $X_n - X_0 = \sum_{k=1}^n (X_k - X_{k-1}) \leq \sum_{k=1}^n \eta_k$ a.s., where $\eta_k \triangleq X_k - \mathbb{E}[X_k | \mathcal{F}_{k-1}]$ is \mathcal{F}_k -measurable. Hence $\mathbb{P}(X_n - X_0 \geq \alpha n) \leq \mathbb{P}(\sum_{k=1}^n \eta_k \geq \alpha n)$ where a.s. $\eta_k \leq d$, $\mathbb{E}[\eta_k | \mathcal{F}_{k-1}] = 0$, and $\text{Var}(\eta_k | \mathcal{F}_{k-1}) \leq \sigma^2$. The continuation of the proof coincides with the proof of Theorem 1 (starting from (17) and (25)). Since $\eta_k \leq d$ a.s. for every $k \in \{1, \dots, n\}$, then $X_n - X_0 \leq nd$ a.s.; hence, if $\delta > 1$ (i.e., $\alpha > d$), then the probability on the left-hand side of (16) is equal to zero. The other inequality for submartingales holds due to the fact that if $\{X_k, \mathcal{F}_k\}$ is a conditionally symmetric submartingale, then $\{-X_k, \mathcal{F}_k\}$ is a conditionally symmetric supermartingale.

E. Proof of Corollary 2

Since $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a supermartingale, then a.s., for every $k \in \mathbb{N}$, $X_k - X_0 \leq \sum_{j=1}^k \eta_j$ where, as before, $\eta_j \triangleq X_j - \mathbb{E}[X_j | \mathcal{F}_{j-1}]$. Consider the martingale $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ where $Y_n \triangleq \sum_{j=1}^n \eta_j$ for every $n \in \mathbb{N}$, and $Y_0 \triangleq 0$ (it is a martingale since a.s. $\mathbb{E}[\eta_j | \mathcal{F}_{j-1}] = 0$). Since $Y_k - Y_{k-1} = \eta_k$ for every $k \in \mathbb{N}$, then the predictable quadratic variation process $\{Q_n\}_{n \in \mathbb{N}_0}$ which corresponds to the martingale $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is, from (10), the same process as the one which corresponds to the supermartingale $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$. Furthermore, $X_k - X_0 \leq \sum_{j=1}^k \xi_j = Y_k - Y_0$ for every $k \in \mathbb{N}$. Hence, it follows that for every $z, r > 0$,

$$\mathbb{P}\left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} (X_k - X_0) \geq z, Q_n \leq r\right) \leq \mathbb{P}\left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} (Y_k - Y_0) \geq z, Q_n \leq r\right).$$

Theorem 5, applied to the martingale $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$, gives the satisfiability of the inequality in (13) and (14). If $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a conditionally symmetric supermartingale, then $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a conditionally symmetric martingale. Theorem 4, applied again to the martingale $\{Y_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$, implies the improved inequality in (11) and (12) under the conditional symmetry assumption of the supermartingale.

III. DISCUSSION OF THE NEW RESULTS AND SOME OTHER KNOWN BOUNDS

We discuss in the following the new results, and also consider some relations with some other known exponential bounds for discrete-time real-valued conditionally symmetric martingales.

A. Discussion of the New Results and Their Proofs

Remark 3: Note that the two exponents in Theorems 2 and 1 are both discontinuous at $\delta = 1$. This is consistent with the assumption of the bounded jumps that implies that $\mathbb{P}(|X_n - X_0| \geq nd\delta)$ is equal to zero if $\delta > 1$.

If $\delta \rightarrow 1^-$ then, from (5), $x \rightarrow \infty$; hence, for every $\gamma \in (0, 1]$,

$$\lim_{\delta \rightarrow 1^-} E(\gamma, \delta) = \lim_{x \rightarrow \infty} [x - \ln(1 + \gamma(\cosh(x) - 1))] = \ln\left(\frac{2}{\gamma}\right). \tag{42}$$

On the other hand, the right limit at $\delta = 1$ is infinity since $E(\gamma, \delta) = +\infty$ for every $\delta > 1$. The same discontinuity also exists for the exponent in Theorem 2 where the right limit at $\delta = 1$ is infinity, and the left limit is equal to

$$\lim_{\delta \rightarrow 1^-} D\left(\frac{\delta + \gamma}{1 + \gamma} \parallel \frac{\gamma}{1 + \gamma}\right) = \ln\left(1 + \frac{1}{\gamma}\right) \tag{43}$$

where the last equality follows from (7). A comparison of the limits in (42) and (43) is consistent with the improvement that is obtained in Theorem 1 as compared to Theorem 2 due to the additional assumption of the conditional symmetry that is relevant if $\gamma \in (0, 1)$. Furthermore, straightforward calculations show that the two exponents indeed coincide if $\gamma = 1$ (which is equivalent to removing the constraint on the conditional variance and staying only with the assumption of the bounded differences), and their common value is equal to

$$f(\delta) = \begin{cases} \ln(2) \left[1 - h_2\left(\frac{1-\delta}{2}\right)\right], & 0 \leq \delta \leq 1 \\ +\infty, & \delta > 1 \end{cases} \tag{44}$$

and $h_2(x) \triangleq -x \log_2(x) - (1-x) \log_2(1-x)$ for $0 \leq x \leq 1$ denotes the binary entropy function to the base 2.

Remark 4: Theorem 2 provides the correct exponential decay for moderate deviations. More explicitly, we get the following:

Proposition 1: let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a discrete-time real-valued martingale, and let $\eta \in [\frac{1}{2}, 1)$ be a fixed number. Then, from Theorem 2, for every $\alpha \geq 0$,

$$\mathbb{P}(|X_n - X_0| \geq \alpha n^\eta) \leq 2 \exp\left(-\frac{\delta^2 n^{2\eta-1}}{2\gamma}\right) \left(1 + O(n^{-(1-\eta)})\right). \tag{45}$$

where γ and δ are introduced in (3). In particular, the following central limit theorem (CLT) holds for every $\alpha \geq 0$

$$\mathbb{P}(|X_n - X_0| \geq \alpha \sqrt{n}) \leq 2 \exp\left(-\frac{\delta^2}{2\gamma}\right) \left(1 + O\left(\frac{1}{\sqrt{n}}\right)\right). \tag{46}$$

Proof: This result follows by re-scaling δ in Theorem 2, where it follows that for every $\alpha \geq 0$,

$$\mathbb{P}(|X_n - X_0| \geq \alpha n^\eta) \leq 2 \exp \left(-n D \left(\frac{\delta_n + \gamma}{1 + \gamma} \parallel \frac{\gamma}{1 + \gamma} \right) \right) \quad (47)$$

where from (3)

$$\delta_n \triangleq \frac{\alpha n^{-(1-\eta)}}{d} = \frac{\delta}{n^{1-\eta}}. \quad (48)$$

From the equality

$$(1 + u) \ln(1 + u) = u + \sum_{k=2}^{\infty} \frac{(-u)^k}{k(k-1)}, \quad -1 < u \leq 1$$

it follows from (7) that for every $n > \left(\frac{\delta}{\gamma}\right)^{\frac{1}{1-\eta}}$

$$nD \left(\frac{\delta_n + \gamma}{1 + \gamma} \parallel \frac{\gamma}{1 + \gamma} \right) = \left(\frac{\delta^2}{2\gamma} \right) n^{2\eta-1} - \left(\frac{\delta^3(1-\gamma)}{6\gamma^2} \right) n^{3\eta-2} + \dots$$

Substituting this into the exponent on the right-hand side of (47) gives that, for every $\alpha \geq 0$, inequality (45) holds. Note that $\eta = \frac{1}{2}$ refers to small deviations, and it gives the CLT-type inequality in (46). ■

The moderate deviations principle provides the exact asymptotic exponent for martingales with bounded jumps (see [7] and [20]). From Proposition 1, it follows that Theorem 2 gives the correct asymptotic exponential decay for small and moderate deviations (it is in contrast to the Azuma-Hoeffding inequality where a factor of $\frac{1}{\gamma}$ is missing in the exponent). Therefore, one cannot achieve an improvement for conditionally symmetric martingales via Theorem 1 when small or moderate deviations are considered, but only for large deviations (scaling linearly with n). In the large deviations regime, one indeed obtains an improvement in the exponent of Theorem 1 as compared to Theorem 2.

Remark 5: Consider the optimization problem

$$\text{maximize } \int_{\mathbb{R}} \exp(\lambda x) p_X(x) dx$$

s.t.

$$\int_{\mathbb{R}} p_X(x) dx = 1$$

$$\int_{\mathbb{R}} x p_X(x) dx = 0$$

$$\int_{\mathbb{R}} x^2 p_X(x) dx \leq \gamma d^2$$

$$p_X(x) \geq 0 \quad \forall x \in \mathbb{R}$$

$$p_X(x) = 0 \quad \forall |x| > d$$

$$p_X(x) = p_X(-x) \quad \forall x \in \mathbb{R},$$

where $\lambda \in \mathbb{R}$, $d \in \mathbb{R}_+$, $\gamma \in [0, 1]$, and the optimization is carried over the *pdf* p_X . This is a linear programming (LP) problem whose solution, according to Corollary 3, is given by the symmetric distribution

$$p_X(x) = (1 - \gamma)\delta_0(x) + \frac{\gamma}{2} \delta_d(x) + \frac{\gamma}{2} \delta_{-d}(x) \quad (49)$$

where $\delta_a(x) \triangleq \delta(x - a)$, for $a \in \mathbb{R}$, denotes the shifted Dirac's delta function. This refers to a discrete random variable X that gets the three values $0, d, -d$ with probabilities $1 - \gamma, \frac{\gamma}{2}, \frac{\gamma}{2}$ respectively. The corresponding solution of this optimization problem is given by the right-hand side of (23). Note that the removal of the last (symmetry) constraint in the above optimization problem gives, according to Bennett's inequality, the solution

$$p_X(x) = \frac{\gamma}{1 + \gamma} \delta_d(x) + \frac{1}{1 + \gamma} \delta_{-\gamma d}(x) \quad (50)$$

with the corresponding maximal value

$$\int_{\mathbb{R}} \exp(\lambda x) p_X(x) dx = \frac{\gamma \exp(\lambda d) + \exp(-\lambda \gamma d)}{1 + \gamma}.$$

The difference between (49) and (50), where the *pdf* that achieves the maximum in the latter case is symmetric if and only if $\gamma = 1$, is the reason for the improved exponent in Theorem 1 as compared to Theorem 2 when $\gamma \in (0, 1)$. This explains the source of the improvement in Theorem 1 for discrete-time real-valued conditionally symmetric martingales with bounded jumps, as compared to Theorem 2 which applies to the extended class of martingales that are not necessarily conditionally symmetric. The enhanced improvement of the exponent in Theorem 1 over the exponent in Theorem 2, as the value of $\gamma \in (0, 1)$ gets smaller (which is also evidenced by a comparison of the limits in (42) and (43)), is consistent with the enhanced asymmetry of the maximizing *pdf* in (50) as compared to the symmetric *pdf* in (49) for all values of γ .

Remark 6: The infimum in Theorem 3 is attained and thus is a minimum. This follows from [21, Remark 13] and because Theorem 3 is a particularization of [21, Theorem 4] for conditionally symmetric martingales with bounded jumps.

Remark 7: The function B in (14) is monotonic decreasing over the interval $(0, \infty)$, and it is bounded between 0 and 1. Also, $\lim_{u \rightarrow 0^+} B(u) = 1$, and $\lim_{u \rightarrow \infty} B(u) = 0$. Similarly to the function B in (14), also the function C in (12) is monotonic decreasing over the interval $(0, \infty)$, and it is bounded between 0 and 1. Calculus shows that

$$\begin{aligned} \lim_{u \rightarrow 0^+} C(u) &= 1, & \lim_{u \rightarrow \infty} C(u) &= 0, \\ \lim_{u \rightarrow 0^+} \frac{C(u)}{B(u)} &= \lim_{u \rightarrow \infty} \frac{C(u)}{B(u)} = 1. \end{aligned}$$

However, Theorem 4 improves the exponent in Theorem 5 due to the additional requirement of the conditional symmetry of the martingale in the former theorem. Although clear from the combined proof of these theorems, one can verify analytically that $C(u) > B(u)$ for every $u > 0$.

Remark 8: An immediate implication of Theorem 5 is given in the following:

Corollary 4: Let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a discrete-time real-valued martingale, and assume that $|X_k - X_{k-1}| \leq d$ a.s. for every $k \in \mathbb{N}$ and for a fixed $d > 0$. Let $\{Q_n\}_{n \in \mathbb{N}_0}$ be the quadratic variation process of the martingale, as is defined in (10). Then, for every $z, r > 0$,

$$\mathbb{P} \left(\max_{1 \leq k \leq n} |X_k - X_0| \geq z, Q_n \leq r \text{ for some } n \in \mathbb{N} \right) \leq 2 \exp \left(-\frac{z^2}{2r} \cdot B \left(\frac{zd}{r} \right) \right) \tag{51}$$

where B is introduced in (14).

A similar result follows from Theorem 4 under the conditional symmetry assumption of the martingale (where the function B in (14) is replaced with the function C in (12)).

B. Discussion of Some Related Known Inequalities

Remark 9: Under the conditions in Theorem 5, the inequality in [17, Theorem 1.2A] is equivalent to the following one:

$$\mathbb{P} \left(\max_{1 \leq k \leq n} (X_k - X_0) \geq z, Q_n \leq r \text{ for some } n \in \mathbb{N} \right) \leq \exp \left(-\frac{z^2}{2r} \cdot \tilde{B} \left(\frac{zd}{r} \right) \right) \tag{52}$$

where

$$\tilde{B}(u) \triangleq \frac{2}{1 + u + \sqrt{1 + 2u}}, \quad \forall u > 0. \tag{53}$$

It is noted that on the left-hand side of inequality (52), the first event is $\max_{1 \leq k \leq n} (X_k - X_0) \geq r$ instead of the loosened condition $X_n - X_0 \geq r$ (as it appears in [17, Eq. (1.2)]) due to the equality in (41). After introducing inequality (52), it was noted in [17, p. 538] that this inequality should be compared to Freedman’s bound in Theorem 5 but this comparison was not available. For such a comparison, it is easy to verify that

$$\frac{1}{1 + u} < \tilde{B}(u) < B(u) < 1, \quad \forall u > 0.$$

This inequality shows that, under the condition of Theorem 5, the exponential bound in [17, Theorem 1.2A] (see (52)) is looser than Freedman's bound in Theorem 5. It is noted however that [17, Theorem 1.2A] holds even under a milder condition where, for a fixed $d > 0$ and for every $j \in \mathbb{N}$ and $k > 2$,

$$\mathbb{E}[|X_j - X_{j-1}|^k | \mathcal{F}_{j-1}] \leq \frac{k!}{2} \cdot \mathbb{E}[(X_j - X_{j-1})^2 | \mathcal{F}_{j-1}] d^{k-2}.$$

This condition is automatically satisfied if $|X_j - X_{j-1}| \leq d$ for every $j \in \mathbb{N}$, which is indeed the condition of bounded jumps (a.s.) for the martingale in Theorem 5. For discrete-time real-valued martingales with bounded jumps (a.s.), we therefore revisited in this subsection only the proof of Theorem 5 in order to improve it in the next subsection for conditionally symmetric martingales (see Theorem 4).

Remark 10: Corollary 4 was generalized in [9, Theorem 3.3]. Applied to a discrete-time and real-valued martingale $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ with $\xi_n \triangleq X_n - X_{n-1}$ for every $n \in \mathbb{N}$, this generalization states that if, for some $a > 0$, the second-order process is defined by

$$H_n^{(a)} \triangleq \sum_{k=1}^n \xi_k^2 I_{\{|\xi_k| > a\}} + \sum_{k=1}^n \mathbb{E}[\xi_k^2 | \mathcal{F}_{k-1}] \quad (54)$$

then, at each finite stopping time τ ,

$$\mathbb{P}\left(\max_{n \leq \tau} |X_n - X_0| \geq z, H_\tau^{(a)} \leq r\right) \leq 2 \exp\left(-\frac{z^2}{2r} \cdot B\left(\frac{za}{r}\right)\right)$$

where B is defined in (14). Note that in the setting considered in Theorem 5 and Corollary 4 where the martingale has bounded jumps a.s. and $|X_k - X_{k-1}| \leq d$ a.s. for some fixed $d > 0$ and every $k \in \mathbb{N}$, then choosing $a = d$ in the above inequality gives the result in Corollary 4 (note that, in this case, the equality $H_n^{(a)} = Q_n$ holds for every $n \in \mathbb{N}_0$ where $\{Q_n\}_{n \in \mathbb{N}_0}$ is the predictable quadratic variation in (10)).

Remark 11: In the setting of Theorem 5, the bound in [17, Theorem 1.5A] is looser. The reason is that, under the condition in Theorem 5, it follows from [17, Theorem 1.5A] and the equality in (41) that for every $z, r > 0$

$$\mathbb{P}\left(\max_{1 \leq k \leq n} (X_k - X_0) \geq z, Q_n \leq r \text{ for some } n \in \mathbb{N}\right) \leq \exp\left(-\frac{z^2}{2r} \cdot B^*\left(\frac{zd}{r}\right)\right)$$

where $B^*(u) \triangleq \frac{1}{u} \sinh^{-1}\left(\frac{u}{2}\right)$ for every $u > 0$. The claim now follows by verifying that $B(u) > B^*(u)$ for every $u > 0$ (the function B is introduced in (14)).

Remark 12: In the following, we consider an inequality that was derived in [9, Section 4] and [17, Section 6], and emphasize its difference from the result in Theorem 4.

In [17, Theorem 6.1], it was stated that if $\{\xi_i\}$ is a sequence adapted to the filtration $\{\mathcal{F}_i\}$, and these random variables are conditionally symmetric, then for all $z, r > 0$

$$\mathbb{P}\left(\sum_{i=1}^n \xi_i \geq z, \sum_{i=1}^n \xi_i^2 \leq r \text{ for some } n \in \mathbb{N}\right) \leq \exp\left(-\frac{z^2}{2r}\right).$$

Its proof is similar in concept to the proof of Theorem 4 with the crucial difference that the two proofs rely on different supermartingales for the derivation of the respective inequalities. More explicitly, the main task in the proof of Theorem 4 was to show that $\{U_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a supermartingale (for a suitable choice of the free parameters λ and θ). The supermartingale that was required for the proof of [17, Theorem 6.1] is provided in [17, Lemma 6.1]. It is noted that the proof of [17, Lemma 6.1] is less involved (since it only required the use of a simple equality that follows from the conditional symmetry of the sequence $\{\xi_i\}$, and the use of the inequality $\cosh(x) \leq \exp(\frac{x^2}{2})$ that holds for every $x \in \mathbb{R}$). As a consequence of [17, Theorem 6.1], the following result holds:

Corollary 5: Let $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ be a discrete-time and real-valued conditionally symmetric martingale. Then, for every $r, z > 0$,

$$\mathbb{P}\left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} |X_k - X_0| \geq z, \sum_{i=1}^n (X_i - X_{i-1})^2 \leq r\right) \leq 2 \exp\left(-\frac{z^2}{2r}\right). \quad (55)$$

Proof: Let $\xi_i \triangleq X_i - X_{i-1}$ for all $i \in \mathbb{N}$, and $\xi_0 \triangleq 0$. Note that $\{\xi_i\}$ is a sequence of conditionally symmetric random variables, and ξ_i is \mathcal{F}_i -measurable (i.e., $\{\xi_i\}$ is adapted to the filtration $\{\mathcal{F}_i\}$). Since $\sum_{i=1}^n \xi_i = X_n - X_0$ then, from [17, Theorem 6.1], for every $r, z > 0$,

$$\mathbb{P} \left(\exists n \in \mathbb{N} : X_n - X_0 \geq z, \sum_{i=1}^n (X_i - X_{i-1})^2 \leq r \right) \leq \exp \left(-\frac{z^2}{2r} \right).$$

Since $Y_n \triangleq \sum_{i=1}^n (X_i - X_{i-1})^2$ is monotonic non-decreasing in n then, similarly to the reasoning of the equality in (41), one can replace the first event $\{X_n - X_0 \geq z\}$ with $\{\max_{1 \leq k \leq n} (X_k - X_0) \geq z\}$. Hence, for every $r, z > 0$,

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} (X_k - X_0) \geq z, \sum_{i=1}^n (X_i - X_{i-1})^2 \leq r \right) \leq \exp \left(-\frac{z^2}{2r} \right).$$

By referring to the martingale $\{-X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ (note that the predictable quadratic variation in (10) is not affected by altering the sign of $\{X_n\}$), then also

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \min_{1 \leq k \leq n} (X_k - X_0) \leq -z, \sum_{i=1}^n (X_i - X_{i-1})^2 \leq r \right) \leq \exp \left(-\frac{z^2}{2r} \right).$$

The proof of the corollary is completed by combining the last two inequalities and using the union bound. \blacksquare

The result in Corollary 5 was further generalized in [9, Theorem 4.1] for square-integrable, conditionally symmetric local martingales.

Note that Theorem 4 and Corollary 4 provide an upper bound on the probability

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} |X_k - X_0| \geq z, \sum_{k=1}^n \mathbb{E}[(X_k - X_{k-1})^2 | \mathcal{F}_{k-1}] \leq r \right) \quad (56)$$

whereas [9, Theorem 4.1], [17, Theorem 6.1], and Corollary 5 provide a bound on the probability

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} |X_k - X_0| \geq z, \sum_{k=1}^n (X_k - X_{k-1})^2 \leq r \right) \quad (57)$$

so the second events in (56) and (57) refer to the predictable and optional quadratic variations of the martingale, respectively, which explains the difference between these two bounds.

IV. AN EXAMPLE IN THE CONTEXT OF GAMBLING

Let $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ be a conditionally symmetric supermartingale. Consider a game where a gambler is allowed to place a wager of size W_n before the n -th toss where the size of the wager may depend on the past elements of the sequence X_0, \dots, X_{n-1} , but not on X_n or the future elements of this sequence. Lets assume that there exists some fixed $W > 0$ such that $0 \leq W_k \leq W$ for every $k \in \mathbb{N}$ (i.e., the size of the wager cannot exceed a fixed amount W at any toss). Hence, $W_k \in L^\infty(\Omega, \mathcal{F}_{k-1}, \mathbb{P})$ for every $k \in \mathbb{N}$. If the n -th toss satisfies $X_n \geq X_{n-1}$, the gambler nets $W_n(X_n - X_{n-1})$, and if $X_n \leq X_{n-1}$ then the gambler losses $-W_n(X_n - X_{n-1})$. The net winnings S_n after n tosses is therefore equal to $S_n = \sum_{k=1}^n W_k(X_k - X_{k-1})$ for every $n \in \mathbb{N}$ with $S_0 = 0$. According to Example 4, $\{S_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a conditionally symmetric supermartingale. Note that since $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ is by assumption a supermartingale then this game is sub-fair from the point of view of the gambler, unless this process is a martingale. Lets assume that there exists a fixed $c > 0$ such that a.s.

$$X_k - \mathbb{E}[X_k | \mathcal{F}_{k-1}] \leq c, \quad \forall k \in \mathbb{N}. \quad (58)$$

Since W_k, S_{k-1} and X_{k-1} are \mathcal{F}_{k-1} -measurable then a.s.

$$\begin{aligned} & S_k - \mathbb{E}[S_k | \mathcal{F}_{k-1}] \\ &= W_k(X_k - \mathbb{E}[X_k | \mathcal{F}_{k-1}]) \\ &\leq cW \triangleq d. \end{aligned} \quad (59)$$

The predictable quadratic variation process $\{Q_n\}_{n \in \mathbb{N}_0}$ that is associated with the supermartingale $\{S_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is

$$\begin{aligned} Q_n &= \sum_{k=1}^n \text{Var}(S_k | \mathcal{F}_{k-1}) \\ &= \sum_{k=1}^n W_k^2 \text{Var}(X_k | \mathcal{F}_{k-1}) \end{aligned} \quad (60)$$

with $Q_0 = 0$. Lets assume that a.s. $\text{Var}(X_k | \mathcal{F}_{k-1}) \leq \sigma_k^2$ for every $k \in \mathbb{N}$, then a.s.

$$Q_n \leq \sum_{k=1}^n \sigma_k^2 W_k^2, \quad \forall n \in \mathbb{N}. \quad (61)$$

Based on Corollary 2 and (59)–(61), it follows that due to the conditional symmetry of the supermartingale

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} S_k \geq z, \sum_{k=1}^n \sigma_k^2 W_k^2 \leq r \right) \leq \exp \left(-\frac{z^2}{2r} \cdot C \left(\frac{dz}{r} \right) \right), \quad \forall r, z > 0 \quad (62)$$

where the function C is defined in (12). The interpretation of this inequality is as follows: Since $\{\sum_{k=1}^n \sigma_k^2 W_k^2\}_{n \in \mathbb{N}}$ is a non-negative and monotonic non-decreasing sequence in n , then the probability of the joint event in (62) refers to the event where the maximal net winnings up to an arbitrary time instance n is larger than or equal to a fixed amount $z > 0$, as long as the total wagers up to this time instance are not too large in the sense that

$$\sum_{k=1}^n \sigma_k^2 W_k^2 \leq r \quad (63)$$

for a fixed value $r > 0$. If the value of r is increased while keeping the value of z fixed then this means that we allow an enlargement of the observation time of the process of net winnings since the constraint in (63), which plays as a criterion for defining the stopping time, becomes less severe so it is expected to be violated at a later instance of time. Therefore, by enlarging the observation time of the process of net winnings, the event that there is an instance during this period of time where the maximal net winnings up to this time instance is larger than or equal to a fixed value $z > 0$ becomes more likely. Hence, the probability of the joint event is increased by increasing the value of r while keeping the value of z fixed. On the other hand, assume that the value of $r > 0$ is kept fixed while increasing the value of $z > 0$. The observation time of the process of net winnings is kept fixed in this case (since the constraint in (63) is not affected by a change of the value of z), but the event of achieving net winnings larger than or equal to z at some point during this observation period of time becomes less likely by increasing the value of z . Therefore, the probability of the joint event in (62) is decreased by increasing the value of z while keeping the value of r fixed. Note that the inequality in (62) holds under the assumption that the supermartingale $\{X_k, \mathcal{F}_k\}_{k \in \mathbb{N}_0}$ is conditionally symmetric. Without this assumption, Corollary 2 gives the loosened upper bound

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} S_k \geq z, \sum_{k=1}^n \sigma_k^2 W_k^2 \leq r \right) \leq \exp \left(-\frac{z^2}{2r} \cdot B \left(\frac{dz}{r} \right) \right), \quad \forall r, z > 0 \quad (64)$$

where the function B is defined in (14). These observations are exemplified in the following.

Example 6: As an example of the considered setting, let $p \geq 2$ be an arbitrary even number, and define a conditional symmetric martingale $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ by

$$X_k = X_{k-1} + U_k, \quad \forall k \in \mathbb{N}$$

where $\{U_k\}_{k \in \mathbb{N}}$ is a sequence of i.i.d. random variables getting the p values from the set $\{\pm 1, \dots, \pm(p-1)\}$ with equal probability ($\frac{1}{p}$). Hence, in this fair game (since $\{X_n, \mathcal{F}_n\}_{n \in \mathbb{N}_0}$ is a martingale), the net winnings after n tosses is given by $S_n = \sum_{k=1}^n W_k U_k$ for every $n \in \mathbb{N}$ with $S_0 = 0$. Since, by assumption, W_k may only depend on X_0, \dots, X_{k-1} but not on the elements X_k, X_{k+1}, \dots then W_k is \mathcal{F}_{k-1} -measurable. For every $k \in \mathbb{N}$,

$$X_k - \mathbb{E}[X_k | \mathcal{F}_{k-1}] = X_k - X_{k-1} = U_k \leq p - 1$$

so, from (58) and (59), set $c = p - 1$ and $d = (p - 1)W$. Furthermore, from (60), the predictable quadratic variation of the net winnings is equal to

$$\begin{aligned} Q_n &= \sum_{k=1}^n W_k^2 \text{Var}(X_k | \mathcal{F}_{k-1}) \\ &= \sum_{k=1}^n W_k^2 \text{Var}(U_k) \\ &= \left(\frac{p^2 - 1}{3}\right) \sum_{k=1}^n W_k^2 \end{aligned} \tag{65}$$

so set $\sigma_k^2 \triangleq \frac{p^2 - 1}{3}$ in (61) for every $k \in \mathbb{N}$. Referring to (62) and (64), lets define new parameters $x, y > 0$ such that

$$z \triangleq Wx, \quad r \triangleq \left(\frac{p^2 - 1}{3}\right) W^2 y \tag{66}$$

then the bounds in (62) and (64) yield that

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} S_k \geq Wx, \sum_{i=1}^n W_i^2 \leq W^2 y \right) \leq \exp \left(-\frac{3}{2(p^2 - 1)} \frac{x^2}{y} \cdot C \left(\frac{3}{p + 1} \frac{x}{y} \right) \right) \tag{67}$$

and

$$\mathbb{P} \left(\exists n \in \mathbb{N} : \max_{1 \leq k \leq n} S_k \geq Wx, \sum_{i=1}^n W_i^2 \leq W^2 y \right) \leq \exp \left(-\frac{3}{2(p^2 - 1)} \frac{x^2}{y} \cdot B \left(\frac{3}{p + 1} \frac{x}{y} \right) \right). \tag{68}$$




Fig. 1. This figure refers to Example 6 with $p = 6$. The left plot shows the upper bounds on the probability of the joint event in (67) (solid line) and (68) (dotted line), which follow respectively from Theorems 4 and 5 (the improvement in Theorem 4 is due to the conditional symmetry of the martingale that is associated with the net winnings). The right plot shows the improved bound in (67) as a function of the parameter x , for some fixed values of the other parameter y .

Fig. 1 presents the bounds in (68) and (67) with $p = 6$. The left plot in this figure shows the upper bounds on the probability of the joint event in (67) (solid line) and (68) (dotted line), which follow respectively from Theorems 4 and 5. This improvement, as shown in the solid line, is obtained due to the conditional symmetry of the martingale that refers to the net winnings (as was discussed earlier). The right plot of Fig. 1 shows the improved bound in (67) as a function of the parameter x , for some fixed values of the parameter y . Due to the linear dependence in (66) between x and z and also between y and r , and in light of the interpretation that was provided before this example, the probability of the joint event in (67) and (68) is

- Monotonic increasing if the parameter $x > 0$ is kept fixed while increasing the value of $y > 0$,
- Monotonic decreasing if the parameter $x > 0$ is increased while the parameter $y > 0$ is kept fixed.

This is indeed reflected from the exponential upper bound in (67), as is shown in the right plot of Fig. 1.

Theorem 1 provides an upper bound on the probability $\mathbb{P}(S_n \geq \alpha n)$ for an arbitrary value of $\alpha \geq 0$ (clearly, if $\alpha > (p-1)W \triangleq d$ then this probability is zero). The exponent of the bound in Theorem 1 is equal to $E(\gamma, \delta)$, as is introduced in (4), where the notation in (3) gives that

$$\gamma = \frac{\sigma^2}{d^2} = \frac{p+1}{3(p-1)}, \quad \delta = \frac{\alpha}{(p-1)W}. \quad (69)$$

Note that γ approaches $\frac{1}{3}$ as $p \gg 1$. For small values of γ , there is a significant advantage of the exponent that follows from Theorem 1 as compared to Theorem 2 (as is also evidenced from the comparison of the limits in (42) and (43), differing especially for values of γ close to zero). This is already the case for $p = 4$ (where, from (69), $\gamma = \frac{5}{9}$). Note that in the case where $p = 2$, which corresponds to the setting of the Saint-Petersburg game, one gets $\gamma = 1$ in (69); therefore, in the latter case, the resulting exponent is given by the function f in (44). One can also apply Theorem 3 to this example. In this case, it follows from (9) that for every $l \in \mathbb{N}$

$$\gamma_{2l} = \frac{2}{p} \sum_{j=1}^{\frac{p}{2}} \left(\frac{2j-1}{p-1} \right)^{2l}$$

and $\gamma_{2l-1} = 0$. Fig. 2 compares, for $p = 6$, the exponents which follow from Theorems 1–3 as a function of




Fig. 2. The figure refers to Example 6 with $p = 6$. It compares the exponents in Theorems 1–3 as a function of $\delta \in [0, 1]$. The left plot shows the exponent in Theorem 3, for various even values of $m \geq 2$. The improvement in the exponent of Theorem 3, which follows by taking into consideration the conditional moments of $S_k - S_{k-1}$ (given \mathcal{F}_{k-1}) up to order m , is shown in the left plot for $m = 2, 4, \dots, 16$ ($m = 2, 4, 6$ refer to the dotted, dashed and dashed-dot lines, respectively; $m = 8, \dots, 16$ refer to the solid lines). The right plot compares the exponent of the Azuma-Hoeffding inequality (dotted line), the exponent in Theorem 2 (dot-dashed line), the improvement for conditionally symmetric martingales in Theorem 1 (dashed line), and the exponents in Theorem 3 (the two solid lines) while taking into consideration either the conditional variance (in the lower solid line) or all the conditional moments of $S_k - S_{k-1}$ (practically, up to order $m = 16$, due to the fast convergence in terms of m as is shown in the left plot of this figure).

$\delta \in [0, 1]$. The left plot of Fig. 2 shows the exponent of Theorem 3, applied to the considered example with $p = 6$ for various even values of $m \geq 2$. The improvement in the exponent of Theorem 3, which follows by taking into consideration the orders of the conditional moments for $S_k - S_{k-1}$ (given \mathcal{F}_{k-1}) up to order m , is shown in the left plot of Fig. 2 for $m = 2, 4, \dots, 16$ ($m = 2$ refers to the dotted line, $m = 4$ to the dashed line, $m = 6$ refers to the dashed-dot line, and $m = 8, \dots, 16$ refers to the solid lines, hence showing the fast convergence in terms of m ; as is shown in the left plot of Fig. 2, the improvement in the exponent that is obtained by taking into consideration the conditional moments of order higher than 10 is marginal). The right plot of Fig. 2 compares the exponents of the Azuma-Hoeffding inequality (dotted line) and Theorem 2 (dot-dashed line), the refined exponent for conditionally symmetric martingales in Theorem 1 (dashed line), and the exponents in Theorem 3 (the two solid lines) while taking into consideration only the conditional variance or alternatively all the conditional moments (practically, up to order 16, as the convergence of these exponents is fast due to the left plot of this figure). Note that according to

[21, Proposition 2], the exponent in Theorem 3 for $m = 2$ is looser than the exponent in Theorem 2; hence, it is also looser than the exponent in Theorem 1. This is indeed shown in the right plot of Fig. 2. However, the exponent which relies on all the conditional moments for this specific example is better than the exponent in Theorem 1 for conditionally symmetric martingales (where the latter exponent only relies on the conditional variance of $\{S_k\}$).

ACKNOWLEDGMENT

I thank Robert Adler and Ofer Zeitouni for their help in improving the presentation of the results, and Ofer Zeitouni for a suggestion that led to a weakening of assumptions in Lemma 1.

REFERENCES

- [1] N. Alon and J. H. Spencer, *The Probabilistic Method*, Wiley Series in Discrete Mathematics and Optimization, Third Edition, 2008.
- [2] K. Azuma, “Weighted sums of certain dependent random variables,” *Tohoku Mathematical Journal*, vol. 19, pp. 357–367, 1967.
- [3] G. Bennett, “Probability inequalities for the sum of independent random variables,” *Journal of the American Statistical Association*, vol. 57, no. 297, pp. 33–45, March 1962.
- [4] D. L. Burkholder, “Explorations in martingale theory and its applications,” *Ecole d’Eté de Probabilités de Saint-Flour XIX–1989, Lecture Notes in Mathematics*, Springer-Verlag, vol. 1464, pp. 1–66, 1991.
- [5] F. Chung and L. Lu, *Complex Graphs and Networks, Regional Conference Series in Mathematics*, vol. 107, 2006. Chapter 2 entitled “Old and new concentration inequalities” is available at: <http://www.math.ucsd.edu/~fan/complex/ch2.pdf>.
- [6] F. Chung and L. Lu, “Concentration inequalities and martingale inequalities: a survey,” *Internet Mathematics*, vol. 3, no. 1, pp. 79–127, March 2006.
- [7] A. Dembo, “Moderate deviations for martingales with bounded jumps,” *Electronic Communications in Probability*, vol. 1, no. 3, pp. 11–17, March 1996.
- [8] A. Dembo and O. Zeitouni, *Large Deviations Techniques and Applications*, Springer, second edition, 1997.
- [9] K. Dzhaparidze and J. H. van Zanten, “On Bernstein-type inequalities for martingales,” *Stochastic Processes and their Applications*, vol. 93, no. 1, pp. 109–117, May 2001.
- [10] D. Freedman, “On tail probabilities for martingales,” *Annals of Probability*, vol. 3, no. 1, pp. 100–118, January 1975.
- [11] I. Grama and E. Haeusler, “Large deviations for martingales via Cramer’s method,” *Stochastic Processes and their Applications*, vol. 85, no. 2, pp. 279–293, February 2000.
- [12] W. Hoeffding, “Probability inequalities for sums of bounded random variables,” *Journal of the American Statistical Association*, vol. 58, no. 301, pp. 13–30, March 1963.
- [13] C. McDiarmid, “On the method of bounded differences,” *Surveys in Combinatorics*, vol. 141, pp. 148–188, Cambridge University Press, Cambridge, 1989.
- [14] C. McDiarmid, “Concentration,” *Probabilistic Methods for Algorithmic Discrete Mathematics*, pp. 195–248, Springer, 1998.
- [15] A. Osekowski, “Weak type inequalities for conditionally symmetric martingales,” *Statistics and Probability Letters*, vol. 80, no. 23–24, pp. 2009–2013, December 2010.
- [16] A. Oskowski, “Sharp ratio inequalities for a conditionally symmetric martingale,” *Bulletin of the Polish Academy of Sciences Mathematics*, vol. 58, no. 1, pp. 65–77, 2010.
- [17] V. H. de la Pena, “A general class of exponential inequalities for martingales and ratios,” *Annals of Probability*, vol. 27, no. 1, pp. 537–564, January 1999.
- [18] V. H. de la Pena, M. J. Klass and T. L. Lai, “Self-normalized processes: exponential inequalities, moment bounds and iterated logarithm laws,” *Annals of Probability*, vol. 32, no. 3, pp. 1902–1933, July 2004.
- [19] I. Pinelis, “Optimum bounds for the distributions of martingales in Banach spaces,” *Annals of Probability*, vol. 22, no. 4, pp. 1679–1706, October 1994.
- [20] A. Rackauskas, “Large deviations for martingales,” *Acta Applicandae Mathematicae*, vol. 38, no. 1, pp. 109–129, January 1995.
- [21] I. Sason, “On refined versions of the Azuma-Hoeffding inequality with applications in information theory,” submitted to the *IEEE Trans. on Information Theory*, July 2011. [Online]. Available at <http://arxiv.org/abs/1111.1977> (version 3).
- [22] W. L. Steiger, “A best possible Kolmogoroff-type inequality for martingales and a characteristic property,” *Annals of Mathematical Statistics*, vol. 40, no. 3, pp. 764–769, June 1969.
- [23] G. Wang, “Sharp maximal inequalities for conditionally symmetric martingales and Brownian motion,” *Proceedings of the American Mathematical Society*, vol. 112, no. 2, pp. 579–586, June 1991.