

*Bernoulli* **19**(5A), 2013, 2000–2009  
DOI: 10.3150/12-BEJ438

# Integrability properties and limit theorems for the exit time from a cone of planar Brownian motion

STAVROS VAKEROUDIS<sup>1,2,\*</sup> and MARC YOR<sup>1,3,\*\*</sup>

<sup>1</sup>*Laboratoire de Probabilités et Modèles Aléatoires (LPMA), CNRS: UMR7599, Université Pierre et Marie Curie – Paris VI, Université Paris-Diderot Paris VII, 4, Place Jussieu, 75252 Paris Cedex 05, France. E-mail: \*stavros.vakeroudis@upmc.fr; url: <http://svakeroudis.wordpress.com>; \*\*yormarc@aol.com*

<sup>2</sup>*Probability and Statistics Group, School of Mathematics, University of Manchester, Alan Turing Building, Oxford Road, Manchester M13 9PL, United Kingdom*

<sup>3</sup>*Institut Universitaire de France, Paris, France*

We obtain some integrability properties and some limit theorems for the exit time from a cone of a planar Brownian motion, and we check that our computations are correct via Bougerol's identity.

*Keywords:* Bougerol's identity; exit time from a cone; planar Brownian motion; skew-product representation

## 1. Introduction

We consider a standard planar Brownian motion  $(Z_t = X_t + iY_t, t \geq 0)$ , starting from  $x_0 + i0, x_0 > 0$ , where  $(X_t, t \geq 0)$  and  $(Y_t, t \geq 0)$  are two independent linear Brownian motions, starting respectively, from  $x_0$  and 0 (when we simply write: Brownian motion, we always mean real-valued Brownian motion, starting from 0; for 2-dimensional Brownian motion, we indicate planar or complex BM).

As is well known Itô and McKean [10], since  $x_0 \neq 0$ ,  $(Z_t, t \geq 0)$  does not visit a.s. the point 0 but keeps winding around 0 infinitely often. In particular, the continuous winding process  $\theta_t = \text{Im}(\int_0^t \frac{dZ_s}{Z_s})$ ,  $t \geq 0$  is well defined. A scaling argument shows that we may assume  $x_0 = 1$ , without loss of generality, since, with obvious notation:

$$(Z_t^{(x_0)}, t \geq 0) \stackrel{(law)}{=} (x_0 Z_{(t/x_0^2)}^{(1)}, t \geq 0). \quad (1)$$

Thus, from now on, we shall take  $x_0 = 1$ .

This is an electronic reprint of the original article published by the ISI/BS in *Bernoulli*, 2013, Vol. 19, No. 5A, 2000–2009. This reprint differs from the original in pagination and typographic detail.

Furthermore, there is the skew product representation:

$$\log |Z_t| + i\theta_t \equiv \int_0^t \frac{dZ_s}{Z_s} = (\beta_u + i\gamma_u) \Big|_{u=H_t=\int_0^t \frac{ds}{|Z_s|^2}}, \quad (2)$$

where  $(\beta_u + i\gamma_u, u \geq 0)$  is another planar Brownian motion starting from  $\log 1 + i0 = 0$ . Thus, the Bessel clock  $H$  plays a key role in many aspects of the study of the winding number process  $(\theta_t, t \geq 0)$  (see, e.g., Yor [21]).

Rewriting (2) as:

$$\log |Z_t| = \beta_{H_t}; \quad \theta_t = \gamma_{H_t}, \quad (3)$$

we easily obtain that the two  $\sigma$ -fields  $\sigma\{|Z_t|, t \geq 0\}$  and  $\sigma\{\beta_u, u \geq 0\}$  are identical, whereas  $(\gamma_u, u \geq 0)$  is independent from  $(|Z_t|, t \geq 0)$ .

We shall also use Bougerol's celebrated identity in law, see, for example, Bougerol [5], Alili, Dufresne and Yor [1] and Yor [24] (page 200), which may be written as:

$$\text{for fixed } t \quad \sinh(\beta_t) \stackrel{(law)}{=} \hat{\beta}_{A_t(\beta)}, \quad (4)$$

where  $(\beta_u, u \geq 0)$  is 1-dimensional BM,  $A_u(\beta) = \int_0^u ds \exp(2\beta_s)$  and  $(\hat{\beta}_v, v \geq 0)$  is another BM, independent of  $(\beta_u, u \geq 0)$ . For the random times  $T_c^{|\theta|} \equiv \inf\{t: |\theta_t| = c\}$ , and  $T_c^{|\gamma|} \equiv \inf\{t: |\gamma_t| = c\}$ , ( $c > 0$ ) by using the skew-product representation (3) of planar Brownian motion Revuz and Yor [15], we obtain:

$$T_c^{|\theta|} = A_{T_c^{|\gamma|}}(\beta) \equiv \int_0^{T_c^{|\gamma|}} ds \exp(2\beta_s) = H_u^{-1} \Big|_{u=T_c^{|\gamma|}}. \quad (5)$$

Moreover, it has been recently shown that, Bougerol's identity applied with the random time  $T_c^{|\theta|}$  instead of  $t$  in (4) yields the following Vakeroudis [18].

**Proposition 1.1.** *The distribution of  $T_c^{|\theta|}$  is characterized by its Gauss-Laplace transform:*

$$E \left[ \sqrt{\frac{2c^2}{\pi T_c^{|\theta|}}} \exp\left(-\frac{x}{2T_c^{|\theta|}}\right) \right] = \frac{1}{\sqrt{1+x}} \varphi_m(x) \quad (6)$$

for every  $x \geq 0$ , with  $m = \frac{\pi}{2c}$ , and:

$$\varphi_m(x) = \frac{2}{(G_+(x))^m + (G_-(x))^m}, \quad G_{\pm}(x) = \sqrt{1+x} \pm \sqrt{x}. \quad (7)$$

The remainder of this article is organized as follows: in Section 2, we study some integrability properties for the exit times from a cone; more precisely, we obtain some new results concerning the negative moments of  $T_c^{|\theta|}$  and of  $T_c^\theta \equiv \inf\{t: \theta_t = c\}$ . In

Section 3, we state and prove some Limit theorems for these random times for  $c \rightarrow 0$  and for  $c \rightarrow \infty$  followed by several generalizations (for extensions of these works to more general planar processes, see, e.g., Doney and Vakeroudis [7]). We use these results in order to obtain (see Remark 3.4) a new simple non-computational proof of Spitzer's celebrated asymptotic theorem Spitzer [16], which states that:

$$\frac{2}{\log t} \theta_t \xrightarrow[t \rightarrow \infty]{(law)} C_1, \quad (8)$$

with  $C_1$  denoting a standard Cauchy variable (for other proofs, see, e.g., Williams [20], Durrett [9], Messulam and Yor [13], Bertoin and Werner [2], Yor [23], Vakeroudis [18]). Finally, in Section 4, we use the Gauss–Laplace transform (6) which is equivalent to Bougerol's identity (4) in order to check our results.

## 2. Integrability properties

Concerning the moments of  $T_c^{|\theta|}$ , we have the following (a more extended discussion is found in, e.g., Matsumoto and Yor [12]).

**Theorem 2.1.** *For every  $c > 0$ ,  $T_c^{|\theta|}$  enjoys the following integrability properties:*

- (i) for  $p > 0$ ,  $E[(T_c^{|\theta|})^p] < \infty$ , if and only if  $p < \frac{\pi}{4c}$ ,
- (ii) for any  $p < 0$ ,  $E[(T_c^{|\theta|})^p] < \infty$ .

**Corollary 2.2.** *For  $0 < c < d$ , the random times  $T_{-d,c}^\theta \equiv \inf\{t: \theta_t \notin (-d, c)\}$ ,  $T_c^{|\theta|}$  and  $T_c^\theta$  satisfy the inequality:*

$$T_c^\theta \geq T_{-d,c}^\theta \geq T_c^{|\theta|}. \quad (9)$$

Thus, their negative moments satisfy:

$$\text{for } p > 0 \quad E\left[\frac{1}{(T_c^\theta)^p}\right] \leq E\left[\frac{1}{(T_{-d,c}^\theta)^p}\right] \leq E\left[\frac{1}{(T_c^{|\theta|})^p}\right] < \infty. \quad (10)$$

### Proofs of Theorem 2.1 and of Corollary 2.2.

(i) The original proof is given by Spitzer [16], followed later by many authors Williams [20], Burkholder [6], Messulam and Yor [13], Durrett [9], Yor [22]. See also Revuz and Yor [15], Ex. 2.21, page 196.

(ii) In order to obtain this result, we might use the representation  $T_c^{|\theta|} = A_{T_c^{|\theta|}\gamma_1}$  together with a recurrence formula for the negative moments of  $A_t$  [8], Theorem 4.2, page 417 (in fact, Dufresne also considers  $A_t^{(\mu)} = \int_0^t ds \exp(2\beta_s + 2\mu s)$ , but we only need to take  $\mu = 0$  for our purpose, and we note  $A_t \equiv A_t^{(0)}$ ), [17]. However, we can also obtain this result by

simply remarking that the RHS of the Gauss–Laplace transform (6) in Proposition 1.1 is an infinitely differentiable function in 0 (see also [19]), thus:

$$E \left[ \frac{1}{(T_c^{|\theta|})^p} \right] < \infty \quad \text{for every } p > 0. \quad (11)$$

Now, Corollary 2.2 follows immediately from Theorem 2.1(ii).  $\square$

### 3. Limit theorems for $T_c^{|\theta|}$

#### 3.1. Limit theorems for $T_c^{|\theta|}$ , as $c \rightarrow 0$ and $c \rightarrow \infty$

The skew-product representation of planar Brownian motion allows to prove the three following asymptotic results for  $T_c^{|\theta|}$ .

**Proposition 3.1.**

(a) For  $c \rightarrow 0$ , we have:

$$\frac{1}{c^2} T_c^{|\theta|} \xrightarrow[c \rightarrow 0]{(law)} T_1^{|\gamma|}. \quad (12)$$

(b) For  $c \rightarrow \infty$ , we have:

$$\frac{1}{c} \log(T_c^{|\theta|}) \xrightarrow[c \rightarrow \infty]{(law)} 2|\beta|_{T_1^{|\gamma|}}. \quad (13)$$

(c) For  $\varepsilon \rightarrow 0$ , we have:

$$\frac{1}{\varepsilon^2} (T_{c+\varepsilon}^{|\theta|} - T_c^{|\theta|}) \xrightarrow[\varepsilon \rightarrow 0]{(law)} \exp(2\beta_{T_1^{|\gamma|}}) T_1^{\gamma'}, \quad (14)$$

where  $\gamma'$  stands for a real Brownian motion, independent from  $\gamma$ , and  $T_1^{\gamma'} = \inf\{t: \gamma'_t = 1\}$ .

**Proof.** We rely upon (5) for the three proofs. By using the scaling property of BM, we obtain:

$$T_c^{|\theta|} = A_{T_c^{|\gamma|}}(\beta) \stackrel{(law)}{=} A_u(\beta)|_{u=c^2 T_1^{|\gamma|}}$$

thus:

$$\frac{1}{c^2} T_c^{|\theta|} \stackrel{(law)}{=} \int_0^{T_1^{|\gamma|}} dv \exp(2c\beta_v). \quad (15)$$

(a) For  $c \rightarrow 0$ , the RHS of (15) converges to  $T_1^{|\gamma|}$ , thus we obtain part (a) of the proposition.

(b) For  $c \rightarrow \infty$ , taking logarithms on both sides of (15) and dividing by  $c$ , on the LHS we obtain  $\frac{1}{c} \log(T_c^{|\theta|}) - \frac{2}{c} \log c$  and on the RHS:

$$\frac{1}{c} \log \left( \int_0^{T_1^{|\gamma|}} dv \exp(2c\beta_v) \right) = \log \left( \int_0^{T_1^{|\gamma|}} dv \exp(2c\beta_v) \right)^{1/c},$$

which, from the classical Laplace argument:  $\|f\|_p \xrightarrow{p \rightarrow \infty} \|f\|_\infty$ , converges for  $c \rightarrow \infty$ , towards:

$$2 \sup_{v \leq T_1^{|\gamma|}} (\beta_v) \stackrel{(law)}{=} 2|\beta|_{T_1^{|\gamma|}}.$$

This proves part (b) of the proposition.

(c)

$$\begin{aligned} T_{c+\varepsilon}^{|\theta|} - T_c^{|\theta|} &= \int_{T_c^{|\gamma|}}^{T_{c+\varepsilon}^{|\gamma|}} du \exp(2\beta_u) \\ &= \int_0^{T_{c+\varepsilon}^{|\gamma|} - T_c^{|\gamma|}} dv \exp(2\beta_{T_c^{|\gamma|}}) \exp(2(\beta_{v+T_c^{|\gamma|}} - \beta_{T_c^{|\gamma|}})) \\ &= \exp(2\beta_{T_c^{|\gamma|}}) \int_0^{T_{c+\varepsilon}^{|\gamma|} - T_c^{|\gamma|}} dv \exp(2B_v), \end{aligned} \quad (16)$$

where  $(B_s \equiv \beta_{s+T_c^{|\gamma|}} - \beta_{T_c^{|\gamma|}}, s \geq 0)$  is a BM independent of  $T_c^{|\gamma|}$ .

We study now  $\tilde{T}_{c,c+\varepsilon}^{|\gamma|} \equiv T_{c+\varepsilon}^{|\gamma|} - T_c^{|\gamma|}$ , the first hitting time of the level  $c + \varepsilon$  from  $|\gamma|$ , starting from  $c$ . Thus, we define:  $\rho_u \equiv |\gamma_u|$ , starting also from  $c$ . Thus,  $\rho_u = c + \delta_u + L_u$ , where  $(\delta_s, s \geq 0)$  is a BM and  $(L_s, s \geq 0)$  is the local time of  $\rho$  at 0. Thus,

$$\begin{aligned} \tilde{T}_{c,c+\varepsilon}^{|\gamma|} &= \inf\{u \geq 0: \rho_u = c + \varepsilon\} \equiv \inf\{u \geq 0: \delta_u + L_u = \varepsilon\} \\ &\stackrel{u=\varepsilon^2 v}{=} \varepsilon^2 \inf\left\{v \geq 0: \frac{1}{\varepsilon} \delta_{v\varepsilon^2} + \frac{1}{\varepsilon} L_{v\varepsilon^2} = 1\right\}. \end{aligned} \quad (17)$$

From Skorokhod's lemma Revuz and Yor [15]:

$$L_u = \sup_{y \leq u} ((-c - \delta_y) \vee 0)$$

we deduce:

$$\frac{1}{\varepsilon} L_{v\varepsilon^2} = \sup_{y \leq v\varepsilon^2} ((-c - \delta_y) \vee 0) \stackrel{y=\varepsilon^2 \sigma}{=} \sup_{\sigma \leq v} \left( \left( -c - \varepsilon \frac{1}{\varepsilon} \delta_{\sigma\varepsilon^2} \right) \vee 0 \right) = 0. \quad (18)$$

Hence, with  $\gamma'$  denoting a new BM independent from  $\gamma$ , (16) writes:

$$T_{c+\varepsilon}^{|\theta|} - T_c^{|\theta|} = \exp(2\beta_{T_c^{|\gamma|}}) \int_0^{\varepsilon^2 T_1^{\gamma'}} dv \exp(2B_v). \quad (19)$$

Thus, dividing both sides of (19) by  $\varepsilon^2$  and making  $\varepsilon \rightarrow 0$ , we obtain part (c) of the proposition.  $\square$

**Remark 3.2.** The asymptotic result (c) in Proposition 3.1 may also be obtained in a straightforward manner from (16) by analytic computations. Indeed, using the Laplace transform of the first hitting time of a fixed level by the absolute value of a linear Brownian motion  $E[e^{-(\lambda^2/2)T_b^{|\gamma|}}] = \frac{1}{\cosh(\lambda b)}$  (see, e.g., Proposition 3.7, page 71 in Revuz and Yor [15]), we have that for  $0 < c < b$ , and  $\lambda \geq 0$ :

$$E[e^{-(\lambda^2/2)(T_b^{|\gamma|} - T_c^{|\gamma|})}] = \frac{\cosh(\lambda c)}{\cosh(\lambda b)}. \quad (20)$$

Using now  $b = c + \varepsilon$ , for every  $\varepsilon > 0$ , the latter equals:

$$\frac{\cosh(\lambda c/\varepsilon)}{\cosh((\lambda/\varepsilon)(c + \varepsilon))} \xrightarrow{\varepsilon \rightarrow 0} e^{-\lambda}.$$

The result follows now by remarking that  $e^{-\lambda}$  is the Laplace transform (for the argument  $\lambda^2/2$ ) of the first hitting time of 1 by a linear Brownian motion  $\gamma'$ , independent from  $\gamma$ .

### 3.2. Generalizations

Obviously, we can obtain several variants of Proposition 3.1, by studying  $T_{-bc,ac}^\theta$ ,  $0 < a, b \leq \infty$ , for  $c \rightarrow 0$  or  $c \rightarrow \infty$ , and  $a, b$  fixed. We define  $T_{-d,c}^\gamma \equiv \inf\{t: \gamma_t \notin (-d, c)\}$  and we have:

- $\frac{1}{c^2} T_{-bc,ac}^\theta \xrightarrow[c \rightarrow 0]{(law)} T_{-b,a}^\gamma$ .
- $\frac{1}{c} \log(T_{-bc,ac}^\theta) \xrightarrow[c \rightarrow \infty]{(law)} 2|\beta|_{T_{-b,a}^\gamma}$ .

In particular, we can take  $b = \infty$ , hence the following corollary.

#### Corollary 3.3.

(a) For  $c \rightarrow 0$ , we have

$$\frac{1}{c^2} T_{ac}^\theta \xrightarrow[c \rightarrow 0]{(law)} T_a^\gamma. \quad (21)$$

(b) For  $c \rightarrow \infty$ , we have

$$\frac{1}{c} \log(T_{ac}^\theta) \xrightarrow{c \rightarrow \infty} 2|\beta|_{T_a^\gamma} \stackrel{(law)}{=} 2|C_a|, \quad (22)$$

where  $(C_a, a \geq 0)$  is a standard Cauchy process.

**Remark 3.4 (Yet another proof of Spitzer's theorem).** Taking  $a = 1$ , from Corollary 3.3(b), we can obtain yet another proof of Spitzer's celebrated asymptotic theorem stated in (8). Indeed, (22) can be equivalently stated as:

$$P(\log T_c^\theta < cx) \xrightarrow{c \rightarrow \infty} P(2|C_1| < x). \quad (23)$$

Now, the LHS of (23) equals:

$$\begin{aligned} P(\log T_c^\theta < cx) &\equiv P(T_c^\theta < \exp(cx)) \equiv P\left(\sup_{u \leq \exp(cx)} \theta_u > c\right) \\ &= P(|\theta_{\exp(cx)}| > c) = P\left(|\theta_t| > \frac{\log t}{x}\right), \end{aligned} \quad (24)$$

with  $t = \exp(cx)$ . Thus, because  $|C_1| \stackrel{(law)}{=} |C_1|^{-1}$ , (23) now writes:

$$\text{for every } x > 0 \text{ given } P\left(|\theta_t| > \frac{\log t}{x}\right) \xrightarrow{t \rightarrow \infty} P\left(|C_1| > \frac{2}{x}\right), \quad (25)$$

which yields precisely Spitzer's theorem (8).

### 3.3. Speed of convergence

We can easily improve upon Proposition 3.1 by studying the speed of convergence of the distribution of  $\frac{1}{c^2} T_c^{|\theta|}$  towards that of  $T_1^{|\gamma|}$ , that is, the following proposition.

**Proposition 3.5.** For any function  $\varphi \in \mathcal{C}^2$ , with compact support,

$$\begin{aligned} &\frac{1}{c^2} \left( E \left[ \varphi \left( \frac{1}{c^2} T_c^{|\theta|} \right) \right] - E[\varphi(T_1^{|\gamma|})] \right) \\ &\xrightarrow{c \rightarrow 0} E \left[ \varphi'(T_1^{|\gamma|}) (T_1^{|\gamma|})^2 + \frac{2}{3} \varphi''(T_1^{|\gamma|}) (T_1^{|\gamma|})^3 \right]. \end{aligned} \quad (26)$$

**Proof.** We develop  $\exp(2c\beta_v)$ , for  $c \rightarrow 0$ , up to the second order term, that is,

$$e^{2c\beta_v} = 1 + 2c\beta_v + 2c^2\beta_v^2 + \dots$$

More precisely, we develop up to the second order term, and we obtain

$$\begin{aligned} E\left[\varphi\left(\frac{1}{c^2}T_c^{|\theta|}\right)\right] &= E\left[\varphi\left(\int_0^{T_1^{|\gamma|}} dv \exp(2c\beta_v)\right)\right] \\ &= E\left[\varphi(T_1^{|\gamma|}) + \varphi'(T_1^{|\gamma|}) \int_0^{T_1^{|\gamma|}} (2c\beta_v + 2c^2\beta_v^2) dv\right] \\ &\quad + \frac{1}{2}E\left[\varphi''(T_1^{|\gamma|})4c^2\left(\int_0^{T_1^{|\gamma|}} \beta_v dv\right)^2\right] + c^2o(c). \end{aligned}$$

We then remark that  $E[\int_0^t \beta_v dv] = 0$ ,  $E[\int_0^t \beta_v^2 dv] = t^2/2$  and  $E[(\int_0^t \beta_v dv)^2] = t^3/3$ , thus we obtain (26).  $\square$

#### 4. Checks via Bougerol's identity

So far, we have not made use of Bougerol's identity (4), which helps us to characterize the distribution of  $T_c^{|\theta|}$  [18]. In this subsection, we verify that writing the Gauss–Laplace transform in (6) as:

$$E\left[\sqrt{\frac{2}{\pi}} \frac{1}{\sqrt{(1/c^2)T_c^{|\theta|}}} \exp\left(-\frac{xc^2}{2T_c^{|\theta|}}\right)\right] = \frac{1}{\sqrt{1+xc^2}} \varphi_m(xc^2), \quad (27)$$

with  $m = \pi/(2c)$ , we find asymptotically for  $c \rightarrow 0$  the Gauss–Laplace transform of  $T_1^{|\gamma|}$ . Indeed, from (27), for  $c \rightarrow 0$ , we obtain:

$$\begin{aligned} E\left[\sqrt{\frac{2}{\pi}} \frac{1}{\sqrt{T_1^{|\gamma|}}} \exp\left(-\frac{x}{2T_1^{|\gamma|}}\right)\right] \\ = \lim_{c \rightarrow 0} \frac{2}{(\sqrt{1+xc^2} + \sqrt{xc^2})^{\pi/(2c)} + (\sqrt{1+xc^2} - \sqrt{xc^2})^{\pi/(2c)}}. \end{aligned} \quad (28)$$

Let us now study:

$$\begin{aligned} (\sqrt{1+xc^2} + \sqrt{xc^2})^{\pi/(2c)} &= \exp\left(\frac{\pi}{(2c)} \log[1 + (\sqrt{1+xc^2} - 1) + \sqrt{xc^2}]\right) \\ &\sim \exp\left(\frac{\pi}{2c} \left[c\sqrt{x} + \frac{xc^2}{2}\right]\right) \xrightarrow{c \rightarrow 0} \exp\left(\frac{\pi\sqrt{x}}{2}\right). \end{aligned}$$

A similar calculation finally gives

$$E\left[\sqrt{\frac{2}{\pi}} \frac{1}{\sqrt{T_1^{|\gamma|}}} \exp\left(-\frac{x}{2T_1^{|\gamma|}}\right)\right] = \frac{1}{\cosh((\pi/2)\sqrt{x})}, \quad (29)$$



a result which is in agreement with the law of  $\beta_{T_1^{|\gamma|}}$ , whose density is

$$E\left[\frac{1}{\sqrt{2\pi T_1^{|\gamma|}}} \exp\left(-\frac{y^2}{2T_1^{|\gamma|}}\right)\right] = \frac{1}{2 \cosh((\pi/2)y)}. \quad (30)$$

Indeed, the law of  $\beta_{T_c^{|\gamma|}}$  may be obtained from its characteristic function which is given by Revuz and Yor [15], page 73:

$$E[\exp(i\lambda\beta_{T_c^{|\gamma|}})] = \frac{1}{\cosh(\lambda c)}.$$

It is well known that Lévy [11], Biane and Yor [4]:

$$\begin{aligned} E[\exp(i\lambda\beta_{T_c^{|\gamma|}})] &= \frac{1}{\cosh(\lambda c)} = \frac{1}{\cosh(\pi\lambda c/\pi)} = \int_{-\infty}^{\infty} e^{i(\lambda c/\pi)y} \frac{1}{2\pi} \frac{1}{\cosh(y/2)} dy \\ &\stackrel{x=cy/\pi}{=} \int_{-\infty}^{\infty} e^{i\lambda x} \frac{1}{2\pi} \frac{\pi/c}{\cosh(x\pi/(2c))} dx = \int_{-\infty}^{\infty} e^{i\lambda x} \frac{1}{2c} \frac{1}{\cosh(x\pi/(2c))} dx. \end{aligned} \quad (31)$$

So, the density  $h_{-c,c}$  of  $\beta_{T_c^{|\gamma|}}$  is:

$$h_{-c,c}(y) = \left(\frac{1}{2c}\right) \frac{1}{\cosh(y\pi/(2c))} = \left(\frac{1}{c}\right) \frac{1}{e^{y\pi/(2c)} + e^{-y\pi/(2c)}}$$

and for  $c = 1$ , we obtain (30).

We recall from Remark 3.2 that (see also Pitman and Yor [14], where further results concerning the infinitely divisible distributions generated by some Lévy processes associated with the hyperbolic functions  $\cosh$ ,  $\sinh$  and  $\tanh$  can also be found):

$$E\left[\exp\left(-\frac{\lambda^2}{2} T_c^{|\gamma|}\right)\right] = \frac{1}{\cosh(\lambda c)}, \quad (32)$$

thus, for  $c = 1$  and  $\lambda = \frac{\pi}{2}\sqrt{x}$ , (29) now writes:

$$E\left[\sqrt{\frac{2}{\pi}} \frac{1}{\sqrt{T_1^{|\gamma|}}} \exp\left(-\frac{x}{2T_1^{|\gamma|}}\right)\right] = E\left[\exp\left(-\frac{x\pi^2}{8} T_1^{|\gamma|}\right)\right], \quad (33)$$

a result which gives a probabilistic proof of the reciprocal relation that was obtained in Biane, Pitman and Yor [3] (using the notation of this article, Table 1, page 442):

$$f_{C_1}(x) = \left(\frac{2}{\pi x}\right)^{3/2} f_{C_1}\left(\frac{4}{\pi^2 x}\right).$$

## References

- [1] ALILI, L., DUFRESNE, D. and YOR, M. (1997). Sur l'identité de Bougerol pour les fonctionnelles exponentielles du mouvement Brownien avec drift. In *Exponential Functionals and Principal Values Related to Brownian Motion* (M. YOR, ed.). *A collection of research papers. Biblioteca de la Revista Matemática, Ibero-Americana* 3–14. Madrid: Revista Matemática Iberoamericana.
- [2] BERTOIN, J. and WERNER, W. (1994). Asymptotic windings of planar Brownian motion revisited via the Ornstein–Uhlenbeck process. In *Séminaire de Probabilités, XXVIII. Lecture Notes in Math.* **1583** 138–152. Berlin: Springer. [MR1329109](#)
- [3] BIANE, P., PITMAN, J. and YOR, M. (2001). Probability laws related to the Jacobi theta and Riemann zeta functions, and Brownian excursions. *Bull. Amer. Math. Soc. (N.S.)* **38** 435–465 (electronic). [MR1848256](#)
- [4] BIANE, P. and YOR, M. (1987). Valeurs principales associées aux temps locaux browniens. *Bull. Sci. Math.* **111** 23–101. [MR0886959](#)
- [5] BOUGEROL, P. (1983). Exemples de théorèmes locaux sur les groupes résolubles. *Ann. Inst. H. Poincaré Sect. B (N.S.)* **19** 369–391. [MR0730116](#)
- [6] BURKHOLDER, D.L. (1977). Exit times of Brownian motion, harmonic majorization, and Hardy spaces. *Advances in Math.* **26** 182–205. [MR0474525](#)
- [7] DONEY, R.A. and VAKEROUDES, S. (2013). Windings of planar stable processes. In *Sém. Prob. XLV, Lect. Notes in Mathematics*. Berlin/Heidelberg/New York: Springer. To appear.
- [8] DUFRESNE, D. (2000). Laguerre series for Asian and other options. *Math. Finance* **10** 407–428. [MR1785163](#)
- [9] DURRETT, R. (1982). A new proof of Spitzer's result on the winding of two-dimensional Brownian motion. *Ann. Probab.* **10** 244–246. [MR0637391](#)
- [10] ITÔ, K. and MCKEAN, H.P. JR. (1965). *Diffusion Processes and Their Sample Paths. Die Grundlehren der Mathematischen Wissenschaften* **125**. Berlin/Heidelberg/New York: Springer. [MR0199891](#)
- [11] LÉVY, P. (1980). *Œuvres de Paul Lévy. Processus stochastiques* **4** Paris: Gauthier-Villars. Published under the direction of Daniel Dugué with the collaboration of Paul Deheuvels and Michel Ibero. [MR0586765](#)
- [12] MATSUMOTO, H. and YOR, M. (2005). Exponential functionals of Brownian motion. I. Probability laws at fixed time. *Probab. Surv.* **2** 312–347. [MR2203675](#)
- [13] MESSULAM, P. and YOR, M. (1982). On D. Williams' "pinching method" and some applications. *J. London Math. Soc. (2)* **26** 348–364. [MR0675178](#)
- [14] PITMAN, J. and YOR, M. (2003). Infinitely divisible laws associated with hyperbolic functions. *Canad. J. Math.* **55** 292–330. [MR1969794](#)
- [15] REVUZ, D. and YOR, M. (1999). *Continuous Martingales and Brownian Motion*, 3rd ed. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]* **293**. Berlin: Springer. [MR1725357](#)
- [16] SPITZER, F. (1958). Some theorems concerning 2-dimensional Brownian motion. *Trans. Amer. Math. Soc.* **87** 187–197. [MR0104296](#)
- [17] VAKEROUDES, S. (2011). Nombres de tours de certains processus stochastiques plans et applications à la rotation d'un polymère. (Windings of some planar stochastic processes and applications to the rotation of a polymer.) PhD thesis, Université Pierre et Marie Curie (Paris VI), Paris.

- [18] VAKERODIS, S. (2011). On hitting times of the winding processes of planar Brownian motion and of Ornstein–Uhlenbeck processes, via Bougerol’s identity. *Teor. Veroyatnost. i Primenen.—SIAM Theory Probab. Appl.* **56** 566–591 (in TVP).
- [19] VAKERODIS, S. and YOR, M. (2012). Some infinite divisibility properties of the reciprocal of planar Brownian motion exit time from a cone. *Electron. Commun. Probab.* **17** 9. [MR2950189](#)
- [20] WILLIAMS, D. (1974). *A simple geometric proof of Spitzer’s winding number formula for 2-dimensional Brownian motion. Preprint.* Swansea: Univ. College.
- [21] YOR, M. (1980). Loi de l’indice du lacet brownien, et distribution de Hartman–Watson. *Z. Wahrsch. Verw. Gebiete* **53** 71–95. [MR0576898](#)
- [22] YOR, M. (1985). Une décomposition asymptotique du nombre de tours du mouvement brownien complexe. *Astérisque* **132** 103–126. Colloquium in honor of Laurent Schwartz, Vol. 2 (Palaiseau, 1983). [MR0816763](#)
- [23] YOR, M. (1997). Generalized meanders as limits of weighted Bessel processes, and an elementary proof of Spitzer’s asymptotic result on Brownian windings. *Studia Sci. Math. Hungar.* **33** 339–343. [MR1454119](#)
- [24] YOR, M. (2001). *Exponential Functionals of Brownian Motion and Related Processes. Springer Finance.* Berlin: Springer. With an introductory chapter by Hélyette Geman, Chapters 1, 3, 4, 8 translated from the French by Stephen S. Wilson. [MR1854494](#)

*Received January 2012*