arXiv:1201.4110v1 [physics.bio-ph] 19 Jan 2012

Contractile units in disordered actomyosin bundles arise from F-actin buckling

Martin Lenz¹, Todd Thoresen², Margaret L. Gardel^{1,2,3}, and Aaron R. Dinner^{1,2,4,*}

¹James Franck Institute, ²Institute for Biophysical Dynamics, ³Department of Physics,

⁴Department of Chemistry, University of Chicago, Chicago IL 60637, USA

Bundles of filaments and motors are central to contractility in cells. The classic example is striated muscle, where actomyosin contractility is mediated by highly organized sarcomeres which act as fundamental contractile units. However, many contractile bundles *in vivo* and *in vitro* lack sarcomeric organization. Here we propose a model for how contractility can arise in actomyosin bundles without sarcomeric organization and validate its predictions with experiments on a reconstituted system. In the model, internal stresses in frustrated arrangements of motors with diverse velocities cause filaments to buckle, leading to overall shortening. We describe the onset of buckling in the presence of stochastic actin-myosin detachment and predict that buckling-induced contractile units" associated with this process. Consistent with these results, our reconstituted actomyosin bundles contract at relatively high motor density, and we observe buckling at the predicted length scale.

Contractility arising from interactions between myosin molecular motors and actin filaments (F-actin) is used ubiquitously by cells to build tension and drive morphological changes [1]. Such force transmission from molecular to cellular length scales is well understood in striated muscle, where it critically relies on highly organized structures known as sarcomeres [Fig. 1(a)] [2]. However, many contractile actomyosin bundles found in vivo, such as smooth muscle fibers [3], graded polarity bundles [4] and the contractile ring [5], lack a sarcomeric organization. Most recently, we have shown that in vitro bundles lacking apparent sarcomeric organization can also contract [6] [e.g., Fig. 1(b)]. In these disparate systems, contraction occurs with a well-defined contraction velocity per unit length, suggesting that contractile bundles can be meaningfully divided into elementary units that are arranged in series [5–7]. The mechanisms giving rise to such units in the absence of sarcomeric organization are not understood.

Much theoretical work on non-sarcomeric actomyosin assemblies posits contractility as a fundamental assumption, and predicts larger-scale effects such as polarity organization [8], the appearance of topological defects [9], active stiffening [10], and oscillatory behavior in cells [11]. Models that address the microscopic origin of contractility assume that myosin motors dwell at the barbed ends of F-actin, thus acting as transient static cross-linkers [12]. This generates sufficient sarcomere-like organization to elicit contraction [13]. Experimental evidence for this behavior is unfortunately lacking [14], and it is thus important to investigate alternative mechanisms.

Here we demonstrate another route to contractility in non-sarcomeric bundles through theory and experiments. We first show experimentally that contraction in reconstituted actomyosin bundles is accompanied by F-actin buckling. Such buckling could provide a symmetry-breaking mechanism necessary for contraction [15] [Fig. 1(c)], and we investigate the general consequences of asymmetric filament response theoretically by considering the build-up of forces in a bundle with randomly arranged motors. We predict that buckling, and thus contraction, can only be achieved in an intermediate range of motor density, due to the stabilizing effects of strong cross-linking at high density and of stochastic motor detachment at low density. This picture yields a characteristic length scale between two buckles, which provides a natural size for a contractile unit. These predictions are consistent with experimental observations, suggesting that buckling is necessary for contractility in non-sarcomeric actomyosin bundles.

To form reconstituted actomyosin bundles, we incubate F-actin with length $\ell_f \simeq 5\,\mu m$ with smooth muscle myosin thick filaments of length $\simeq 300 \,\mathrm{nm}$ in buffer lacking ATP such that thick filaments cross-link F-actin with high affinity. While flexible motors have been considered as a basis for contraction [16], this is unlikely to apply here as thick filaments are significantly more rigid than F-actin. The bundle lengths range from 10 to $100 \,\mu\text{m}$ with 4-6 F-actin per bundle cross-section, and no sarcomeric organization is observed [6]. By varying the concentration of myosin filaments, the average spacing ℓ_0 between two consecutive myosin filaments can be varied from 390 nm to $5.3 \,\mu m$ [6]. Perfusing buffer containing 1 mM ATP causes bundles formed with high myosin density ($\ell_0 = 540 \,\mathrm{nm}$) to shorten by $\simeq 10\%$ rapidly $(100-600 \text{ nm} \cdot \text{s}^{-1})$ [Fig. 1(d) and Ref. [14]; Movie S1]. In contrast, contraction does not occur at low myosin density $(\ell_0 = 1.5 \,\mu\text{m})$ [Fig. 1(e) and Movie S1]. A sharp transition between those two behaviors is observed at $\ell_0 = 1.3 \,\mu \text{m}$ [Fig. 1(f)].

In considering these observations, it is important to recognize that actomyosin interactions can *a priori* elicit extension just as well as contraction. As shown in Fig. 1(c), elementary bundles comprised of two F-actin ("filaments") and one myosin thick filament ("motor") containing numerous myosin heads contract when the




FIG. 1. Contraction in actomyosin bundles. (a) Sarcomeric structure as in striated muscle, where passive cross-linkers are attached near F-actin barbed ends. Motors move towards F-actin barbed ends, causing each contractile unit (sarcomere) to contract. (b) Bundle devoid of sarcomeric organization or passive cross-linkers, as in our experiments. (c) Motors and polar filaments induce local contraction or extension depending on the geometry of their assembly. Overall contractility requires breaking the balance between these two effects. (d) Time-lapse images of a bundle comprised of Factin and fluorescent myosin thick filaments (inverted contrast) with $\ell_0 = 540$ nm. The initially wavy bundle becomes taut following the addition of 1 mM ATP at t = 0 s, indicating contraction. Scale bar, $5\,\mu m$. (e) Similar experiment with $\ell_0 = 1.5 \,\mu\text{m}$, showing no contraction. Scale bar as in (a). See also Movie S1. (f) Bundle contraction as a function of ℓ_0 . Bars indicate standard deviation $(n \ge 25)$.

motor is located in the vicinity of the pointed ends, but extend when it is close to the barbed ends. In sarcomeres, myosin is restricted towards the pointed ends of actin filaments, which favors contraction over extension [Fig. 1(a)]. By contrast, we show in Refs. [14, 15] that these two tendencies compensate and prevent overall contraction in non-sarcomeric bundles unless two specific conditions are fulfilled. First, the unloaded motor velocities need to have a certain dispersion among the motor population. Otherwise, motors merely induce filament translation without inducing overall bundle contraction [13]. Such dispersion has been observed experimentally [17], and in our system likely arises from the variation of number of myosin heads in the thick filaments. In its presence, stresses build in the bundle as motors translate the filaments with different preferred velocities. While these stresses present a potential for bundle deformation, the disordered nature of the bundle implies that they have equal chances of being compressive or extensile. Thus the second condition is an asymmetric response of the filaments to such stresses, which breaks the symmetry




FIG. 2. Buckling in non-sarcomeric contractile actomyosin bundles. (a) Time-lapse images of fluorescent actin (inverted contrast) showing F-actin buckling (arrowheads) following the addition of 1 mM ATP at t = 0s. Scale bar, 5μ m. See also Movie S2. (b) Relative contraction (filled squares) and number of F-actin buckles (open circles) as a function of time. Data shows mean \pm sd averaged over n = 3 bundles with $\ell_0 \simeq 1 \mu$ m. (c) The presence of fast (grey) and slow (white) motors generically induce compressive (red) and extensile (blue) stresses in filaments. (d) Buckling of the compressed filaments leads to an overall shortening of the bundle.

between contraction and extension. Contraction occurs when filaments yield under compression while resisting extension [14, 15].

We next look for evidence of this behavior in our experiments. In this respect, we observe F-actin buckling coinciding with contraction [Fig. 2(a); Movie S2]. This constitutes an extreme form of asymmetric response of the filaments, and thus enables contractility. Prior to ATP addition, compact bundles with aligned F-actin are observed. Upon ATP addition, the frequency of buckles increases rapidly during contraction, and then diminishes once contraction stops [Fig. 2(b)]. These F-actin buckles are dynamic, with their amplitude, curvature and location changing over time.

Qualitatively, the relationship between buckling and contraction can be understood as follows. Consider two antiparallel filaments interacting through several different motors with distinct speeds [Fig. 2(c)]. As motors start to move relative to the filaments, stresses build in sections of the filament flanked by motors with different speeds. When the flanking motor proximal to the barbed end is faster than that proximal to the pointed end, compression arises. When it is slower, tension arises. Following buckling of the compressed filament sections, fast motors are free to move quickly while the others move slowly. This results in the growth of the compressed sections and shrinkage of the extended ones, and thus in overall bundle contraction [Fig. 2(d)]. The region centered around each buckle thus plays the role of a contractile unit, whose typical size is equal to the distance ℓ_B between two buckles.

In this picture, the contractile behavior of the bundle hinges on the ability of the motors to induce filament buckling. At high motor density, we expect the bundle to be so strongly cross-linked that buckling becomes impossible despite the sizable stresses induced by a large number of motors. At low motor density, we expect that stochastic detachment of the motors undermines stress build-up and thus prevents buckling. Here we present a mathematical model to predict the range of myosin densities enabling contraction and the contractile unit length ℓ_B . These results are then compared with the observations in Figs. 1 and 2 to validate the proposed contraction mechanism.

The key assumptions of our model are that (1) motors have a dispersion in their unloaded velocities, (2) a section of filament between two motors buckles above a certain threshold force F_B , and (3) motors intermittently detach from the filaments, thus allowing local stress relaxation. We consider a bundle of weakly deformed filaments and ask whether the forces developing within it are sufficient to induce buckling [Fig. 3(a)].

To this end, we focus on a single filament of length ℓ_f and approximate its surroundings by an effective medium composed of immobile [14] point-like motors separated by a distance $\ell_0 \ll \ell_f$ [Fig. 3(b)]. This divides the filament into discrete sections, which we label by $i = 0, \ldots, \ell_f/\ell_0$. We take into account the possibility that the filaments are not straight, but bend away from the x-axis, implying that the contour length L_i of filament section i can be larger than ℓ_0 . Defining f_i as the tension of filament section i ($f_i < 0$ for a compressed filament section), we expand its force-extension relationship for small deformations:

$$L_i = L_i(f_i = 0) - cf_i,$$
 (1)

where c > 0 is the filament compliance. We refer to the motor flanked by filament sections i - 1 and i as "motor i", and describe its operation by the simplified force-velocity relationship

$$f_{i-1} - f_i = F_i - \chi v_i.$$
 (2)

Here v_i denotes the local velocity of the filament at the location of motor i and $\chi > 0$ is the motor susceptibility. Eqs. (1) and (2) yield a local relaxation time scale $\tau_r = \chi c/2$. The time-independent stall force of motor iis denoted by F_i in Eq. (2), and is drawn from a random distribution satisfying

$$\overline{F_i} = F_S$$
 and $\overline{F_i F_j} - \overline{F_i} \overline{F_j} = \delta F_S^2 \delta_{ij},$ (3)

where bars denote averages over the motor distribution. As a result, different motors have different unloaded velocities F_i/χ as required for contraction. Owing to the




FIG. 3. Stress build-up in bundles with non-identical motors. (a) In a bundle with motors having non-identical velocities (shades of grey), filaments of lengths $\approx \ell_f$ are subjected to random motor forces at points $\approx \ell_0$ apart distributed throughout their length. (b) Prior to buckling, the environment of a filament of interest (red) can be approximated by a collection of immobile motors (shades of grey) [14].

conservation of filament mass:

$$\frac{\mathrm{d}L_i}{\mathrm{d}t} = v_i - v_{i+1}.\tag{4}$$

Finally, a motor bound to several filaments as in Fig. 3(a) can transiently detach from one while still holding onto the others [14]. We thus let each motor *i* randomly detach from the filament with a constant rate $1/\tau_d$. Following detachment, local filament stresses relax instantaneously, yielding $f_i = f_{i-1} = (f_i + f_{i-1})/2$. The motor then reattaches after a time much shorter than τ_r and τ_d [14]. We denote by $\langle \ldots \rangle$ the average over the Poisson process of motor detachment.

We obtain the space and time evolution of the filament tension f(x,t) in the continuum limit $i \to x/\ell_0$ by combining Eqs. (1-4) and averaging over motor detachment [14]:

$$\partial_t \langle f \rangle - D \partial_x^2 \langle f \rangle = (\ell_0 / 2\tau_r) \partial_x F, \tag{5}$$

where $D = \ell_0^2(\tau_r^{-1} + \tau_d^{-1})/2$. The right-hand-side of Eq. (5) involves the spatial gradient of the stall force F(x), reflecting the fact that non-identical motors lead to force build-up. This effect competes with the relaxation of filament forces through motor detachment, which enters through the diffusion term $D\partial_r^2 \langle f \rangle$.

An initially relaxed filament [f(x, t = 0) = 0] experiences a vanishing average force $\langle f \rangle(x, t) = 0$ throughout its dynamics. To quantify the magnitude of the motorinduced stress, we thus calculate the rms filament force $(\overline{\langle f^2 \rangle})^{1/2}$ [14] and find that it increases monotonically from zero at t = 0 to

$$f_{\infty} = \frac{1}{2\sqrt{3}} \left(\frac{\ell_f}{\ell_0}\right)^{1/2} \frac{\delta F_S}{1 + \tau_r / \tau_d}.$$
 (6)

at $t = \infty$ [Fig. 4(a-b)]. We next estimate the dependence of the ratio τ_r/τ_d on the experimentally accessible




FIG. 4. Model predictions for filament force build-up. (a) Black line: Steady-state filament force f_{∞} as a function of motor spacing ℓ_0 as in Eq. (6). For $\ell_0 \ll \ell_0^*$ and $\ell_0 \gg \ell_0^*$, $f_{\infty} \propto \ell_0^{-1/2}$ and $\ell_0^{-9/2}$, respectively. Colored lines: buckling force $F_B \propto \ell_0^{-2}$. (b) Typical filament force $(\langle \overline{f^2} \rangle)^{1/2}$ as a function of time [14]. (c) Contractile unit size ℓ_B as a function of ℓ_0 as in Eq. (8) ($\ell_f \simeq 5 \,\mu$ m).

parameter ℓ_0 . A Worm-Like Chain model for filament elasticity yields $c \approx \ell_0^4/k_B T \ell_p^2$, where ℓ_p is the filament persistence length [18], and we approximate $\chi \approx F_S/v$, where v is a characteristic motor velocity. This implies $\tau_r/\tau_d \approx (\ell_0/\ell_0^{-1})^4$, with $\ell_0^* = (k_B T \ell_p^2 v \tau_d/F_S)^{1/4}$. We can thus distinguish two regimes for the steady-state force f_∞ [Fig. 4(a)]. For $\ell_0 \ll \ell_0^*$, detachment events are rare compared to the time τ_r needed for the force to recover from such an event, and f_∞ is not affected by them. For $\ell_0 \gg \ell_0^*$, f_∞ quickly decreases with increasing ℓ_0 as detachment becomes much faster than recovery.

Up to a prefactor of order one, contraction proceeds as in Fig. 2(c-d) if $f_{\infty} > F_B \approx k_B T \ell_p / \ell_0^2$ [18]. Comparing f_{∞} to F_B as in Fig. 4(a), we find a threshold stiffness above which buckling cannot occur (as exampled by the *blue line*). Reasonable values for our actomyosin system are $\ell_p \simeq 10 \,\mu\text{m}, v \simeq 200 \,\text{nm} \cdot \text{s}^{-1},$ $\delta F_S \approx F_S \simeq 1 \,\text{pN}$ and $\tau_d \simeq 200 \,\text{ms}$ based on the typical time scales involved in the myosin mechanochemical cycle [14]. These values put us in the soft filament regime defined by $\ell_p \ll \delta F_S^4 L_f^2 (v \tau_d)^{3/2} / k_B T^{5/2} F_S^{3/2} \simeq 20 \,\text{cm}$ (*red line*). In this regime, the lines representing F_B and f_{∞} intersect at

$$\ell_0^- = (k_B T \ell_p / \delta F_S \ell_f^{1/2})^{2/3} \simeq 70 \,\mathrm{nm}, \tag{7a}$$

$$\ell_0^+ = (\ell_f^{1/2} v \tau_d \ell_p \delta F_S / F_S)^{2/5} \simeq 1 \,\mu \mathrm{m},$$
 (7b)

meaning that buckling and contraction occur for $\ell_0^- < \ell_0 < \ell_0^+$. This range reflects the fact that strong crosslinking ($\ell_0 < \ell_0^-$) suppresses buckling while sparse motors ($\ell_0 > \ell_0^+$) are undermined by stochastic detachment. While the regime $\ell_0 \simeq \ell_0^-$ is not accessible experimentally, the predicted value for ℓ_0^+ is strikingly similar to the motor spacing at which the breakdown of contraction is observed in Fig. 1(f) (1.3 μ m), suggesting that the proposed mechanism is a good description of our experiments.

To characterize the contractile units resulting from this mechanism when $\ell_0^- < \ell_0 < \ell_0^+$, we turn to the transient regime leading up to filament buckling. The filament force profile as a function of x is initially flat, and subsequently coarsens into a random walk for $t = +\infty$. According to Eq. (5), this coarsening occurs diffusively with diffusion coefficient D. The typical filament forces at time $t \ll \ell_f^2/D$ are thus of order $f_\infty (\sqrt{Dt}/\ell_f)^{1/2}$. We denote the time that this force reaches the buckling threshold F_B by t_B , following which contraction proceeds as in Fig. 2(c-d) and the coarsening dynamics is interrupted. The distance between buckles at t_B thus yields the contractile unit size

$$\ell_B \approx \sqrt{Dt_B} \approx \frac{\ell_p^2}{\ell_0^3} \left(\frac{k_B T}{\delta F_S}\right)^2 \left(1 + \frac{\tau_r}{\tau_d}\right)^2.$$
(8)

As illustrated in Fig. 4(c), ℓ_B is typically in the micrometer range, in agreement with the observations of Fig. 2(a) and the findings of Ref. [6].

Because of compensating effects between contractile and extensile motor-filament configurations, the familiar framework involving rigid filaments and identical motors commonly used to describe striated muscle contraction is not suited to study actomyosin bundles lacking sarcomeric organization. Here, we put forward an alternative mechanism based on our observation of buckling. The buckling arises from the nonlinear elastic response of Factin [19] and dispersion in the speeds of myosin motors [20]. F-actin buckling has previously been invoked to explain contraction qualitatively [21]. Addition of passive cross-linkers, which are formally equivalent to immobile motors, would reinforce a dispersion of motor velocities and promote contraction.

The order-of-magnitude agreement between theory and experiments with respect to the size of contractile units and the critical myosin concentration required for contraction suggests that our current analysis offers a good description of the onset of bundle contractility. Our conclusions are robust to inclusion of features such as inhomogeneous motor spacings ℓ_0 and force dependence of the motor detachment rate [14]. Our mechanism is a general one and applies to any one-dimensional system of polar filaments and motors. Further experiments and theory are needed to better understand the molecular basis for motor inhomogeneities and filament asymmetric response in the myriad of non-sarcomeric organizations found *in vivo*.

We thank Yitzhak Rabin and Tom Witten for useful discussions. This work was supported by NSF dmr-mrsec 0820054, NIH P50 GM081892 and NIH DP10D00354.

^{*} dinner@uchicago.edu

- J. Stricker, T. Falzone, and M. L. Gardel. J. Biomech., 43(1):9, 2010.
- [2] B. Alberts et al. Garland, New-York, 1998.
- [3] F. S. Fay et al. J. Cell Biol., 96(3):783, 1983.
- [4] L. P. Cramer, M. Siebert, and T. J. Mitchison. J. Cell Biol., 136(6):1287, 1997.
- [5] A. Carvalho, A. Desai, and K. Oegema. *Cell*, 137(5):926, 2009.
- [6] T. Thoresen, M. Lenz, and M. L. Gardel. Biophys. J., 100(11):2698, 2011.
- [7] W. M. Bement and D. G. Capco. Cell Motil. Cytoskeleton, 20(2):145, 1991; A. M. Herrera et al. J. Cell Sci., 118(11):2381, 2005.
- [8] K. Kruse, A. Zumdieck, and F. Jülicher. Europhys. Lett., 64(5):716, 2003; N. Yoshinaga et al. Phys. Rev. Lett., 105(23):238103, 2010.
- K. Kruse et al. Phys. Rev. Lett., 92:078101, 2004;
 K. Kruse et al. Eur. Phys. J. E, 16(1):5, 2005.
- [10] F. C. MacKintosh and A. J. Levine. Phys. Rev. Lett., 100(1):018104, 2008.

- [11] G. Salbreux et al. Phys. Biol., 4(4):268, 2007.
- [12] K. Kruse and F. Jülicher. *Phys. Rev. Lett.*, 85(8):1778, 2000; K. Kruse and K. Sekimoto. *Phys. Rev. E*, 66(3):031904, 2002; K. Kruse and F. Jülicher. *Phys. Rev. E*, 67(5):051913, 2003; T. B. Liverpool and M. C. Marchetti. *Phys. Rev. Lett.*, 90(13):138102, 2003; R. Peter *et al. New J. Phys.*, 10(3):035002, 2008.
- [13] A. Zemel and A. Mogilner. Phys. Chem. Chem. Phys., 11(24):4821, 2009.
- [14] Supporting material.
- [15] M. Lenz and A. Dinner. arXiv:1101.1058, 2011.
- [16] T. B. Liverpool et al. Europhys. Lett., 85:18007, 2009.
- [17] A. Yamada, N. Ishii, and K. Takahashi. J. Biochem., 108(3):341, 1990.
- [18] T. Odijk. Macromolecules, 28(20):7016, 1995.
- [19] J. Berro et al. Biophys. J., 92(7):2546, 2007.
- [20] Y. Tanaka, A. Ishijima, and S. Ishiwata. *Biochim. Bio-phys. Acta*, 1159(1):94, 1992.
- [21] M. Soares e Silva et al. Proc. Natl. Acad. Sci. U.S.A., 108(23):9408, 2011.