

FAMILIES AND SPRINGER'S CORRESPONDENCE

G. LUSZTIG

INTRODUCTION

0.1. Let G be a connected reductive algebraic group over an algebraically closed field \mathbf{k} of characteristic p . Let W be the Weyl group of G ; let $\text{Irr}W$ be a set of representatives for the isomorphism classes of irreducible representations of W over $\bar{\mathbf{Q}}_l$, an algebraic closure of the field of l -adic numbers (l is a fixed prime number $\neq p$).

Now $\text{Irr}W$ is partitioned into subsets called *families* as in [L1, Sec.9], [L3, 4.2]. Moreover to each family \mathcal{F} in $\text{Irr}W$, a certain set $\mathbf{X}_{\mathcal{F}}$, a pairing $\{, \} : \mathbf{X}_{\mathcal{F}} \times \mathbf{X}_{\mathcal{F}} \rightarrow \bar{\mathbf{Q}}_l$, and an imbedding $\mathcal{F} \rightarrow \mathbf{X}_{\mathcal{F}}$ was canonically attached in [L1], [L3, Ch.4]. (The set $\mathbf{X}_{\mathcal{F}}$ with the pairing $\{, \}$, which can be viewed as a nonabelian analogue of a symplectic vector space, plays a key role in the classification of unipotent representations of a finite Chevalley group [L3] and in that of unipotent character sheaves on G). In [L1], [L3] it is shown that $\mathbf{X}_{\mathcal{F}} = M(\mathcal{G}_{\mathcal{F}})$ where $\mathcal{G}_{\mathcal{F}}$ is a certain finite group associated to \mathcal{F} and, for any finite group Γ , $M(\Gamma)$ is the set of all pairs (g, ρ) where g is an element of Γ defined up to conjugacy and ρ is an irreducible representation over $\bar{\mathbf{Q}}_l$ (up to isomorphism) of the centralizer of g in Γ ; moreover $\{, \}$ is given by the "nonabelian Fourier transform matrix" of [L1, Sec.4] for $\mathcal{G}_{\mathcal{F}}$.

In the remainder of this paper we assume that either $p = 0$ or that p is a good prime for G . In this case a uniform definition of the group $\mathcal{G}_{\mathcal{F}}$ was proposed in [L3, 13.1] in terms of special unipotent classes in G and the Springer correspondence, but the fact that this leads to a group isomorphic to $\mathcal{G}_{\mathcal{F}}$ as defined in [L3, Ch.4] was stated in [L3, (13.1.3)] without proof. One of the aims of this paper is to supply the missing proof.

To state the results of this paper we need some definitions. For $E \in \text{Irr}W$ let $a_E \in \mathbf{N}, b_E \in \mathbf{N}$ be as in [L3, 4.1]. As noted in [L2], for $E \in \text{Irr}W$ we have

(a) $a_E \leq b_E$;

we say that E is *special* if $a_E = b_E$.

Let C be a unipotent conjugacy class in G and let $u \in C$. Let $Z(u)$ be the centralizer of u in G and let $A(u)$ be the group of connected components of $Z(u)$. Let \mathcal{B}_u be the variety of Borel subgroups of G that contain u ; this is a nonempty

Supported in part by National Science Foundation grant DMS-0758262.

variety of dimension, say, e_C . The conjugation action of $Z(u)$ on \mathcal{B}_u induces an action of $A(u)$ on $\mathbf{S}_u := H^{2e_C}(\mathcal{B}_u, \mathbf{Q}_l)$. Now W acts on \mathbf{S}_u by Springer's representation [Spr]; however here we adopt the definition of the W -action on \mathbf{S}_u given in [L4] which differs from Springer's original definition by tensoring by sign. The W -action on \mathbf{S}_u commutes with the $A(u)$ -action. Hence we have canonically $\mathbf{S}_u = \oplus_{E \in \text{Irr} W} E \otimes \mathcal{V}_E$ (as $W \times A(u)$ -modules) where \mathcal{V}_E are finite dimensional \mathbf{Q}_l -vector spaces with $A(u)$ -action. Let $\text{Irr}_C W = \{E \in \text{Irr} W; \mathcal{V}_E \neq 0\}$; this set does not depend on the choice of u in C . By [Spr], the sets $\text{Irr}_C W$ (for C variable) form a partition of $\text{Irr} W$; also, if $E \in \text{Irr}_C W$ then \mathcal{V}_E is an irreducible $A(u)$ -module and, if $E \neq E'$ in $\text{Irr}_C W$, then the $A(u)$ -modules $\mathcal{V}_E, \mathcal{V}_{E'}$ are not isomorphic. By [BM] we have

(b) $e_C \leq b_E$ for any $E \in \text{Irr}_C W$

and the equality $b_E = e_C$ holds for exactly one $E \in \text{Irr}_C W$ which we denote by E_C (for this E , \mathcal{V}_E is the unit representation of $A(u)$).

Following [L3, (13.1.1)] we say that C is *special* if E_C is special. (This concept was introduced in [L2, Sec.9] although the word "special" was not used there.) From (b) we see that C is special if and only if $a_{E_C} = e_C$.

Now assume that C is special. We denote by $\mathcal{F} \subset \text{Irr} W$ the family that contains E_C . (Note that $C \mapsto \mathcal{F}$ is a bijection from the set of special unipotent classes in G to the set of families in $\text{Irr} W$.) We set $\text{Irr}_C^* W = \{E \in \text{Irr}_C W; E \in \mathcal{F}\}$ and

$$\mathcal{K}(u) = \{a \in A(u); a \text{ acts trivially on } \mathcal{V}_E \text{ for any } E \in \text{Irr}_C^* W\}.$$

This is a normal subgroup of $A(u)$. We set $\bar{A}(u) = A(u)/\mathcal{K}(u)$, a quotient group of $A(u)$. Now, for any $E \in \text{Irr}_C^* W$, \mathcal{V}_E is naturally an (irreducible) $\bar{A}(u)$ -module. Another definition of \bar{A}_u is given in [L3, (13.1.1)]. In that definition $\text{Irr}_C^* W$ is replaced by $\{E \in \text{Irr}_C W; a_E = e_C\}$ and $\mathcal{K}(u), \bar{A}(u)$ are defined as above but in terms of this modified $\text{Irr}_C^* W$. However the two definitions are equivalent in view of the following result.

Proposition 0.2. *Assume that C is special. Let $E \in \text{Irr}_C W$.*

- (a) *We have $a_E \leq e_C$.*
- (b) *We have $a_E = e_C$ if and only if $E \in \mathcal{F}$.*

This follows from [L6, 10.9]. Note that (a) was stated without proof in [L3, (13.1.2)] (the proof I had in mind at the time of [L3] was combinatorial).

0.3. The following result is equivalent to a result stated without proof in [L3, (13.1.3)].

Theorem 0.4. *Let C be a special unipotent class of G , let $u \in C$ and let \mathcal{F} be the family that contains E_C . Then we have canonically $\mathbf{X}_{\mathcal{F}} = M(\bar{A}(u))$ so that the pairing $\{, \}$ on $\mathbf{X}_{\mathcal{F}}$ coincides with the pairing $\{, \}$ on $M(\bar{A}(u))$. Hence $\mathcal{G}_{\mathcal{F}}$ can be taken to be $\bar{A}(u)$.*

This is equivalent to the corresponding statement in the case where G is adjoint, which reduces immediately to the case where G is adjoint simple. It is then enough

to prove the theorem for one G in each isogeny class of semisimple, almost simple algebraic groups; this will be done in §3 after some combinatorial preliminaries in §1, §2. The proof uses the explicit description of the Springer correspondence: for type A_n, G_2 in [Spr]; for type B_n, C_n, D_n in [S1] (as an algorithm) and in [L4] (by a closed formula); for type F_4 in [S2]; for type E_n in [AL], [Sp1].

An immediate consequence of (the proof of) Theorem 0.4 is the following result which answers a question of R. Bezrukavnikov.

Corollary 0.5. *In the setup of 0.4 let $E \in \text{Irr}_C^* W$ and let \mathcal{V}_E be the corresponding $A(u)$ -module viewed as an (irreducible) $\bar{A}(u)$ -module. The image of E under the canonical imbedding $\mathcal{F} \rightarrow \mathbf{X}_{\mathcal{F}} = M(\bar{A}(u))$ is represented by the pair $(1, \mathcal{V}_E) \in M(\bar{A}(u))$. Conversely, if $E \in \mathcal{F}$ and the image of E under $\mathcal{F} \rightarrow \mathbf{X}_{\mathcal{F}} = M(\bar{A}(u))$ is represented by the pair $(1, \rho) \in M(\bar{A}(u))$ where ρ is an irreducible representation of $\bar{A}(u)$, then $E \in \text{Irr}_C^* W$ and $\rho \cong \mathcal{V}_E$.*

0.6. Corollary 0.5 has the following interpretation. Let Y be a (unipotent) character sheaf on G whose restriction to the regular semisimple elements is an irreducible local system; assume that in the usual parametrization of unipotent character sheaves by $\sqcup_{\mathcal{F}} \mathbf{X}_{\mathcal{F}}$, Y corresponds to $(1, \rho) \in M(\bar{A}(u))$ where C is the special unipotent class corresponding to a family \mathcal{F} , $u \in C$ and ρ is an irreducible representation of $\bar{A}(u)$. Then $Y|_C$ is (up to shift) the irreducible local system on C defined by ρ ; moreover, if C' is a unipotent class of dimension $< \dim C$ then $Y|_{C'} = 0$. A similar interpretation of the parametrization of unipotent character sheaves on G by $\sqcup_{\mathcal{F}} \mathbf{X}_{\mathcal{F}}$ in terms of restrictions to various subsets of G is expected in general and is illustrated in [L5, 4.7(a)].

0.7. Notation. If A, B are subsets of \mathbf{N} we denote by $A \dot{\cup} B$ the union of A and B regarded as a multiset (each element of $A \cap B$ appears twice). For any set \mathcal{X} , we denote by $\mathcal{P}(\mathcal{X})$ the set of subsets of \mathcal{X} viewed as an F_2 -vector space with sum given by the symmetric difference. If $\mathcal{X} \neq \emptyset$ we note that $\{\emptyset, \mathcal{X}\}$ is a line in $\mathcal{P}(\mathcal{X})$ and we set $\bar{\mathcal{P}}(\mathcal{X}) = \mathcal{P}(\mathcal{X}) / \{\emptyset, \mathcal{X}\}$, $\mathcal{P}_{ev}(\mathcal{X}) = \{L \in \mathcal{P}(\mathcal{X}); |L| \equiv 0 \pmod{2}\}$; let $\bar{\mathcal{P}}_{ev}(\mathcal{X})$ be the image of $\mathcal{P}_{ev}(\mathcal{X})$ under the obvious map $\mathcal{P}(\mathcal{X}) \rightarrow \bar{\mathcal{P}}(\mathcal{X})$ (thus $\bar{\mathcal{P}}_{ev}(\mathcal{X}) = \bar{\mathcal{P}}(\mathcal{X})$ if $|\mathcal{X}|$ is odd and $\bar{\mathcal{P}}_{ev}(\mathcal{X})$ is a hyperplane in $\bar{\mathcal{P}}(\mathcal{X})$ if $|\mathcal{X}|$ is even). Now if $\mathcal{X} \neq \emptyset$, the assignment $L, L' \mapsto |L \cap L'| \pmod{2}$ defines a symplectic form on $\mathcal{P}_{ev}(\mathcal{X})$ which induces a nondegenerate symplectic form $(,)$ on $\bar{\mathcal{P}}_{ev}(\mathcal{X})$ via the obvious linear map $\mathcal{P}_{ev}(\mathcal{X}) \rightarrow \bar{\mathcal{P}}_{ev}(\mathcal{X})$.

For $z \in (1/2)\mathbf{Z}$ we set $\lfloor z \rfloor = z$ if $z \in \mathbf{Z}$ and $\lfloor z \rfloor = z - (1/2)$ if $z \in \mathbf{Z} + (1/2)$.

Erratum to [L3]. On page 86, line -6 delete: " $b' < b$ " and on line -4 before "In the language..." insert: "The array above is regarded as identical to the array obtained by interchanging its two rows."

On page 343, line -5, after "respect to M " insert: "and where the group $\mathcal{G}_{\mathcal{F}}$ defined in terms of (u', M) is isomorphic to the group $\mathcal{G}_{\mathcal{F}}$ defined in terms of (u, G) ".

Erratum to [L4]. In the definition of A_{α}, B_{α} in [L4, 11.5], the condition $I \in \alpha$ should be replaced by $I \in \alpha'$ and the condition $I \in \alpha'$ should be replaced by $I \in \alpha$.

1. COMBINATORICS

1.1. Let N be an even integer ≥ 0 . Let $a := (a_0, a_1, a_2, \dots, a_N) \in \mathbf{N}^{N+1}$ be such that $a_0 \leq a_1 \leq a_2 \leq \dots \leq a_N$, $a_0 < a_2 < a_4 < \dots$, $a_1 < a_3 < a_5 < \dots$. Let $\mathcal{J} = \{i \in [0, N]; a_i \text{ appears exactly once in } a\}$. We have $\mathcal{J} = \{i_0, i_1, \dots, i_{2M}\}$ where $M \in \mathbf{N}$ and $i_0 < i_1 < \dots < i_{2M}$ satisfy $i_s = s \pmod{2}$ for $s \in [0, 2M]$. Hence for any $s \in [0, 2M-1]$ we have $i_{s+1} = i_s + 2m_s + 1$ for some $m_s \in \mathbf{N}$. Let \mathcal{E} be the set of $b := (b_0, b_1, b_2, \dots, b_N) \in \mathbf{N}^{N+1}$ such that $b_0 < b_2 < b_4 < \dots$, $b_1 < b_3 < b_5 < \dots$ and such that $[b] = [a]$ (we denote by $[b], [a]$ the multisets $\{b_0, b_1, \dots, b_N\}, \{a_0, a_1, \dots, a_N\}$). We have $a \in \mathcal{E}$. For $b \in \mathcal{E}$ we set

$$\hat{b} = (\hat{b}_0, \hat{b}_1, \hat{b}_2, \dots, \hat{b}_N) = (b_0, b_1+1, b_2+1, b_3+2, b_4+2, \dots, b_{N-1}+(N/2), b_N+(N/2)). \blacksquare$$

Let $[\hat{b}]$ be the multiset $\{\hat{b}_0, \hat{b}_1, \hat{b}_2, \dots, \hat{b}_N\}$.

For $s \in \{1, 3, \dots, 2M-1\}$ we define $a^{\{s\}} = (a_0^{\{s\}}, a_1^{\{s\}}, a_2^{\{s\}}, \dots, a_N^{\{s\}}) \in \mathcal{E}$ by

$$\begin{aligned} & (a_{i_s}^{\{s\}}, a_{i_s+1}^{\{s\}}, a_{i_s+2}^{\{s\}}, a_{i_s+3}^{\{s\}}, \dots, a_{i_s+2m_s}^{\{s\}}, a_{i_s+2m_s+1}^{\{s\}}) \\ &= (a_{i_s+1}, a_{i_s}, a_{i_s+3}, a_{i_s+2}, \dots, a_{i_s+2m_s+1}, a_{i_s+2m_s}) \end{aligned}$$

and $a_i^{\{s\}} = a_i$ if $i \in [0, N] - [i_s, i_{s+1}]$. More generally for $X \subset \{1, 3, \dots, 2M-1\}$ we define $a^X = (a_0^X, a_1^X, a_2^X, \dots, a_N^X) \in \mathcal{E}$ by $a_i^X = a_i^{\{s\}}$ if $s \in X$, $i \in [i_s, i_{s+1}]$, and $a_i^X = a_i$ for all other $i \in [0, N]$. Note that $[a^X] = [\hat{a}]$. Conversely, we have the following result.

Lemma 1.2. *Let $b \in \mathcal{E}$ be such that $[\hat{b}] = [\hat{a}]$. There exists $X \subset \{1, 3, \dots, 2M-1\}$ such that $b = a^X$.*

The proof is given in 1.3-1.5.

1.3. We argue by induction on M . We have

$$a = (y_1 = y_1 < y_2 = y_2 < \dots < y_r = y_r < a_{i_0} < \dots)$$

for some r . Since $[b] = [a]$, we must have

$$(b_0, b_2, b_4, \dots) = (y_1, y_2, \dots, y_r, \dots), (b_1, b_3, b_5, \dots) = (y_1, y_2, \dots, y_r, \dots).$$

Thus,

$$(a) \ b_i = a_i \text{ for } i < i_0.$$

We have $a = (\dots > a_{2M} > y'_1 = y'_1 < y'_2 = y'_2 < \dots < y'_{r'} = y'_{r'})$ for some r' . Since $[b] = [a]$, we must have

$$(b_0, b_2, b_4, \dots) = (\dots, y'_1, y'_2, \dots, y'_{r'}), (b_1, b_3, b_5, \dots) = (\dots, y'_1, y'_2, \dots, y'_{r'}).$$

Thus,

$$(b) \ b_i = a_i \text{ for } i > i_{2M}.$$

If $M = 0$ we see that $b = a$ and there is nothing further to prove. In the rest of the proof we assume that $M \geq 1$.

1.4. From 1.3 we see that

$$(a_0, a_1, a_2, \dots, a_{i_{2M}}) = (\dots, a_{i_{2M}-1} < x_1 = x_1 < x_2 = x_2 < \dots < x_q = x_q < a_{i_{2M}})$$

(for some q) has the same entries as $(b_0, b_1, b_2, \dots, b_{i_{2M}}$ (in some order)). Hence the pair

$$(\dots, b_{i_{2M}-5}, b_{i_{2M}-3}, b_{i_{2M}-1}), (\dots, b_{i_{2M}-4}, b_{i_{2M}-2}, b_{i_{2M}})$$

must have one of the following four forms.

$$\begin{aligned} & (\dots, a_{i_{2M}-1}, x_1, x_2, \dots, x_q), (\dots, x_1, x_2, \dots, x_q, a_{i_{2M}}), \\ & (\dots, x_1, x_2, \dots, x_q, a_{i_{2M}}), (\dots, a_{i_{2M}-1}, x_1, x_2, \dots, x_q), \\ & (\dots, x_1, x_2, \dots, x_q), (\dots, a_{i_{2M}-1}, x_1, x_2, \dots, x_q, a_{i_{2M}}), \\ & (\dots, a_{i_{2M}-1}, x_1, x_2, \dots, x_q, a_{i_{2M}}), (\dots, x_1, x_2, \dots, x_q). \end{aligned}$$

Hence $(\dots, b_{i_{2M}-2}, b_{i_{2M}-1}, b_{i_{2M}})$ must have one of the following four forms.

$$\begin{aligned} \text{(I)} & (\dots, a_{i_{2M}-1}, x_1, x_1, x_2, x_2, \dots, x_q, x_q, a_{i_{2M}}), \\ \text{(II)} & (\dots, x_1, a_{i_{2M}-1}, x_2, x_1, x_3, x_2, \dots, x_q, x_{q-1}, a_{i_{2M}}, x_q), \\ \text{(III)} & (\dots, a_{i_{2M}-1}, z, x_1, x_1, x_2, x_2, \dots, x_q, x_q, a_{i_{2M}}), \\ \text{(IV)} & (\dots, a_{i_{2M}-1}, z', x_1, z'', x_2, x_1, x_3, x_2, \dots, x_q, x_{q-1}, a_{i_{2M}}, x_q), \end{aligned}$$

where $a_{i_{2M}-1} > z$, $a_{i_{2M}-1} > z'' \geq z'$ and all entries in \dots are $< a_{i_{2M}-1}$. Correspondingly, $(\dots, \hat{b}_{i_{2M}-2}, \hat{b}_{i_{2M}-1}, \hat{b}_{i_{2M}})$ must have one of the following four forms.

$$\begin{aligned} \text{(I)} & (\dots, a_{i_{2M}-1} + h - q, x_1 + h - q, x_1 + h - q + 1, x_2 + h - q + 1, x_2 + h - q + 2, \dots, x_q + h - 1, x_q + h, a_{i_{2M}} + h), \\ \text{(II)} & (\dots, x_1 + h - q, a_{i_{2M}-1} + h - q, x_2 + h - q + 1, x_1 + h - q + 1, x_3 + h - q + 2, x_2 + h - q + 1, \dots, x_q + h - 1, x_{q-1} + h - 1, a_{i_{2M}} + h, x_q + h), \\ \text{(III)} & (\dots, a_{i_{2M}-1} + h - q - 1, z + h - q, x_1 + h - q, x_1 + h - q + 1, x_2 + h - q + 1, x_2 + h - q + 2, \dots, x_q + h - 1, x_q + h, a_{i_{2M}} + h), \\ \text{(IV)} & (\dots, a_{i_{2M}-1} + h - q - 1, z' + h - q - 1, x_1 + h - q, z'' + h - q, x_2 + h - q + 1, x_1 + h - q + 1, x_3 + h - q + 2, x_2 + h - q + 1, \dots, x_q + h - 1, x_{q-1} + h - 1, a_{i_{2M}} + h, x_q + h) \end{aligned}$$

where $h = i_{2M}/2$ and in case (III) and (IV), $a_{i_{2M}-1} + h - q$ is not an entry of $(\dots, \hat{b}_{i_{2M}-2}, \hat{b}_{i_{2M}-1}, \hat{b}_{i_{2M}})$.

Since $(\dots, \hat{a}_{i_{2M}-2}, \hat{a}_{i_{2M}-1}, \hat{a}_{i_{2M}})$ is given by (I) we see that $a_{i_{2M}-1} + h - q$ is an entry of $(\dots, \hat{a}_{i_{2M}-2}, \hat{a}_{i_{2M}-1}, \hat{a}_{i_{2M}})$. Using 1.3(b) we see that

$$\{\dots, \hat{a}_{i_{2M}-2}, \hat{a}_{i_{2M}-1}, \hat{a}_{i_{2M}}\} = (\dots, b_{i_{2M}-2}, b_{i_{2M}-1}, b_{i_{2M}})$$

as multisets. We see that cases (III) and (IV) cannot arise. Hence we must be in case (I) or (II). Thus

we have either

$$\begin{aligned} & (b_{i_{2M}-1}, b_{i_{2M}-1}+1, \dots, b_{i_{2M}-2}, b_{i_{2M}-1}, b_{i_{2M}}) \\ \text{(a)} \quad & = (a_{i_{2M}-1}, a_{i_{2M}-1}+1, \dots, a_{i_{2M}-2}, a_{i_{2M}-1}, a_{i_{2M}}) \end{aligned}$$

or

$$\begin{aligned} & (b_{i_{2M}-1}, b_{i_{2M}-1}+1, \dots, b_{i_{2M}-2}, b_{i_{2M}-1}, b_{i_{2M}}) \\ \text{(b)} \quad & = (a_{i_{2M}-1}+1, a_{i_{2M}-1}, a_{i_{2M}-1}+3, a_{i_{2M}-1}+2, \dots, a_{i_{2M}}, a_{i_{2M}-1}). \end{aligned}$$

1.5. Let $a' = (a_0, a_1, a_2, \dots, a_{i_{2M-1}-1})$, $b' = (b_0, b_1, b_2, \dots, b_{i_{2M-1}-1})$,
 $\hat{a}' = (a_0, a_1 + 1, a_2 + 1, a_3 + 2, a_4 + 2, \dots, a_{i_{2M-1}-1} + (i_{2M-1} - 1)/2)$,
 $\hat{b}' = (b_0, b_1 + 1, b_2 + 1, b_3 + 2, b_4 + 2, \dots, b_{i_{2M-1}-1} + (i_{2M-1} - 1)/2)$,

From $[\hat{b}] = [\hat{a}]$ and 1.3(b), 1.4(a),(b) we see that the multiset formed by the entries of \hat{a}' coincides with the multiset formed by the entries of \hat{b}' . Using the induction hypothesis we see that there exists $X' \subset \{1, 3, \dots, 2M - 3\}$ such that $b' = a'^{X'}$ where $a'^{X'}$ is defined in terms of a' , X' in the same way as a^X was defined (see 1.1) in terms of a , X . We set $X = X'$ if we are in case 1.4(a) and $X = X' \cup \{2M - 1\}$ if we are in case 1.4(b). Then we have $b = a^X$ (see 1.4(a),(b)), as required. This completes the proof of Lemma 1.2.

1.6. We shall use the notation of 1.1. Let \mathfrak{T} be the set of all unordered pairs $(\mathfrak{A}, \mathfrak{B})$ where $\mathfrak{A}, \mathfrak{B}$ are subsets of $\{0, 1, 2, \dots\}$ and $\mathfrak{A} \dot{\cup} \mathfrak{B} = (a_0, a_1, a_2, \dots, a_N)$ as multisets. For example, setting $\mathfrak{A}_\emptyset = (a_0, a_2, a_4, \dots, a_N)$, $\mathfrak{B}_\emptyset = (a_1, a_3, \dots, a_{N-1})$, we have $(\mathfrak{A}_\emptyset, \mathfrak{B}_\emptyset) \in \mathfrak{T}$. For any subset \mathfrak{a} of \mathcal{J} we consider

$$\mathfrak{A}_\mathfrak{a} = ((\mathcal{J} - \mathfrak{a}) \cap \mathfrak{A}_\emptyset) \cup (\mathfrak{a} \cap \mathfrak{B}_\emptyset) \cup (\mathfrak{A}_\emptyset \cap \mathfrak{B}_\emptyset),$$

$$\mathfrak{B}_\mathfrak{a} = ((\mathcal{J} - \mathfrak{a}) \cap \mathfrak{B}_\emptyset) \cup (\mathfrak{a} \cap \mathfrak{A}_\emptyset) \cup (\mathfrak{A}_\emptyset \cap \mathfrak{B}_\emptyset).$$

Then $(\mathfrak{A}_\mathfrak{a}, \mathfrak{B}_\mathfrak{a}) \in \mathfrak{T}$ and the map $\mathfrak{a} \mapsto (\mathfrak{A}_\mathfrak{a}, \mathfrak{B}_\mathfrak{a})$ induces a bijection $\bar{\mathcal{P}}(\mathcal{J}) \leftrightarrow \mathfrak{T}$. (Note that if $\mathfrak{a} = \emptyset$ then $(\mathfrak{A}_\mathfrak{a}, \mathfrak{B}_\mathfrak{a})$ agrees with the earlier definition of $(\mathfrak{A}_\emptyset, \mathfrak{B}_\emptyset)$.)

Let \mathfrak{T}' be the set of all $(\mathfrak{A}, \mathfrak{B}) \in \mathfrak{T}$ such that $|\mathfrak{A}| = |\mathfrak{A}_\emptyset|$, $|\mathfrak{B}| = |\mathfrak{B}_\emptyset|$.

Let $\mathcal{P}(\mathcal{J})_0$ be the subspace of $\mathcal{P}_{ev}(\mathcal{J})$ spanned by the 2-element subsets

$$\{a_{i_0}, a_{i_1}\}, \{a_{i_2}, a_{i_3}\}, \dots, \{a_{i_{2M-2}}, a_{i_{2M-1}}\}$$

of \mathcal{J} . Let $\mathcal{P}(\mathcal{J})_1$ be the subspace of $\mathcal{P}_{ev}(\mathcal{J})$ spanned by the 2-element subsets

$$\{a_{i_1}, a_{i_2}\}, \{a_{i_3}, a_{i_4}\}, \dots, \{a_{i_{2M-1}}, a_{i_{2M}}\}$$

of \mathcal{J} .

Let $\bar{\mathcal{P}}(\mathcal{J})_0$ (resp. $\bar{\mathcal{P}}(\mathcal{J})_1$) be the image of $\mathcal{P}(\mathcal{J})_0$ (resp. $\mathcal{P}(\mathcal{J})_1$) under the obvious map $\mathcal{P}(\mathcal{J}) \rightarrow \bar{\mathcal{P}}(\mathcal{J})$. Note that

(a) $\bar{\mathcal{P}}(\mathcal{J})_0$ and $\bar{\mathcal{P}}(\mathcal{J})_1$ are opposed Lagrangian subspaces of the symplectic vector space $\bar{\mathcal{P}}(\mathcal{J})$, (\cdot, \cdot) , (see 0.7); hence (\cdot, \cdot) defines an identification $\bar{\mathcal{P}}(\mathcal{J})_0 = \bar{\mathcal{P}}(\mathcal{J})_1^*$ where $\bar{\mathcal{P}}(\mathcal{J})_1^*$ is the vector space dual to $\bar{\mathcal{P}}(\mathcal{J})_1$.

Let \mathfrak{T}_0 (resp. \mathfrak{T}_1) be the subset of \mathfrak{T} corresponding to $\bar{\mathcal{P}}(\mathcal{J})_0$ (resp. $\bar{\mathcal{P}}(\mathcal{J})_1$) under the bijection $\bar{\mathcal{P}}(\mathcal{J}) \leftrightarrow \mathfrak{T}$. Note that $\mathfrak{T}_0 \subset \mathfrak{T}'$, $\mathfrak{T}_1 \subset \mathfrak{T}'$, $|\mathfrak{T}_0| = |\mathfrak{T}_1| = 2^M$.

For any $X \subset \{1, 3, \dots, 2M - 1\}$ we set $\mathfrak{a}_X = \cup_{s \in X} \{a_{i_s}, a_{i_{s+1}}\} \in \mathcal{P}(\mathcal{J})$. Then $(\mathfrak{A}_{\mathfrak{a}_X}, \mathfrak{B}_{\mathfrak{a}_X}) \in \mathfrak{T}_1$ is related to a^X in 1.1 as follows:

$$\mathfrak{A}_{\mathfrak{a}_X} = \{a_0^X, a_2^X, a_4^X, \dots, a_N^X\}, \mathfrak{B}_{\mathfrak{a}_X} = \{a_1^X, a_3^X, \dots, a_{N-1}^X\}.$$

1.7. We shall use the notation of 1.1. Let T be the set of all ordered pairs (A, B) where A is a subset of $\{0, 1, 2, \dots\}$, B is a subset of $\{1, 2, 3, \dots\}$, A contains no consecutive integers, B contains no consecutive integers, and $A \dot{\cup} B = (\hat{a}_0, \hat{a}_1, \hat{a}_2, \dots, \hat{a}_N)$ as multisets. For example, setting $A_\emptyset = (\hat{a}_0, \hat{a}_2, \hat{a}_4, \dots, \hat{a}_N)$, $B_\emptyset = (\hat{a}_1, \hat{a}_3, \dots, \hat{a}_{N-1})$, we have $(A_\emptyset, B_\emptyset) \in T$.

For any $(A, B) \in T$ we define (A^-, B^-) as follows: A^- consists of $x_0 < x_1 - 1 < x_2 - 2 < \dots < x_p - p$ where $x_0 < x_1 < \dots < x_p$ are the elements of A ; B^- consists of $y_1 - 1 < y_2 - 2 < \dots < y_q - q$ where $y_1 < y_2 < \dots < y_q$ are the elements of B .

We can enumerate the elements of T as in [L4, 11.5]. Let J be the set of all $c \in \mathbf{N}$ such that c appears exactly once in the sequence

$$(\hat{a}_0, \hat{a}_1, \hat{a}_2, \dots, \hat{a}_N) = (a_0, a_1 + 1, a_2 + 1, a_3 + 2, a_4 + 2, \dots, a_{N-1} + (N/2), a_N + (N/2)).$$

A nonempty subset I of J is said to be an interval if it is of the form $\{i, i + 1, i + 2, \dots, j\}$ with $i - 1 \notin J, j + 1 \notin J$ and with $i \neq 0$. Let \mathcal{I} be the set of intervals of J . For any $s \in \{1, 3, \dots, 2M - 1\}$, the set $I_s := \{\hat{a}_{i_s}, \hat{a}_{i_s+1}, \hat{a}_{i_s+2}, \dots, \hat{a}_{i_s+2m_s+1}\}$ is either a single interval or a union of intervals $I_s^1 \sqcup I_s^2 \sqcup \dots \sqcup I_s^{t_s}$ ($t_s \geq 2$) where $\hat{a}_{i_s} \in I_s^1$, $\hat{a}_{i_s+2m_s+1} \in I_s^{t_s}$, $|I_s^1|, |I_s^{t_s}|$ are odd, $|I_s^h|$ are even for $h \in [2, t_s - 1]$ and any element in I_s^e is $<$ than any element in $I_s^{e'}$ for $e < e'$. Let \mathcal{I}_s be the set of all $I \in \mathcal{I}$ such that $I \subset I_s$. We have a partition $\mathcal{I} = \sqcup_{s \in \{1, 3, \dots, 2M-1\}} \mathcal{I}_s$. Let H be the set of elements of $c \in J$ such that $c < a_{i_1}$ (that is such that c does not belong to any interval). For any subset $\alpha \subset \mathcal{I}$ we consider

$$A_\alpha = \cup_{I \in \mathcal{I} - \alpha} (I \cap A_\emptyset) \cup \cup_{I \in \alpha} (I \cap B_\emptyset) \cup (H \cap A_\emptyset) \cup (A_\emptyset \cap B_\emptyset),$$

$$B_\alpha = \cup_{I \in \mathcal{I} - \alpha} (I \cap B_\emptyset) \cup \cup_{I \in \alpha} (I \cap A_\emptyset) \cup (H \cap B_\emptyset) \cup (A_\emptyset \cap B_\emptyset).$$

Then $(A_\alpha, B_\alpha) \in T$ and the map $\alpha \mapsto (A_\alpha, B_\alpha)$ is a bijection $\mathcal{P}(\mathcal{I}) \leftrightarrow T$. (Note that if $\alpha = \emptyset$ then (A_α, B_α) agrees with the earlier definition of $(A_\emptyset, B_\emptyset)$.)

Let $T' = \{(A, B) \in T; |A| = |A_\emptyset|, |B| = |B_\emptyset|\}$, $T_1 = \{(A, B) \in T'; A^- \dot{\cup} B^- = A_\emptyset^- \dot{\cup} B_\emptyset^-\}$. Let $\mathcal{P}(\mathcal{I})'$ (resp. $\mathcal{P}(\mathcal{I})_1$) be the subset of $\mathcal{P}(\mathcal{I})$ corresponding to T' (resp. T_1) under the bijection $\mathcal{P}(\mathcal{I}) \leftrightarrow T$.

Now let X be a subset of $\{1, 3, \dots, 2M - 1\}$. Let $\alpha_X = \cup_{s \in X} \mathcal{I}_s \in \mathcal{P}(\mathcal{I})$. From the definitions we see that

$$(a) \ A_{\alpha_X}^- = \mathfrak{A}_{\alpha_X}, \ B_{\alpha_X}^- = \mathfrak{B}_{\alpha_X}$$

(notation of 1.6). In particular we have $(A_{\alpha_X}, B_{\alpha_X}) \in T_1$. Thus $|T_1| \geq 2^M$. Using Lemma 1.2 we see that

$$(b) \ |T_1| = 2^M \text{ and } T_1 \text{ consists of the pairs } (A_{\alpha_X}, B_{\alpha_X}) \text{ with } X \subset \{1, 3, \dots, 2M - 1\}.$$

Using (a), (b) we deduce:

$$(c) \ \text{The map } T_1 \rightarrow \mathfrak{T}_1 \text{ given by } (A, B) \mapsto (A^-, B^-) \text{ is a bijection.}$$

2. COMBINATORICS (CONTINUED)

2.1. Let $N \in \mathbf{N}$. Let

$$a := (a_0, a_1, a_2, \dots, a_N) \in \mathbf{N}^{N+1}$$

be such that $a_0 \leq a_1 \leq a_2 \leq \dots \leq a_N$, $a_0 < a_2 < a_4 < \dots$, $a_1 < a_3 < a_5 < \dots$ and such that the set $\mathcal{J} := \{i \in [0, N]; a_i \text{ appears exactly once in } a\}$ is nonempty. Now \mathcal{J} consists of $\mu + 1$ elements $i_0 < i_1 < \dots < i_\mu$ where $\mu \in \mathbf{N}$, $\mu = N \bmod 2$. We have $i_s = s \bmod 2$ for $s \in [0, \mu]$. Hence for any $s \in [0, \mu - 1]$ we have $i_{s+1} = i_s + 2m_s + 1$ for some $m_s \in \mathbf{N}$. Let \mathcal{E} be the set of $b := (b_0, b_1, b_2, \dots, b_N) \in \mathbf{N}^{N+1}$ such that $b_0 < b_2 < b_4 < \dots$, $b_1 < b_3 < b_5 < \dots$ and such that $[b] = [a]$ (we denote by $[b], [a]$ the multisets $\{b_0, b_1, \dots, b_N\}$, $\{a_0, a_1, \dots, a_N\}$). We have $a \in \mathcal{E}$. For $b \in \mathcal{E}$ we set

$$\overset{\circ}{b} = (\overset{\circ}{b}_0, \overset{\circ}{b}_1, \overset{\circ}{b}_2, \dots, \overset{\circ}{b}_N) = (b_0, b_1, b_2 + 1, b_3 + 1, b_4 + 2, b_5 + 2, \dots) \in \mathbf{N}^{N+1}.$$

Let $\overset{\circ}{[b]}$ be the multiset $\{\overset{\circ}{b}_0, \overset{\circ}{b}_1, \overset{\circ}{b}_2, \dots, \overset{\circ}{b}_N\}$. For any $s \in [0, \mu - 1] \in 2\mathbf{N}$ we define $a^{\{s\}} = (a_0^{\{s\}}, a_1^{\{s\}}, a_2^{\{s\}}, \dots, a_N^{\{s\}}) \in \mathcal{E}$ by

$$\begin{aligned} & (a_{i_s}^{\{s\}}, a_{i_s+1}^{\{s\}}, a_{i_s+2}^{\{s\}}, a_{i_s+3}^{\{s\}}, \dots, a_{i_s+2m_s}^{\{s\}}, a_{i_s+2m_s+1}^{\{s\}}) \\ & = (a_{i_s+1}, a_{i_s}, a_{i_s+3}, a_{i_s+2}, \dots, a_{i_s+2m_s+1}, a_{i_s+2m_s}) \end{aligned}$$

and $a_i^{\{s\}} = a_i$ if $i \in [0, N] - [i_s, i_{s+1}]$. More generally for a subset X of $[0, \mu - 1] \cap 2\mathbf{N}$ we define $a^X = (a_0^X, a_1^X, a_2^X, \dots, a_N^X) \in \mathcal{E}$ by $a_i^X = a_i^{\{s\}}$ if $s \in X$, $i \in [i_s, i_{s+1}]$, and $a_i^X = a_i$ for all other $i \in [0, N]$. Note that $[\overset{\circ}{a}^X] = [\overset{\circ}{a}]$. Conversely, we have the following result.

Lemma 2.2. *Let $b \in \mathcal{E}$ be such that $[\overset{\circ}{b}] = [\overset{\circ}{a}]$. Then there exists $X \subset [0, \mu - 1] \cap 2\mathbf{N}$ such that $b = a^X$.*

The proof is given in 2.3-2.5.

2.3. We argue by induction on μ . By the argument in 1.3 we have

- (a) $b_i = a_i$ for $i < i_0$,
- (b) $b_i = a_i$ for $i > i_\mu$.

If $\mu = 0$ we see that $b = a$ and there is nothing further to prove. In the rest of the proof we assume that $\mu \geq 1$.

2.4. From 2.3 we see that $(a_{i_0}, a_{i_0+1}, \dots, a_N) = (a_{i_0} < x_1 = x_1 < x_2 = x_2 < \dots < x_p = x_p < a_{i_1} < \dots)$ (for some p) has the same entries as $(b_{i_0}, b_{i_0+1}, \dots, b_N)$ (in some order). Hence the pair $(b_{i_0}, b_{i_0+2}, b_{i_0+4}, \dots), (b_{i_0+1}, b_{i_0+3}, b_{i_0+5}, \dots)$ must have one of the following four forms.

$$\begin{aligned} & (a_{i_0}, x_1, x_2, \dots, x_p, \dots), (x_1, x_2, \dots, x_p, a_{i_1}, \dots), \\ & (x_1, x_2, \dots, x_p, a_{i_1}, \dots), (a_{i_0}, x_1, x_2, \dots, x_p, \dots), \\ & (a_{i_0}, x_1, x_2, \dots, x_p, a_{i_1}, \dots), (x_1, x_2, \dots, x_p, \dots), \\ & (x_1, x_2, \dots, x_p, \dots), (a_{i_0}, x_1, x_2, \dots, x_p, a_{i_1}, \dots). \end{aligned}$$

Hence $(b_{i_0}, b_{i_0+1}, b_{i_0+2}, \dots, b_N)$ must have one of the following four forms.

$$(I) (a_{i_0}, x_1, x_1, x_2, x_2, \dots, x_p, x_p, a_{i_1}, \dots),$$

- (II) $(x_1, a_{i_0}, x_2, x_1, x_3, x_2, \dots, x_p, x_{p-1}, a_{i_1}, x_p, \dots)$,
 (III) $(a_{i_0}, x_1, x_1, x_2, x_2, \dots, x_p, x_p, z, a_{i_1}, \dots)$,
 (IV) $(x_1, a_{i_0}, x_2, x_1, x_3, x_2, \dots, x_p, x_{p-1}, z', x_p, z'', a_{i_1}, \dots)$

where $a_{i_1} < z$, $a_{i_1} < z' \leq z''$ and all entries in \dots are $> a_{i_1}$. Correspondingly, $(\overset{\circ}{b}_{i_0}, \overset{\circ}{b}_{i_0+1}, \overset{\circ}{b}_{i_0+2}, \dots, \overset{\circ}{b}_N)$ must have one of the following four forms.

- (I) $(a_{i_0} + h, x_1 + h, x_1 + h + 1, x_2 + h + 1, x_2 + h + 2, \dots, x_p + h + p - 1, x_p + h + p, a_{i_1} + h + p, \dots)$,
 (II) $(x_1 + h, a_{i_0} + h, x_2 + h + 1, x_1 + h + 1, x_3 + h + 2, x_2 + h + 2, \dots, x_p + h + p - 1, x_{p-1} + h + p - 1, a_{i_1} + h + p, x_p + h + p, \dots)$,
 (III) $(a_{i_0} + h, x_1 + h, x_1 + h + 1, x_2 + h + 1, x_2 + h + 2, \dots, x_p + h + p - 1, x_p + h + p, z + p, a_{i_1} + h + p + 1, \dots)$,

- $(x_1 + h, a_{i_0} + h, x_2 + h + 1, x_1 + h + 1, x_3 + h + 2, x_2 + h + 2, \dots, x_p + h + p - 1,$
 (IV) $x_{p-1} + h + p - 1, z' + h + p, x_p + h + p, z'' + h + p + 1, a_{i_1} + h + p + 1, \dots)$

where $h = i_0/2$ and in case (III) and (IV) $a_{i_1} + h + p$ is not an entry of $(\overset{\circ}{b}_{i_0}, \overset{\circ}{b}_{i_0+1}, \overset{\circ}{b}_{i_0+2}, \dots)$.

Since $(\overset{\circ}{a}_{i_0}, \overset{\circ}{a}_{i_0+1}, \overset{\circ}{a}_{i_0+2}, \dots)$ is given by (I) we see that $a_{i_1} + h + p$ is an entry of $(\overset{\circ}{a}_{i_0}, \overset{\circ}{a}_{i_0+1}, \overset{\circ}{a}_{i_0+2}, \dots)$. Using 2.3 we see that

$$\{\overset{\circ}{a}_{i_0}, \overset{\circ}{a}_{i_0+1}, \overset{\circ}{a}_{i_0+2}, \dots\} = \{\overset{\circ}{b}_{i_0}, \overset{\circ}{b}_{i_0+1}, \overset{\circ}{b}_{i_0+2}, \dots\}$$

as multisets. We see that cases (III) and (IV) cannot arise. Hence we must be in case (I) or (II). Thus

we have either

$$(a) \quad (b_{i_0}, b_{i_0+1}, b_{i_0+2}, \dots, b_{i_1}) = (a_{i_0}, a_{i_0+1}, a_{i_0+2}, \dots, a_{i_1})$$

or

$$(b) \quad (b_{i_0}, b_{i_0+1}, b_{i_0+2}, \dots, b_{i_1}) = (a_{i_0+1}, a_{i_0}, a_{i_0+3}, a_{i_0+2}, \dots, a_{i_1}, a_{i_1-1}).$$

. From 2.3 and (a),(b) we see that if $\mu = 1$ then Lemma 2.2 holds. Thus in the rest of the proof we can assume that $\mu \geq 2$.

2.5. Let $a' = (a_{i_1+1}, a_{i_1+2}, \dots, a_N)$, $b' = (b_{i_1+1}, b_{i_1+2}, \dots, b_N)$,

$$\overset{\circ}{a}' = (a_{i_1+1}, a_{i_1+2}, a_{i_1+3} + 1, a_{i_1+4} + 1, a_{i_1+5} + 2, a_{i_1+6} + 2, \dots),$$

$$\overset{\circ}{b}' = (b_{i_1+1}, b_{i_1+2}, b_{i_1+3} + 1, b_{i_1+4} + 1, b_{i_1+5} + 2, b_{i_1+6} + 2, \dots).$$

From $\overset{\circ}{[b]} = \overset{\circ}{[a]}$ and 2.3(a), 2.4(a), (b) we see that the multiset formed by the entries of $\overset{\circ}{a}'$ coincides with the multiset formed by the entries of $\overset{\circ}{b}'$. Using the induction hypothesis we see that there exists $X' \subset [2, \mu - 1] \cap 2\mathbf{N}$ such that $b' = a'^{X'}$ where $a'^{X'}$ is defined in terms of a' , X' in the same way as a^X (see 2.1) was defined in terms of a , X . We set $X = X'$ if we are in case 2.4(a) and $X = \{0\} \cup X'$ if we are in case 2.4(b). Then we have $b = a^X$ (see 2.4(a), (b)), as required. This completes the proof of Lemma 2.2.

2.6. We shall use the notation of 2.1. Let \mathfrak{T} be the set of all unordered pairs $(\mathfrak{A}, \mathfrak{B})$ where $\mathfrak{A}, \mathfrak{B}$ are subsets of $\{0, 1, 2, \dots\}$ and $\mathfrak{A} \dot{\cup} \mathfrak{B} = (a_0, a_1, a_2, \dots, a_N)$ as multisets. For example, setting $\mathfrak{A}_\emptyset = \{a_i; i \in [0, N] \cap 2\mathbf{N}\}$, $\mathfrak{B}_\emptyset = \{a_i; i \in [0, N] \cap (2\mathbf{N} + 1)\}$, we have $(\mathfrak{A}_\emptyset, \mathfrak{B}_\emptyset) \in \mathfrak{T}$. For any subset \mathfrak{a} of \mathcal{J} we consider

$$\mathfrak{A}_\mathfrak{a} = ((\mathcal{J} - \mathfrak{a}) \cap \mathfrak{A}_\emptyset) \cup (\mathfrak{a} \cap \mathfrak{B}_\emptyset) \cup (\mathfrak{A}_\emptyset \cap \mathfrak{B}_\emptyset),$$

$$\mathfrak{B}_\mathfrak{a} = ((\mathcal{J} - \mathfrak{a}) \cap \mathfrak{B}_\emptyset) \cup (\mathfrak{a} \cap \mathfrak{A}_\emptyset) \cup (\mathfrak{A}_\emptyset \cap \mathfrak{B}_\emptyset).$$

Then $(\mathfrak{A}_\mathfrak{a}, \mathfrak{B}_\mathfrak{a}) = (\mathfrak{A}_{\mathcal{J}-\mathfrak{a}}, \mathfrak{B}_{\mathcal{J}-\mathfrak{a}}) \in \mathfrak{T}$ and the map $\mathfrak{a} \mapsto (\mathfrak{A}_\mathfrak{a}, \mathfrak{B}_\mathfrak{a})$ induces a bijection $\bar{\mathcal{P}}(\mathcal{J}) \leftrightarrow \mathfrak{T}$. (Note that if $\mathfrak{a} = \emptyset$ then $(\mathfrak{A}_\mathfrak{a}, \mathfrak{B}_\mathfrak{a})$ agrees with the earlier definition of $(\mathfrak{A}_\emptyset, \mathfrak{B}_\emptyset)$.)

Let \mathfrak{T}' be the set of all $(\mathfrak{A}, \mathfrak{B}) \in \mathfrak{T}$ such that $|\mathfrak{A}| = |\mathfrak{A}_\emptyset|$, $|\mathfrak{B}| = |\mathfrak{B}_\emptyset|$. Let $\mathcal{P}(\mathcal{J})_1$ be the subspace of $\mathcal{P}(\mathcal{J})$ spanned by the following 2-element subsets of \mathcal{J} :

$$\{a_{i_1}, a_{i_2}\}, \{a_{i_3}, a_{i_4}\}, \dots, \{a_{i_{\mu-2}}, a_{i_{\mu-1}}\} \text{ (if } N \text{ is odd)}$$

or

$$\{a_{i_1}, a_{i_2}\}, \{a_{i_3}, a_{i_4}\}, \dots, \{a_{i_{\mu-1}}, a_{i_\mu}\} \text{ (if } N \text{ is even)}.$$

Let $\mathcal{P}(\mathcal{J})_0$ be the subspace of $\mathcal{P}(\mathcal{J})$ spanned by the following 2-element subsets of \mathcal{J} :

$$\{a_{i_0}, a_{i_1}\}, \{a_{i_2}, a_{i_3}\}, \dots, \{a_{i_{\mu-1}}, a_{i_\mu}\} \text{ (if } N \text{ is odd)}$$

or

$$\{a_{i_0}, a_{i_1}\}, \{a_{i_2}, a_{i_3}\}, \dots, \{a_{i_{\mu-2}}, a_{i_{\mu-1}}\} \text{ (if } N \text{ is even)}.$$

Let $\bar{\mathcal{P}}(\mathcal{J})_0$ (resp. $\bar{\mathcal{P}}(\mathcal{J})_1$) be the image of $\mathcal{P}(\mathcal{J})_0$ (resp. $\mathcal{P}(\mathcal{J})_1$) under the obvious map $\mathcal{P}(\mathcal{J}) \rightarrow \bar{\mathcal{P}}(\mathcal{J})$.

Note that

(a) $\bar{\mathcal{P}}(\mathcal{J})_0$ and $\bar{\mathcal{P}}(\mathcal{J})_1$ are opposed Lagrangian subspaces of the symplectic vector space $\bar{\mathcal{P}}_{ev}(\mathcal{J})$, $(,)$, (see 0.7); hence $(,)$ defines an identification $\bar{\mathcal{P}}(\mathcal{J})_1 = \bar{\mathcal{P}}(\mathcal{J})_0^*$ where $\bar{\mathcal{P}}(\mathcal{J})_0^*$ is the vector space dual to $\bar{\mathcal{P}}(\mathcal{J})_0$.

Let \mathfrak{T}_0 (resp. \mathfrak{T}_1) be the subset of \mathfrak{T} corresponding to $\bar{\mathcal{P}}(\mathcal{J})_0$ (resp. $\bar{\mathcal{P}}(\mathcal{J})_1$) under the bijection $\bar{\mathcal{P}}(\mathcal{J}) \leftrightarrow \mathfrak{T}$. Note that $\mathfrak{T}_0 \subset \mathfrak{T}'$, $\mathfrak{T}_1 \subset \mathfrak{T}'$, $|\mathfrak{T}_0| = |\mathfrak{T}_1| = 2^{\lfloor \mu/2 \rfloor}$.

For any $X \subset [0, \mu - 1] \cap 2\mathbf{N}$ we set $\mathfrak{a}_X = \cup_{s \in X} \{a_{i_s}, a_{i_{s+1}}\} \in \mathcal{P}(\mathcal{J})$. Then $(\mathfrak{A}_{\mathfrak{a}_X}, \mathfrak{B}_{\mathfrak{a}_X})$ is related to a^X in 2.1 as follows:

$$\mathfrak{A}_{\mathfrak{a}_X} = \{a_i^X; i \in [0, N] \cap 2\mathbf{N}\}, \mathfrak{B}_{\mathfrak{a}_X} = \{a_i^X; i \in [0, N] \cap (2\mathbf{N} + 1)\}.$$

2.7. We shall use the notation of 2.1. Let T be the set of all unordered pairs (A, B) where A is a subset of $\{0, 1, 2, \dots\}$, B is a subset of $\{1, 2, 3, \dots\}$, A contains no consecutive integers, B contains no consecutive integers, and $A \dot{\cup} B = (\overset{\circ}{a}_0, \overset{\circ}{a}_1, \overset{\circ}{a}_2, \dots, \overset{\circ}{a}_N)$ as multisets. For example, setting

$$A_\emptyset = \{\overset{\circ}{a}_i; i \in [0, N] \cap 2\mathbf{N}\}, B_\emptyset = (\overset{\circ}{a}_i; i \in [0, N] \cap (2\mathbf{N} + 1)),$$

we have $(A_\emptyset, B_\emptyset) \in T$.

For any $(A, B) \in T$ we define (A^-, B^-) as follows: A^- consists of $x_1 < x_2 - 1 < x_3 - 2 < \dots < x_p - p + 1$ where $x_1 < x_2 < \dots < x_p$ are the elements of A ; B^- consists of $y_1 < y_2 - 1 < \dots < y_q - q + 1$ where $y_1 < y_2 < \dots < y_q$ are the elements of B .

We can enumerate the elements of T as in [L4, 11.5]. Let J be the set of all $c \in \mathbf{N}$ such that c appears exactly once in the sequence

$$(\overset{\circ}{a}_0, \overset{\circ}{a}_1, \overset{\circ}{a}_2, \dots, \overset{\circ}{a}_N) = (a_0, a_1, a_2 + 1, a_3 + 1, a_4 + 2, a_5 + 2, \dots).$$

A nonempty subset I of J is said to be an interval if it is of the form $\{i, i + 1, i + 2, \dots, j\}$ with $i - 1 \notin J, j + 1 \notin J$. Let \mathcal{I} be the set of intervals of J . For any $s \in [0, \mu - 1] \cap 2\mathbf{N}$, the set $I_s := \{\overset{\circ}{a}_{i_s}, \overset{\circ}{a}_{i_s+1}, \overset{\circ}{a}_{i_s+2}, \dots, \overset{\circ}{a}_{i_s+2m_s+1}\}$ is either a single interval or a union of intervals $I_s^1 \sqcup I_s^2 \sqcup \dots \sqcup I_s^{t_s}$ ($t_s \geq 2$) where $\overset{\circ}{a}_{i_s} \in I_s^1$, $\overset{\circ}{a}_{i_s+2m_s+1} \in I_s^{t_s}$, $|I_s^1|, |I_s^{t_s}|$ are odd, $|I_s^h|$ are even for $h \in [2, t_s - 1]$ and any element in I_s^e is $<$ than any element in $I_s^{e'}$ for $e < e'$. Let \mathcal{I}_s be the set of all $I \in \mathcal{I}$ such that $I \subset I_s$. We have a partition $\mathcal{I} = \sqcup_{s \in [0, \mu - 1] \cap 2\mathbf{N}} \mathcal{I}_s$. For any subset $\alpha \subset \mathcal{I}$ we consider

$$A_\alpha = \cup_{I \in \mathcal{I} - \alpha} (I \cap A_\emptyset) \cup \cup_{I \in \alpha} (I \cap B_\emptyset) \cup (A_\emptyset \cap B_\emptyset),$$

$$B_\alpha = \cup_{I \in \mathcal{I} - \alpha} (I \cap B_\emptyset) \cup \cup_{I \in \alpha} (I \cap A_\emptyset) \cup (A_\emptyset \cap B_\emptyset).$$

Then $(A_\alpha, B_\alpha) \in T$ and the map $\alpha \mapsto (A_\alpha, B_\alpha)$ is a bijection $\bar{\mathcal{P}}(\mathcal{I}) \leftrightarrow T$. (Note that if $\alpha = \emptyset$ then (A_α, B_α) agrees with the earlier definition of $(A_\emptyset, B_\emptyset)$.)

Let $T' = \{(A, B) \in T; |A| = |A_\emptyset|, |B| = |B_\emptyset|\}$, $T_1 = \{(A, B) \in T'; A^- \dot{\cup} B^- = A_\emptyset^- \dot{\cup} B_\emptyset^-\}$. Let $\bar{\mathcal{P}}(\mathcal{I})'$ (resp. $\bar{\mathcal{P}}(\mathcal{I})_1$) be the subset of $\bar{\mathcal{P}}(\mathcal{I})$ corresponding to T' (resp. T_1) under the bijection $\bar{\mathcal{P}}(\mathcal{I}) \leftrightarrow T$.

Now let X be a subset of $[0, \mu - 1] \cap 2\mathbf{N}$. Let $\alpha_X = \cup_{s \in X} \mathcal{I}_s \in \mathcal{P}(\mathcal{I})$. From the definitions we see that

$$(a) \ A_{\alpha_X}^- = \mathfrak{A}_{\alpha_X}, B_{\alpha_X}^- = \mathfrak{B}_{\alpha_X}$$

(notation of 2.6). In particular we have $(A_{\alpha_X}, B_{\alpha_X}) \in T_1$. Thus $|T_1| \geq 2^{\lfloor \mu/2 \rfloor}$. Using Lemma 2.2 we see that

(b) $|T_1| = 2^{\lfloor \mu/2 \rfloor}$ and T_1 consists of the pairs $(A_{\alpha_X}, B_{\alpha_X})$ with $X \subset [0, \mu - 1] \cap 2\mathbf{N}$.

Using (a), (b) we deduce:

(c) The map $T_1 \rightarrow \mathfrak{T}_1$ given by $(A, B) \mapsto (A^-, B^-)$ is a bijection.

3. PROOF OF THEOREM 0.4 AND OF COROLLARY 0.5

3.1. If G is simple adjoint of type A_n , $n \geq 1$, then 0.4 and 0.5 are obvious: we have $A(u) = \{1\}$, $\bar{A}(u) = \{1\}$.

3.2. Assume that $G = Sp_{2n}(\mathbf{k})$ where $n \geq 2$. Let N be a sufficiently large even integer. Now $u : \mathbf{k}^{2n} \rightarrow \mathbf{k}^{2n}$ has i_e Jordan blocks of size e ($e = 1, 2, 3, \dots$). Here i_1, i_3, i_5, \dots are even. Let $\Delta = \{e \in \{2, 4, 6, \dots\}; i_e \geq 1\}$. Then $A(u)$ can be identified in the standard way with $\mathcal{P}(\Delta)$. Hence the group of characters $\hat{A}(u)$ of $A(u)$ (which may be canonically identified with the F_2 -vector space dual to $\mathcal{P}(\Delta)$) may be also canonically identified with $\mathcal{P}(\Delta)$ itself (so that the basis given by the one element subsets of Δ is self-dual).

To the partition $1i_1 + 2i_2 + 3i_3 + \dots$ of $2n$ we associate a pair (A, B) as in [L4, 11.6] (with $N, 2m$ replaced by $2n, N$). We have $A = (\hat{a}_0, \hat{a}_2, \hat{a}_4, \dots, \hat{a}_N)$, $B = (\hat{a}_1, \hat{a}_3, \dots, \hat{a}_{N-1})$, where $\hat{a}_0 \leq \hat{a}_1 \leq \hat{a}_2 \leq \dots \leq \hat{a}_N$ is obtained from a sequence $a_0 \leq a_1 \leq a_2 \leq \dots \leq a_N$ as in 1.1. (Here we use that C is special.) Now the definitions and results in §1 are applicable. As in [L3, 4.5] the family \mathcal{F} is in canonical bijection with \mathfrak{T}' in 1.6.

We arrange the intervals in \mathcal{I} in increasing order $I_{(1)}, I_{(2)}, \dots, I_{(f)}$ (any element in $I_{(1)}$ is smaller than any element in $I_{(2)}$, etc.). We arrange the elements of Δ in increasing order $e_1 < e_2 < \dots < e_{f'}$; then $f = f'$ and we have a bijection $\mathcal{I} \leftrightarrow \Delta$, $I_{(h)} \leftrightarrow e_h$; moreover we have $|I_{(h)}| = i_{e_h}$ for $h \in [1, f]$; see [L4, 11.6]. Using this bijection we see that $A(u)$ and $\hat{A}(u)$ are identified with the F_2 -vector space $\mathcal{P}(\mathcal{I})$ with basis given by the one element subsets of \mathcal{I} . Let $\pi : \mathcal{P}(\mathcal{I}) \rightarrow \mathcal{P}(\mathcal{I})_1^*$ (with $\mathcal{P}(\mathcal{I})_1^*$ as in 1.7(c)) be the (surjective) F_2 -linear map which to $X \subset \mathcal{I}$ associates the linear form $L \mapsto |X \cap L| \pmod{2}$ on $\mathcal{P}(\mathcal{I})_1$. We will show that

(a) $\ker \pi = \mathcal{K}(u)$ ($\mathcal{K}(u)$ as in 0.1).

We identify $\text{Irr}_C W$ with T' (see 1.7) via the restriction of the bijection in [L4, (12.2.4)] (we also use the description of the Springer correspondence in [L4, 12.3]). Under this identification the subset $\text{Irr}_C^* W$ of $\text{Irr}_C W$ becomes the subset T_1 (see 1.7) of T' . Via the identification $\mathcal{P}(\mathcal{I})' \leftrightarrow T'$ in 1.7 and $\hat{A}(u) \leftrightarrow \mathcal{P}(\mathcal{I})$ (see above), the map $E \mapsto \mathcal{V}_E$ from T' to $\hat{A}(u)$ becomes the obvious imbedding $\mathcal{P}(\mathcal{I})' \rightarrow \mathcal{P}(\mathcal{I})$ (we use again [L4, 12.3]). By definition, $\mathcal{K}(u)$ is the set of all $X \in \mathcal{P}(\mathcal{I})$ such that for any $L \in \mathcal{P}(\mathcal{I})_1$ we have $|X \cap L| = 0 \pmod{2}$. Thus, (a) holds.

Using (a) we have canonically $\bar{A}(u) = \mathcal{P}(\mathcal{I})_1^*$ via π . We define an F_2 -linear map $\mathcal{P}(\mathcal{I})_1 \rightarrow \bar{\mathcal{P}}(\mathcal{J})_1$ (see 1.6) by $I_s \mapsto \{a_{i_s}, a_{i_s+1}\}$ for $s \in \{1, 3, \dots, 2M-1\}$ (I_s as in 1.7). This is an isomorphism; it corresponds to the bijection 1.7(c) under the identification $T_1 \leftrightarrow \mathcal{P}(\mathcal{I})_1$ in 1.7 and the identification $\mathfrak{T}_1 \leftrightarrow \bar{\mathcal{P}}(\mathcal{J})_1$ in 1.6. Hence we can identify $\mathcal{P}(\mathcal{I})_1^*$ with $\bar{\mathcal{P}}(\mathcal{J})_1^*$ and with $\bar{\mathcal{P}}(\mathcal{J})_0$ (see 1.6(a)). We obtain an identification $\bar{A}(u) = \bar{\mathcal{P}}(\mathcal{J})_0$.

By [L3, 4.5] we have $\mathbf{X}_{\mathcal{F}} = \bar{\mathcal{P}}(\mathcal{J})$. Using 1.6(a) we see that $\bar{\mathcal{P}}(\mathcal{J}) = M(\bar{\mathcal{P}}(\mathcal{J})_0) = M(\bar{A}(u))$ canonically so that 0.4 holds in our case. From the arguments above we see that in our case 0.5 follows from 1.7(c).

3.3. Assume that $G = SO_n(\mathbf{k})$ where $n \geq 7$. Let N be a sufficiently large integer

such that $N = n \bmod 2$. Now $u : \mathbf{k}^n \rightarrow \mathbf{k}^n$ has i_e Jordan blocks of size e ($e = 1, 2, 3, \dots$). Here i_2, i_4, i_6, \dots are even. Let $\Delta = \{e \in \{1, 3, 5, \dots\}; i_e \geq 1\}$. If $\Delta = \emptyset$ then $A(u) = \{1\}$, $\hat{A}(u) = \{1\}$ and $\mathcal{G}_{\mathcal{F}} = \{1\}$ so that the result is trivial.

In the remainder of this subsection we assume that $\Delta \neq \emptyset$. Then $A(u)$ can be identified in the standard way with the F_2 -subspace $\mathcal{P}_{ev}(\Delta)$ of $\mathcal{P}(\Delta)$ and the group of characters $\hat{A}(u)$ of $A(u)$ (which may be canonically identified with the F_2 -vector space dual to $A(u)$) becomes $\bar{\mathcal{P}}(\Delta)$; the obvious pairing $A(u) \times \hat{A}(u) \rightarrow F_2$ is induced by the inner product $L, L' \mapsto |L \cap L'| \bmod 2$ on $\mathcal{P}(\Delta)$.

To the partition $1i_1 + 2i_2 + 3i_3 + \dots$ of n we associate a pair (A, B) as in [L4, 11.7] (with N, M replaced by n, N). We have $A = \{\overset{\circ}{a}_i; i \in [0, N] \cap 2\mathbf{N}\}$, $B = \{\overset{\circ}{a}_i; i \in [0, N] \cap (2\mathbf{N} + 1)\}$ where $\overset{\circ}{a}_0 \leq \overset{\circ}{a}_1 \leq \overset{\circ}{a}_2 \leq \dots \leq \overset{\circ}{a}_N$ is obtained from a sequence $a_0 \leq a_1 \leq a_2 \leq \dots \leq a_N$ as in 2.1. (Here we use that C is special.) Now the definitions and results in §2 are applicable. As in [L3, 4.5] (if N is even) or [L3, 4.6] (if N is odd) the family \mathcal{F} is in canonical bijection with \mathfrak{T}' in 2.6.

We arrange the intervals in \mathcal{I} in increasing order $I_{(1)}, I_{(2)}, \dots, I_{(f)}$ (any element in $I_{(1)}$ is smaller than any element in $I_{(2)}$, etc.). We arrange the elements of Δ in increasing order $e_1 < e_2 < \dots < e_{f'}$; then $f = f'$ and we have a bijection $\mathcal{I} \leftrightarrow \Delta$, $I_{(h)} \leftrightarrow e_h$; moreover we have $|I_{(h)}| = i_{e_h}$ for $h \in [1, f]$; see [L4, 11.7]. Using this bijection we see that $A(u)$ is identified with $\mathcal{P}_{ev}(\mathcal{I})$ and $\hat{A}(u)$ is identified with $\bar{\mathcal{P}}(\mathcal{I})$. For any $X \in \mathcal{P}_{ev}(\mathcal{I})$, the assignment $L \mapsto |X \cap L| \bmod 2$ can be viewed as an element of $\bar{\mathcal{P}}(\mathcal{I})_1^*$ (the dual space of $\bar{\mathcal{P}}(\mathcal{I})_1$ in 2.7 which by 2.7(b) is an F_2 -vector space of dimension $2^{\lfloor \mu/2 \rfloor}$). This induces a (surjective) F_2 -linear map $\pi : \mathcal{P}_{ev}(\mathcal{I}) \rightarrow \bar{\mathcal{P}}(\mathcal{I})_1^*$. We will show that

(a) $\ker \pi = \mathcal{K}(u)$ ($\mathcal{K}(u)$ as in 0.1).

We identify $\text{Irr}_C W$ with T' (see 2.7) via the restriction of the bijection in [L4, (13.2.5)] if N is odd or [L4, (13.2.6)] if N is even (we also use the description of the Springer correspondence in [L4, 13.3]). Under this identification the subset $\text{Irr}_C^* W$ of $\text{Irr}_C W$ becomes the subset T_1 (see 2.7) of T' . Via the identification $\bar{\mathcal{P}}(\mathcal{I})' \leftrightarrow T'$ in 2.7 and $\hat{A}(u) \leftrightarrow \bar{\mathcal{P}}(\mathcal{I})$ (see above), the map $E \mapsto \mathcal{V}_E$ from T' to $\hat{A}(u)$ becomes the obvious imbedding $\bar{\mathcal{P}}(\mathcal{I})_0 \rightarrow \bar{\mathcal{P}}(\mathcal{I})$ (we use again [L4, 13.3]). By definition, $\mathcal{K}(u)$ is the set of all $X \in \mathcal{P}_{ev}(\mathcal{I})$ such that for any $L \in \mathcal{P}(\mathcal{I})$ representing a vector in $\bar{\mathcal{P}}(\mathcal{I})_1$ we have $|X \cap L| = 0 \bmod 2$. Thus, (a) holds.

Using (a) we have canonically $\hat{A}(u) = \bar{\mathcal{P}}(\mathcal{I})_1^*$ via π . We have an F_2 -linear map $\bar{\mathcal{P}}(\mathcal{I})_1 \rightarrow \bar{\mathcal{P}}(\mathcal{J})_0$ (see 2.6) induced by $I_s \mapsto \{a_{i_s}, a_{i_{s+1}}\}$ for $s \in [0, \mu - 1] \cap 2\mathbf{N}$ (I_s as in 2.7). This is an isomorphism; it corresponds to the bijection 2.7(c) under the identification $T_1 \leftrightarrow \bar{\mathcal{P}}(\mathcal{I})_1$ in 2.7 and the identification $\mathfrak{T}_1 \leftrightarrow \bar{\mathcal{P}}(\mathcal{J})_0$ in 2.6. Hence we can identify $\bar{\mathcal{P}}(\mathcal{I})_1^*$ with $\bar{\mathcal{P}}(\mathcal{J})_0^*$ and with $\bar{\mathcal{P}}(\mathcal{J})_1$ (see 2.6(a)). We obtain an identification $\hat{A}(u) = \bar{\mathcal{P}}(\mathcal{J})_1$.

By [L3, 4.6] we have $\mathbf{X}_{\mathcal{F}} = \bar{\mathcal{P}}_{ev}(\mathcal{J})$. Using 2.6(a) we see that $\bar{\mathcal{P}}(\mathcal{J}) = M(\bar{\mathcal{P}}(\mathcal{J})_1) = M(\hat{A}(u))$ canonically so that 0.4 holds in our case. From the arguments above we see that in our case 0.5 follows from 2.7(c). ■

3.4. In 3.5-3.9 we consider the case where G is simple adjoint of exceptional type.

In each case we list the elements of the set $\text{Irr}_C W$ for each special unipotent class C of G ; the elements of $\text{Irr}_C W - \text{Irr}^* C W$ are enclosed in $[\]$. (The notation for the various C is as in [Sp2]; the notation for the objects of $\text{Irr} W$ is as in [Sp2] (for type E_n) and as in [L3, 4.10] for type F_4 .) In each case the structure of $A(u)$, $\bar{A}(u)$ (for $u \in C$) is indicated; here S_n denotes the symmetric group in n letters. The order in which we list the objects in $\text{Irr}_C W$ corresponds to the following order of the irreducible representations of $A(u) = S_n$:

$1, \epsilon$ ($n = 2$); $1, r, \epsilon$ ($n = 3, G \neq G_2$); $1, r$ ($n = 3, G = G_2$); $1, \lambda^1, \lambda^2, \sigma$ ($n = 4$);
 $1, \nu, \lambda^1, \nu', \lambda^2, \lambda^3$ ($n = 5$)

(notation of [L3, 4.3]). Now 0.4 and 0.5 follow in our case from the tables in 3.5-3.9 and the definitions in [L3, 4.8-4.13].

3.5. Assume that G is of type E_8 .

$\text{Irr}_{E_8} W = \{1_0\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{E_8(a_1)} W = \{8_1\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{E_8(a_2)} W = \{35_2\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{E_7 A_1} W = \{112_3, 28_8\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{D_8} W = \{210_4, 160_7\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{E_7(a_1) A_1} W = \{560_5, [50_8]\}$; $A(u) = S_2, \bar{A}(u) = \{1\}$
 $\text{Irr}_{E_7(a_1)} W = \{567_6\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{D_8(a_1)} W = \{700_6, 300_8\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{E_7(a_2) A_1} W = \{1400_7, 1008_9, 56_{19}\}$; $A(u) = S_3, \bar{A}(u) = S_3$
 $\text{Irr}_{A_8} W = \{1400_8, 1575_{10}, 350_{14}\}$; $A(u) = S_3, \bar{A}(u) = S_3$
 $\text{Irr}_{D_7(a_1)} W = \{3240_9, [1050_{10}]\}$; $A(u) = S_2, \bar{A}(u) = \{1\}$
 $\text{Irr}_{D_8(a_3)} W = \{2240_{10}, [175_{12}], 840_{13}\}$; $A(u) = S_3, \bar{A}(u) = S_2$
 $\text{Irr}_{D_6 A_1} W = \{2268_{10}, 1296_{13}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{E_6(a_1) A_1} W = \{4096_{11}, 4096_{12}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{E_6} W = \{525_{12}\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{D_7(a_2)} W = \{4200_{12}, 3360_{13}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{E_6(a_1)} W = \{2800_{13}, 2100_{16}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{D_5 A_2} W = \{4536_{13}, [840_{14}]\}$; $A(u) = S_2, \bar{A}(u) = \{1\}$
 $\text{Irr}_{D_6(a_1) A_1} W = \{6075_{14}, [700_{16}]\}$; $A(u) = S_2, \bar{A}(u) = \{1\}$
 $\text{Irr}_{A_6 A_1} W = \{2835_{14}\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{A_6} W = \{4200_{15}\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{D_6(a_1)} W = \{5600_{15}, 2400_{17}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{2A_4} W = \{4480_{16}, 4536_{18}, 5670_{18}, 1400_{20}, 1680_{22}, 70_{32}\}$; $A(u) = S_5, \bar{A}(u) = S_5$
 $\text{Irr}_{D_5} W = \{2100_{20}\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{(A_5 A_1)''} W = \{5600_{21}, 2400_{23}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{D_4 A_2} W = \{4200_{15}, [168_{24}]\}$; $A(u) = S_2, \bar{A}(u) = \{1\}$
 $\text{Irr}_{A_4 A_2 A_1} W = \{2835_{22}\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{A_4 A_2} W = \{4536_{23}\}$; $A(u) = \{1\}, \bar{A}(u) = \{1\}$
 $\text{Irr}_{D_5(a_1)} W = \{2800_{25}, 2100_{28}\}$; $A(u) = S_2, \bar{A}(u) = S_2$
 $\text{Irr}_{A_4 2A_1} W = \{4200_{24}, 3360_{25}\}$; $A(u) = S_2, \bar{A}(u) = S_2$

$$\begin{aligned}
\text{Irr}_{D_4} W &= \{525_{36}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_4 A_1} W &= \{4096_{26}, 4096_{27}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{A_4} W &= \{2268_{30}, 1296_{33}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{D_4(a_1)A_2} &= \{2240_{28}, 840_{31}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{A_3 A_2} W &= \{3240_{31}, [972_{32}]\}; A(u) = S_2, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_4(a_1)A_1} W &= \{1400_{32}, 1575_{34}, 350_{38}\}; A(u) = S_3, \bar{A}(u) = S_3 \\
\text{Irr}_{D_4(a_1)} W &= \{1400_{37}, 1008_{39}, 564_9\}; A(u) = S_3, \bar{A}(u) = S_3 \\
\text{Irr}_{2A_2} W &= \{700_{42}, 300_{44}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{A_3} W &= \{567_{46}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2 2A_1} W &= \{560_{47}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2 A_1} W &= \{210_{52}, 160_{55}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{A_2} W &= \{112_{63}, 28_{68}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{2A_1} W &= \{35_{74}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_1} W &= \{8_{91}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{\emptyset} W &= \{1_{120}\}; A(u) = \{1\}, \bar{A}(u) = \{1\}
\end{aligned}$$

3.6. Assume that G is adjoint of type E_7 .

$$\begin{aligned}
\text{Irr}_{E_7} W &= \{1_0\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{E_7(a_1)} W &= \{7_1\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{E_7(a_2)} W &= \{27_2\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_6 A_1} W &= \{56_3, 21_6\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{E_6} W &= \{21_3\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{E_6(a_1)} W &= \{120_4, 105_5\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{D_6(a_1)A_1} W &= \{189_5, [15_7]\}; A(u) = S_2, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_6(a_1)} W &= \{210_6\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_6} W &= \{105_6\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_5 A_1} W &= \{168_6\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_5} W &= \{189_7\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_6(a_2)A_1} W &= \{315_7, 280_9, 35_{13}\}; A(u) = S_3, \bar{A}(u) = S_3 \\
\text{Irr}_{(A_5 A_1)'} W &= \{405_8, 189_{10}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{D_5(a_1)A_1} W &= \{378_9\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_4 A_2} W &= \{210_{10}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_5(a_1)} W &= \{420_{10}, 336_{11}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{A_5''} W &= \{105_{12}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_4 A_1} W &= \{512_{11}, 512_{12}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{D_4} W &= \{105_{15}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_4} W &= \{420_{13}, 336_{14}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{A_3 A_2 A_1} W &= \{210_{13}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_3 A_2} W &= \{378_{14}, [84_{15}]\}; A(u) = S_2, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_4(a_1)A_1} W &= \{405_{15}, 189_{17}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{D_4(a_1)} W &= \{315_{16}, 280_{18}, 35_{22}\}; A(u) = S_3, \bar{A}(u) = S_3 \\
\text{Irr}_{(A_3 A_1)''} W &= \{189_{20}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{2A_2} W &= \{168_{21}\}; A(u) = \{1\}, \bar{A}(u) = \{1\}
\end{aligned}$$

$$\begin{aligned}
\text{Irr}_{A_2 3 A_1} W &= \{105_{21}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_3} W &= \{210_{21}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2 2 A_1} W &= \{189_{22}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2 A_1} W &= \{120_{25}, 105_{26}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{3 A_1'} W &= \{21_{36}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2} W &= \{56_{30}, 21_{33}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{2 A_1} W &= \{27_{37}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_1} W &= \{7_{46}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{\emptyset} W &= \{1_{63}\}; A(u) = \{1\}, \bar{A}(u) = \{1\}
\end{aligned}$$

3.7. Assume that G is adjoint of type E_6 .

$$\begin{aligned}
\text{Irr}_{E_6} W &= \{1_0\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{E_6(a_1)} W &= \{6_1\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_5} W &= \{20_2\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_5 A_1} W &= \{30_3, 15_5\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{D_5(a_1)} W &= \{64_4\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_4 A_1} W &= \{60_5\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_4} W &= \{81_6\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_4} W &= \{24_6\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{D_4(a_1)} W &= \{80_7, 90_8, 20_{10}\}; A(u) = S_3, \bar{A}(u) = S_3 \\
\text{Irr}_{2 A_2} W &= \{24_{12}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_3} W &= \{81_{10}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2 2 A_1} W &= \{60_{11}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2 A_1} W &= \{64_{13}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2} W &= \{30_{15}, 15_{17}\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{2 A_1} W &= \{20_{20}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_1} W &= \{6_{25}\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{\emptyset} W &= \{1_{36}\}; A(u) = \{1\}, \bar{A}(u) = \{1\}
\end{aligned}$$

3.8. Assume that G is of type F_4 .

$$\begin{aligned}
\text{Irr}_{F_4} W &= \{1_1\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{F_4(a_1)} W &= \{4_2, 2_3\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{F_4(a_2)} W &= \{9_1\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{B_3} W &= \{8_1\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{C_3} W &= \{8_3\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{F_4(a_3)} W &= \{12_1, 9_3, 6_2, 1_3\}; A(u) = S_4, \bar{A}(u) = S_4 \\
\text{Irr}_{\tilde{A}_2} W &= \{8_2\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_2} W &= \{8_4, [1_2]\}; A(u) = S_2, \bar{A}(u) = \{1\} \\
\text{Irr}_{A_1 \tilde{A}_1} W &= \{9_4\}; A(u) = \{1\}, \bar{A}(u) = \{1\} \\
\text{Irr}_{\tilde{A}_1} W &= \{4_5, 2_2\}; A(u) = S_2, \bar{A}(u) = S_2 \\
\text{Irr}_{\emptyset} W &= \{1_4\}; A(u) = \{1\}, \bar{A}(u) = \{1\}
\end{aligned}$$

3.9. Assume that G is of type G_2 .

$\text{Irr}_{G_2}W$ is the unit representation; $A(u) = \{1\}, \bar{A}(u) = \{1\}$

$\text{Irr}_{G_2(a_1)}W$ consists of the reflection representation and the one dimensional representation on which the reflection with respect to a long (resp.short) simple coroot acts nontrivially (resp. trivially); $A(u) = S_3, \bar{A}(u) = S_3$

$\text{Irr}_\emptyset W$ is the sign representation; $A(u) = \{1\}, \bar{A}(u) = \{1\}$

3.10. This completes the proof of Theorem 0.4 and that of Corollary 0.5.

We note that the definition of $\mathcal{G}_{\mathcal{F}}$ given in [L3] (for type C_n, B_n) is $\bar{\mathcal{P}}(\mathcal{J})_1$ (in the setup of 3.2) and $\bar{\mathcal{P}}(\mathcal{J})_0$ (in the setup of 3.3) which is noncanonically isomorphic to $\bar{A}(u)$, unlike the definition adopted here that is, $\bar{\mathcal{P}}(\mathcal{J})_0$ (in the setup of 3.2) and $\bar{\mathcal{P}}(\mathcal{J})_1$ (in the setup of 3.3) which makes $\mathcal{G}_{\mathcal{F}}$ canonically isomorphic to $\bar{A}(u)$.

REFERENCES

- [AL] D. Alvis and G. Lusztig, *On Springer's correspondence for simple groups of type E_n ($n = 6, 7, 8$)*, Math. Proc. Camb. Phil. Soc. **92** (1982), 65-72.
- [BM] J. Borho and R. MacPherson, *Représentations des groupes de Weyl et homologie d'intersection pour les variétés nilpotents*, C.R. Acad. Sci. Paris **292** (1981), 707-710.
- [L1] G. Lusztig, *Unipotent representations of a finite Chevalley group of type E_8* , Quart.J. Math. Oxford **30** (1979), 315-338.
- [L2] G. Lusztig, *A class of irreducible representations of a Weyl group*, Proc. Kon. Nederl. Akad.(A) **82** (1979), 323-335.
- [L3] G. Lusztig, *Characters of reductive groups over a finite field*, Ann. Math. Studies, vol. 107, Princeton Univ.Press, 1984.
- [L4] G. Lusztig, *Intersection cohomology complexes on a reductive group*, Invent.Math. **75** (1984), 205-272.
- [L5] G. Lusztig, *Introduction to character sheaves*, Proc. Symp. Pure Math. **47** (1987), Amer. Math. Soc., 165-189.
- [L6] G. Lusztig, *A unipotent support for irreducible representations*, Adv. Math **94** (1992), 139-179.
- [S1] T. Shoji, *On the Springer representations of the Weyl groups of classical algebraic groups*, Comm.in Alg. **7** (1979), 1713-1745; **7** (1979), 2027-2033.
- [S2] T. Shoji, *On the Springer representations of Chevalley groups of type F_4* , Comm.in Alg. **8** (1980), 409-440.
- [Sp1] J. Spaltenstein, *Appendix*, Math. Proc. Camb. Phil. Soc. **92** (1982), 73-78.
- [Sp2] J. Spaltenstein, *On the generalized Springer correspondence for exceptional groups*, Algebraic groups and related topics, Adv.Studies in Pure Math., vol. 6, 1985, pp. 317-338.
- [Spr] T. A. Springer, *Trigonometric sums, Green functions of finite groups and representations of Weyl groups*, Invent.Math. **36** (1976), 173-207.

DEPARTMENT OF MATHEMATICS, M.I.T., CAMBRIDGE, MA 02139