# 1 Genetic diversity and fitness in small populations of partially asexual, self-

2	incompatible plants
3	
4	Miguel Navascués <sup>1,2</sup> , Solenn Stoeckel <sup>3</sup> and Stéphanie Mariette <sup>4</sup>
5	
6	<sup>1</sup> Unidad de Genética, Centro de Investigación Forestal, INIA, Carretera de La Coruña km 7.5,
7	28040 Madrid, Spain
8	<sup>2</sup> Équipe Éco-évolution Mathématique, CNRS, UMR 7625 Écologie et Évolution, Université
9	Pierre et Marie Curie & École Normale Supérieure, 46 rue d'Ulm 75230 Paris Cedex 05,
10	France
11	<sup>3</sup> UMR 1099 BIO3P (Biology of Organisms and Populations applied to Plant Protection),
12	INRA Agrocampus Rennes, F-35653 Le Rheu, France
13	<sup>4</sup> Unité de Recherche sur les Espèces Fruitières, INRA, Domaine de la Grande Ferrade, 71
14	avenue Edouard Bourlaux, BP 81, 33883 Villenave d'Ornon Cedex, France
15	Author for correspondence: Stéphanie Mariette, Tel: (33) 557 122 383, Fax: (33) 557 122
16	439, E-mail: smariett@bordeaux.inra.fr
17	
18	Keywords
19	Self-incompatibility, asexual reproduction, number of S-alleles, linkage disequilibrium,
20	inbreeding depression, mutation load
21	Running title: Asexuality and self-incompatibility in small populations
22	Word count: 5433
23	

## Summary

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

How self-incompatibility systems are maintained in plant populations is still a debated issue. Theoretical models predict that self-incompatibility systems break down according to the intensity of inbreeding depression and number of S-alleles. Other studies have explored the role of asexual reproduction in the maintenance of self-incompatibility. However, the population genetics of partially asexual, self-incompatible populations are poorly understood and previous studies have failed to consider all possible effects of asexual reproduction or could only speculate on those effects. In the present study, we investigated how partial asexuality may affect genetic diversity at the S-locus and fitness in small self-incompatible populations. A genetic model including an S-locus and a viability locus was developed to perform forward simulations of the evolution of populations of various sizes. Drift combined with partial asexuality produced a decrease in the number of alleles at the S-locus. In addition, an excess of heterozygotes was present in the population, causing an increase in mutation load. This heterozygote excess was enhanced by the self-incompatibility system in small populations. In addition, in highly asexual populations, individuals produced asexually had some fitness advantages over individuals produced sexually (due to the increased heterozygosity, sex produces the homozygosis of deleterious alleles). Our results suggest that future research on the role of asexuality for the maintenance of self-incompatibility will need to (1) account for whole genome fitness (mutation load generated by asexuality, selfincompatibility and drift) and (2) acknowledge that the maintenance of self-incompatibility may not be independent of the maintenance of sex itself.

## Introduction

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

Hermaphroditic plant species reproduce with variable rates of selfing, ranging from strict selfing to strict outcrossing (Barrett, 2002). Self-incompatibility (SI) is a reproductive system that prevents self-fertilization. In the case of heteromorphic self-incompatibility, distinct morphologies result in distinct compatibility groups, whereas in the case of homomorphic self-incompatibility, compatible individuals cannot be distinguished by their morphology (de Nettancourt, 1977). Most SI systems depend on physiological mechanisms that prevent pollen germination or pollen tube growth. In sporophytic self-incompatibility (SSI) systems, the compatibility of a pollen grain depends on the diploid genotype of the plant that produced it. In gametophytic self-incompatibility (GSI) systems, the compatibility of a pollen grain depends on its haploid genotype. GSI is more widespread than SSI (Glémin et al., 2001). Fisher (1941) showed that self-fertilization should have a selective advantage because a selfing genotype will transmit more copies of its genome than a non-selfing genotype (this has been termed the automatic advantage of selfing). However, numerous studies have shown that inbred offspring are less fit than outbred offspring. The relative decrease in the mean fitness of selfed versus outcrossed individuals (inbreeding depression) is generally recognized as the only main factor that counterbalances the selective advantage of selfing (Charlesworth and Charlesworth, 1987). Consequently, the level of inbreeding depression in populations should play a determining role in the evolution of SI systems. Inbreeding depression decreases as population size becomes smaller due to reduced polymorphism for selection to act on (Bataillon and Kirkpatrick, 2000). Charlesworth and Charlesworth (1979) showed that the number of Salleles also plays a role in maintaining SI, since a low number of alleles will limit the number of compatible crosses in the population. A decrease in population size can also cause a reduction in the number of S-alleles (Brennan et al., 2003), and a self-compatible mutant can

be positively selected for (Reinartz and Les, 1994). Thus, small populations may be particularly prone to the breakdown of SI due to weak inbreeding depression and low numbers of S-alleles. However, small self-incompatible populations may maintain high levels of inbreeding depression due to a sheltered load of deleterious alleles linked to the S-locus (Glémin et al., 2001). The existence of a sheltered load has been demonstrated experimentally in Solanum carolinense by Stone (2004) and Mena-Alí et al. (2009). Overall, under a wide range of conditions, SI can evolve to self-compatibility. In effect, the loss of SI systems is very frequent in plant evolution (Igic et al., 2008). However, the reasons for which some species maintain a SI system while other species lose it are not completely understood. It has been suggested that asexual reproduction, "when an individual produces new individuals that are genetically identical to the ancestor at all loci in the genome, except at those sites that have experienced somatic mutations" (de Meeûs et al., 2007), plays a role in the maintenance or breakdown of SI. Two studies suggest that asexuality could relieve the main selective pressures that favour the breakdown of SI. First, Chen et al. (1997) showed in Australian Droseraceae that all self-incompatible taxa have effective forms of asexual reproduction, whereas the obligatory sexual annual taxon, Drosera glanduligera, is selfcompatible. Their interpretation is that self-incompatible forms accumulate recessive lethal polymorphisms, especially in association with biparental inbreeding generated by elevated levels of asexual reproduction. The hypothetical high genetic load constitutes the selective pressure that maintains the outbreeding mechanisms. Second, Vallejo-Marín and O'Brien (2007) hypothesized that asexuality could provide reproductive assurance in cross-fertilizing species subject to pollen limitation. They predicted that cross-fertilizing species subject to pollen limitation would often have some means of asexual reprduction, and they found a strong correlation between self-incompatibility and asexuality in Solanum (Solanaceae). Conversely, Young et al. (2002) developed a contrasting hypothesis for the case of Rutidosis

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

*leiolepis* in which they found SSI and high rates of asexual reproduction. They suggested that increased asexual reproduction causes mate limitation by reducing genotype diversity at the *S*-locus, favouring a breakdown of self-incompatibility.

The net effect of asexual reproduction on the maintenance or breakdown of SI systems is therefore still unknown. Moreover, it is unclear to what extent the effects of asexual reproduction discussed by the various authors are actually present in partially asexual, self-incompatible populations. Theoretical studies have described the effects of partial asexuality on the inbreeding depression of self-compatible populations (Muirhead and Lande, 1997) and the number of S-alleles (Vallejo-Marín and Uyenoyama, 2008). However, no model has been developed so far to study the joint effect of asexuality, self-incompatibility and drift on the evolution of diversity and fitness parameters for a hermaphroditic species. Using individual-based simulations, we investigated the effect of (i) partial asexual reproduction and drift on diversity at the S-locus, (ii) combined asexuality, self-incompatibility and drift on the frequency of deleterious alleles. Thus, our main goal was to characterise the dynamics of two key factors (number of S-alleles and inbreeding depression) for the maintenance of SI in partially asexual, self-incompatible populations. It was not the aim of this work to estimate the probability of invasion of self-compatible alleles or other modifiers of reproduction.

## **Materials and Methods**

Genetic model

The model considered in this study was based on the model developed by Glémin *et al.* (2001). This model consists of a population of N (four population sizes evaluated: 25, 50, 100 and 1000) diploid hermaphroditic individuals with a GSI system. In addition, our model also included asexual reproduction at rate c (probability that an individual is generated by asexual reproduction, seven values evaluated: 0, 0.5, 0.8, 0.9, 0.99, 0.999 and 1). We considered an

asexual reproduction event as the production of a new independent individual that is an exact copy of the parental individual (or only different by somatic mutation) (e.g. de Meeûs et al., 2007). As in Glémin et al. (2001), each individual genome possessed the S-locus, which regulated SI, and a viability locus, whose state determined individual fitness. Two neutral loci were also included for reference. The four loci were considered to be physically unlinked and were inherited independently through sexual reproduction. Sexual reproduction events were controlled by the S-locus: crosses between individuals were only possible when the S-allele carried by the pollen grain was different from both S-alleles of the pistil (i.e. at least three different S-alleles were necessary in the population for sexual reproduction to occur). To focus on effects specifically attributable to asexuality on the maintenance and breakdown of SI, we assumed unlimited pollen availability and that sexual crosses were always fruitful. Thus, sexual reproduction was only limited when asexuality and genetic drift reduced the number of S-alleles to less than three within a population. In this case, independent of the original rate of asexual reproduction, all individuals reproduced asexually (c=1) until mutation introduced a third S-allele. Then, c was reset to its original value. Individual fitness, f, was determined by the viability locus, which had two alleles A and a. Strength of the dominance of allele A over deleterious allele a was regulated by the dominance coefficient h, and strength of the selection against a was regulated by the selection coefficient s. Thus, relative fitness of genotypes were:  $f_{AA}=1$ ,  $f_{Aa}=1$ -hs and  $f_{aa}=1$ -s (0<s<=1 and 0 < h <= 1/2). Three selection regimes at the viability locus were considered: (i) a neutral case, to be able to study the effects of the interaction between SI and asexuality without the interference of selection, where parameters s and h were set to zero; (ii) a case of mildly deleterious and partially recessive mutations, for which values assigned to s and h were

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

respectively 0.1 and 0.2; and (*iii*) a case of highly recessive lethal mutations, values assigned to *s* and *h* were respectively 1 and 0.02.

Mutations were allowed in all loci. Neutral loci mutated at rate  $\mu_N=10^{-3}$ , following a *k*-allele model (*k*=100). Mutation at the *S*-locus followed an infinite-allele model (Kimura and Crow, 1964) with rate  $\mu_S$ ; three values were considered for this rate ( $\mu_S=10^{-3}$ ,  $10^{-4}$  and  $10^{-5}$  as in Glémin *et al.*, 2001). Mutations at the viability locus occurred at rate  $\mu_1=10^{-3}$  from *A* to *a* and

at rate  $\mu_2=10^{-4}$  from a to A.

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

144

145

146

147

148

149

150

## Individual-based simulations

Forward-time simulations of a population of N diploid individuals with no overlapping generations were performed. Simulations started with all individuals carrying two unique alleles at the S-locus (2N different S-alleles in the whole population), alleles A or a randomly assigned (with equal probability, 0.5) to the viability locus and any of the k alleles (k=100) randomly assigned (with equal probability, 1/k) to the two neutral loci. At each generation, the number of individuals generated by asexual reproduction, x, was drawn from a Poisson distribution with mean Nc, and the remaining N-x individuals were generated by sexual reproduction (parents contributing to these two types of offspring were not determined in this step). Genotypes of the N individuals were generated one by one. If a given individual originated by asexual reproduction, a genotype i was randomly drawn from all individuals of the previous generation. That genotype was assigned to the individual with a probability equal to the fitness of genotype i; this will be called the selection step. If the genotype was not successfully assigned, the genotype draft and the selection step was repeated until a genotype passed the selection step. If the individual originated by sexual reproduction, the simulation followed a different procedure. First, an individual from previous generation was randomly drawn to act as a mother and the ovule haploid genotype was

randomly generated. Then, a second individual was drawn to act as a father and the pollen haploid genotype was randomly generated. If the *S*-allele at the pollen grain was not different from both *S*-alleles of the mother (i.e. incompatible pollen), the pollen was discarded. New fathers and pollen grain genotypes were generated until a compatible cross was obtained (i.e. pollen availability was unlimited). Then, a diploid genotype i was generated from the haploid genotypes of the ovule and the pollen. The selection step was carried out on genotype i as in the case of asexual reproduction; if the genotype was not assigned the process was repeated by drawing two new parents until a genotype was able to pass the selection step. Once all N genotypes were selected, mutations were applied to each individual. The numbers of mutants at the S-locus and at the two neutral loci were drawn from Poisson distributions with means  $2N\mu_S$  and  $2N\mu_N$  respectively. At the viability locus, a mutant was formed with a probability  $\mu_1$  if the allele was A and a probability  $\mu_2$  if the allele was A. The rate of asexual reproduction was a fixed parameter of our model and did not change with the selection process.

## Monitored genetic indices

Simulations were run for 50 000 generations before beginning to monitor the genetic indices, this allowed monitored indices to become stable and avoided the influence of initial conditions in their values (Glémin *et al.*, 2001). From generation 50 000, indices were calculated every 3 000 generations until 107 000 generations (i.e. 20 times). For the same combination of parameters, 50 replicates of the simulations were performed providing 1 000 (indices calculated 20 times in each of the 50 replicates) evaluations of the monitored variables for each set of parameter values. Indices were calculated from all genotypes of the population (i.e. they were not estimates, but the true values of the population).

frequencies, effective number of alleles ( $n_e = 1 / \sum_{i=1}^{n_a} p_i^2$ , where  $p_i$  was the frequency of allele),

Genetic diversity indices: For each locus we calculated the number of alleles  $(n_a)$ , allele

- 194 expected heterozygosity  $(H_e = 1 1/n_e)$ , observed heterozygosity  $(H_o, proportion of$
- heterozygous individuals) and the inbreeding coefficient  $(F_{IS} = (H_e H_o)/H_e)$ . Theoretical
- 196 predictions for some of these indices were obtained from the work of several authors for some
- of the scenarios considered:
- 198 a) Vallejo-Marín and Uyenoyama (2008) studied the effects of partial asexuality in the
- number of alleles of the S-locus using a diffusion approximation and found that the number of
- 200 common alleles ( $n_c$ , which is approximately equivalent to the effective number of S-alleles,
- 201  $n_e$ , Takahata, 1990) is determined by:

$$202 (1) 1 = \theta \frac{2Mn_c}{2Mn_c - \theta(n_c - 1)(n_c - 2)} e^{\frac{2Mn_c}{(n_c - 1)(n_c - 2)} - \theta} \sqrt{\frac{2\pi(n_c - 1)}{Mn_c + \theta(n_c - 1)}} \left\{ \left[ 1 + \frac{\theta(n_c - 1)}{Mn_c} \right] \frac{n_c - 2}{n_c} \right\}^{\theta + Mn_c/(n_c - 1)}$$

- where M = N(1-c) and  $\theta = 2N\mu_s$ . This equation was solved numerically to obtain the
- 204 expected effective number of S-alleles in the simulations without selection on the viability
- 205 locus.
- b) Glémin *et al.* (2001) obtained the expected  $F_{IS}$  for a neutral locus linked to the S-locus as:

207 (2) 
$$F_{IS} = -\frac{(1+2f_S)(n_c-1)}{1+n_c^2+2f_S(n_c-1)(4Nr'+n_c-1)}$$

- where  $f_S$  is the scaling factor of the genealogy of the S-locus and r' is the net recombination
- 209 rate between the two loci. For the case of partial asexuality, the number of common alleles
- 210 was given by Equation 1 and the scaling factor was  $f_s = \frac{n_c^2(n_c 1)(n_c 2)}{4\theta[2Mn_c \theta(n_c 1)(n_c 2)]}$
- 211 (Vallejo-Marín and Uyenoyama, 2008). To correct for the reduction of recombination due to
- as a sexuality, we used r'=r(1-c), where r was the recombination rate during sexual
- 213 reproduction (i.e. r=0.5 for unlinked loci). This approach ignored mutation on the neutral
- 214 locus.

- c) Finally, an expression of the expected  $F_{IS}$  of a neutral locus in a self-compatible, partially
- asexual population was derived following Balloux et al. (2003):

217 (3) 
$$F_{IS} = \frac{\gamma[(1-c)Ns - 1]}{2N + \gamma[1+cN(s-2) - Ns]}$$

- where  $\gamma = (1 \mu_N)^2$  and s is the selfing rate. Selfing rate in Balloux et al. (2003) model is
- 219 defined as the rate of self-fertilization of individuals; thus a fertilization event between two
- 220 individuals from the same genet is considered an outcrossing event. Under random mating,
- 221 s=1/N and for negligible mutation rates,  $\gamma=1$ .
- 222 <u>Linkage disequilibrium:</u> The linkage disequilibrium between pairs of loci (S-locus-viability
- locus and the two neutral loci) was studied with the correlation coefficient between two loci,
- $R_{GGD}$ , developed by Garnier-Géré and Dillmann (1992) which is particularly appropriate for
- partially asexual diploids (de Meeûs and Balloux, 2004).
- 226 Fitness: Mean fitness of the population ( $\overline{W}$ ) was calculated from the genotype frequencies as
- 227  $\overline{W} = f(AA) \times 1 + f(Aa) \times (1 hs) + f(aa) \times (1 s)$ , and the average fitness of individuals
- potentially produced by selfing  $(W_s)$ , outcrossing  $(W_o)$  and asexuality  $(W_a)$  were calculated as

229 
$$W_s = f(AA) \times 1 + f(Aa) \times \left[ \frac{1}{4} \times 1 + \frac{1}{2} \times (1 - hs) + \frac{1}{4} \times (1 - s) \right] + f(aa) \times (1 - s),$$

- 230  $W_o = 1 \times p_A^2 + (1 hs) \times 2 \times p_A \times p_a + (1 s) \times p_a^2$  (where  $p_A$  and  $p_a$  are the frequencies of alleles
- 231 A and a) and  $W_a = \overline{W}$ . Inbreeding depression ( $\delta$ ) and mutation load (L) were calculated as
- 232  $\delta = 1 (W_s/W_o)$  and  $L = 1 \overline{W}$ . In addition, we compared the fitness of individuals produced
- by asexual reproduction  $(W_a)$  to the fitness of individuals produced sexually, using the ratio
- $W_o/W_a$ .  $W_s$ ,  $W_o$  and  $W_a$  were calculated as the fitness of potential offspring if all individuals
- were produced by selfing, by outcrossing and asexually, respectively. However, these modes
- of reproduction might not be actually present in the different scenarios considered (e.g. in

self-incompatible populations, no selfing occurs, but potential  $W_s$  and inbreeding depression can be calculated).

Fixation probabilities: To interpret the results, we calculated the probabilities of four events:

<u>Fixation probabilities:</u> To interpret the results, we calculated the probabilities of four events: fixation of alleles A and a in the viability locus, 'fixation' of viability locus heterozygote (Aa) and fixation of a genotype in the S-locus (reduction of number of S-alleles to two, this was only monitored in self-incompatible populations). These fixation probabilities were estimated on 1 000 evaluations of the monitored variables as the proportion of times in which  $p_a$ =0,

# 246 Results

Effect of asexuality and drift on linkage disequilibrium between loci

 $p_a$ =1 and  $H_o$ =1 for the viability locus, or n=2 for the S-locus.

Probably the most obvious effect of partial asexual reproduction was the reduction of recombination. Asexuality generated identical genotypes in offspring, so allele associations were transmitted in the same way that they would be transmitted if they were physically linked throughout the whole genome. In fully asexual populations, maximum linkage disequilibrium was expected (equivalent to the linkage disequilibrium between fully linked loci) and in partially asexual populations, recombination was reduced proportionally to the rate of asexual reproduction. Figure 1 shows how non-random associations of alleles increased with asexual reproduction rates. Linkage disequilibrium levels also depended on population size because drift produced a departure from the expected values of frequencies of allele combinations (Figure 1). Strong drift might have even caused some combinations of alleles to be absent in the population.

In small populations, some generations with complete allelic association started to appear at asexual reproduction rates of 0.8 and were predominant at rates of 0.99 and higher. In larger populations, complete allelic association was only found in the fully asexual scenario. These

262 associations of alleles were observed between loci subject to selection and between neutral 263 loci, which showed that linkage disequilibrium was not a product of selective processes 264 (Figure 1a and Figure 1b). 265 Physical linkage between the S-locus and loci with deleterious alleles can decrease the 266 number of S-alleles (Uyenoyama, 2003). It can also dramatically increase inbreeding 267 depression in small populations (Glémin et al., 2001). Therefore, for the present model, it was 268 important to discern whether the linkage disequilibrium generated by asexual reproduction is 269 sufficient for the selective forces of the S-locus to interact with the viability locus. The results 270 are presented below taking this perspective into account. 271 272 Effect of asexuality and drift on diversity parameters: number of S-alleles and heterozygote 273 excess 274 The effective number of S-alleles decreased with asexuality (Figure 2) as shown analytically 275 by Vallejo-Marín and Uyenoyama (2008). Balancing selection on the S-locus only occurred 276 during sexual reproduction events. Since these events were scarce at high rates of asexual 277 reproduction, the influence of balancing selection (which promotes high allelic diversity) on 278 the population diminished. This can temporarily stop sexual reproduction by reducing the 279 number of S-alleles to two (i.e. fixation of an S-locus genotype, Table 1). 280 No significant differences in the number of S-alleles were found among the three different 281 selection regimes for the viability locus (frequency distributions of the number of S-alleles 282 among different selection regimes were undistinguishable in a Kolmogorov-Smirnov test, P-283 value>0.98). It must be noted that small populations (with high linkage disequilibrium) were

fixed for the wild type allele A most of the time (Table 2b) and this may explain the weak

influence of deleterious alleles on the S-locus.

284

Another general effect of partial asexual reproduction in diploids was the reduction of allele segregation. This can increase heterozygosity by the independent accumulation of mutations on the two alleles of an asexual lineage (Pamilo, 1987; Birky, 1996). To quantify this effect, we measured the inbreeding coefficient  $F_{IS}$ , which compares observed and expected heterozygosities. Theoretical equilibrium values for  $F_{IS}$  at a neutral locus (unlinked to the S-locus) were very similar in self-compatible and self-incompatible populations (Figure 3), and, for large population sizes, the simulated populations followed the same pattern (Figure 3b). However, for small population sizes, drift generated strong linkage disequilibrium between the S-locus and other loci. Thus, balancing selection on the S-locus had a hitchhiking effect on other loci, increasing their heterozygosity. This explained the contrasting  $F_{IS}$  values on simulated selfcompatible and self-incompatible populations (Figure 3a). Analytical predictions do not reflect this dramatic difference because they do not account for linkage disequilibrium.  $F_{IS}$ decreased with asexuality and this decrease was enhanced by self-incompatibility in small populations, particularly noticeable at asexual reproduction rates higher than 0.8. This decrease in  $F_{IS}$  observed in neutral loci also occurred at the viability locus (both for the mildly deleterious and lethal recessive cases), which implied an increase in the frequency of the deleterious allele (Figure 4).

304

305

306

307

308

309

310

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

Effect of asexuality and drift on fitness

For both self-compatible and self-incompatible populations, mutation load globally increased with asexuality and population size, but a greater variance among generations and populations was observed for very small populations (Figure 5). As mentioned above, the frequency of the deleterious allele increased with asexuality in small self-incompatible populations, due to a hitchhiking effect of the *S*-locus over the viability locus. In the self-incompatibility system,

when the population size was small enough, the mutation load increased with asexual reproduction rates; this load resulted from the increase of the deleterious allele frequency (Figure 5a and 5c). Under similar conditions in the self-compatible system, mutation loads were lower than the mutation load observed in the self-incompatible system (Figure 5b and 5d). In contrast, inbreeding depression had similar values in self-incompatible and self-compatible populations (Figure 6), showing some increases with asexual reproduction rate for lethal recessive mutations that was more apparent in large populations.

The average fitness was higher in sexual populations than in asexual populations due to Mendelian segregation (e.g. Figure 5; Chasnov, 2000; Kirkpatrick and Jenkins, 1989). The relative fitness between individuals produced by sexual and asexual reproduction was influenced very little by partial asexuality in a self-compatible population (Figure 7b and 7d). However, in small self-incompatible populations, the asexual reproduction rate increased the advantage of asexual compared to sexual reproduction (Figure 7a and 7c). These populations consisted mainly of heterozygous individuals (Figure 3). Therefore, asexual reproduction produced mainly *Aa* individuals, with only slightly lower fitness than the fittest *AA* haplotype. However, sexual reproduction produced a high proportion of *aa* individuals, reducing the average fitness of offspring, a reduction that was not compensated by the production of the fittest *AA* individuals. This could potentially lead to a positive feedback effect on the asexual reproduction rate, but this was not studied in our model where the asexual reproduction rate was a fixed parameter.

#### Discussion

The number of S-alleles and inbreeding depression are considered to be the main factors that influence the maintenance or breakdown of self-incompatibility systems (Charlesworth and Charlesworth, 1979). A decrease in the number of S-alleles is expected to favour the

breakdown of self-incompatibility since self-compatible mutants can invade the population, whereas the maintenance of inbreeding depression within populations prevents the breakdown of self-incompatibility. In the present study, we examined how drift and partial asexuality may modify the dynamics of these two key parameters in a self-incompatible population.

Drift, combined with partial asexuality, dramatically reduced the number of S-alleles

As already predicted by Yokoyama and Hetherington (1982), we showed that the number of *S*-alleles decreases with the effective size of the population. Thus, in small populations, repeated bottlenecks or founder events may lead to the breakdown of the self-incompatibility system. However, Karron (1987) found no differences in mating systems between rare and widespread congeners across several families (but see also a study on some Brassicaceae species, Kunin and Shmida, 1997). As for the impact of asexuality, our simulation study confirmed a recent result demonstrated analytically by Vallejo-Marín and Uyenoyama (2008). The number of *S*-alleles decreased with asexuality due to the weakened influence of balancing selection in a partially asexual population (Figure 2). In contrast with other effects of asexuality that were only observed for very high rates of asexuality, intermediate asexual reproduction rates were sufficient to significantly reduce the number of *S*-alleles. High rates of asexual reproduction combined with drift produced extremely low numbers of *S*-alleles. This effect was strong enough to even reduce the number of *S*-alleles to two (i.e. fixation of an *S*-locus genotype, Table 1), which stopped all possibility of sexual reproduction until a

new S-allele arose from mutation or migration. Therefore, in such extreme cases, this drastic

effect of asexuality on the number of S-alleles should favour the breakdown of self-

incompatibility. However, paradoxically, the number of S-alleles may not be very relevant in

a system where asexuality serves to provide reproductive assurance.

Drift, combined with partial asexuality and self-incompatibility, led to an increase in mutation load Drift influenced the level of mutation load. As population size decreased, the frequency of the deleterious allele decreased and mutation load decreased, due to more effective purging. We also observed a much greater variance in mutation load in very small populations (Table 2 and Figure 5, N=25). In this case, selection was overwhelmed by drift and this caused a higher probability of fixation of the deleterious allele, and an increase in mutation load. Similar results have been described by Bataillon and Kirkpatrick (2000), Glémin (2003) and Haag and Roze (2007). Drift is also a key component of mutation load in asexual populations (Haag and Roze, 2007). Indeed, large asexual and sexual populations showed comparable frequencies of the deleterious allele and comparable mutation loads (Figure 5; see also Haag and Roze, 2007). However, mutation load was greater in small asexual populations than in small sexual ones. This could be explained by the absence of segregation in asexual populations, in which heterozygous individuals are present in high frequencies (Table 2 and Figure 5; see also Haag and Roze, 2007). In this study, we additionally showed that self-incompatibility played a role in increasing mutation load in partially asexual, self-incompatible populations. Partial asexual reproduction and small population sizes produced strong associations (i.e. linkage disequilibrium) between deleterious alleles (on any locus of the genome) and S-alleles. Glémin et al. (2001) have already shown that deleterious mutations on loci linked to the S-locus may be sheltered by balancing selection acting on rare S-alleles. Asexuality and self-incompatibility both favoured the increase of the heterozygote genotype Aa within the population, leading to an increase of mutation load in small self-incompatible, partially asexual populations.

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

Inbreeding depression remained low in small, partially asexual, self-incompatible populations Inbreeding depression followed different patterns than the ones observed for mutation load. For very small populations, inbreeding depression is expected to decrease (Bataillon and Kirkpatrick, 2000), due to the absence of polymorphism at the viability locus. This was observed in small self-compatible and self-incompatible populations (Glémin et al., 2001, and our own simulations, Figure 6). As for the impact of asexuality, our results showed no large differences between the self-compatible and the self-incompatible cases for small populations. The effect of strong drift, in the presence of strong asexuality, was to reduce genotype diversity. In most cases, populations were fixed at one allele, but sometimes the population was fixed at the heterozygote genotype Aa. This mainly occurred in self-incompatible populations (see Table 2). With no or very low genotype diversity, inbreeding depression had very low values, even if deleterious alleles were present at high frequencies in the population. Chen et al. (1997) proposed that asexuality would favour the maintenance of SI. Their reasoning was that asexual reproduction would increase a population's genetic load due to lethal recessive mutations, which was confirmed by our simulation results. However, mutation load and inbreeding depression did not show parallel trends, and the high level of mutation load due to the combination of drift, asexuality and SI was accompanied by low levels of inbreeding depression. Nevertheless, inbreeding depression generally tended to increase with asexuality due to the accumulation of deleterious alleles and this should favour the maintenance of SI.

407

408

409

410

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

Evolution of partially asexual, self-incompatible populations: a third possible outcome

Until now, the evolutionary interest of partially asexual, self-incompatible populations has
been focused on two possible outcomes: (1) the system is stable and SI is maintained or (2)

the system is unstable and SI disappears (Vallejo-Marín, 2007; Vallejo-Marín and O'Brien, 2007; Chen et al., 1997; Young et al., 2002). However, a third possible outcome should be considered: being the unstable system, sexual reproduction completely disappears. This type of event was observed in our simulations due to a reduction in the number of S-alleles to two (see above). However, we also studied how selection on the viability locus could contribute to the potential loss of sexuality. Our results showed that the fitness of individuals produced by as exual reproduction  $(W_a)$  was greater than that of individuals produced by random sexual mating  $(W_o)$  in small populations (Figure 7). Fitness of individuals (potentially) produced by selfing  $(W_s)$  was also lower than  $W_a$  in small self-incompatible populations, suggesting that asexuality may have an immediate advantage over selfing under such circumstances. Therefore, we speculate that asexual reproduction could take over sexual reproduction more easily than self-compatibility over self-incompatibility when low numbers of S-alleles reduce mate availability. Despite the effect of asexuality on inbreeding depression and S-locus diversity, asexual reproduction may offer an alternative means of reproduction under these adverse conditions and self-incompatibility may be temporarily dormant instead of breaking down. Asexual populations could subsequently recover a sexual self-incompatible reproductive system when new S-alleles are introduced in the population (for instance, by migration). However, prolonged periods of asexual reproduction may facilitate the accumulation of mutations affecting sexual traits, including self-incompatibility, since selective pressures are ineffective for these traits (Eckert, 2001). Thus, alternative periods of sexual and asexual reproduction may be necessary for such dynamics, which is probably the case for plants combining asexual and sexual reproduction (Bengtsson and Ceplitis, 2000). Nevertheless, the scenarios described here need to be confirmed by further and more complex simulations and other factors may favour self-compatible sexual reproduction, such as the more effective purging of deleterious mutations.

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

## Perspectives

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

In this study, we used a model with physically unlinked loci and a single viability locus. However, because the linkage disequilibrium produced by asexuality affected the whole genome, population genomic models may be necessary to fully study the effect of deleterious alleles on the S-locus for small, partially asexual populations. In addition, physically linked loci with deleterious alleles (which were not studied here) may reduce the number of S-alleles (Uyenoyama, 2003) and the strength of that effect may be influenced by the rate of asexuality as discussed by Vallejo-Marín and Uyenoyama (2008). Moreover, in the present study, it was not possible to investigate the evolutionary dynamics of mating system and asexuality modifiers since these were assumed to be fixed parameters. It would be then interesting to develop a model to test the invasion propensity of self-compatible or asexual mutants. Finally, models incorporating migration or population structure are necessary to study more realistic dynamics of partially asexual, self-incompatible populations. Population structure is expected to affect the genetic diversity of viability and the S-locus, for instance by introducing a third allele via migration within subpopulations fixed at two S-alleles. These models should also take into account the specificities of the different types of asexual reproduction where, for instance, the asexual propagules may have a lower dispersal capacity (e.g. vegetative reproduction) and may create spatial clusters of individuals with the same genotype. Recent theoretical work (Vallejo-Marín and Uyenoyama, 2008) and our own model lead to several predictions about S-locus diversity, mutation load and inbreeding depression within partially asexual self-incompatible populations. However, we need more experimental observations to test and validate these theoretical advances. First, as outlined by Vallejo-Marín and O'Brien (2007), if asexuality relieves selective pressures favouring the breakdown of SI, the co-occurrence of asexuality and self-incompatibility should be frequent. Association of self-incompatibility and asexuality has been studied in only a few monospecific studies (Rutidosis leiolepis, Young et al., 2002; Eucalyptus morrisbyi, Jones et al., 2005; Prunus avium, Stoeckel et al., 2006). The association was found at a multispecific level by Chen et al. (1997) in Australian Droseraceae and by Vallejo-Marín and O'Brien (2007) in Solanaceae. More reports on species combining self-incompatibility and asexuality would enhance our understanding of pressures explaining their maintenance. Second, experimental measurements of asexual reproduction rates and number of S-alleles in species showing various levels of asexual reproduction would be useful to test predictions from our simulations. Although experimental measurements of inbreeding depression in self-incompatible species are available (Mena-Alí et al., 2008), studies comparing inbreeding depression from different populations with contrasting sizes and rates of asexual reproduction are lacking.

## Conclusions

This study investigated for the first time the effect of partial asexual reproduction on the fitness of self-incompatible populations. For loci unlinked to the S-locus and large populations, fitness values were similar to those of a self-compatible population. However, in small populations, the combination of drift, asexuality and SI increased mutation load due to the accumulation of deleterious alleles. This increase in the mutation load was accompanied by a selective advantage of asexually produced offspring compared to sexually produced offspring. Therefore, future studies addressing the maintenance of SI in partially asexual populations (by studying the invasibility of self-compatible genotypes) will need to (1) account for whole genome fitness and (2) acknowledge that the process may not be independent of the maintenance of sex itself (i.e. will need to establish whether modifiers of asexual reproduction rates invade more easily than self-compatible alleles).

## Acknowledgements

Collaboration between MN and SM was promoted through the implementation of the REPROFOR project, financed by the Spanish Ministry for Education and Science. We would like to thank Santiago C. González-Martínez and Ricardo Alía for helping to develop this collaboration. E. Porcher is acknowledged for helpful discussions on self-incompatibility evolution. We are also very thankful to P. Garnier-Géré for a detailed discussion on linkage disequilibrium estimators.

## 493 References

- 494 Balloux F, Lehmann L, de Meeûs T (2003). The population genetics of clonal and partially
- 495 clonal diploids. *Genetics* **164**: 1635-1644.
- Barrett SCH (2002). The evolution of plant sexual diversity. *Nat Rev Genet* **3**: 274-284.
- 497 Bataillon T, Kirkpatrick M (2000). Inbreeding depression due to mildly deleterious mutations
- in finite populations: size does matter. *Genet Res* **75**: 75-81.
- Bengtsson BO, Ceplitis A (2000). The balance between sexual and asexual reproduction in
- plants living in variable environments. *J Evolution Biol* **13**: 415-422.
- 501 Birky CW (1996). Heterozygosity, Heteromorphy, and Phylogenetic Trees in Asexual
- 502 Eukaryotes. *Genetics* **144**, 427-437.
- Brennan AC, Harris SA, Hiscock SJ (2003). The population genetics of sporophytic self-
- incompatibility in Senecio squalidus L. (Asteraceae): avoidance of mating constraints
- 505 imposed by low S-allele number. *Philos T R Soc B* **358**: 1047-1050.
- 506 Charlesworth D, Charlesworth B (1987). Inbreeding depression and its evolutionary
- 507 consequences. *Annu Rev EcolEvol S* **18**: 237-268.
- 508 Charlesworth D, Charlesworth B (1979). The evolution and breakdown of S-allele systems.
- 509 *Heredity* **43**: 41-55.
- 510 Chasnov JR (2000). Mutation-selection balance, dominance and the maintenance of sex.
- 511 *Genetics* **156**: 1419-1425.
- 512 Chen L, Stace HM, James SH (1997). Self-incompatibility, seed abortion and clonality in the
- 513 breeding systems of several Western Australian Drosera species (Droseraceae) . Aust J
- 514 *Bot* **45**: 191-201.
- Eckert CG (2001). The loss of sex in clonal plants. Evol Ecol 15: 501-520.
- Fisher RA (1941). Average excess and average effect of a gene substitution. *Ann Eugen* 11:
- 517 53-63.

518 Garnier-Géré P, Dillmann C (1992). A computer program for testing pairwise linkage 519 disequilibria in subdivided populations. J Hered 83: 239. 520 Glémin S (2003). How are deleterious mutations purged? Drift versus nonrandom mating. 521 Evolution **57**: 2678—2687. 522 Glémin S, Bataillon T, Ronfort J, Mignot A, Olivieri I (2001). Inbreeding depression in small 523 populations of self-incompatible plants. *Genetics* **159**: 1217-1229. 524 Haag CR, Roze D (2007). Genetic load in sexual and asexual diploids: segregation, 525 dominance and genetic drift. Genetics 176: 1663-1678. 526 Igic B, Lande R, Kohn JR (2008). Loss of self-incompatibility and its evolutionary 527 consequences. Int J Plant Sci 169: 93-104. 528 Jones RC, McKinnon GE, Potts BM, Vaillancourt RE (2005). Genetic diversity and mating 529 system of an endangered tree Eucalyptus morrisbyi. Aust J Bot 53: 367-377. Karron JD (1987). A comparison of levels of genetic polymorphism and self-compatibility in 530 531 geographically restricted and widespread plant congeners. Evol Ecol 1: 47-58. 532 Kimura M, Crow JF (1964). The number of alleles that can be maintained in a finite 533 population. Genetics 49: 725-738. 534 Kirkpatrick M, Jenkins CD (1989). Genetic segregation and the maintenance of sexual 535 reproduction. Nature, 339: 300-301. 536 Kunin WE, Shmida A (1997). Plant reproductive traits as a function of local, regional, and 537 global abundance. Conserv Biol 11: 183-192. 538 de Meeûs T, Balloux F (2004). Clonal reproduction and linkage disequilibrium in diploids: a 539 simulation study. Infect Genet Evol 4: 345—351. 540 de Meeûs T, Prugnolle F, Agnew P (2007). Asexual reproduction: genetics and evolutionary

aspects. Cell Mol Life Sci 64: 1355-1372.

542 Mena-Alí J, Keser L, Stephenson A (2008). Inbreeding depression in Solanum carolinense 543 (Solanaceae), a species with a plastic self-incompatibility response. BMC Evol Biol 8: 544 10. 545 Mena-Alí J, Keser L, Stephenson A (2009). The effect of sheltered load on reproduction in 546 Solanum carolinense, a species with variable self-incompatibility. Sex Plant Reprod 547 **22**: 63-71. 548 Muirhead CA, Lande R (1997). Inbreeding depression under joint selfing, outcrossing, and 549 asexuality. *Evolution* **51**: 1409-1415. 550 de Nettancourt D (1977). Incompatibility in Angiosperms, New York: Springer, Berlin 551 Heidelberg. 552 Pamilo P (1987). Heterozygosity in apomictic organisms. *Hereditas* **107**: 95-101. 553 Reinartz JA, Les DH (1994). Bottleneck-induced dissolution of self-incompatibility and 554 breeding system consequences in Aster furcatus (Asteraceae). Am J Bot 81: 446-455. 555 Stoeckel S, Grange J, Fernandez-Manjarres JF, Bilger I, Frascaria-Lacoste N, Mariette S 556 (2006). Heterozygote excess in a self-incompatible and partially clonal forest tree 557 species - Prunus avium L. Mol Ecol 15: 2109-2118. 558 Stone JL (2004). Sheltered load associated with S-alleles in Solanum carolinense. Heredity 559 **92**: 335-342. 560 Takahata N (1990). A simple genealogical structure of strongly balanced allelic lines and 561 trans-species evolution of polymorphism. PNAS 87: 2419-2423. Uyenoyama MK (2003). Genealogy-dependent variation in viability among self-562 563 incompatibility genotypes. Theor Popul Biol 63: 281-293. 564 Vallejo-Marín M (2007). The paradox of clonality and the evolution of self-incompatibility. 565 Plant Signal Behav 2: 265-266.

566	Vallejo-Marin M, O'Brien HE (2007). Correlated evolution of self-incompatibility and clonal
567	reproduction in Solanum (Solanaceae). New Phytol 173: 415-421.
568	Vallejo-Marín M, Uyenoyama MK (2008). On the evolutionary modification of self-
569	incompatibility: implications of partial clonality for allelic diversity and genealogical
570	structure. In Self-Incompatibility in Flowering Plants. Evolution, Diversity, and
571	Mechanisms . pp. 53-71.
572	Yokoyama S, Hetherington LE (1982). The expected number of self-incompatibility alleles in
573	finite plant populations. <i>Heredity</i> <b>48</b> : 299-303.
574	Young AG, Hill JH, Murray BG, Peakall R (2002). Breeding system, genetic diversity and
575	clonal structure in the sub-alpine forb Rutidosis leiolepis F. Muell. (Asteraceae). Biol
576	Conserv 106: 71-78.

## Figure legends

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

Figure 1 Linkage disequilibrium ( $R_{GDD}$  index, Garnier-Géré & Dillmann 1992) at increasing rates of asexual reproduction. Box-plots represent median (black line), first and third quartiles (box) and 5% and 95% percentiles (whiskers) of 1 000 observations from simulations performed at seven rates of asexual reproduction (c=0, 0.5, 0.8, 0.9, 0.99, 0.999 and 1) and four population sizes (N=25, 50, 100 and 1000). (a)  $R_{GDD}$  index between S-locus and viability locus (s=0.1, h=0.2); (b)  $R_{GDD}$  index between two neutral loci. **Figure 2** Effective number of S-alleles at increasing rates of asexual reproduction. Box-plots represent median (black line), first and third quartiles (grey box) and 5% and 95% percentiles (whiskers) of 1 000 observations from the simulations performed at six rates of asexuality (c=0, 0.5, 0.8, 0.9, 0.99 and 1), population size N=25 (a) and N=100 (b), mutation rate  $\mu_S=10^{\circ}$  $^{5}$ , s=0 and h=0 for the viability locus. Grey line represents the theoretical expected number of common S-alleles obtained numerically from Equation 1 (Equation 3.7 in Vallejo-Marín & Uyenoyama, 2008). Figure 3 Inbreeding coefficient  $F_{IS}$  at a neutral locus at increasing rates of asexual reproduction. Box-plots represent median (black line), first and third quartiles (white and grey boxes, for self-compatible and self-incompatible populations respectively) and 5% and 95% percentiles (whiskers) of 1 000 observations from the simulations performed at six rates of asexuality (c=0, 0.5, 0.8, 0.9, 0.99 and 1), mutation rates  $\mu_S$ =10<sup>-5</sup> and  $\mu_N$ =10<sup>-3</sup>, and population size (a) N=25 and (b) N=100. Theoretical equilibrium values for  $F_{IS}$  at a neutral locus in a self-compatible, random-mating population (dotted black line, from Equation 3, neglecting mutation) and in a self-incompatible population (continuous grey line, from Equation 2) are also represented. Figure 4 Frequency of deleterious alleles at increasing rates of asexual reproduction under two selection regimes. Box-plots represent median (black line), first and third quartiles (white

603 and 95% percentiles (whiskers) of 1000 observations from the simulations performed at seven rates of asexuality (c=0, 0.5, 0.8, 0.9, 0.99, 0.999 and 1), mutation rates  $\mu_S$ =10<sup>-5</sup> and  $\mu_N$ =10<sup>-3</sup>. 604 605 Population size and viability locus coefficients: (a) N=25, s=0.1, h=0.2, (b) N=25, s=1, 606 h=0.02, (c) N=100, s=0.1, h=0.2, and (d) N=100, s=1, h=0.02. 607 Figure 5 Mutation load at increasing rates of asexual reproduction under two different 608 selection regimes according to reproductive system. Box-plots represent median (black line), 609 first and third quartiles (box) and 5% and 95% percentiles (whiskers) of 1 000 observations 610 from simulations performed at seven rates of asexual reproduction (c=0, 0.5, 0.8, 0.9, 0.99, 611 0.999 and 1) and four population sizes (N=25, 50, 100 and 1000). (a) Self-incompatible 612 population with a viability locus with mildly deleterious allele (s=0.1, h=0.2), (b) self-613 compatible population with a viability locus with mildly deleterious allele (s=0.1, h=0.2), (c) 614 self-incompatible population with a viability locus with highly recessive lethal allele (s=1, 615 h=0.02), and (d) self-compatible population with a viability locus with highly recessive lethal 616 allele (s=1, h=0.02). 617 Figure 6 Inbreeding depression at increasing rates of asexual reproduction and under two 618 different selection regimes according to reproductive system. Box-plots represent median 619 (black line), first and third quartiles (box) and 5% and 95% percentiles (whiskers) of 1000 620 observations from simulations performed at seven rates of asexual reproduction (c=0, 0.5, 0.8, 621 0.9, 0.99, 0.999 and 1) and four population sizes (N=25, 50, 100 and 1000). (a) Self-622 incompatible population with a viability locus with mildly deleterious allele (s=0.1, h=0.2), 623 (b) self-compatible population with a viability locus with mildly deleterious allele (s=0.1, 624 h=0.2), (c) self-incompatible population with a viability locus with highly recessive lethal 625 allele (s=1, h=0.02), and (d) self-compatible population with a viability locus with highly

and grey boxes, for self-compatible and self-incompatible populations respectively) and 5%

602

626

recessive lethal allele (s=1, h=0.02).

**Figure 7** Fitness ratio between sexually and asexually produced individuals at increasing rates of asexual reproduction under two different selection regimes according to reproductive system. Box-plots represent median (black line), first and third quartiles (box) and 5% and 95% percentiles (whiskers) of 1 000 observations from simulations performed at seven rates of asexual reproduction (c=0, 0.5, 0.8, 0.9, 0.99, 0.999 and 1) and four population sizes (N=25, 50, 100 and 1000). (a) Self-incompatible population with a viability locus with mildly deleterious allele (s=0.1, h=0.2), (b) self-compatible population with a viability locus with mildly deleterious allele (s=0.1, h=0.2), (c) self-incompatible population with a viability locus with highly recessive lethal allele (s=1, h=0.02).

**Table 1** Genotype fixation probability at the *S*-locus under two different selection regimes at the viability locus

Population	Selection regime	Rate of asexual reproduction							
size		c=0	c=0.5	c=0.8	c=0.9	c=0.99	c=0.999	c=1	
N=25	Mildly deleterious <sup>1</sup>	0.000	0.000	0.036	0.325	0.784	0.807	0.997	
<i>N</i> =25	Lethal recessive <sup>2</sup>	0.000	0.000	0.018	0.315	0.790	0.805	0.994	
<i>N</i> =50	Mildly deleterious	0.000	0.000	0.000	0.002	0.072	0.117	0.990	
<i>N</i> =50	Lethal recessive	0.000	0.000	0.000	0.001	0.075	0.090	0.989	
<i>N</i> =100	Mildly deleterious	0.000	0.000	0.000	0.000	0.001	0.001	0.983	

The probability of genotype fixation at the *S*-locus (only two alleles in the population) was estimated as the proportion of observations in which this event was recorded. Results in this table are for populations with a mutation rate of  $\mu_S$ =10<sup>-5</sup> at the *S*-locus.

<sup>&</sup>lt;sup>1</sup> Viability locus selection and dominance coefficients: s=0.1 and h=0.2

<sup>644</sup> Viability locus selection and dominance coefficients: s=1 and h=0.02

Table 2 Fixation probabilities at the viability locus in a self-incompatible population under two different selection regimes.

Population Selection regime Rate of asexual reproduction								
size		c=0	c=0.5	c=0.8	c=0.9	c=0.99	c=0.999	<i>c</i> =1
(a) Fixation	of deleterious allele a							
N=25	Mildly deleterious <sup>1</sup>	0.066	0.049	0.056	0.050	0.085	0.081	0.067
<i>N</i> =50	Mildly deleterious	0.000	0.000	0.001	0.000	0.001	0.001	0.001
<i>N</i> =100	Mildly deleterious	0.000	0.000	0.000	0.000	0.000	0.000	0.000
(b) Fixation of allele A								
N=25	Mildly deleterious	0.670	0.714	0.663	0.519	0.264	0.279	0.082
<i>N</i> =25	Lethal recessive <sup>2</sup>	0.831	0.783	0.767	0.581	0.293	0.317	0.085
<i>N</i> =50	Mildly deleterious	0.583	0.541	0.574	0.507	0.474	0.466	0.139
<i>N</i> =50	Lethal recessive	0.661	0.635	0.596	0.564	0.518	0.493	0.150
<i>N</i> =100	Mildly deleterious	0.314	0.306	0.320	0.294	0.274	0.290	0.187
<i>N</i> =100	Lethal recessive	0.437	0.361	0.318	0.313	0.269	0.296	0.189
(c) Fixation of genotype Aa								
N=25	Mildly deleterious	0.000	0.000	0.023	0.184	0.472	0.467	0.723
<i>N</i> =25	Lethal recessive	0.000	0.000	0.013	0.212	0.551	0.569	0.879
<i>N</i> =50	Mildly deleterious	0.000	0.000	0.000	0.000	0.026	0.050	0.585
<i>N</i> =50	Lethal recessive	0.000	0.000	0.000	0.000	0.032	0.041	0.721
<i>N</i> =100	Mildly deleterious	0.000	0.000	0.000	0.000	0.000	0.000	0.230
<i>N</i> =100	Lethal recessive	0.000	0.000	0.000	0.000	0.000	0.000	0.317

The probability of fixation at the viability locus (fixation of genotypes, AA, aa or Aa) was estimated as the proportion of observations in which fixation was recorded.

648

649

651

<sup>650</sup> Viability locus selection and dominance coefficients: s=0.1 and h=0.2

<sup>&</sup>lt;sup>2</sup> Viability locus selection and dominance coefficients: s=1 and h=0.02















