

EINSTEIN HERMITIAN METRICS OF NON NEGATIVE SECTIONAL CURVATURE

EZIO DE ARAUJO COSTA

Abstract

In this paper we will prove that if M is a compact simply connected Hermitian Einstein 4-manifold with non negative sectional curvature then M is isometric to complex projective space $\mathbb{C}\mathbb{P}^2$ with the Fubini-Study metric or M is isometric to a product of two two-spheres $\mathbb{S}^2 \times \mathbb{S}^2$, with their canonical metrics.

1. Introduction

Let $M = M^4$ be a 4-manifold. A Riemannian metric g on M is called *Einstein* if M has constant Ricci curvature and called *Hermitian* if $g(J\cdot, J\cdot) = g(\cdot, \cdot)$ for a complex structure J on M . In [5], C. LeBrun proved the following

Theorem 1. (LeBrun)- *Let $(M = M^4, g, J)$ be a compact connected complex surface with metric g and complex structure J . If g is Einstein and Hermitian with respect to J then only one of the following holds:*

- (1) g is Kaehler-Einstein with positive Ricci curvature.
- (2) M is isometric to $\mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$ and g is the Page metric.
- (3) M is isometric to $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ and g is the Chen-LeBrun-Weber metric.

Using the previous theorem, C. Koca proved in [4] the following:

Theorem 2. (Koca)-*Let $(M = M^4, g, J)$ be a compact complex surface with metric g and complex structure J . If g is Einstein and Hermitian with respect to J and g has positive sectional curvature then M is isometric to complex projective space $\mathbb{C}\mathbb{P}^2$ with the Fubini-Study metric.*

Now, consider M a compact simply connected Kaehler-Einstein 4-manifold with non negative sectional curvature. In this case, M. Berger proved in [1], that M is isometric to complex projective space $\mathbb{C}\mathbb{P}^2$ with the Fubini-Study metric or isometric to a product of two spheres $\mathbb{S}^2 \times \mathbb{S}^2$, with their canonical metrics.

In the next two sections we will prove that the Page metric and the Chen-LeBrun-Weber metrics no has non negative sectional curvature. This will conclude the proof our main result:

Theorem 3. *Let $(M = M^4, g, J)$ be a compact simply connected complex surface with metric g and complex structure J . If g is Einstein and Hermitian with respect to J and g has non negative sectional curvature then M is isometric to complex projective space $\mathbb{C}\mathbb{P}^2$ with the Fubini-Study metric or M is isometric to a product of two spheres $\mathbb{S}^2 \times \mathbb{S}^2$, with their canonical metrics.*

2. Page metric

The Page metric (see [6]) lives in connected sum $\mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$, where $\overline{\mathbb{C}\mathbb{P}^2}$ is the complex projective space $\mathbb{C}\mathbb{P}^2$ with opposite orientation. In [4], C. Koca showed that the Page metric no has non negative sectional curvature using a computer program like *Maple*. In this section we will prove this result using a different argument. For this consider the page metric g as in Koca [4] :

$$g = W^2(x)dx^2 + g^2(x)(\sigma_1^2 + \sigma_2^2) + \frac{D^2}{W(x)}\sigma_3^2$$

where $x \in (-1, 1)$,

$$W(x) = \sqrt{\frac{1 - a^2x^2}{[3 - a^2 - a^2(1 + a^2)x^2](1 - x^2)}},$$

$$g(x) = \frac{2}{\sqrt{3 + 6a^2 - a^4}} \sqrt{1 - a^2x^2},$$

$D = \frac{2}{3+a^2}$ and a is the unique positive root of equation $f(x) = x^4 + 4x^3 - 6x^2 + 12x - 3 = 0$. Notice that $a < 1$, since that $f(0) = -3$ and $f(1) = 8$.

In accord with Koca, there exists a two-plane where the sectional curvature satisfies

$$K_{01} = 2\left[\frac{g'W' - g''W}{gW^3}\right].$$

Then we have

$$K_{01} = -\frac{2}{gW}F',$$

where

$$F = \frac{g'}{W}.$$

Claim: There exist $c \in [0, 1)$ such that $F'(c) > 0$.

Proof of Claim :

Notice that $g' = -Ax(1 - a^2x^2)^{-1/2}$, where $A = \frac{2a}{\sqrt{3+6a^2-a^4}} > 0$. Moreover,

$$F = -Ax\sqrt{[3 - a^2 - a^2(1 + a^2)x^2](1 - x^2)}(1 - a^2x^2)^{-1},$$

where $1 - a^2x^2 > 0$. Assumes that $F'(x) \leq 0$ for all $x \in [0, 1)$.

Then F is a increasing function in $[0, 1)$ and follows of this that $F(y) \leq F(x) \leq F(0) = 0$, for all $y > x \in [0, 1)$. Since that F is continuo in $[0, 1]$, we have that $0 = F(1) \leq F(x) \leq 0$, for all $x \in [0, 1)$. So $F = 0$ in $[0, 1)$ (contradiction)

This proves that there exists points where $K_{01} < 0$.

3. Chen-LeBrun-Weber metric

In [2], Chen, LeBrun and Weber proved that $M = \mathbb{C}\mathbb{P}^2 \# 2\mathbb{C}\mathbb{P}^2$ admits an Hermitian non-Kaehler Einstein metric g . In particular, there exists a Kaehler metric h on M of positive scalar curvature s such that $g = s^2h$. Now consider W_h^+ and W_g^+ the self-dual Weyl part of the Weyl tensor W of the respective metrics h and g . Since that h and g are conformally related we have $W_g^+ = \frac{1}{s^2}W_h^+$. On the other hand, the self-dual tensor W_h^+ of the Kaehler metric h has exactly two different eigenvalues and so W_g^+ has also two different eigenvalues. By Proposition 4 of Derdzinski [3], M admits a non trivial Killing vector field with respect to metric g . Assumes that M has non negative sectional curvature with respect to metric g . Then M is a compact simply connected 4-manifold with non negative sectional curvature and with a non trivial Killing vector field. By Theorem 1 of Searle and Yang in [7], we have that the Euler characteristic of M satisfies $\chi(M) \leq 4$ which contradicts the fact of that $\chi(\mathbb{C}\mathbb{P}^2 \# 2\mathbb{C}\mathbb{P}^2) = 5$.

REFERENCES

- [1] Berger, M. *Les varietes Kaehleriennes compacts d'Einstein de dimension quatre a courbure positive*. Tensor. **13**(1963) 71-74.
- [2] Chen, X. LeBrun, C. and Weber B. *On Conformally Kahler Einstein Manifolds*. Journ. Amer. mth. Soc. **21**(2008) 1137-1168.
- [3] Derdzinsky, A. *Self-dual Kaehler manifolds and Einstein manifolds of dimension four*. Comp. Math. **49**(1983) 405-433.
- [4] Koca, C. *Einstein Hermitian Metrics of Positive Sectional Curvature*. arXiv:1112.4181v1 [math.DG] 18 Dec 2011.
- [5] LeBrun, C. *On Einstein Hermitian 4-manifold*. arXiv : 1010.0238v1 [math.DG] 10 Oct 2010.
- [6] Page, D. *A compact rotating gravitational instanton*. Phys. Lett. B, **79** (1979) 235-238.
- [7] Searle, C. and Yang, D. *On the topology of nonnegatively curved simply connected 4-manifolds with continous symmetry*. Duke Math J., **74**(1994), 547-556.

Author's address:

Mathematics Department, Federal University of Bahia,

zipcode: 40170110- Salvador -Bahia-Brazil

Author's email

ezio@ufba.br