# A NOTE ON A BRUNN-MINKOWSKI INEQUALITY FOR THE GAUSSIAN MEASURE

#### PIOTR NAYAR AND TOMASZ TKOCZ

ABSTRACT. We give the counter-examples related to a Gaussian Brunn-Minkowski inequality and the (B) conjecture.

### 1. INTRODUCTION AND NOTATION

Let  $\gamma_n$  be the standard Gaussian distribution on  $\mathbb{R}^n$ , i.e. the measure with the density

$$g_n(x) = \frac{1}{(2\pi)^{n/2}} e^{-|x|^2/2},$$

where  $|\cdot|$  stands for the standard Euclidean norm. A powerful tool in convex geometry is the Brunn-Minkowski inequality for Lebesgue measure (see [Sch] for more information). Concerning the Gaussian measure, the following question has recently been posed.

**Question** (R. Gardner and A. Zvavitch, [GZ]). Let  $0 < \lambda < 1$  and let A and B be closed convex sets in  $\mathbb{R}^n$  such that  $o \in A \cap B$ . Is it true that

(GBM) 
$$\gamma_n(\lambda A + (1-\lambda)B)^{1/n} \ge \lambda \gamma_n(A)^{1/n} + (1-\lambda)\gamma_n(B)^{1/n}?$$

A counter-example is given in this note. However, we believe that this question has an affirmative answer in the case of o-symmetric convex sets, i.e. the sets satisfying K = -K.

In [CFM] it is proved that for an o-symmetric convex set K in  $\mathbb{R}^n$  the function

(1) 
$$\mathbb{R} \ni t \mapsto \gamma_n(e^t K),$$

is log-concave. This was conjectured by W. Banaszczyk and popularized by R. Latała [Lat]. It turns out that the (B) conjecture cannot be extended to the class of sets which are not necessarily *o*-symmetric yet contain the origin, as one of the sets provided in our counter-example shows.

As for the notation, we frequently use the function

$$T(x) = \frac{1}{\sqrt{2\pi}} \int_x^\infty e^{-t^2/2} \mathrm{d}t.$$

<sup>2010</sup> Mathematics Subject Classification. Primary 52A40; Secondary 60G15.

Key words and phrases. Convex body, Gauss measure, Brunn-Minkowski inequality, B-conjecture.

Research of the first author partially supported by NCN Grant no. 2011/01/N/ST1/01839.

Research of the second author partially supported by NCN Grant no.  $2011/01/\mathrm{N}/\mathrm{ST1}/05960.$ 

### 2. Counter-examples

Now we construct the convex sets  $A, B \subset \mathbb{R}^2$  containing the origin such that inequality (GBM) does not hold. Later on we show that for the set B the (B) conjecture is not true.

Fix  $\alpha \in (0, \pi/2)$  and  $\varepsilon > 0$ . Take

$$A = \{ (x, y) \in \mathbb{R}^2 \mid y \ge |x| \tan \alpha \},\$$
  
$$B = B_{\varepsilon} = \{ (x, y) \in \mathbb{R}^2 \mid y \ge |x| \tan \alpha - \varepsilon \} = A - (0, \varepsilon).$$

Clearly, A, B are convex and  $0 \in A \cap B$ . Moreover, from convexity of A we have  $\lambda A + (1 - \lambda)A = A$  and therefore

$$\lambda A + (1 - \lambda)B = \lambda A + (1 - \lambda)(A - (0, \varepsilon)) = A - (1 - \lambda)(0, \varepsilon).$$

Observe that

$$\gamma_2(A) = \frac{1}{2} - \frac{\alpha}{\pi},$$
  

$$\gamma_2(B) = 2 \int_0^{+\infty} T(x \tan \alpha - \varepsilon) \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \, \mathrm{d}x,$$
  

$$\gamma_2(\lambda A + (1 - \lambda)B) = 2 \int_0^{+\infty} T(x \tan \alpha - \varepsilon (1 - \lambda)) \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \, \mathrm{d}x$$

and that these expressions are analytic functions of  $\varepsilon$ . We will expand these functions in  $\varepsilon$  up to the order 2. Let

$$a_k = \int_0^{+\infty} T^{(k)}(x \tan \alpha) \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \, \mathrm{d}x,$$

for k = 0, 1, 2, where  $T^{(k)}$  is the k-th derivative of T (we adopt the standard notation  $T^{(0)} = T$ ). We get

$$\begin{split} \gamma_2(A) &= 2a_0, \\ \gamma_2(B) &= 2a_0 - 2\varepsilon a_1 + \varepsilon^2 a_2 + o(\varepsilon^2), \\ \gamma_2(\lambda A + (1-\lambda)B) &= 2a_0 - 2\varepsilon (1-\lambda)a_1 + \varepsilon^2 (1-\lambda)^2 a_2 + o(\varepsilon^2). \end{split}$$

Thus

$$\sqrt{\gamma_2(B)} = \sqrt{2a_0} - \frac{a_1}{\sqrt{2a_0}}\varepsilon + \left(\frac{a_2}{2\sqrt{2a_0}} - \frac{a_1^2}{2(2a_0)^{3/2}}\right)\varepsilon^2 + o(\varepsilon^2).$$

Taking  $\varepsilon(1-\lambda)$  instead of  $\varepsilon$  we obtain

$$\sqrt{\gamma_2(\lambda A + (1-\lambda)B)} = \sqrt{2a_0} - \frac{a_1}{\sqrt{2a_0}}(1-\lambda)\varepsilon + \left(\frac{a_2}{2\sqrt{2a_0}} - \frac{a_1^2}{2(2a_0)^{3/2}}\right)(1-\lambda)^2\varepsilon^2 + o(\varepsilon^2).$$

Since

$$\sqrt{\gamma_2(\lambda A + (1-\lambda)B)} - \lambda\sqrt{\gamma_2(A)} - (1-\lambda)\sqrt{\gamma_2(B)}$$
$$= -\lambda(1-\lambda)\frac{1}{2(2a_0)^{3/2}}(2a_0a_2 - a_1^2)\varepsilon^2 + o(\varepsilon^2),$$

we will have a counter-example if we find  $\alpha \in (0, \pi/2)$  such that

$$2a_0a_2 - a_1^2 > 0$$

Recall that  $a_0 = \frac{1}{2}\gamma_2(A) = \frac{1}{2}\left(\frac{1}{2} - \frac{\alpha}{\pi}\right)$ . The integrals that define the  $a_k$ 's can be calculated. Namely,

$$a_{1} = \int_{0}^{\infty} T'(x \tan \alpha) \frac{e^{-x^{2}/2}}{\sqrt{2\pi}} dx = -\frac{1}{\sqrt{2\pi}} \frac{1}{2} \int_{\mathbb{R}} e^{-(1+\tan^{2}\alpha)x^{2}/2} \frac{dx}{\sqrt{2\pi}}$$
$$= -\frac{1}{\sqrt{2\pi}} \frac{1}{2\sqrt{1+\tan^{2}\alpha}},$$
$$a_{2} = \int_{0}^{\infty} T''(x \tan \alpha) \frac{e^{-x^{2}/2}}{\sqrt{2\pi}} dx = \frac{1}{\sqrt{2\pi}} \int_{0}^{\infty} (x \tan \alpha) e^{-(1+\tan^{2}\alpha)x^{2}/2} \frac{dx}{\sqrt{2\pi}}$$
$$= \frac{1}{2\pi} \frac{\tan \alpha}{1+\tan^{2}\alpha}.$$

Therefore,

$$2a_0a_2 - a_1^2 = 2\left(\frac{1}{2}\left(\frac{1}{2} - \frac{\alpha}{\pi}\right) \cdot \frac{1}{2\pi} \frac{\tan\alpha}{1 + \tan^2\alpha}\right) - \frac{1}{2\pi} \cdot \frac{1}{4(1 + \tan^2\alpha)}$$
$$= \frac{1}{8\pi} \frac{1}{1 + \tan^2\alpha} \left(\tan\alpha\left(2 - \frac{4\alpha}{\pi}\right) - 1\right),$$

which is positive for  $\alpha$  close to  $\pi/2$ .

Now we turn our attention to the (B) conjecture. We are to check that for the set  $B = B_{\varepsilon}$  the function  $\mathbb{R} \ni t \mapsto \gamma_n(e^t B)$  is not log-concave, provided that  $\varepsilon$  is sufficiently small. Since

$$e^t B = \{(x, y) \in \mathbb{R}^2 \mid y \ge \tan \alpha |x| - \varepsilon e^t\}$$

we get

$$\ln \gamma_2(e^t B) = \ln \left( 2 \int_0^\infty T(x \tan \alpha - e^t \varepsilon) \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right)$$
$$= \ln \left( 2 \int_0^\infty T(x \tan \alpha) \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx \right) - \varepsilon e^t \frac{\int_0^\infty T'(x \tan \alpha) e^{-x^2/2} dx}{\int_0^\infty T(x \tan \alpha) e^{-x^2/2} dx} + o(\varepsilon).$$

This produces the desired counter-example for sufficiently small  $\varepsilon$  as the function  $t \mapsto \beta e^t$ , where

$$\beta = -\frac{\int_0^\infty T'(x\tan\alpha)e^{-x^2/2}\mathrm{d}x}{\int_0^\infty T(x\tan\alpha)e^{-x^2/2}\mathrm{d}x} > 0,$$

is convex.

*Remark.* The set  $B_{\varepsilon}$  which serves as a counter-example to the (B) conjecture in the nonsymmetric case works when the parameter  $\alpha = 0$  as well (and  $\varepsilon$ is sufficiently small). Since  $B_{\varepsilon}$  is simply a halfspace in this case, it shows that symmetry of K is required for log-concavity of (1) even in the onedimensional case.

## Acknowledgements

The authors would like to thank Professors R. Gardner and A. Zvavitch for pointing out that the constructed set may also serve as a counter-example to the (B) conjecture in the non-symmetric case. An anonymous referee deserves thanks for the remark.

#### References

- [CFM] D. Cordero-Erausquin, M. Fradelizi, and B. Maurey, The (B) conjecture for the Gaussian measure of dilates of symmetric convex sets and related problems, J. Funct. Anal. 214 (2004), no. 2, 410–427. MR2083308 (2005g:60064)
- [GZ] R. J. Gardner, A. Zvavitch, Gaussian Brunn-Minkowski-type inequalities, Trans. Amer. Math. Soc. 360 (2010), no. 10, 5333–5353. MR2657682 (Review)
- [Lat] R. Latała, On some inequalities for Gaussian measures, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 813–822, Higher Ed. Press, Beijing, 2002. MR1957087 (2004b:60055)
- [Sch] R. Schneider, Convex bodies: the Brunn-Minkowski theory. Encyclopedia of Mathematics and its Applications, 44. Cambridge University Press, Cambridge, 1993. MR1216521 (94d:52007)

PIOTR NAYAR, INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097 WARSZAWA, POLAND.

*E-mail address*: nayar@mimuw.edu.pl

TOMASZ TKOCZ, INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097 WARSZAWA, POLAND.

*E-mail address*: t.tkocz@mimuw.edu.pl