

RIEMANN-ROCH THEORY ON FINITE SETS

RODNEY JAMES AND RICK MIRANDA

ABSTRACT. In [1] M. Baker and S. Norine developed a theory of divisors and linear systems on graphs, and proved a Riemann-Roch Theorem for these objects (conceived as integer-valued functions on the vertices). In [2] and [3] the authors generalized these concepts to real-valued functions, and proved a corresponding Riemann-Roch Theorem in that setting, showing that it implied the Baker-Norine result. In this article we prove a Riemann-Roch Theorem in a more general combinatorial setting that is not necessarily driven by the existence of a graph.

1. INTRODUCTION

Baker and Norine showed in [1] that a Riemann-Roch formula holds for an analogue of linear systems defined on the vertices of finite connected graphs. There, the image of the graph Laplacian induces an equivalence relation on the group of *divisors* of the graph, which are integer-valued functions defined on the set of vertices. This equivalence relation is the analogue of linear equivalence in the classical algebro-geometric setting.

We showed in [2] that the Baker-Norine result implies a generalization of the Riemann-Roch formula to edge-weighted graphs, where the edge weights can be R -valued, where R is an arbitrary subring of the reals; the equivalence relation induced by image of the edge-weighted graph Laplacian applies equally well to divisors which are R -valued functions defined on the set of vertices. In [3], we proved our version of the R -valued Riemann-Roch theorem from first principles; this gave an independent proof of the Baker-Norine result as well.

The notion of linear equivalence above is induced by the appropriate graph Laplacian acting on the group of divisors, which may be viewed as points in \mathbb{Z}^n (the Baker-Norine case) or more generally R^n , where n is the number of vertices of the graph. In this paper, we propose a generalization of this Riemann-Roch formula for graphs where linear equivalence is induced by a group action on the points of R^n . The setup we will use is as follows.

Choose a subring R of the reals, and fix a positive integer n . Let V be the group of points in R^n under component-wise addition. If $x \in V$, we will use the functional notation $x(i)$ to denote the i -th component of x .

For any $x \in V$, define the *degree* of x as

$$\deg(x) = \sum_{i=1}^n x(i).$$

For any $d \in R$, define the subset $V_d \subset V$ to be

$$V_d = \{x \in V \mid \deg(x) = d\}.$$

Note that the subset V_0 is a subgroup; for any d , V_d is a coset of V_0 in V .

Let $H \subset V_0$ be a subgroup of V_0 and consider the action on V by H by translation: if $h \in H$ and $x \in V$, then $(h, x) \mapsto h + x$. This action of H on V induces the equivalence relation $x \sim y$ if and only if $x - y \in H$; or equivalently, $x \sim y$ if and only if there is a $h \in H$ such that $x = (h, y)$.

Fix the parameter $g \in R$, which we call the *genus*, and choose a set $\mathcal{N} \subset V_{g-1}$. For $x \in V$, define

$$\begin{aligned} x^+ &= \max(x, 0) \\ x^- &= \min(x, 0) \end{aligned}$$

where \max and \min are evaluated at each coordinate. It follows that $x = x^+ + x^-$ and $x^+ = -(-x)^-$. We then define the *dimension* of $x \in V$ to be

$$\ell(x) = \min_{\nu \in \mathcal{N}} \{\deg((x - \nu)^+)\}.$$

We will discuss the motivation for this definition of dimension in the next section.

We can now state our main result.

Theorem 1.1. *Let V be the additive group of points in R^n for a subring $R \subset \mathbb{R}$, and let H be a subgroup of V_0 that acts by translation on V . Fix $g \in R$. Suppose $\kappa \in V_{2g-2}$, and $\mathcal{N} \subset V_{g-1}$, satisfying the symmetry condition*

$$\nu \in \mathcal{N} \iff \kappa - \nu \in \mathcal{N}.$$

Then for every $x \in V$,

$$\ell(x) - \ell(\kappa - x) = \deg(x) - g + 1$$

We will give a proof of Theorem 1.1 in §2. In §3, we will give examples of H , κ , and \mathcal{N} (coming from the graph setting) which satisfy the conditions of Theorem 1.1, and show how this Riemann-Roch formulation is equivalent to that given in [3]. Finally in §4, we give examples that do not arise from graphs.

2. PROOF OF RIEMANN-ROCH FORMULA

The dimension of $x \in V$

$$\ell(x) = \min_{\nu \in \mathcal{N}} \{\deg((x - \nu)^+)\}$$

can be written as

$$\ell(x) = \min_{\nu \in \mathcal{N}} \left\{ \sum_{i=1}^n \max\{x(i) - \nu(i), 0\} \right\}.$$

If $x(i) \geq \nu(i)$ for each i , $\sum_{i=1}^n \max\{x(i) - \nu(i), 0\}$ is the *taxicab* distance from x to ν . Thus, $\ell(x)$ is the taxicab distance from x to the portion of the set \mathcal{N} such that $x \geq \mathcal{N}$, where the inequality is evaluated at each component.

We will now proceed with the proof of the Riemann-Roch formula.

Proof. (Theorem 1.1)

Suppose that $\mathcal{N} \subset V_{g-1}$ and $\kappa \in V$ satisfy the symmetry condition. We can then write

$$\begin{aligned} \ell(\kappa - x) &= \min_{\nu \in \mathcal{N}} \{\deg((\kappa - x - \nu)^+)\} \\ &= \min_{\nu \in \mathcal{N}} \{\deg(((\kappa - \nu) - x)^+)\} \\ &= \min_{\mu \in \mathcal{N}} \{\deg((\mu - x)^+)\}. \end{aligned}$$

Using the identities $x = x^+ + x^-$ and $x^+ = -(x^-)$, we have

$$\begin{aligned} \min_{\mu \in \mathcal{N}} \{\deg((\mu - x)^+)\} &= \min_{\mu \in \mathcal{N}} \{\deg((\mu - x)) - \deg((\mu - x)^-)\} \\ &= \min_{\mu \in \mathcal{N}} \{\deg((\mu - x)) + \deg((x - \mu)^+)\}. \end{aligned}$$

Since $\mu \in \mathcal{N}$ we know that $\deg(\mu) = g - 1$, thus $\deg(\mu - x) = g - 1 - \deg(x)$ and thus

$$\begin{aligned} \ell(\kappa - x) &= \deg((\mu - x)) + \min_{\mu \in \mathcal{N}} \{\deg((x - \mu)^+)\} \\ &= g - 1 - \deg(x) + \ell(x). \end{aligned}$$

□

Note that $\kappa \in V$ is the analogue to the canonical divisor in the classical Riemann-Roch formula.

3. GRAPH EXAMPLES

Let Γ be a finite, edge-weighted connected simple graph with n vertices. We will assume that Γ has no loops. Let $w_{ij} \in R$ with $w_{ij} \geq 0$ be the weight of the edge connecting vertices v_i and v_j . The no loops assumption is also applied to the edge weights so that $w_{ii} = 0$ for each i . We showed in [3] that such a graph satisfies an equivalent Riemann-Roch formula as in Theorem 1.1.

In this setting, $H = \langle h_1, h_2, \dots, h_{n-1} \rangle$ where each $h_i \in R^n$ is defined as

$$h_i(j) = \begin{cases} \deg(v_i) & \text{if } i = j \\ -w_{ij} & \text{if } i \neq j. \end{cases}$$

(Here $\deg(v)$ for a vertex v is the sum of the weights of the edges incident to v .) Note that H is the edge-weighted Laplacian of Γ .

As shown in [3], the set $\mathcal{N} \subset V_{g-1}$ is generated by a set $\{\nu_1, \dots, \nu_s\}$ as follows. Fix a vertex v_k and let (j_1, \dots, j_n) be a permutation of $(1, \dots, n)$ such that $j_1 = k$. There are then $(n-1)!$ such permutations; for each permutation, we compute a $\nu \in V_{g-1}$ defined by

$$\nu(j_l) = \begin{cases} -1 & \text{if } l = 1 \\ -1 + \sum_{i=1}^{l-1} w_{j_i j_l} & \text{if } l > 1. \end{cases}$$

Each such ν may not be unique; set s to be the number of unique ν 's and index this set $\{\nu_1, \dots, \nu_s\}$. We then define the set \mathcal{N} as

$$\mathcal{N} = \{x \in V \mid x \sim \nu_i \text{ for some } i = 1, \dots, s\}.$$

The canonical element κ is defined by $\kappa(j) = \deg(v_j) - 2$, and the genus $g = 1 + \sum_{i < j} w_{ij} - n$.

As an example, consider a two vertex graph Γ with edge weight $w_{12} = p > 0$. Then $g = p - 1$ and $H = \langle (p, -p) \rangle = \mathbb{Z}(p, -p)$. The set $\mathcal{N} \subset V_{g-1}$ is $\mathcal{N} = \{\nu \mid \nu \sim (p-1, -1)\}$ and $\kappa = (p-2, p-2)$. Figure 1 shows the divisors $x \in \mathbb{R}^2$ for this graph in the plane. The shaded region, which is bounded by the corner points in the set \mathcal{N} , represent points x with $\ell(x) = 0$.

We can show directly that \mathcal{N} and κ for the two-vertex graph example satisfy the necessary condition for Theorem 1.1 to hold. If $\kappa - x \in \mathcal{N}$, then $\kappa - x =$




FIGURE 1. Divisors in \mathbb{R}^2 for a two-vertex graph with p edges. The shaded region represents points $x \in \mathbb{R}^2$ with $\ell(x) = 0$; for a general point $x \in \mathbb{R}^2$, $\ell(x)$ is the taxicab distance to the shaded region.

$m(p, -p) + (p - 1, -1)$ for some $m \in \mathbb{Z}$. Solving for x , we have

$$\begin{aligned} x &= (p - 2, p - 2) - m(p, -p) - (p - 1, -1) \\ &= (-mp - 1, mp + p - 1) \\ &= (p - 1, -1) - (m + 1)(p, -p) \end{aligned}$$

and thus $x \in \mathcal{N}$. Similarly, if $\nu \in \mathcal{N}$, it easily follows that $\kappa - \nu \in \mathcal{N}$.

Now consider a three vertex graph with edge weights $w_{12} = p$, $w_{13} = q$ and $w_{23} = r$. The set \mathcal{N} can be generated by $\nu_1 = (-1, p - 1, q + r - 1)$ and $\nu_2 = (-1, p + r - 1, q - 1)$; H can be generated by $h_1 = (p + q, -p, -q)$ and $h_2 = (-p, p + r, -r)$. In Figure 2, the region representing points $x \in \mathbb{R}^3$ such that $\ell(x) = 0$ is shown for a three vertex graph with edge weights $p = 1$, $q = 3$ and $r = 4$.

4. NON-GRAPH EXAMPLES

The main result of this paper would not be interesting if there were no examples of subgroups $H \subset V_0$ with \mathcal{N} and κ that were not derived from graphs.

Theorem 4.1. *There are subgroups $H \subset V_0$ with $\kappa \in V$ and $\mathcal{N} \subset V_{g-1}$ such that Theorem 1.1 holds where H is not the Laplacian of a finite connected graph.*

Proof. Let $n = 2$ and choose $H = \langle (-4, 4) \rangle$, $\mathcal{N} = \{\nu \in G \mid \nu \sim (2, -2)\}$ with $\kappa = (0, 0)$ and $g = 1$. If H were generated from a two vertex graph, using the notation from the previous section we would have $p = 4$. This would require $\kappa = (2, 2)$ with \mathcal{N} generated by $\nu_1 = (3, -1)$.

Since there is no integer m such that $\kappa = (0, 0) = (2, 2) + m(-4, 4)$ (and likewise there is no m such that $\nu_1 = (2, -2) = (3, -1) + m(-4, 4)$), H cannot be generated from a two vertex graph.




FIGURE 2. Divisors in \mathbb{R}^3 for a three-vertex graph edge weights $w_{12} = 1$, $w_{13} = 3$ and $w_{23} = 4$. The solid region represents points $x \in \mathbb{R}^3$ with $\ell(x) = 0$; for a general point $x \in \mathbb{R}^3$, $\ell(x)$ is the taxicab distance to the surface.

Now suppose that $\nu \in \mathcal{N}$. Then $\nu = (2, -2) + m(-4, 4)$ for some integer m , and

$$\begin{aligned} \kappa - \nu &= (0, 0) - (2, -2) - m(-4, 4) \\ &= (2, -2) - (m - 1)(-4, 4) \end{aligned}$$

thus $\kappa - \nu \in \mathcal{N}$.

Similarly, if $\kappa - \nu \in \mathcal{N}$, $\kappa - \nu = (2, -2) + m(-4, 4)$ for some integer m , and

$$\begin{aligned} \nu &= \kappa - (2, -2) - m(-4, 4) \\ &= (4m - 2, -4m + 2) \\ &= (2, -2) - (m - 1)(-4, 4) \end{aligned}$$

thus $\nu \in \mathcal{N}$ and H, κ, \mathcal{N} satisfies Theorem 1.1. \square

We include in Figure 3 a representation of the divisors $x \in \mathbb{R}^2$ with $\ell(x) = 0$ for the example used in the proof of Theorem 4.1. The plot is identical to that of a two vertex graph with $p = 4$ but is shifted by -1 in each direction.

It is also possible to produce non-graph examples by using more generators for \mathcal{N} . In Figure 4, the divisors $x \in \mathbb{R}^2$ with $\ell(x) = 0$ are shown where \mathcal{N} is generated by two points $\nu_1 = (0, 4)$ and $\nu_2 = (1, 3)$, using $H = \langle (-3, 3) \rangle$ and $\kappa = (0, 0)$.




FIGURE 3. Divisors $x \in \mathbb{R}^2$ with $\ell(x) = 0$ for the non-graph example in the proof of Theorem 4.1. Note that this example is identical to the two vertex graph example in Figure 1 with $p = 4$, but shifted by $(-1, -1)$.




FIGURE 4. Divisors $x \in \mathbb{R}^2$ with $\ell(x) = 0$ for a non-graph example with \mathcal{N} generated by $(0, 4)$ and $(1, 3)$, $H = \langle (-3, 3) \rangle$ and $\kappa = (0, 0)$.

REFERENCES

- [1] Matthew Baker and Serguei Norine, Riemann-Roch and Abel-Jacobi Theory on a Finite Graph, *Advances in Mathematics* 215 (2007) 766-788.
- [2] Rodney James and Rick Miranda, A Riemann-Roch theorem for edge-weighted graphs, arXiv:0908.1197v2 [math.AG], 2009.
- [3] Rodney James and Rick Miranda, Linear Systems on Edge-Weighted Graphs, arXiv:1105.0227v1 [math.AG], 2011.

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF COLORADO,
DENVER, CO, USA

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO, USA