

# DISCRIMINANTS, SYMMETRIZED GRAPH MONOMIALS, AND SUMS OF SQUARES CERTIFICATES

PER ALEXANDERSSON

ABSTRACT. Here we present certificates for 5 classes of 6-edged multigraphs whose symmetrized graph monomials may be represented as sum of squares, but not as linear combinations of partition square graphs. This is a complement to the results presented in [1].

## 1. BACKGROUND

In [1], we have the following definition:

**Definition 1.** Let  $g$  be a directed graph, with vertices  $x_1, \dots, x_n$  and adjacency matrix  $(a_{ij})$ , being non-negative integers. Define first its graph monomial  $P_g$  as follows

$$P_g(x_1, \dots, x_n) := \prod_{1 \leq i, j \leq n} (x_i - x_j)^{a_{ij}},$$

where  $a_{ij}$  is the number of directed edges joining  $x_i$  with  $x_j$ .

The symmetrized graph monomial of  $g$  is defined as

$$\tilde{g}(\mathbf{x}) = \sum_{\sigma \in S_n} P_g(\sigma \mathbf{x}), \quad \mathbf{x} = x_1, \dots, x_n.$$

Two multigraphs are *equivalent* if the first corresponding symmetrized graph monomial is a constant multiple of the other.

We study the set of symmetrized graph monomials obtained from multigraphs with 6 edges. There are 212 such multigraphs.

One main result in [1] is the following proposition:

**Proposition 1.** (i) 102 graphs with 6 edges have identically vanishing symmetrized graph monomial. (ii) The remaining 110 graphs are divided into 27 equivalence classes. (iii) 12 of these classes can be expressed as non-negative linear combinations of square graphs, i.e. lie in the convex cone spanned by the square graphs. (iv) Of the remaining 15 classes, symmetrized graph monomial of 7 of them change sign. (v) Of the remaining 8 classes (which are presented in Fig. 1) the first 5 are sums of squares.

Proposition (i),(ii), (iii) and (iv) follows from direct computation.

## 2. CERTIFICATES

Here we give certificates that the symmetrized graph monomial for the first 5 classes given in Fig. 1 are sum of squares, corresponding to the last case in Prop. 1. The graphs in each row yield the same polynomial, up to a constant. The symmetrized graph monomial from row  $i$  is a constant multiple of the polynomial

$v_i Q_i v_i^T$ , where  $v_i$  is the coefficient vector and  $Q_i$  is the corresponding symmetric positive semi-definite matrix, given below. It is well-known that this certifies that the first 5 classes of graphs are sum of squares.

It is relatively straightforward to verify that the polynomials indeed are non-negative, using methods similarly to [2]. By using Matlab together with Yalmip, one may verify that the last three classes cannot be expressed as sums of squares.

$$v_1 = \{x_1^2 x_2, x_1 x_2^2, x_1^2 x_3, x_2^2 x_3, x_1 x_3^2, x_2 x_3^2, x_1^2 x_4, x_2^2 x_4, x_3^2 x_4, x_1 x_4^2, x_2 x_4^2, x_3 x_4^2, \\ x_1^2 x_5, x_2^2 x_5, x_3^2 x_5, x_4^2 x_5, x_1 x_5^2, x_2 x_5^2, x_3 x_5^2, x_4 x_5^2, x_5 x_6^2, x_5^2 x_6, x_4 x_6^2, x_4^2 x_6, \\ x_3 x_6^2, x_3^2 x_6, x_2 x_6^2, x_2^2 x_6, x_1 x_6^2, x_1^2 x_6, x_1 x_2 x_3, x_1 x_2 x_4, x_1 x_3 x_4, x_2 x_3 x_4, x_1 x_2 x_5, \\ x_1 x_3 x_5, x_2 x_3 x_5, x_1 x_4 x_5, x_2 x_4 x_5, x_3 x_4 x_5, x_4 x_5 x_6, x_3 x_5 x_6, x_3 x_4 x_6, x_2 x_5 x_6, x_2 x_4 x_6, x_2 x_3 x_6, \\ x_1 x_5 x_6, x_1 x_4 x_6, x_1 x_3 x_6, x_1 x_2 x_6\}$$

$$v_2 = \{x_5 x_6^2, x_5^2 x_6, x_4 x_6^2, x_4 x_5 x_6, x_4 x_5^2, x_4^2 x_6, x_4^2 x_5, x_3 x_6^2, x_3 x_5 x_6, x_3 x_5^2, x_3 x_4 x_6, \\ x_3 x_4 x_5, x_3 x_4^2, x_3^2 x_6, x_3^2 x_5, x_3^2 x_4, x_2 x_6^2, x_2 x_5 x_6, x_2 x_5^2, x_2 x_4 x_6, x_2 x_4 x_5, x_2 x_4^2, \\ x_2 x_3 x_6, x_2 x_3 x_5, x_2 x_3 x_4, x_2 x_3^2, x_2^2 x_6, x_2^2 x_5, x_2^2 x_4, x_2^2 x_3, x_1 x_6^2, x_1 x_5 x_6, x_1 x_5^2, \\ x_1 x_4 x_6, x_1 x_4 x_5, x_1 x_4^2, x_1 x_3 x_6, x_1 x_3 x_5, x_1 x_3 x_4, x_1 x_3^2, x_1 x_2 x_6, x_1 x_2 x_5, x_1 x_2 x_4, \\ x_1 x_2 x_3, x_1 x_2^2, x_1^2 x_6, x_1^2 x_5, x_1^2 x_4, x_1^2 x_3, x_1^2 x_2\}$$

$$v_3 = \{x_3 x_4^2, x_3^2 x_4, x_2 x_4^2, x_2 x_3 x_4, x_2 x_3^2, x_2^2 x_4, x_2^2 x_3, x_1 x_4^2, x_1 x_3 x_4, x_1 x_3^2, x_1 x_2 x_4, \\ x_1 x_2 x_3, x_1 x_2^2, x_1^2 x_4, x_1^2 x_3, x_1^2 x_2\}$$

$$v_4 = \{x_4^3, x_3 x_4^2, x_3^2 x_4, x_3^3, x_2 x_4^2, x_2 x_3 x_4, x_2 x_3^2, x_2^2 x_4, x_2^2 x_3, x_2^3, x_1 x_4^2, x_1 x_3 x_4, x_1 x_3^2, \\ x_1 x_2 x_4, x_1 x_2 x_3, x_1 x_2^2, x_1^2 x_4, x_1^2 x_3, x_1^2 x_2, x_1^3\}$$

$$v_5 = \{x_5 x_6^2, x_5^2 x_6, x_4 x_6^2, x_4 x_5 x_6, x_4 x_5^2, x_4^2 x_6, x_4^2 x_5, x_3 x_6^2, x_3 x_5 x_6, x_3 x_5^2, x_3 x_4 x_6, \\ x_3 x_4 x_5, x_3 x_4^2, x_3^2 x_6, x_3^2 x_5, x_3^2 x_4, x_2 x_6^2, x_2 x_5 x_6, x_2 x_5^2, x_2 x_4 x_6, x_2 x_4 x_5, x_2 x_4^2, \\ x_2 x_3 x_6, x_2 x_3 x_5, x_2 x_3 x_4, x_2 x_3^2, x_2^2 x_6, x_2^2 x_5, x_2^2 x_4, x_2^2 x_3, x_1 x_6^2, x_1 x_5 x_6, x_1 x_5^2, \\ x_1 x_4 x_6, x_1 x_4 x_5, x_1 x_4^2, x_1 x_3 x_6, x_1 x_3 x_5, x_1 x_3 x_4, x_1 x_3^2, x_1 x_2 x_6, x_1 x_2 x_5, x_1 x_2 x_4, \\ x_1 x_2 x_3, x_1 x_2^2, x_1^2 x_6, x_1^2 x_5, x_1^2 x_4, x_1^2 x_3, x_1^2 x_2\}$$




FIGURE 1. 8 equivalence classes of all non-negative graphs with 6 edges.




FIGURE 2.  $Q_1$




FIGURE 3.  $Q_2$




FIGURE 4.  $Q_3$




FIGURE 5.  $Q_4$




FIGURE 6.  $Q_5$

## REFERENCES

- [1] P. Alexandersson, B. Shapiro, Discriminants, symmetrized graph monomials, and sums of squares, arXiv:1104.0589v1 [math.AG]
- [2] A. Lax, P. Lax, On sums of squares, Linear Algebra and its Applications. 20 (1978) 71–75.

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, S-10691, STOCKHOLM, SWEDEN  
*E-mail address:* `per@math.su.se`