Extension of a radiative neutrino mass model based on a cosmological view point

Daijiro Suematsu¹

Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan

Abstract

We consider an extension of the radiative neutrino mass model at TeV regions so as to give the origin for inflation of the universe. This extension also gives a consistent explanation for both the origin of baryon number asymmetry and dark matter. A small scalar coupling which plays a crucial role in the neutrino mass generation in the original model may be related to parameters which are control inflation.

 $^{^{1}\}mathrm{e\text{-}mail:}$ suematsu@hep.s.kanazawa-u.ac.jp

1 Introduction

The radiative neutrino mass model proposed in [1] is a simple but interesting possibility which can take the place of the canonical seesaw model for neutrino masses. It could give the explanation for not only the neutrino mass problem but also the origin of dark matter (DM) [2, 3, 4].² A Z_2 symmetry to forbid the tree level neutrino mass also guarantees the stability of the lightest Z_2 odd field. As a result, the reason of the smallness of neutrino masses and the existence of DM are closely related in this model. Although the model has these noticeable features, baryon number asymmetry in the universe seems not to be easily explained in this framework unfortunately. The ordinary thermal leptogenesis [7, 8] seems not to work for the generation of sufficient lepton number asymmetry in a consistent way with the relic abundance of DM without some modification [9].

In this model the lightest right-handed neutrino is a most promising DM candidate. In that case both its relic abundance and small neutrino masses require O(1) neutrino Yukawa couplings in general [2, 3, 4]. This allows the model to cause the large CP asymmetry in the decay of the right-handed neutrinos even though their masses are of O(1) TeV. However, the same neutrino Yukawa couplings could cause a large washout of the generated lepton number asymmetry through the lepton number violating scattering processes. As a result, the thermal leptogenesis is not easy to generate sufficient lepton number asymmetry in a consistent way with the neutrino oscillation data and the DM abundance at least in the simplest form of the model. Non-thermal leptogenesis might give a consistent scenario for the origin of the baryon number asymmetry in this model [10].

In this paper we propose a simple extension to solve this fault of the model based on a cosmological viewpoint. We extend the model to make it connect with inflation of the universe and also generate the lepton number asymmetry through the inflaton decay.³ Since the lepton number violating effect could be separated from the neutrino mass generation, both the reduction of DM relic abundance and the washout of lepton number asymmetry are reconciled for the same neutrino Yukawa couplings. As a bonus, we have an explanation for the smallness of a scalar coupling which is crucial for this

²The supersymmetric extension of the model is proposed in [5, 6].

³The similar idea has been proposed in the inflation scenario in which the right-handed sneutrino plays a role of inflaton [11].

radiative neutrino mass generation.

2 Extension of the model

The model considered here is a simple model for the radiative neutrino mass generation [1]. This model is an extension of the standard model (SM) with three right-handed singlet fermion N_{R_i} and an inert doublet scalar η . These new fields are supposed to have odd parity of an assumed Z_2 symmetry, under which all SM contents have even parity. Invariant Yukawa couplings and scalar potential related to these new fields are summarized as

$$-\mathcal{L}_{y} = h_{ij}\bar{N}_{R_{i}}\eta^{\dagger}\ell_{L_{j}} + h_{ij}^{*}\bar{\ell}_{L_{i}}\eta N_{R_{j}} + \frac{1}{2}\left(M_{i}\bar{N}_{R_{i}}N_{R_{i}}^{c} + M_{i}\bar{N}_{R_{i}}^{c}N_{R_{i}}\right),$$

$$V = m_{\phi}^{2}\phi^{\dagger}\phi + m_{\eta}^{2}\eta^{\dagger}\eta + \lambda_{1}(\phi^{\dagger}\phi)^{2} + \lambda_{2}(\eta^{\dagger}\eta)^{2} + \lambda_{3}(\phi^{\dagger}\phi)(\eta^{\dagger}\eta) + \lambda_{4}(\eta^{\dagger}\phi)(\phi^{\dagger}\eta)$$

$$+ \left[\frac{\lambda_{5}}{2}(\phi^{\dagger}\eta)^{2} + \text{h.c.}\right],$$
(1)

where ℓ_{L_i} is a left-handed lepton doublet and ϕ is an ordinary Higgs doublet. Interaction terms are written by using the basis under which both matrices for Yukawa couplings of charged leptons and masses of the singlet neutrinos are real and diagonal. Since the new doublet scalar η is assumed to have no vacuum expectation value, the Z_2 symmetry forbids neutrino masses at tree level. We note that the lightest field with its odd parity is stable and then its thermal relic behaves as DM in the universe.

In this model we can define lepton number L in two different ways, since N_{R_i} and η are introduced as the fields which couple with ℓ_{L_i} . In fact, if either $M_i \bar{N}_{R_i} N_{R_i}^c$ or $\lambda_5 (\phi^{\dagger} \eta)^2$ decouples, the conserved lepton number can be defined for each case, respectively, such as

(i)
$$L(\eta) = 0$$
, $L(N_{R_i}) = 1$, (ii) $L(\eta) = 1$, $L(N_{R_i}) = 0$. (2)

The case (i) corresponds to the canonical seesaw model and the ordinary thermal leptogenesis is considered in this framework. The case (ii) is a new possibility allowed in this model. If we take account of sphaleron interaction, both the baryon number B and the lepton number L are violated but B-L is conserved as usual [12]. In each case, we find that B-L is related to B through the sphaleron interaction as

(i)
$$B = \frac{8}{23}(B - L)$$
, (ii) $B = \frac{7}{19}(B - L)$. (3)

These are derived by using chemical equilibrium conditions [13].

Now we consider the extension of the model at high energy regions by introducing canonically normalized real singlet scalars S_i to which the odd parity of Z_2 is assigned. Their potential and interaction terms are assumed to be given by

$$-\mathcal{L}_{S} = \sum_{i=1}^{K} \left(\frac{1}{2} m_{i}^{2} S_{i}^{2} + \mu_{i} S_{i} \phi^{\dagger} \eta + \text{h.c.} \right), \tag{4}$$

where μ_i is supposed to be complex.⁴ It is easy to find that the model defined by eq. (1) can be obtained as the effective theory with $\lambda_5 = \sum_i \frac{\mu_i^2}{m_i^2}$ after singlet scalars S_i are integrated out. Thus, if we adopt the definition (ii) for the lepton number in this extension, the origin of the lepton number violating λ_5 term in eq. (1) is found in \mathcal{L}_S . As long as this λ_5 term is out of thermal equilibrium at low energy regions, the B-L is found to be a good symmetry there. We easily find that the lepton number violation due to λ_5 is crucial for the neutrino mass generation at weak scale.

Neutrino masses are generated through the one-loop effect by picking up this lepton number violation associated with the electroweak symmetry breaking. They can be expressed as

$$\mathcal{M}_{ij}^{\nu} = \sum_{k=1}^{3} h_{ik} h_{jk} \left[\frac{\lambda_5 \langle \phi \rangle^2}{8\pi^2 M_k} \frac{M_k^2}{M_{\eta}^2 - M_k^2} \left(1 + \frac{M_k^2}{M_{\eta}^2 - M_k^2} \ln \frac{M_k^2}{M_{\eta}^2} \right) \right]$$

$$\simeq \left(\frac{\lambda_5}{10^{-10}} \right) \left(\frac{\bar{h}}{1.0} \right)^2 \left(\frac{1 \text{ TeV}}{M} \right) \times O\left(10^{-1} \right) \text{ eV}, \tag{5}$$

where $M_{\eta}^2 = m_{\phi}^2 + (\lambda_3 + \lambda_4) \langle \phi \rangle^2$. In the second line we put $h_{ik} \sim \bar{h}$ and $M_i, M_{\eta} \sim M$ for the rough estimation. If we remember that neutrino oscillation data requires that the neutrino masses are $O(10^{-1})$ eV or less,⁵ we find that λ_5 should be extremely small for the values of \bar{h} and M implicated in the second line of eq. (5). The explanation of the smallness of λ_5 seems to be important for this neutrino mass generation mechanism to be natural. Several solutions for this question have been proposed in [3, 6]. As we will see below, the present extension may give an alternative solution for this problem if $|\mu_i| \ll m_i$ is satisfied. The effective coupling constant λ_5 is found to be closely related

⁴The Z_2 symmetry can not forbid quartic terms of S_i . However, we only assume that there are no such terms here since the radiatively generated one can be neglected for sufficiently small μ_i .

⁵It should be noted that one of eigenvalues of this mass matrix is zero and also the cosmological upper bound for the neutrino masses is 0.58 eV [14].

to the cosmological issues such as the inflation and the baryon number asymmetry in the universe.

3 Cosmological features of the model

At first, we discuss the features of this extended model as an inflation model. If we assume that S_i takes a large initial value such as $O(M_{\rm pl})$ where $M_{\rm pl} = (8\pi G)^{-1/2}$ is the reduced Planck mass, the model behaves as the chaotic inflation model with multi-component inflaton. Various features for this kind of model have been discussed in a lot of articles [15]. We follow the results given in [16].

The number of e-folds brought by S_i is given as

$$N = M_{\rm pl}^{-2} \sum_{i=1}^{K} \int_{S_i^{\rm end}}^{S_i^*} \left(\frac{V_i}{V_i'}\right) dS_i \simeq \sum_{i=1}^{K} \left(\frac{S_i^*}{M_{\rm pl}}\right)^2, \tag{6}$$

where $V_i = \frac{1}{2}m_i^2 S_i^2$. S_i^{end} and S_i^* stand for a field value at the end of slow-roll inflation and a value at the time when cosmological scales $k = a_0 H_0$ leave the horizon, respectively. Since $S_i^* \gg S_i^{\text{end}}$ is satisfied, the number of e-folds N is determined only by S_i^* and it does not depend on the inflaton mass m_i . Using this e-folds N, the tilt of the scalar perturbation spectrum is given as

$$1 - n_s = \frac{1}{N} + \frac{\sum_{i=1}^K m_i^4 S_i^4}{\left(\sum_{i=1}^K m_i^2 S_i^2\right)^2}.$$
 (7)

If we note that $(\sum_i m_i^2 S_i^2)^2 > \sum_i m_i^4 S_i^4$ is satisfied for $K \geq 2$, n_s is found to become larger in the multi-component case compared with the single inflaton case. The ratio of tensor to scalar perturbation is given as

$$r = \frac{8}{N}. (8)$$

We note that r is independent of the number of inflaton components and their masses. If all the masses of S_i are equal, both n_s and r reduce to the ones of the single inflaton model [16]. Both n_s and r is now constrained by the 7-year WMAP data for the CMB observation as found in [14]. Although the m^2S^2 type chaotic inflation with a single inflaton S is marginally allowed for reasonable N such as N = 50 - 60, the situation becomes worse for the multi-component case since n_s becomes larger compared to the single inflaton case as discussed above (See Fig. 5 in the first paper of [14]). Taking account of this, we confine our following study to the two component case with $m_1 = m_2$ in eq. (4).

The decay of inflaton S_i occurs at $H \sim \Gamma_{S_i} < m_i$ through three scalars interaction $\mu_i S_i \phi^{\dagger} \eta$ in eq. (4) where H is the Hubble parameter and Γ_{S_i} stands for the decay width of S_i . Since this decay width is given by $\Gamma_{S_i} = \frac{1}{4\pi} \frac{|\mu_i|^2}{m_i}$, the decay $S_i \to \eta \phi^{\dagger}$ yields the thermal plasma with reheating temperature [17]

$$T_R \simeq 0.5 g_*^{-1/2} \left(\frac{\mu_1}{m_1}\right) (m_1 M_{\rm pl})^{1/2},$$
 (9)

where we suppose $|\mu_1| \gg |\mu_2|$. In the present model we have $g_* = 116$ as the relativistic degrees of freedom. If we note that this decay violates the lepton number defined as (ii) in eq. (2), we find that the lepton number asymmetry can be induced in this process through the cross term between tree and one-loop diagrams as long as μ_i is complex. The CP asymmetry induced in this process can be estimated as

$$\varepsilon \equiv \sum_{i=1}^{2} \frac{\Gamma(S_i \to \eta \phi^{\dagger}) - \Gamma(S_i \to \eta^{\dagger} \phi)}{\Gamma(S_i \to \eta \phi^{\dagger}) + \Gamma(S_i \to \eta^{\dagger} \phi)} = \sum_{i,j} \frac{1}{8\pi} \frac{\operatorname{Im}(\mu_i^* \mu_j)^2}{m_i^2 |\mu_i|^2} \ln \frac{m_i^2 + m_j^2}{m_j^2}$$

$$\simeq \frac{|\lambda_5| \ln 2}{8\pi} \sin 2(\theta_1 - \theta_2), \tag{10}$$

where $\theta_i = \arg(\mu_i)$ and we use $|\lambda_5| \simeq \frac{|\mu_1|^2}{m_1^2}$.

The lepton number asymmetry generated in the decay product η is kept in thermal plasma if the effective λ_5 term in eq. (1) is out of thermal equilibrium until the sphaleron decoupling temperature. This condition is written as

$$|\lambda_5| < 2 \times 10^{-7} g_*^{1/4} \left(\frac{T}{1 \text{ TeV}}\right).$$
 (11)

Thus, when $|\lambda_5|$ is in this range, the lepton number asymmetry can be estimated as the one induced through the decay of S_1 to $\eta\phi^{\dagger}$. It may be expressed as

$$Y_L = \varepsilon Y_\eta(T_R) \simeq \varepsilon \frac{m_1 S_1^{\text{end}}}{s(T_R)} \simeq 10\varepsilon \frac{T_R}{m_1},$$
 (12)

where Y_L and Y_{η} stand for the lepton number asymmetry and the η number in the comoving volume. They are expressed as $Y_{L,\eta} = \frac{n_{L,\eta}}{s}$ by using the entropy density $s = \frac{2\pi^2}{45}g_*T^3$. Since the washout of this generated lepton number asymmetry is caused by the

processes whose amplitudes are proportional to the tiny coupling λ_5 , it could be safely neglected.

Combining eqs. (3), (9), (10) and (12), we obtain the baryon number asymmetry Y_B as

$$|Y_B| \simeq 5 \times 10^{-11} \left(\frac{\lambda_5}{10^{-10}}\right)^2 \left(\frac{10^5 \text{ GeV}}{T_R}\right) \sin 2(\theta_1 - \theta_2).$$
 (13)

Since the observed baryon number asymmetry in the universe is $Y_B = (0.7 - 0.9) \times 10^{-10}$ [18], it can be generated for the suitable values of λ_5 and T_R . For example, if we impose $\lambda_5 = 10^{-10}$ and $T_R = 10^5$ GeV so as to generate suitable baryon number for the maximum value of $\sin 2(\theta_1 - \theta_2)$, we have

$$m_1 = 2 \times 10^4 \text{ GeV}, \qquad |\mu_1| = 0.2 \text{ GeV}.$$
 (14)

These results suggests that in this model the inflation should occur at a scale which is not far from the weak scale. In this extension the smallness of λ_5 which is crucial for the small neutrino mass generation is explained as the nature of the inflaton sector.

The decay of η also produces N_{R_i} as the component of thermal plasma. Since the lightest singlet fermion N_{R_1} is stable due to Z_2 symmetry, it behaves as DM. Thus, its relic abundance should satisfy $\Omega_{N_{R_1}}h^2=0.11$ which is obtained from observations of the WMAP [14]. The relic abundance is fixed through the $N_{R_1}N_{R_1}$ annihilation caused by the η exchange. This cross section can be expressed as [2, 4]

$$\langle \sigma v^2 \rangle \simeq \frac{1}{12\pi} \frac{M_1^2 (M_1^4 + M_\eta^4)}{(M_1^2 + M_\eta^2)^4} \sum_{i=e,\mu,\tau} |h_{i1}|^4 \frac{6T}{M_1}.$$
 (15)

By using this cross section, the relic abundance can be approximately estimated as [19]

$$\Omega_{N_{R_1}} h^2 = \frac{2.14 \times 10^9 z_f}{g_*^{1/2} m_{\rm pl} (\text{GeV}) \langle \sigma v \rangle}, \quad z_f = \ln \frac{0.038 g m_{\rm pl} M_1 \langle \sigma v \rangle}{g_*^{1/2} z_f^{1/2}}, \tag{16}$$

where z_f is defined as $z_f = M_1/T_f$ by using the freeze-out temperature T_f of N_{R_1} . If this $\Omega_{N_{R_1}}h^2$ has the required value for the λ_5 estimated above, the model could give a consistent explanation for both the baryon number asymmetry and the DM abundance.

In order to proceed this quantitative analysis, we need to introduce an additional assumption for the neutrino Yukawa couplings. Here we assume the following flavor structure for them [4]:

$$h_{ei} = 0, \quad h_{\mu i} = h_{\tau i} = h_i \quad (i = 1, 2); \qquad h_{e3} = h_{\mu 3} = -h_{\tau 3} = h_3.$$
 (17)




Fig. 1 The left frame shows the N_{R_1} relic abundance as a function of M_1 . The right frame shows the neutrino Yukawa coupling $|h_1|$ as a function of M_1 . Each line is plotted for $M_{\eta} = 2$, 2.5 and 3 TeV.

If we impose this structure, tri-bimaximal mixing is automatically induced⁶ and then the neutrino oscillation data can be explained as long as the following conditions are satisfied

$$|h_1|^2 \Lambda_1 + |h_2|^2 \Lambda_2 \simeq \frac{\sqrt{\Delta m_{\text{atm}}^2}}{2}, \qquad |h_3|^2 \Lambda_3 \simeq \frac{\sqrt{\Delta m_{\text{sol}}^2}}{3},$$
 (18)

where $\Delta m_{\rm atm}^2$ and $\Delta m_{\rm sol}^2$ represent the squared neutrino mass difference required by the atmospheric neutrinos and solar neutrinos [20]. Λ_i is defined as

$$\Lambda_i = \frac{\lambda_5 \langle \phi \rangle^2}{8\pi^2 M_k} \frac{M_k^2}{M_\eta^2 - M_k^2} \left(1 + \frac{M_k^2}{M_\eta^2 - M_k^2} \ln \frac{M_k^2}{M_\eta^2} \right). \tag{19}$$

If we use λ_5 derived from m_1 and μ_1 given in eq. (14) and impose the neutrino mass constraints (18), we can estimate the DM abundance. Here we proceed this analysis only for typical parameters such as

$$|\lambda_5| = 2.4 \times 10^{-10}, \quad M_2 = 4 \text{ TeV}, \quad M_3 = 6 \text{ TeV}.$$
 (20)

In the left frame of Fig. 1, we plot the N_{R_1} relic abundance $\Omega_{N_{R_1}}h^2$ as a function of M_1 for some values of M_η . In this calculation we assume $h_2 = h_1$, for simplicity. If a value of λ_5 becomes smaller, the neutrino Yukawa coupling $|h_1|$ becomes larger as expected from eq. (5) and then $\Omega_{N_{R_1}}h^2$ takes a smaller value. We can check that λ_5 should satisfy $|\lambda_5| \lesssim 2.6 \times 10^{-10}$ to reduce $\Omega_{N_{R_1}}h^2$ to the observed value for the present

⁶Recent T2K and Double Chooz results suggest nonzero $\sin \theta_{13}$ [21]. If we introduce a nonzero value for h_{ei} as a small perturbation to this flavor structure, nonzero $\sin \theta_{13}$ could be obtained [9]. In the present study we ignore this effect.

$M_{\eta} \; ({\rm TeV})$	$M_1 \text{ (TeV)}$	$ h_1 $	$Br(\mu \to e\gamma)$	$Br(\tau \to \mu \gamma)$
2	0.38	2.02	0.36×10^{-12}	9.5×10^{-8}
2	1.55	1.47	1.5×10^{-12}	0.34×10^{-8}
2.5	0.49	2.25	0.42×10^{-12}	10×10^{-8}
2.5	1.9	1.65	0.42×10^{-12}	0.46×10^{-8}
3	0.59	2.47	0.46×10^{-12}	11×10^{-8}
3	2.3	1.82	0.46×10^{-12}	0.55×10^{-8}

Table 1 Predicted values of the neutrino Yukawa coupling $|h_1|$ and the branching ratio of the lepton flavor violating processes by imposing the neutrino oscillation data and the DM relic abundance on the model.

parameter setting. As long as this bound is satisfied, the required N_{R_1} relic abundance can be obtained consistently with the neutrino mass constraints. In right frame of Fig. 1, we plot the neutrino Yukawa coupling $|h_1|$ as a function of M_1 by applying the condition in eq. (18) for the same parameters used in the left frame of Fig. 1.

Combining these results, we can find the values of Yukawa couplings h_1 and h_3 which give the required value of $\Omega_{N_{R_1}}h^2$. These values are shown in Table 1. Since the neutrino Yukawa couplings are large, the model could be severely constrained by the lepton flavor violation processes such as $\tau \to \mu \gamma$ and $\mu \to e \gamma$ [2]. The branching ratio of these processes are given by

$$Br(\tau \to \mu \gamma) \simeq \frac{0.51\alpha}{64\pi (G_F M_\eta^2)^2} \left[|h_1|^2 \left\{ F_2 \left(\frac{M_1^2}{M_\eta^2} \right) + F_2 \left(\frac{M_2^2}{M_\eta^2} \right) \right\} - |h_3|^2 F_2 \left(\frac{M_3^2}{M_\eta^2} \right) \right]^2,$$

$$Br(\mu \to e\gamma) \simeq \frac{3\alpha}{64\pi (G_F M_\eta^2)^2} \left[|h_3|^2 F_2 \left(\frac{M_3^2}{M_\eta^2} \right) \right]^2,$$
(21)

where $F_2(r)$ is defined as

$$F_2(r) = \frac{1 - 6r + 3r^2 + 2r^3 - 6r^2 \ln r}{6(1 - r)^4}.$$
 (22)

The predicted values of these branching ratio for the obtained parameters are also given in Table 1. The present upper bounds for $Br(\mu \to e\gamma)$ and $Br(\tau \to \mu\gamma)$ are 2.4×10^{-12} and 4.4×10^{-8} , respectively [22]. One may consider that the former gives much severe constraint on the model. However, the situation is different here. Although the neutrino Yukawa coupling $|h_1|$ have to take large values to reduce the N_{R_1} relic abundance, the

assumed flavor structure (17) can successfully suppresses $\mu \to e\gamma$ [3]. As found from eq. (21), this process depends only on $|h_3|$ which can be much smaller than $|h_1|$. On the other hand, since $\tau \to \mu\gamma$ depends on $|h_1|$, it could impose severer constraint on this model. Anyway, we can find consistent parameters for which these bounds are satisfied successfully as shown in Table 1.

Finally, we should remark on other phenomenological problems which could appear in relation to the present neutrino mass generation. First, we note that there is one-loop contribution to the muon g-2 through the interactions given in eq. (1) [4]. By using eq. (22), it can be written as [23]

$$\delta a_{\mu} = \sum_{k=1}^{3} \frac{|h_{\mu k}|^{2}}{(4\pi)^{2}} \frac{m_{\mu}^{2}}{M_{\eta}^{2}} F_{2} \left(\frac{M_{k}^{2}}{M_{\eta}^{2}}\right) \simeq 7.1 \times 10^{-11} \left(\frac{2 \text{ TeV}}{M_{\eta}}\right)^{2} \left(\frac{|h_{1}|}{2}\right)^{2} \mathcal{O},$$

$$\mathcal{O} = F_{2} \left(\frac{M_{1}^{2}}{M_{\eta}^{2}}\right) + F_{2} \left(\frac{M_{2}^{2}}{M_{\eta}^{2}}\right) + \frac{|h_{3}|^{2}}{|h_{1}|^{2}} F_{2} \left(\frac{M_{3}^{2}}{M_{\eta}^{2}}\right),$$
(23)

where we use eq. (17) in this derivation. Since \mathcal{O} is less than 0.5 here, the predicted value of δa_{μ} is found to be two orders of magnitude smaller than the present experimental value $\delta a_{\mu} = (30.2 \pm 8.7) \times 10^{-10}$ [24]. We need other contributions to saturate this discrepancy if we take it seriously.

Second, the model could have a problem relevant to the CP phases, which eventually appears when we consider the leptogenesis. As in case of the muon g-2, the interactions introduced to generate the neutrino masses could also contribute to the electric dipole moment of an electron (EDME) through loop diagrams if they violate the CP invariance. As found from the neutrino mass formula (5), the phase of μ_1 contributes to the MNS matrix U as the overall Majorana phase. We note that in the EDME loop diagrams the MNS matrix elements appear at lepton verteces with a W^{\pm} line and also an η_0 line in pairs as $U_{ij}U_{i'j'}^*$. This means that the CP phase of μ_1 which is relevant to the leptogenesis does not contribute to the EDME. Although the EDME could be induced by the diagrams with more than two loops as a result of the CP phases of neutrino Yukawa couplings, it could satisfy the present experimental upper bound [25] easily by fixing the parameters irrelevant to the leptogenesis suitably. The constraint from the EDME does not contradict with the present leptogenesis scenario.

4 Summary

We have considered an extension of the radiative neutrino mass model as an inflation model. In this extension the original model is obtained as the effective theory by integrating out the inflaton field. The lepton number violating term which plays a crucial role in the radiative neutrino mass generation appears in relation to the inflaton sector. As a result, the neutrino masses are closely related to the reheating temperature and also the baryon number asymmetry in the universe through the parameter λ_5 . The model might also give a unified picture for the explanation of the origin of DM other than these. Since the resulting reheating temperature is low enough to escape the gravitino problem [26], the supersymmetric extension of the present model along the framework given in [5, 6] could be an interesting subject. It will be discussed elsewhere.

Acknowledgement

This work is partially supported by a Grant-in-Aid for Scientific Research (C) from Japan Society for Promotion of Science (No.21540262) and also a Grant-in-Aid for Scientific Research on Priority Areas from The Ministry of Education, Culture, Sports, Science and Technology (No.22011003).

References

- [1] E. Ma, Phys. Lett. **B625** (2005) 76.
- [2] E. Ma, Phys. Rev. D73 (2006) 077301; J. Kubo, E. Ma and D. Suematsu, Phys. Lett. B642 (2006) 18; D. Aristizabal Sierra, J. Kubo, D. Suematsu, D. Restrepo and O. Zapata, Phys. Rev. D79 (2009) 013011.
- [3] J. Kubo and D. Suematsu, Phys. Lett. **B643** (2006) 336.
- [4] D. Suematsu, T. Toma and T. Yoshida, Phys. Rev. **D79** (2009) 093004; D. Suematsu,
 T. Toma and T. Yoshida, Phys. Rev. **D82** (2010) 013012.
- [5] E. Ma, Annals Fond.Broglie 31 (2006) 285; H. Fukuoka, J. Kubo and D. Suematsu, Phys. Lett. B678 (2009) 401; H. Fukuoka, D. Suematsu and T. Toma, JCAP 07 (2011) 001.
- [6] D. Suematsu and T. Toma, Nucl. Phys. **B847** (2011) 567.
- [7] M. Fukugita and T. Yanagida, Phys. Lett. **B174** (1986) 45.
- [8] M. Plümacher, Nucl. Phys. B530 (1998) 207; W. Buchmüller and M. Plümacher, Int. J. Mod. Phys. A15 (2000) 5047; W. Buchmüller, P. Di Bari, and M. Plümacher, Phys. Lett. B547 (2002) 128; Nucl. Phys. B643 (2002) 367; Nucl. Phys. B665 (2003) 445; G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Struma, Nucl. Phys. B685 (2004) 89; W. Buchmüller, R. D. Peccei and T. Yanagida, Ann. Rev. Nucl. Part. Sci. 55 (2005) 311.
- [9] D. Suematsu, arXiv:1103.0857[hep-ph].
- [10] D. Suematsu, Eur. Phys. J.C56 (2008) 379; H. Higashi, T. Ishima and D. Suematsu, Int. J. Mod. Phys. A26 (2001) 995.
- [11] H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, Phys. Rev. Lett. 70 (1993)1912; D. Suematsu and Y. Yamagishi, Mod. Phys. Lett. 38 (1995) 2923.
- [12] V. A. Kuzumin, V. A. Rubakov, M. E. Shaposhnikov, Phys. Lett. **B155** (1985) 36.

- [13] S. Yu Khlebnikov and M. E. Shaposhnikov, Nucl. Phys. B308 (1988) 885; J. S. Harvey and M. S. Turner, Phys. Rev. D42 (1990) 3344.
- [14] E. Komatsu, et al. Astrophys. J. Suppl. 180 (2009) 330; E. Komatsu, et al. Astrophys.
 J. Suppl. 192 (2011) 18.
- [15] For a review, see D. H. Lyth and A. Riotto, Phys. Rept. 314 (1999) 1; A. R. Liddle and D. H. Lyth, Cosmological Inflation and Large Scale Structure, (CUP, Cambridge, 2000).
- [16] L. Alabidi and D. H. Lyth, JCAP **05** (2006) 016.
- [17] E. W. Kolb and M. S. Turner, The Early Universe (Addison-Wesley, Redwood City, CA, 1990).
- [18] A. Riotto and M. Trodden, Ann. Rev. Nucl. Part. Sci. 49 (1999) 35; W. Bemrenther, Lect. Notes Phys. 591 (2002) 237; M. Dine and A. Kusenko, Rev. Mod. Phys. 76 (2004) 1.
- [19] K. Griest, M. Kamionkowski and M. S. Turner, Phys. Rev. D 41 (1990) 3565.
- [20] SNO Collaboration, Q. R. Ahmad, et al., Phys. Rev. Lett. 89 (2002) 011301; Super-Kamiokande Collaboration, Y. Fukuda, et al., Phys. Rev. Lett. 81 (1998) 1562; KamLAND Collaboration, K. Eguchi, et al., Phys. Rev. Lett. 90 (2003) 021802; K2K Collaboration, M. H. Ahn, et al., Phys. Rev. Lett. 90 (2003) 041801.
- [21] T2K Collaboration, K. Abe, et al., arXiv:1106.2822; H. D. Kerret, talk given at LowNu11.
- [22] MEG Collaboration, J. Adam, et al., Phys. Rev. Lett. 107 (2011) 171801; Belle Collaboration, K. Hayasaka, et al., Phys. Lett. B666 (2008) 16.
- [23] E. Ma and M. Raidal, Phys. Rev. Lett. 87 (2001) 011802.
- [24] K. Hagiwara, A. D. Martin, D. Nomura and T. Teubner, Phys. Lett. B649 (2007) 173.
- [25] B. C. Regan, E. D. Commins, C. J. Schmidt and D. DeMille, Phys. Rev. Lett. 88 (2002) 071805.

[26] M. Yu. Khlopov and A. D. Linde, Phys. Lett. B138 (1984) 265; J. Ellis, J. E. Kim and D.V. Nanopoulos, Phys. Lett. B145 (1984) 181; J. Ellis, K. A. Olive and S. J. Rey, Astropart. Phys. 4 (1996) 371; M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. D71 (2005) 083502; for a review, see S. Sarkar, Rep. prog. Phys. 59 (1996) 1493; T. Moroi, AIP conf. Proc. 805 (2006) 37.