

Arithmetic area for m planar Brownian paths

Jean Desbois¹ and Stéphane Ouvry²

Université Paris-Sud, Laboratoire de Physique Théorique et Modèles Statistiques³
91405 Orsay, France

Abstract

We pursue the analysis made in [1] on the arithmetic area enclosed by m closed Brownian paths. We pay a particular attention to the random variable $S_{n_1, n_2, \dots, n_m}(m)$ which is the arithmetic area of the set of points, also called winding sectors, enclosed n_1 times by path 1, n_2 times by path 2, \dots , n_m times by path m . Various results are obtained in the asymptotic limit $m \rightarrow \infty$. A key observation is that, since the paths are independent, one can use in the m paths case the SLE information, valid in the 1-path case, on the 0-winding sectors arithmetic area.

1 Introduction

In [1], the asymptotic behavior of the average arithmetic area enclosed by the external frontier of m independent closed Brownian planar paths, of same length t and starting from and ending at the same point, has been obtained using a path integral approach [2, 3].

In the one path case, the random variable of interest happens to be the arithmetic area S_n of the n -winding sectors enclosed by the path, from which the total arithmetic area $S = \sum_n S_n$ can be computed. A n -winding sector is by definition a set of points enclosed n times by the path. Path integral technics [3] give $\langle S_n \rangle = t/(2\pi n^2)$ but end up being somehow less adapted for $\langle S_0 \rangle$, the average arithmetic area of the 0-winding sectors inside the path, i.e. of the set of points enclosed an equal number of times clockwise and anti-clockwise by the path. Indeed path integral cannot distinguish 0-winding sectors inside the path from the outside of the path, which is also 0-winding. Other technics have to be used, in the case at hand SLE technics [4], to get $\langle S \rangle = \pi t/5$ from which $\langle S_0 \rangle = \pi t/30$ can be derived. It means that $\langle S_0 \rangle = q \langle S - S_0 \rangle$ with $q = 1/5$.

For m independant paths, the same path integral technics used in the one path case have now to focus [1] on the random variable $S_n(m)$, the arithmetic area of the n -winding sectors enclosed by the m paths, from which the total area $S(m) = \sum_n S_n(m)$ can in principle be computed. A n -winding sector is again defined as a set of points enclosed n times by the m paths as illustrated in Figure 1 in the $m = 2$ paths case. One has found that the leading asymptotic term scales like $\ln m$, namely that $\langle S(m) - S_0(m) \rangle \sim \frac{\pi t}{2} \ln m$ with, as already stressed, no information on $\langle S_0(m) \rangle$ inside the m paths.

In [5] on the other hand, the asymptotic behavior of the arithmetic area enclosed by the convex envelop of m closed paths was found to be $\langle S(m) \rangle_{\text{convex}} \sim \frac{\pi t}{2} \ln m$, i.e. the same scaling as $\langle S(m) - S_0(m) \rangle$. One then concluded [1] that, necessarily, $\langle S(m) \rangle \sim \frac{\pi t}{2} \ln m$ and that $\langle S_0(m) \rangle$ is subleading.

This particular scaling might be of interest for spatial ecology considerations where one asks about the increase in size of a natural reserve with the animal population (assuming of course that food is homogeneously and sufficiently available). By identifying an animal looking for food to a random walker one thereby obtains the $\log m$ scaling rather than the naive geometrical \sqrt{m} scaling.

¹jean.desbois@u-psud.fr

²stephane.ouvry@u-psud.fr

³Unité Mixte de Recherche CNRS-Paris Sud, UMR 8626




Figure 1: On a square lattice two independent closed walks 1 and 2 with 38 steps each, starting from and returning at the origin. The winding sectors of each walk are labelled by their winding numbers $\{n_1\}$ (for walk 1) and $\{n_2\}$ (for walk 2). The winding sectors enclosed by the external frontier of the two walks 1 + 2 are labelled i) by their joint winding numbers $\{n_1, n_2\}$, ii) by their total winding number $\{n_1 + n_2 \equiv n\}$. The $n = 0$ -winding sectors of interest, namely inside the external frontier of 1 + 2, correspond either to $\{0, 0\}$ with at least one of the 0-winding sectors inside one of the paths, or to $\{n_1, n_2\}$ with $n_1 + n_2 = 0$ and both n_1 and $n_2 \neq 0$.

In the present work we will give a more detailed analysis of the random variable $S_n(m)$. In particular we will pay attention to the random variable $S_{n_1, n_2, \dots, n_m}(m)$, the arithmetic area of the set of points enclosed n_1 times by path 1, n_2 times by path 2, \dots , n_m times by path m , with $\sum_i n_i = n$. Happily enough, this variable can be tackled by path integral technics analogous to the one used in [1, 3] provided that at least one of the $n_i \neq 0$. Note, when $\sum_i n_i = 0$, that $S_{n_1, n_2, \dots, n_m}(m)$ contains a part of the inside 0-winding sector area $S_0(m)$. To get a hand on the other part $S_{0, 0, \dots, 0}(m)$, with at least one of the $n_i = 0$ -winding sectors inside the corresponding path i (otherwise one would be trivially outside the external frontier of the m paths), one key observation is that since the paths are independent one can use in the m paths case the SLE information $\langle S_0 \rangle = q \langle S - S_0 \rangle$ valid in the one path case.

From these considerations will follow $\langle S(m) \rangle$ and $\langle S_0(m) \rangle$. We will show in particular that, when $m \rightarrow \infty$, the subleading $\langle S_0(m) \rangle$ remains finite. Also some information on the asymptotics of $\langle S_{n_1, n_2, 0, \dots, 0}(m) \rangle$ when both n_1 and $n_2 \rightarrow \infty$ will be obtained. Finally $\langle S_0(2) \rangle$ and the overlap between 2 paths, $\langle 2S(1) - S(2) \rangle$, will be considered. This quantity might have some interest in polymer physics where polymers are modelised by Brownian paths.

In view of unifying notations between [1] and [3], we always denote the n -winding sectors arithmetic area and total arithmetic area for m paths by $S_n(m)$ and $S(m)$, which means that, from now on, $S_n(1)$ and $S(1)$ stand for S_n and S , the n -winding sectors arithmetic area and total arithmetic area in the one path case.

2 Winding sectors

2.1 Winding angle and propagator

As stated in the Introduction, the arithmetic area of the winding sectors enclosed by planar Brownian paths can be obtained from their winding properties. Consider a path of length t , starting from and ending at \vec{r} and the angle θ wound by the path around the origin O . The average of the random variable $e^{i\alpha\theta}$ over the set of such paths is

$$\langle e^{i\alpha\theta} \rangle = \frac{\mathcal{G}_\alpha(\vec{r}, \vec{r})}{\mathcal{G}_0(\vec{r}, \vec{r})} \quad (1)$$

where

$$\mathcal{G}_\alpha(\vec{r}, \vec{r}) = \int_{\vec{r}(0)=\vec{r}}^{\vec{r}(t)=\vec{r}} \mathcal{D}\vec{r}(\tau) e^{-\frac{1}{2} \int_0^t \dot{\vec{r}}^2(\tau) d\tau + i\alpha \int_0^t \dot{\theta}(\tau) d\tau} \quad (2)$$

is the quantum propagator of a charged particle coupled to a vortex at location O . By symmetry, it depends only on r

$$\mathcal{G}_\alpha(\vec{r}, \vec{r}) = \frac{1}{2\pi t} e^{-\frac{r^2}{t}} \sum_{k=-\infty}^{+\infty} I_{|k-\alpha|} \left(\frac{r^2}{t} \right) \quad (3)$$

where $I_{|k-\alpha|}$ is a modified Bessel function and

$$\mathcal{G}_0(\vec{r}, \vec{r}) = \frac{1}{2\pi t} \quad (4)$$

Additional symmetry and periodicity considerations lead to $\mathcal{G}_\alpha = \mathcal{G}_{\alpha+1} = \mathcal{G}_{1-\alpha}$ so that α can be restricted to $0 \leq \alpha \leq 1$. We also set $\frac{r^2}{t} \equiv x$ so that

$$\langle e^{i\alpha\theta} \rangle = e^{-x} \sum_{k=-\infty}^{+\infty} I_{|k-\alpha|}(x) \equiv G_\alpha(x)$$

with $G_0(x) = 1$. Clearly areas are proportional to t , so we can, without loss of generality, set $t = 1$.

Let us rewrite $G_\alpha(x)$ in a more suitable form : one has

$$G_\alpha(x) = e^{-x} \sum_{k=0}^{+\infty} I_{k+\alpha}(x) + \{\alpha \rightarrow 1 - \alpha\} \quad (5)$$

Observing that $\frac{d}{dx} \left(e^{-x} \sum_{k=0}^{+\infty} I_{k+\alpha}(x) \right) = \frac{1}{2} e^{-x} (I_{\alpha-1}(x) - I_\alpha(x))$, we get

$$\frac{dG_\alpha(x)}{dx} = \frac{1}{2} e^{-x} (I_{-\alpha}(x) - I_\alpha(x) + I_{\alpha-1}(x) - I_{1-\alpha}(x)) = \frac{\sin(\alpha\pi)}{\pi} e^{-x} (K_\alpha(x) + K_{1-\alpha}(x)) \quad (6)$$

Using the integral representation [6] of $K_\alpha(x)$

$$K_\alpha(x) = \frac{1}{2} \int_{-\infty}^{\infty} du e^{-x \cosh u} \cosh(\alpha u) \quad (7)$$

we deduce

$$G_\alpha(x) = \frac{\sin(\alpha\pi)}{\pi} \int_{-\infty}^{\infty} du \frac{1 - e^{-x(1+\cosh u)}}{1 + \cosh u} \cosh \frac{u}{2} \cosh\left(\left(\alpha - \frac{1}{2}\right)u\right) \quad (8)$$

Clearly

$$\lim_{x \rightarrow \infty} G_\alpha(x) = \frac{\sin(\alpha\pi)}{\pi} \int_{-\infty}^{\infty} du \frac{1}{1 + \cosh u} \cosh \frac{u}{2} \cosh\left(\left(\alpha - \frac{1}{2}\right)u\right) = 1 \quad (9)$$

so that

$$1 - G_\alpha(x) = \frac{\sin(\alpha\pi)}{\pi} \int_{-\infty}^{\infty} du \frac{e^{-x(1+\cosh u)}}{1 + \cosh u} \cosh \frac{u}{2} \cosh\left(\left(\alpha - \frac{1}{2}\right)u\right) \quad (10)$$

Eq. (10) will be extensively used in the following.

2.2 Average area $\langle S_n(m) \rangle$ of n -winding sectors

Let us first consider the average arithmetic area $\langle S_n(m) \rangle$ of the n -winding sectors labelled by their winding number n . For m brownian paths of same length unity, starting from and ending at the same point, one has [1]

$$Z_\alpha(m) \equiv \pi \int_0^\infty dx (1 - (G_\alpha(x))^m) = \sum_{n \neq 0} \langle S_n(m) \rangle (1 - e^{i2\pi\alpha n}) \quad (11)$$

Rewriting $G_\alpha(x) = 1 - (1 - G_\alpha(x))$ leads, for $n \neq 0$, to

$$\langle S_n(m) \rangle = - \int_0^1 d\alpha Z_\alpha(m) \cos(2\pi\alpha n) = \sum_{j=1}^m (-1)^{j+1} \binom{m}{j} \left(-\pi \int_0^1 d\alpha \int_0^\infty dx (1 - G_\alpha(x))^j \cos(2\pi\alpha n) \right) \quad (12)$$

where $\binom{m}{j}$ is the binomial coefficient. Thus

$$\langle S_n(m) \rangle = \sum_{j=1}^m (-1)^{j+1} \binom{m}{j} I_{n,j} \quad (13)$$

with

$$I_{n,j} = -\pi \int_0^1 d\alpha \int_0^\infty dx (1 - G_\alpha(x))^j \cos(2\pi\alpha n) \quad (14)$$

Using (10) one rewrites $I_{n,j}$ as

$$I_{n,j} = -\pi \left(\prod_{i=1}^j \int_{-\infty}^{\infty} \frac{du_i}{2\pi \cosh \frac{u_i}{2}} \right) \frac{\Phi_{n,j}}{j + \sum_{i=1}^j \cosh u_i} \quad (15)$$

where the $\Phi_{n,j}$'s follow from the integration over α

- j odd = $2k + 1$

$$\Phi_{n,j} = \frac{(-1)^{k+1}}{2^{2k+1}} 2\pi \cosh \left(\frac{\sum_{i=1}^j u_i}{2} \right) \sum_{N=n-k}^{n+k+1} (-1)^{N+k-n} \binom{2k+1}{N+k-n} \frac{2N-1}{(\pi(2N-1))^2 + (\sum_{i=1}^j u_i)^2} \quad (16)$$

- j even = $2k$

$$\Phi_{n,j} = \frac{(-1)^k}{2^{2k}} 2 \left(\sum_{i=1}^j u_i \right) \sinh \left(\frac{\sum_{i=1}^j u_i}{2} \right) \sum_{N=n-k}^{n+k} (-1)^{N+k-n} \binom{2k}{N+k-n} \frac{1}{(2\pi N)^2 + (\sum_{i=1}^j u_i)^2} \quad (17)$$

It is possible to compute $I_{n,1}$ exactly [4]

$$I_{n,1} = \frac{\pi}{2} \int_{-\infty}^{\infty} \frac{du}{1 + \cosh u} \left(\frac{2N-1}{(\pi(2N-1))^2 + u^2} - \frac{2N+1}{(\pi(2N+1))^2 + u^2} \right) = \frac{1}{2\pi n^2} \quad (18)$$

On the other hand when $n \rightarrow \infty$

- j odd

$$I_{n,j} \sim \frac{(-1)^{\frac{j-1}{2}} j!}{(2\pi)^j n^{j+1}} c_j \quad (19)$$

$$c_j = \int_0^\infty dx e^{-jx} (K_0(x))^j \quad (20)$$

- j even

$$I_{n,j} \sim \frac{(-1)^{(\frac{j}{2}+1)} j(j+1)!}{(2\pi)^{j+1} n^{j+2}} d_j \quad (21)$$

$$d_j = \int_0^\infty dx e^{-jx} (K_0(x))^{j-1} \int_0^\infty du e^{-x \cosh u} u \tanh \frac{u}{2} \quad (22)$$

We finally obtain

$$\begin{aligned} \langle S_n(1) \rangle &= I_{n,1} = \frac{1}{2\pi n^2} \\ \langle S_n(2) \rangle &= \frac{2}{2\pi n^2} - I_{n,2} \underset{n \rightarrow \infty}{=} \frac{2}{2\pi n^2} - \frac{3}{2\pi^3 n^4} d_2 + O\left(\frac{1}{n^6}\right) \end{aligned}$$

and in the general case

$$\begin{aligned}\langle S_n(m) \rangle &= \frac{m}{2\pi n^2} - \binom{m}{2} I_{n,2} + \binom{m}{3} I_{n,3} - \dots \\ &= \lim_{n \rightarrow \infty} \frac{m}{2\pi n^2} - \frac{3}{4\pi^3 n^4} \left(2 \binom{m}{2} d_2 + \binom{m}{3} c_3 \right) + O\left(\frac{1}{n^6}\right)\end{aligned}$$

with the convention that $\binom{m}{j} = 0$ if $j > m$, $d_2 \simeq 2.84$ and $c_3 \simeq 5.73$. Clearly, and as discussed in the Introduction, we have no information so far on the 0-winding sector inside the m paths. To make some progress on this issue, we have to turn to $\langle S_{n_1, n_2, \dots, n_m}(m) \rangle$, the average arithmetic area of the sectors enclosed n_1 times by path 1, n_2 times by path 2, \dots , and n_m times by path m .

2.3 Average area $\langle S_{n_1, n_2, \dots, n_m}(m) \rangle$

Winding sectors can as well be labelled by the set $\{n_1, n_2, \dots, n_m\}$ of the individual winding numbers n_i enclosed by each path i . In line with section (2.2), one has

$$Z_{\alpha_1, \alpha_2, \dots, \alpha_m}(m) \equiv \pi \int_0^\infty dx \left(1 - \prod_{i=1}^m G_{\alpha_i}(x) \right) = \sum' \langle S_{n_1, n_2, \dots, n_m}(m) \rangle \left(1 - e^{2i\pi(\sum_{i=1}^m \alpha_i n_i)} \right) \quad (23)$$

where in \sum' the set $n_1 = n_2 = \dots = n_m = 0$ is excluded from the sum. So

$$\begin{aligned}\langle S_{n_1, n_2, \dots, n_m}(m) \rangle &= - \int_0^1 d\alpha_1 \dots \int_0^1 d\alpha_m Z_{\alpha_1, \alpha_2, \dots, \alpha_m}(m) \cos \left(2\pi \left(\sum_{i=1}^m \alpha_i n_i \right) \right) \\ &= - \int_0^1 d\alpha_1 \dots \int_0^1 d\alpha_m Z_{\alpha_1, \alpha_2, \dots, \alpha_m}(m) \prod_{i=1}^m \cos(2\pi \alpha_i n_i)\end{aligned} \quad (24)$$

where we have used $\int_0^1 d\alpha \sin(\pi\alpha) \sin(2\pi\alpha n) \cosh \left((\alpha - \frac{1}{2})u \right) = 0$. Now $\langle S_{n_1, n_2, \dots, n_m}(m) \rangle$ is invariant by permutation on the n_i 's, so one can focus without loss of generality on $\langle S_{n_1, \dots, n_j, 0, \dots, 0}(m) \rangle$, $1 \leq j \leq m$, with $n_1, \dots, n_j \neq 0$.

Rewriting $G_{\alpha_i}(x) = 1 - (1 - G_{\alpha_i}(x))$ in (23,24) leads to consider when $n_i \neq 0$

$$\int_0^1 d\alpha_i (1 - G_{\alpha_i}(x)) \cos(2\pi \alpha_i n_i) = \int_{-\infty}^\infty du_i e^{-x(1+\cosh u_i)} P(u_i, n_i) \quad (25)$$

with

$$P(u_i, n_i) = \frac{u_i^2 + (1 - 4n_i^2)\pi^2}{(u_i^2 + (\pi(2n_i + 1))^2)(u_i^2 + (\pi(2n_i - 1))^2)} \sim_{n_i \rightarrow \infty} -\frac{1}{4\pi^2 n_i^2} \quad (26)$$

It follows that

$$\langle S_{n_1, \dots, n_j, 0, \dots, 0}(m) \rangle = \pi(-1)^j \int_0^\infty dx \left(\prod_{i=1}^j \int_{-\infty}^\infty du_i e^{-x(1+\cosh u_i)} P(u_i, n_i) \right) (1 - f(x))^{m-j} \quad (27)$$

where

$$f(x) = \int_0^1 d\alpha (1 - G_\alpha(x)) = \int_{-\infty}^\infty du \frac{e^{-x(1+\cosh u)}}{u^2 + \pi^2} \quad (28)$$

For example when $m = 1, 2$ one gets

$$\begin{aligned}\langle S_{n_1}(1) \rangle &= -\pi \int_0^\infty dx \int_{-\infty}^\infty du e^{-x(1+\cosh u)} P(u, n_1) = \frac{1}{2\pi n_1^2} \\ \langle S_{n_1, n_2}(2) \rangle &= \pi \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{du_1 du_2}{2 + \cosh u_1 + \cosh u_2} P(u_1, n_1) P(u_2, n_2) \\ \langle S_{n_1, 0}(2) \rangle &= \frac{1}{2\pi n_1^2} + \pi \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{du_1 du_2}{2 + \cosh u_1 + \cosh u_2} P(u_1, n_1) P(u_2, 0)\end{aligned}$$

When $n_i \rightarrow \infty$ one obtains

$$\langle S_{n_1, \dots, n_j, 0, \dots, 0}(m) \rangle \sim \frac{1}{2^j \pi^{2j-1}} \frac{1}{n_1^2 \dots n_j^2} c_{j,m}$$

where the constant

$$c_{j,m} = \int_0^\infty dx e^{-jx} (K_0(x))^j (1 - f(x))^{m-j}$$

has to be evaluated numerically to the exception of

$$c_{2,2} = \frac{3}{2}\zeta(2) - 1 + \frac{3}{2}\zeta(2) \sum_{k=1}^\infty \prod_{i=1}^k \left(1 - \frac{1}{2i}\right)^2 - \sum_{k=1}^\infty \prod_{i=1}^k \left(1 - \frac{1}{2i+1}\right)^2$$

3 Arithmetic area enclosed by m Brownian paths

We are now in position to compute $\langle S(m) \rangle$, the average arithmetic area enclosed by the m paths. Obviously

$$\langle S(m) \rangle = \sum' \langle S_{n_1, \dots, n_m}(m) \rangle + \langle S_{0, \dots, 0}(m) \rangle \quad (29)$$

where $\langle S_{0, \dots, 0}(m) \rangle$ is the area of the finite $\{n_1 = 0, n_2 = 0, \dots, n_m = 0\}$ winding sectors inside, at least, one of the paths⁴. From (23), we get

$$\sum' \langle S_{n_1, \dots, n_m}(m) \rangle = \int_0^1 d\alpha_1 \dots \int_0^1 d\alpha_m Z_{\alpha_1, \dots, \alpha_m}(m) \quad (30)$$

so that, using (28),

$$\sum' \langle S_{n_1, \dots, n_m}(m) \rangle = \pi \int_0^\infty dx (1 - (1 - f(x))^m) \quad (31)$$

In order to find $\langle S_{0, \dots, 0}(m) \rangle$ first consider

$$\sum_{n_1 \neq 0} \langle S_{n_1, 0, \dots, 0}(m) \rangle = \pi \int_0^\infty dx f(x) (1 - f(x))^{m-1}$$

and, the sum

$$\begin{aligned}A_1 &\equiv \sum_{n_1 \neq 0} (\langle S_{n_1, 0, \dots, 0}(m) \rangle + \langle S_{0, n_1, 0, \dots, 0}(m) \rangle + \dots + \langle S_{0, \dots, 0, n_1}(m) \rangle) \\ &= \binom{m}{1} \pi \int_0^\infty dx f(x) (1 - f(x))^{m-1}\end{aligned}$$

⁴Remember that $\langle S_0(m) \rangle = \langle S_{0, \dots, 0}(m) \rangle + \sum' \langle S_{n_1, \dots, n_m}(m) \rangle$, with $\sum n_i = 0$ and at least one of the $n_i \neq 0$.

where one has taken into account permutation invariance. Similarly one can consider

$$\begin{aligned} A_2 &\equiv \sum_{n_1, n_2 \neq 0} (\langle S_{n_1, n_2, 0, \dots, 0}(m) \rangle + \dots + \langle S_{0, \dots, 0, n_1, 0, \dots, 0, n_2, 0, \dots, 0}(m) \rangle + \dots + \langle S_{0, \dots, 0, n_1, n_2}(m) \rangle) \\ &= \binom{m}{2} \pi \int_0^\infty dx f(x)^2 (1 - f(x))^{m-2} \end{aligned}$$

and in general

$$A_m \equiv \sum_{n_1, n_2, \dots, n_m \neq 0} \langle S_{n_1, n_2, \dots, n_m}(m) \rangle = \binom{m}{m} \pi \int_0^\infty dx f(x)^m (1 - f(x))^{m-m}$$

with, obviously,

$$\sum_{i=1}^m A_i = \pi \int_0^\infty dx (1 - (1 - f(x))^m) = \sum \langle S_{n_1, n_2, \dots, n_m}(m) \rangle \quad (32)$$

We are interested in $\langle S_{0, \dots, 0}(m) \rangle$: in the case of one closed path one knows from SLE [4] that

$$q = \frac{\langle S_0(1) \rangle}{\sum_{n \neq 0} \langle S_n(1) \rangle} = \frac{1}{5} \quad (33)$$

This means that for any point **inside** the path, with winding number n , q is the ratio of the probability to have a 0-winding to the probability to have a $n \neq 0$ -winding.

A_1 counts the points with only one non-zero winding number. It follows that the corresponding contribution to $\langle S_{0, \dots, 0}(m) \rangle$ is necessarily $q A_1$. Similarly, A_2 counts the points with only two non-zero winding numbers. Since the m paths are independent, it follows that the corresponding contribution to $\langle S_{0, \dots, 0}(m) \rangle$ is $q^2 A_2$. This line of reasoning generalizes to A_k : the contribution to $\langle S_{0, \dots, 0}(m) \rangle$ is $q^k A_k$. Finally

$$\langle S_{0, \dots, 0}(m) \rangle = \sum_{i=1}^m q^i A_i = \Phi_0(m) - \Phi_q(m) \quad (34)$$

with

$$\Phi_q(m) = \pi \int_0^\infty dx \left(1 - (1 - (1 - q)f(x))^m \right) \quad (35)$$

Clearly $\sum_{i=1}^m A_i$ in (32) coincides with $\Phi_0(m)$. It follows that the average arithmetic area enclosed by the m paths is

$$\langle S(m) \rangle = \sum_{i=1}^m (1 + q^i) A_i = 2\Phi_0(m) - \Phi_q(m) \quad (36)$$

When $m \rightarrow \infty$ (see eq.(46) in the appendix) one obtains

$$\langle S(m) \rangle = \frac{\pi}{2} \ln m - \frac{\pi}{4} \ln \ln m + \frac{\pi}{2} \left(\ln \sqrt{\frac{25}{4\pi^3}} + C \right) + o\left(\frac{1}{\sqrt{\ln m}}\right) \quad (37)$$

where C is the Euler constant. In Figure 2, numerical simulations for $\langle S(m) \rangle$ show that the agreement with eq.(37) is quite correct, even for not large values of m .




Figure 2: The average arithmetic area $\langle S(m) \rangle$: the crosses are numerical simulations (10000 events) of closed random walks (10^6 steps for each one) on the 2D square lattice; $m = 4, 8, 16, \dots, 1024$; the line is the analytical result eq.(37). The agreement is quite correct as soon as $m \geq 16$.

Moreover, the asymptotic [1] of $\langle S(m) - S_0(m) \rangle$ when $m \rightarrow \infty$ is known to be

$$\langle S(m) - S_0(m) \rangle = \frac{\pi}{2} \ln m - \frac{\pi}{4} \ln \ln m + \frac{\pi}{2} \left(-\ln \sqrt{4\pi} + C \right) + o\left(\frac{1}{\sqrt{\ln m}}\right) \quad (38)$$

Comparing Eqs.(37) and (38), one deduces that the subleading 0-winding sector average arithmetic area

$$\lim_{m \rightarrow \infty} \langle S_0(m) \rangle = \frac{\pi}{2} \ln \frac{5}{\pi} \quad (39)$$

remains finite in the $m \rightarrow \infty$ limit.

Finally, as another illustration of the path integral formalism, consider the average overlap $\langle 2S(1) - S(2) \rangle$ of the arithmetic areas of two paths and $\langle S_0(2) \rangle$, the average 0-winding sectors arithmetic area of two paths. One has

$$\begin{aligned} \langle S(1) \rangle &= \pi \int_0^\infty dx (1+q) f(x) = \frac{\pi}{5} \\ \langle S(2) \rangle &= \pi \int_0^\infty dx \left(2(1+q)f(x) + f(x)^2((1-q)^2 - 2) \right) \end{aligned}$$

so that

$$\begin{aligned} \langle 2S(1) - S(2) \rangle &= \pi(2 - (1-q)^2) \int_0^\infty dx f(x)^2 \\ &= \frac{34\pi}{25} \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{du_1 du_2}{(2 + \cosh u_1 + \cosh u_2)(\pi^2 + u_1^2)(\pi^2 + u_2^2)} \end{aligned} \quad (40)$$

Numerically

$$\frac{\langle 2S(1) - S(2) \rangle}{\langle S(1) \rangle} \approx 0.286$$

is close to what one would obtain if the paths attached in O were two circles of radius R : the overlap in unit of πR^2 would then be

$$\frac{1}{2} - \frac{2}{\pi^2} \approx 0.297$$

Also, as far as $\langle S_0(2) \rangle$ is concerned, rewrite

$$\langle S_0(2) \rangle = \langle S(2) \rangle - \sum_{n \neq 0} \langle S_n(2) \rangle = 2\langle S(1) \rangle - \langle 2S(1) - S(2) \rangle - \sum_{n \neq 0} \langle S_n(2) \rangle$$

From Section 2.2 one has

$$\sum_{n \neq 0} \langle S_n(2) \rangle = \frac{\pi}{3} - \pi \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} du_1 du_2 \frac{\tanh \frac{u_1}{2} + \tanh \frac{u_2}{2}}{(2 + \cosh u_1 + \cosh u_2)(u_1 + u_2)((2\pi)^2 + (u_1 + u_2)^2)}$$

so that, using $\langle S_0(1) \rangle = \frac{\pi}{30}$ and eq. (40), one finds $\langle 2S_0(1) - S_0(2) \rangle$ to be

$$\pi \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{du_1 du_2}{(2 + \cosh u_1 + \cosh u_2)} \left(\frac{\frac{34}{25}}{(\pi^2 + u_1^2)(\pi^2 + u_2^2)} - \frac{\tanh \frac{u_1}{2} + \tanh \frac{u_2}{2}}{(u_1 + u_2)((2\pi)^2 + (u_1 + u_2)^2)} \right) \quad (41)$$

If the paths were not overlapping, then one would necessarily have $\langle 2S_0(1) - S_0(2) \rangle = 0$. The non vanishing result (41) clearly indicates that the two paths do, on average, overlap as already seen in (40). Note that the 0-winding sectors "overlap" $\langle 2S_0(1) - S_0(2) \rangle$ is different in nature from the arithmetic area overlap $\langle 2S(1) - S(2) \rangle$: the latter is a purely geometric overlap, whereas the former is more subtle since the superposition of the two paths destroy some original 0-winding sectors in each of the paths and create new 0-windings sectors for the two paths.

4 Conclusion

Path integral technics have been extensively used to tackle the issue of n -winding sectors arithmetic area of m Brownian paths. Some information stemming from SLE technics valid only in the one path case have also proved useful in the m paths case, merely because the paths are independent.

Eqs.(37) and (39) are the main results of this paper. In particular the subleading 0-winding arithmetic area has been shown to remain finite in the asymptotic limit. Numerical simulations have nicely confirmed these asymptotics results. The overlap between two paths is also computed numerically. Applications to polymer physics will be studied in a forthcoming publication.

5 Appendix

In the appendix we derive the $m \rightarrow \infty$ asymptotic limit of $\Phi_q(m)$ defined in (35)

$$\Phi_q(m) = \pi \int_0^{\infty} dx (1 - (1 - (1 - q)f(x))^m)$$

with $f(x)$ given in (28)

$$f(x) = \int_{-\infty}^{\infty} du \frac{e^{-x(1+\cosh u)}}{\pi^2 + u^2}$$

Setting $x = \frac{y \ln m}{2}$, $\Phi_q(m)$ becomes

$$\begin{aligned}\Phi_q(m) &= \frac{\pi \ln m}{2} \int_0^\infty dy \left(1 - (1 - (1 - q)f(\frac{y \ln m}{2}))^m\right) \\ &\approx \frac{\pi \ln m}{2} \int_0^\infty dy \left(1 - e^{-m(1-q)f(\frac{y \ln m}{2})}\right)\end{aligned}\quad (42)$$

Using $f(x) \simeq \frac{e^{-2x}}{\pi^2} \sqrt{\frac{2\pi}{x}}$ when $x \rightarrow \infty$, the integrand in (42)

$$1 - e^{-m(1-q)f(\frac{y \ln m}{2})} \approx 1 - e^{-(1-q)\frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{y \ln m}}}$$

behaves, in the $m \rightarrow \infty$ limit, like

$$\begin{array}{lll} 0 & \text{if} & y > 1 \\ 1 & \text{if} & y < 1 \end{array}$$

and so, at leading order, $\Phi_q(m) = \frac{\pi \ln m}{2}$.

Focusing now on the subleading correction at order $\frac{1}{\sqrt{\ln m}}$, namely

$$\frac{\pi \ln m}{2} \left(\int_1^\infty dy \left(1 - e^{-(1-q)\frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{y \ln m}}}\right) - \int_0^1 dy e^{-(1-q)\frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{y \ln m}}}\right) \quad (43)$$

let us first consider the $y > 1$ integration. One has to compute

$$a \approx \frac{\pi \ln m}{2} \int_1^\infty dy (1 - q) \frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{y \ln m}} \approx \frac{\pi \ln m}{2} \int_1^\infty dy (1 - q) \frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{\ln m}}$$

where one has used that, because of the m^{1-y} factor, y is peaked to 1 when $m \rightarrow \infty$. One obtains

$$a = \frac{1 - q}{\sqrt{\pi \ln m}} + o\left(\frac{1}{\sqrt{\ln m}}\right) \quad (44)$$

Considering next the $y < 1$ integration one has to compute

$$\begin{aligned} b &\approx -\frac{\pi \ln m}{2} \int_0^1 dy e^{-(1-q)\frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{y \ln m}}} \\ &\approx -\frac{\pi \ln m}{2} \int_0^1 dy e^{-(1-q)\frac{m^{1-y}}{\pi^2} \sqrt{\frac{4\pi}{\ln m}}} \\ &= -\frac{\pi \ln m}{2} \int_0^1 dy e^{-a' m^{1-y}} \quad \text{with} \quad a' = \frac{1 - q}{\pi^2} \sqrt{\frac{4\pi}{\ln m}} \end{aligned}$$

Setting $a' m^{1-y} = z$ with $-dy \ln m = \frac{dz}{z}$ leads to

$$b \approx -\frac{\pi}{2} \int_{a'}^{a'm} e^{-z} \frac{dz}{z} = -\frac{\pi}{2} \left([\ln z e^{-z}]_{a'}^{a'm} + \int_{a'}^{a'm} \ln z e^{-z} dz \right)$$

At order $\frac{1}{\sqrt{\ln m}}$, one has $[\ln z e^{-z}]_{a'}^{a'm} \approx -\ln a' (1 - a')$ and $\int_{a'}^{a'm} \ln z e^{-z} dz \approx \int_0^\infty \ln z e^{-z} dz - \int_0^{a'} \ln z e^{-z} dz \approx C - \int_0^{a'} \ln z (1 - z) dz$ where C is the Euler constant. The last remaining integral is straightforward and finally

$$b \approx \frac{\pi}{2} (\ln a' + C - a') \quad (45)$$

Noticing that $a' \frac{\pi}{2} = \frac{1 - q}{\sqrt{\pi \ln m}} \approx a$, we are left with $a + b \approx \frac{\pi}{2} (\ln a' + C)$ so that

$$\Phi_q(m) = \frac{\pi}{2} \ln m - \frac{\pi}{4} \ln \ln m + \frac{\pi}{2} \left(\ln(1 - q) + \ln \sqrt{\frac{4}{\pi^3} + C} \right) + o\left(\frac{1}{\sqrt{\ln m}}\right) \quad (46)$$

References

- [1] "Algebraic and arithmetic are for m planar Brownian paths" J. Desbois and S. Ouvry, arXiv: 1101.4135, J. Stat. Mech. (2011) P05024
- [2] R. P. Feynman and A. R. Hibbs, Quantum Mechanics and Path Integrals, New York, Mc Graw-Hill (1965)
- [3] A. Comtet, J. Desbois and S. Ouvry, J. Phys. **A 23** (1990) 3563
- [4] C. Garban and J. A. Trujillo Ferreras, Commun. Math. Phys. **264** (2006) 797
- [5] A. Comtet, S.N. Majumdar and J. Randon-Furling, Phys. Rev. Lett., v-103, 140602 (2009).
- [6] M. Abramowitz and I. Stegun, Handbook of mathematical functions, New York, Dover Publications (1965).