

LOW-DIMENSIONAL FREE AND LINEAR REPRESENTATIONS OF $\text{Out}(F_3)$

Dawid Kielak¹
University of Bonn
kielak@math.uni-bonn.de
November 14, 2018

ABSTRACT. We study homomorphisms from $\text{Out}(F_3)$ to $\text{Out}(F_5)$, and $\text{GL}_m(\mathbb{K})$ for $m \leq 6$, where \mathbb{K} is a field of characteristic other than 2 or 3. We conclude that all \mathbb{K} -linear representations of dimension at most 6 of $\text{Out}(F_3)$ factor through $\text{GL}_3(\mathbb{Z})$, and that all homomorphisms $\text{Out}(F_3) \rightarrow \text{Out}(F_5)$ have finite image.

1 Introduction

This paper constitutes a part of a project of understanding homomorphisms $\text{Out}(F_n) \rightarrow \text{GL}_m(\mathbb{K})$ and $\text{Out}(F_n) \rightarrow \text{Out}(F_m)$, i.e. the \mathbb{K} -linear representation theory and the *free representation theory* of $\text{Out}(F_n)$ respectively (see [13]). The first problem, as well as linear representations of $\text{Aut}(F_n)$, has been studied for example by Potapchik–Rapinchuk [15] and Grunewald–Lubotzky [9]; the latter has been addressed for example by Khramtsov [12], Bogopolski–Puga [3], Bridson–Vogtmann [5, 6], and Aramayona–Leininger–Souto [1].

The results obtained so far have a tendency of working only for large n ; in particular the case of $\text{Out}(F_3)$ has not been studied extensively. It was known (thanks to a result of Khramtsov [12]) that there are no embeddings

$$\text{Out}(F_3) \hookrightarrow \text{Out}(F_4)$$

It was also known that there is an embedding

$$\text{Out}(F_3) \hookrightarrow \text{Out}(F_{55})$$

(this is due to Bogopolski–Puga [3]).

Our attempt to expand the understanding of free representations of $\text{Out}(F_3)$ follows the same general route as our proof in [13]. First we investigate the linear representation theory of $\text{Out}(F_3)$, and we prove

¹The author was supported by the EPSRC of the United Kingdom

Theorem 3.19. *Let ϕ be a \mathbb{K} -linear, six-dimensional representation of $\text{Out}(F_3)$, where \mathbb{K} is a field of characteristic other than 2 or 3. Then ϕ factors as*

$$\begin{array}{ccc} \text{Out}(F_3) & \xrightarrow{\phi} & \text{GL}_6(\mathbb{K}) \\ & \searrow \pi & \uparrow \\ & & \text{GL}_3(\mathbb{Z}) \end{array}$$

where π is the natural projection $\text{Out}(F_3) \rightarrow \text{Out}(H_1(F_3, \mathbb{Z})) \cong \text{GL}_3(\mathbb{Z})$.

It is worth noting here that Turchin–Willwacher [16] constructed a 7-dimensional \mathbb{Q} -linear representation of $\text{Out}(F_3)$ which does not factor through π . Thus our result completes the search for the smallest dimension in which such a representation occurs (at least for fields of characteristic zero).

The second part of this paper deals with the action of two finite subgroups of $\text{Out}(F_3)$ on graphs of rank 5. This was also our strategy in [13], yet the situation here is quite different. The finite groups under consideration do not contain (comparatively) large simple groups, as is the case in the higher rank case; on the other hand, the groups are of order 48, and hence are rather tangible. Our considerations yield

Theorem 4.10. *Suppose $\phi : \text{Out}(F_3) \rightarrow \text{Out}(F_5)$ is a homomorphism. Then the image of ϕ is finite.*

The general question of finding the smallest $m > 3$ such that there is a homomorphism $\text{Out}(F_3) \rightarrow \text{Out}(F_m)$ that is injective, or at least has infinite image, remains open.

Acknowledgements. The author wishes to thank Martin R. Bridson for all his help.

2 Notation and preliminaries

Let us first establish some conventions and definitions.

Definition 2.1. We say that X is a *graph* if and only if it is a 1-dimensional CW complex. The closed 1-cells of X will be called *edges*, the 0-cells will be called *vertices*. The sets of vertices and edges of a graph will be denoted by $V(X)$ and $E(X)$ respectively. The points of intersection of an edge with the vertex set are referred to as *endpoints* of the edge.

We will equip X with the standard path metric in which the length of each edge is 1.

Given two graphs X and Y , a function $f : X \rightarrow Y$ is a *morphism of graphs* if and only if f is a continuous map sending $V(X)$ to $V(Y)$, and sending each open edge in X either to a vertex in Y or isometrically onto an open edge in Y . Note that a morphism can invert edges.

When we say that a group G *acts on a graph* X , we mean that it acts by graph morphisms.

We say that a graph X is *directed* if and only if it comes equipped with a map $o : E(X) \rightarrow X$ such that $o(e)$ is a point on the interior of e of distance $\frac{1}{3}$ from one of its endpoints. We also define $\iota, \tau : E(X) \rightarrow V(X)$ by setting $\tau(e)$

to be the endpoint of e closest to $o(e)$, and $\iota(e)$ to be the endpoint of e farthest from $o(e)$. Note that we allow $\iota(e) = \tau(e)$.

The *rank* of a connected graph is defined to be the size of a minimal generating set of its fundamental group (which is a free group).

Let us also define two families of graphs.

Definition 2.2. The graph with one vertex and n edges will be referred to as the n -*rose*.

The graph with two vertices and n edges, such that each edge has two distinct endpoints, will be referred to as the n -*cage*.

Notation 2.3. Let G be a group. We will adopt the following notation:

- for two elements $g, h \in G$, we define $g^h = h^{-1}gh$;
- for two elements $g, h \in G$, we define $[g, h] = ghg^{-1}h^{-1}$;

We will also use \mathbb{Z}_k to denote the cyclic group of order k .

Definition 2.4. Let us introduce the following notation for elements of $\text{Aut}(F_n)$, the automorphism group of F_n , where F_n is the free group on $\{a_1, \dots, a_n\}$:

$$\begin{aligned} \epsilon_i &: \begin{cases} a_i \mapsto a_i^{-1}, \\ a_j \mapsto a_j, \end{cases} & j \neq i & \quad \sigma_{ij} &: \begin{cases} a_i \mapsto a_j, \\ a_j \mapsto a_i, \\ a_k \mapsto a_k, \end{cases} & k \notin \{i, j\} \\ \rho_{ij} &: \begin{cases} a_i \mapsto a_i a_j, \\ a_k \mapsto a_k, \end{cases} & k \neq i & \quad \lambda_{ij} &: \begin{cases} a_i \mapsto a_j a_i, \\ a_k \mapsto a_k, \end{cases} & k \neq i \end{aligned}$$

Let us also define $\Delta = \prod_{i=1}^n \epsilon_i$ and

$$\sigma_{i(n+1)} : \begin{cases} a_i \mapsto a_i^{-1}, \\ a_j \mapsto a_j a_i^{-1}, \end{cases} \quad j \neq i$$

Below we give an explicit presentation of $\text{Out}(F_n)$, the outer automorphism group of F_n :

Theorem 2.5 (Gersten's presentation [8]). *Suppose $n \geq 3$. The group $\text{Out}(F_n)$ is generated by $\{\epsilon_1, \rho_{ij}, \lambda_{ij} \mid i, j = 1, \dots, n, i \neq j\}$, with relations*

- $[\rho_{ij}, \rho_{kl}] = [\lambda_{ij}, \lambda_{kl}] = 1$ for $k \notin \{i, j\}, l \neq i$;
- $[\lambda_{ij}, \rho_{kl}] = 1$ for $k \neq j, l \neq i$;
- $[\rho_{ij}^{-1}, \rho_{jk}^{-1}] = [\rho_{ij}, \lambda_{jk}] = [\rho_{ij}^{-1}, \rho_{jk}]^{-1} = [\rho_{ij}, \lambda_{jk}^{-1}]^{-1} = \rho_{ik}^{-1}$ for $k \notin \{i, j\}$;
- $[\lambda_{ij}^{-1}, \lambda_{jk}^{-1}] = [\lambda_{ij}, \rho_{jk}] = [\lambda_{ij}^{-1}, \lambda_{jk}]^{-1} = [\lambda_{ij}, \rho_{jk}^{-1}]^{-1} = \lambda_{ik}^{-1}$ for $k \notin \{i, j\}$;
- $\rho_{ij} \rho_{ji}^{-1} \lambda_{ij} = \lambda_{ij} \lambda_{ji}^{-1} \rho_{ij}, (\rho_{ij} \rho_{ji}^{-1} \lambda_{ij})^4 = 1$;
- $[\epsilon_1, \rho_{ij}] = [\epsilon_1, \lambda_{ij}] = 1$ for $i, j \neq 1$;
- $\rho_{12}^{\epsilon_1} = \lambda_{12}^{-1}, \rho_{21}^{\epsilon_1} = \rho_{21}^{-1}$;
- $\epsilon_1^2 = 1$;

- $\prod_{i \neq j} \rho_{ij} \lambda_{ij}^{-1} = 1$ for each fixed j .

Note the action of $\text{Aut}(F_n)$ on F_n and $\text{Out}(F_n)$ on the conjugacy classes of F_n is **on the left**.

Definition 2.6. Let us define some finite subgroups of $\text{Out}(F_n)$:

$$\begin{aligned} S_n &\cong \langle \{\sigma_{ij} \mid i, j = 1, \dots, n, i \neq j\} \rangle \\ S_{n+1} &\cong \langle \{\sigma_{ij} \mid i, j = 1, \dots, n+1, i \neq j\} \rangle \\ \mathbb{Z}_2^n \rtimes S_n \cong W_n &= \langle \{\epsilon_1, \sigma_{ij} \mid i, j = 1 \dots, n, i \neq j\} \rangle \\ \mathbb{Z}_2 \times S_{n+1} \cong G_n &= \langle \{\Delta, \sigma_{ij} \mid i, j = 1 \dots, n+1, i \neq j\} \rangle \end{aligned}$$

We do not give distinctive names to the first two (symmetric) groups; instead, we will usually refer to them as respectively $S_n < W_n$ and $S_{n+1} < G_n$. More generally, whenever we mention S_n or S_{n+1} as subgroups of $\text{Out}(F_n)$, we mean these two groups.

In the case of $\text{Out}(F_3)$, define V_4 and A_4 to be the Klein 4-group and the alternating group of degree 4 satisfying

$$V_4 < A_4 < S_4 < G_3 < \text{Out}(F_3).$$

Note that we abuse notation by also using S_n to denote the abstract symmetric group of degree n , and $A_n < S_n$ to denote its maximal alternating subgroup.

Note that, if $i, j \leq n$, we have

$$\epsilon_i \sigma_{ij} = \lambda_{ij} \lambda_{ji}^{-1} \rho_{ij} = \rho_{ij} \rho_{ji}^{-1} \lambda_{ij}$$

and

$$\epsilon_1 \sigma_{i(n+1)} = \prod_{j \neq i} \rho_{ji} = \prod_{j \neq i} \lambda_{ji}$$

and the subgroup $S_n < \text{Out}(F_n)$ defined above acts on the sets

$$\begin{aligned} &\{\epsilon_i \mid i = 1 \dots, n\}, \\ &\{\rho_{ij} \mid i, j = 1 \dots, n, i \neq j\}, \text{ and} \\ &\{\lambda_{ij} \mid i, j = 1 \dots, n, i \neq j\} \end{aligned}$$

by permuting the indices in the natural way.

3 Representations of $\text{Out}(F_3)$ in dimensions five and six

Before we start investigating 5- and 6-dimensional representations of $\text{Out}(F_3)$, let us first prove the following.

Proposition 3.1. *Suppose $\phi : \text{Out}(F_3) \rightarrow G$ is a group homomorphism such that its kernel contains V_4 . Then ϕ factors as*

$$\phi : \text{Out}(F_3) \rightarrow \mathbb{Z}_2 \rightarrow G$$

and the map is determined by the image of ϵ_1 .

Proof. Since V_4 lies in the kernel, we have $\phi(\sigma_{14}) = \phi(\sigma_{23})$. Hence

$$\phi(\rho_{21}) = \phi(\rho_{21}^{\rho_{21}^{21}\rho_{31}}) = \phi(\rho_{21}^{\epsilon_1\sigma_{14}}) = \phi(\rho_{21})^{\phi(\epsilon_1)\phi(\sigma_{23})} = \phi(\rho_{31}^{-1})$$

and so

$$\phi(\rho_{21}) = \phi(\rho_{21}^{\epsilon_3}) = \phi(\rho_{31}^{-1})^{\phi(\epsilon_3)} = \phi(\lambda_{31})$$

Now

$$\phi(\rho_{31}^{-1}) = \phi([\rho_{32}^{-1}, \rho_{21}^{-1}]) = [\phi(\rho_{32}^{-1}), \phi(\lambda_{31}^{-1})] = 1$$

Thus ρ_{31} lies in the kernel of ϕ . We can however conjugate ρ_{31} to each ρ_{ij} using S_3 , and so all elements ρ_{ij} lie in the kernel. The result follows. \square

Let us recall some basic terminology of the representation theory of symmetric groups.

Remark 3.2. Let S_n be a symmetric group of rank n , and let \mathbb{K} be a field with $\text{char}(\mathbb{K}) = 0$ or $\text{char}(\mathbb{K}) > n$. Then there is a one-to-one correspondence between irreducible \mathbb{K} -linear representations of S_n and partitions of n . In particular the representation corresponding to

- (n) is called *trivial*;
- (1^n) is called *determinant*;
- $(n-1, 1)$ is called *standard*;
- $(2, 1^{n-1})$ is called *signed standard*.

Moreover, a direct sum of the standard and trivial representation is known as the *permutation* representation, and a direct sum of the signed standard and determinant representation is known as the *signed permutation* representation.

Now let us turn our attention to representations of the group W_n .

Definition 3.3. Let V be a representation of W_n . Let $N = \{1, \dots, n\}$. Define

- for each $I \subseteq N$, $E_I = \{v \in V \mid \epsilon_i v = (-1)^{\chi_I(i)} v\}$, where χ_I is the characteristic function of I ;
- $V_i = \bigoplus_{|I|=i} E_I$

We will slightly abuse notation, and often omit parentheses and write E_1 for $E_{\{1\}}$, etc.

Lemma 3.4. *Let V be a representation of W_n . Then $\dim V_i = \binom{n}{i} \dim E_1$ where $|I| = i$.*

Proof. The symmetric group $S_n < W_n$ acts on $\{\epsilon_1, \dots, \epsilon_n\}$ by permuting the indices in the natural way. Hence its action on V_i will permute subspaces E_I by permuting subsets of N of size i . Thus each E_I , for a fixed size of I , has the same dimension. The result follows. \square

An immediate consequence of the above is the following.

Lemma 3.5. *Let V be a 2-dimensional \mathbb{K} -linear representation of $\text{Out}(F_3)$, where $\text{char}(\mathbb{K}) \neq 2$. Then the representation factors as*

$$\text{Out}(F_3) \rightarrow \mathbb{Z}_2 \rightarrow \text{GL}_2(\mathbb{K})$$

and is determined by the image of ϵ_1 .

Proof. Let $\phi : \text{Out}(F_3) \rightarrow \text{GL}_2(\mathbb{K})$ be the representation. Lemma 3.4 tells us that (with the notation of Definition 3.3) $V = V_0 \oplus V_3$.

Since $S_4 < G_3$ commutes with Δ , $V = V_0 \oplus V_3$ is a decomposition of S_4 -modules. Now each of these submodules has dimension at most 2. There are at most three irreducible \mathbb{K} -linear representations of S_4 of dimension at most 2: the trivial representation (corresponding to partition (4)), the determinant representation (corresponding to partition (1⁴)), and the one given by a partition (2, 2) (note that the latter might not be irreducible when $\text{char}(\mathbb{K}) = 3$). In all three cases, the action of $V_4 < S_4$ is trivial. This implies that we have satisfied all the requirements of Proposition 3.1, and the result follows. \square

Lemma 3.6. *Let V be a \mathbb{K} -linear representation of $\text{Out}(F_n)$, where \mathbb{K} is a field of characteristic other than 2. Then, with the notation above, we have*

$$V = \bigoplus_{i=0}^n V_i$$

and for each $i \neq j$, $J \subseteq N \setminus \{i, j\}$ we have

$$\rho_{ij}(E_J \oplus E_{J \cup \{i\}} \oplus E_{J \cup \{j\}} \oplus E_{J \cup \{i, j\}}) = E_J \oplus E_{J \cup \{i\}} \oplus E_{J \cup \{j\}} \oplus E_{J \cup \{i, j\}}.$$

An identical statement holds for λ_{ij} .

Proof. The first statement follows directly from the fact that we can simultaneously diagonalise commuting involutions ϵ_i , since we are working over a field \mathbb{K} whose characteristic is not 2.

For the second statement, let us note that $[\rho_{ij}, \epsilon_k] = 1$ for each $k \notin \{i, j\}$. Hence for each $I \subseteq N$:

$$\rho_{ij}(E_I) \leq \bigoplus_{J \Delta I \subseteq \{i, j\}} E_J$$


where $A \Delta B$ denotes the symmetric difference of two sets A and B . An identical argument works for λ_{ij} . \square

To help us visualise the combinatorics of representations of $\text{Out}(F_n)$ we are going to use the following diagrams.

Definition 3.7. Suppose V is a finite dimensional, \mathbb{K} -linear representation of $\text{Out}(F_n)$ over any field \mathbb{K} , and let $x \in \text{Out}(F_n)$. Let us use the notation of Definition 3.3. We define *the minimal diagram for x over V* (often abbreviated to the minimal diagram for x) to be a directed graph D with the vertex set equal to a subset S of the power set of $N = \{1, 2, \dots, n\}$, where $I \in S$ if and only if $E_I \neq \{0\}$, and the edge set given by the following rule: there is a directed edge from I to J if and only if $p_J(x(E_I)) \neq \{0\}$, where $p_J : \bigoplus_{K \subseteq N} E_K \rightarrow E_J$ is the natural projection.

We also say that a graph D' is *a diagram for x over V* if and only if the minimal diagram D for x is a subgraph of D' .

In practice, when realising these diagrams in terms of actual pictures, we are going to align vertices corresponding to subsets of N of the same cardinality in horizontal lines; each such line will correspond to some V_i . We are also going to represent edges as follows: if two vertices are joined by two directed edges, we are going to draw one edge without any arrowheads between them; we are not going to draw edges from a vertex to itself – instead, if a vertex does not have such a loop, then all edges emanating from it will be drawn with a tail (see example below);




To get a firmer grip on these diagrams, let us have a look at a number of facts one can easily deduce from (not necessarily minimal) diagrams.

Example 3.8. Let Γ_0 be a connected component of Γ , a diagram for x . Let $v \in \bigoplus_{I \in V(\Gamma_0)} E_I$ be a vector. Then $v = \sum v_I$ where $v_I \in E_I$. Let $J \notin V(\Gamma_0)$. Note that there are no edges between J and $V(\Gamma_0)$, and so $p_J(x(v_I)) = 0$ for all $I \in V(\Gamma_0)$. Hence $x(v_I) \in \bigoplus_{I \in V(\Gamma_0)} E_I$ and therefore


$$x\left(\bigoplus_{I \in V(\Gamma_0)} E_I\right) = \bigoplus_{I \in V(\Gamma_0)} E_I$$

The following illustrates the relationship between our diagrams and matrices.

Example 3.9. Suppose we have a diagram for x with two vertices, I and J say, such that the union of the connected components containing these vertices does not contain any other vertex. Fix a basis for E_I and E_J . The following illustrates the way the x action on $E_I \oplus E_J$ (seen as a matrix) depends on the diagram:



Example 3.10. Suppose we have a diagram Γ for x such that Γ has a connected component with only two vertices, I and J say, as depicted below.



Exercise 3.8 tells us that $E_I \oplus E_J$ is x -invariant, and thus $x|_{E_I \oplus E_J}$ is an isomorphism. Let $\{v_1, \dots, v_k\}$ be a basis for E_I . Our diagram tells us that $p_I(x(v_i)) = 0$ for each i , and so $x(v_i) \in E_J$. Since x is an isomorphism, we immediately see that

$$\{x(v_1), \dots, x(v_k)\}$$

is a linearly independent set, and hence $\dim E_J \geq \dim E_I$.

Example 3.11. Let Γ be a diagram for x . Note that $p_J(x^{\epsilon_i}(v)) = \pm p_J(x(v))$ whenever $v \in E_I$ for some I . Therefore $p_J(x^{\epsilon_i}(E_I)) = \{0\}$ if and only if $p_J(x(E_I)) = \{0\}$, and so Γ is also a diagram for x^{ϵ_i} .


Now consider $\sigma \in S_n$. We have

$$p_J(x^\sigma(E_I)) = \sigma^{-1}\left(p_{\sigma(J)}(x(E_{\sigma(I)}))\right)$$


and so the image of Γ under the graph morphism induced by $I \mapsto \sigma(I)$ is a diagram for x^σ .

We will use the last example very often, for example to relate diagrams for ρ_{21} with ones for $\rho_{21}^{-1} = \rho_{21}^{\epsilon_1}$ or $\rho_{31} = \rho_{21}^{\sigma_{23}}$.

Lemma 3.12. Let $\phi : \text{Out}(F_3) \rightarrow V$ be a representation such that we have a diagram for ρ_{21} of the form



where $\dim E_i = 1$ for all i . Then ρ_{21} has a diagram




Note that the above diagrams are not necessarily minimal; this means that the lines in the top diagram between E_1 and E_2 , and between V_3 and E_3 , may represent the real situation, or it may be possible to replace each by an edge directed from the right to the left, or one directed from the left to the right, or no edge at all. Lemma 3.12 says that the line between E_2 and E_1 can indeed be replaced by either one of the directed edges or no edge at all.

Proof. Firstly, note that the left-hand side diagram corresponds to

$$p_2(\rho_{21}(E_1)) = \{0\}$$

and the right-hand side diagram to $p_1(\rho_{21}(E_2)) = \{0\}$. Suppose for a contradiction that we have neither of the diagrams, that is that $p_1(\rho_{21}(E_2)) \neq \{0\}$ and $p_2(\rho_{21}(E_1)) \neq \{0\}$. We claim that then ρ_{21} has a diagram




Once we have proven the above claim, take $x \in E_2 \setminus \{0\}$. Then, by the assumptions of our claim, $p_1(\rho_{21}(x)) \neq 0$, since E_2 is 1 dimensional, and so spanned by x . Thus we have

$$\rho_{21}(x) = x_1 + x_2$$


with $x_1 \in E_1 \setminus \{0\}$ and $x_2 \in E_2$. Now

$$p_3(\rho_{31}\rho_{21}(x)) = p_3(\rho_{31}(x_1)) + p_3(\rho_{31}(x_2))$$

By the conclusion of our claim, the following is a diagram for ρ_{21}



By Exercise 3.11 (using the action of σ_{23}) the following is then a diagram for ρ_{31}



Therefore

$$p_3(\rho_{31}(x_2)) = 0$$

and so

$$p_3(\rho_{31}\rho_{21}(x)) = p_3(\rho_{31}(x_1))$$

Note that we can apply the assumptions of our claim to ρ_{31} as well, using the relation $\rho_{31} = \rho_{21}^{\sigma_{23}}$ and Example 3.11; specifically we may assume that $p_3(\rho_{31}(E_1)) \neq \{0\}$. Now, observing that x_1 spans E_1 (which is 1 dimensional), we conclude that

$$p_3(\rho_{31}\rho_{21}(x)) \neq 0$$

But $\rho_{31}(x) \in E_2$ by the diagram above (recall that $x \in E_2$), and thus $\rho_{21}\rho_{31}(x) \in E_2 \oplus E_1$, which in turn implies that

$$p_3(\rho_{21}\rho_{31}(x)) = 0$$

This contradicts the relation $[\rho_{21}, \rho_{31}] = 1$, and our proof is complete.

Now, to prove the claim, let $v_1 \in E_1 \setminus \{0\}$. Then $v_3 = \sigma_{23}\rho_2(\rho_{21}(v_1))$ generates E_3 . Now if

$$p_{1,2,3}(\rho_{21}(v_3)) = 0$$

then we have proven our claim. If not, let $U = \langle u \rangle$ be a subspace of V_3 of dimension 1, where $u = p_{1,2,3}(\rho_{21}(v_3))$. Note that $\rho_{21}v_3 \in u + E_3$, and so

$$\rho_{21}^{-1}(-v_3) = \epsilon_1\rho_{21}\epsilon_1(-v_3) \in u + E_3.$$

In particular, $\rho_{21}^{-1}(-v_3) - v'_3 = u$ for some $v'_3 \in E_3$. Now

$$\rho_{21}u = \rho_{21}(\rho_{21}^{-1}(-v_3) - v'_3) \in E_3 \oplus U,$$

and hence $U \oplus E_3$ is ρ_{21} -invariant.

Let us rewrite $\rho_{21}\rho_{31} = \rho_{31}\rho_{21}$ as

$$\rho_{21}\sigma_{23}\rho_{21}\sigma_{23} = \sigma_{23}\rho_{21}\sigma_{23}\rho_{21}$$

which yields $[\rho_{21}\sigma_{23}\rho_{21}, \sigma_{23}] = 1$.

Now E_1 is σ_{23} -invariant and one dimensional, and so each non-zero vector in E_1 is an eigenvector of σ_{23} with eigenvalue $\mu = \pm 1$. In particular v_1 lies in the μ -eigenspace of σ_{23} , and hence so does $\rho_{21}\sigma_{23}\rho_{21}(v_1)$, since σ_{23} and $\rho_{21}\sigma_{23}\rho_{21}$ commute. Since V_3 and V_1 are σ_{23} -invariant (in fact S_3 -invariant), we also have $p_{1,2,3}(\rho_{21}\sigma_{23}\rho_{21}(v_1))$ lying in the μ -eigenspace of σ_{23} .

Now let

$$\rho_{21}(v_1) = y_1 + y_2$$

with $y_i \in E_i$. We have

$$\begin{aligned} p_{1,2,3}(\rho_{21}\sigma_{23}\rho_{21}(v_1)) &= p_{1,2,3}(\rho_{21}\sigma_{23}(y_1 + y_2)) \\ &= p_{1,2,3}(\rho_{21}(\mu y_1 + \sigma_{23}(y_2))) \\ &= p_{1,2,3}(\rho_{21}\sigma_{23}(y_2)) \end{aligned}$$

since $\rho_{21}(\mu y_2) \in E_1 \oplus E_2$ by assumption of our lemma, and thus lies in the kernel of $p_{1,2,3}$.

By definition we have

$$v_3 = \sigma_{23}p_2\rho_{21}(v_1) = \sigma_{23}(y_2)$$

and so we deduce

$$p_{1,2,3}(\rho_{21}\sigma_{23}\rho_{21}(v_1)) = p_{1,2,3}(\rho_{21}(v_3)) = u$$

where the last equality is the definition of u . Therefore U lies in the μ -eigenspace of σ_{23} .

Note that the eigenspaces of Δ are $V_0 \oplus V_2$ and $V_1 \oplus V_3$, and on each Δ acts as ± 1 . Hence, since $V_0 \oplus V_2 = \{0\}$, we see that $[\phi(\rho_{21}), \phi(\Delta)] = 1$. Therefore, since $\rho_{ij}^\Delta = \lambda_{ij}$, the elements ρ_{ij} and λ_{ij} act identically for each i and j . This in turn implies that $[\phi(\rho_{21}), \phi(\rho_{23})] = 1$ since $[\rho_{21}, \lambda_{23}] = 1$. Rewriting the first relation as before we get $[\phi(\rho_{21}\sigma_{13}\rho_{21}), \phi(\sigma_{13})] = 1$.

Let $v_2 \in E_2 \setminus \{0\}$. We have $\rho_{21}(v_2) = z_1 + z_2$, with $z_i \in E_i$, and with $z_1 \neq 0$. Thus z_1 is a non-zero multiple of v_1 , since E_1 is 1 dimensional. Arguing as before we now get

$$\langle p_{1,2,3}(\rho_{21}\sigma_{13}\rho_{21}(v_2)) \rangle = \langle p_{1,2,3}(\rho_{21}\sigma_{13}(z_1)) \rangle = U$$

The group S_3 can act on V_1 in two ways: via the permutation or the signed permutation representation. In each case however, if E_1 is in the μ -eigenspace of σ_{23} , then E_2 is in the μ -eigenspace of σ_{13} . Thus $\sigma_{13}(v_2) = \mu v_2$ and so U lies in the μ -eigenspace of σ_{13} , arguing as before. Therefore U also lies in the μ -eigenspace of $\sigma_{12} = \sigma_{23}^{\sigma_{13}}$, and so does E_3 . This shows that

$$\phi(\rho_{12})|_{E_3 \oplus U} = \phi(\rho_{21}^{\sigma_{12}})|_{E_3 \oplus U} = \phi(\rho_{21})|_{E_3 \oplus U}$$

and therefore that

$$\phi(\rho_{21})|_{E_3 \oplus U} = \phi(\lambda_{21})|_{E_3 \oplus U} = \phi(\rho_{21}\rho_{12}^{-1}\lambda_{21})|_{E_3 \oplus U} = \phi(\epsilon_2\sigma_{12})|_{E_3 \oplus U}$$


But $\epsilon_2\sigma_{12}(E_3) = E_3$ and so ρ_{21} has a diagram of the form claimed. \square

We shall now focus on five- and six-dimensional representations of $\text{Out}(F_3)$.


Lemma 3.13. *Let V be a \mathbb{K} -linear, six-dimensional representation of $\text{Out}(F_3)$, where $\text{char}(\mathbb{K}) \neq 2$. Suppose that, with notation of Definition 3.3,*

$$\dim V_1 \oplus V_2 \leq 3.$$

Then, if $V_2 = \{0\}$, we have a (not necessarily minimal) diagram for ρ_{21} of the form



and if $V_1 = \{0\}$ of the form




In both cases at least one of $V_0 \oplus V_2$ and $V_1 \oplus V_3$ is $\text{Out}(F_3)$ -invariant.

Proof. Lemma 3.4 tells us that the dimensions of V_1 and V_2 are divisible by 3. Hence, by assumption, at least one of V_1 and V_2 is trivial. If both of them are trivial then Lemma 3.6 immediately tells us that the decomposition $V = V_0 \oplus V_3$ is preserved by each ρ_{ij} and λ_{ij} . Thus the minimal diagram for ρ_{21} is a subdiagram of all the above.


Suppose one of V_1, V_2 is non-trivial. Without loss of generality let us assume $\dim V_1 \neq 0$. Again by assumption we see that $\dim V_1 = 3$, and hence $\dim E_i = 1$ for all i .

Our strategy here is to start with the most general possible diagram for ρ_{21} , and then gradually add restriction until we arrive at one of the diagrams described above.

Lemma 3.6 allows us to conclude that we have the following diagrams for ρ_{21} and ρ_{31} respectively:




The element Δ lies in the centre of G_3 , and so in particular $[\Delta, \epsilon_1 \sigma_{14}] = 1$. This implies that $\epsilon_1 \sigma_{14}$ preserves the eigenspaces of Δ , which happen to be the direct sums of all subspaces V_i with the index i of a given parity (even for the $(+1)$ - and odd for the (-1) -eigenspace). Hence the following is a diagram for $\epsilon_1 \sigma_{14}$:



But, in $\text{Out}(F_3)$, we have $\epsilon_1 \sigma_{14} = \rho_{31} \rho_{21}$, and Example 3.8 tells us that

$$\rho_{31} \left(p_2(\rho_{21}(V_0)) \right) \leq E_2 \oplus V_3$$


We can therefore conclude that $p_2(\rho_{21}(V_0)) = \{0\}$, and so that we have a diagram for ρ_{21} as follows:




Again, by Example 3.8, $\rho_{31}|_{E_2 \oplus V_3}$ is an isomorphism. Hence there exists $v \in E_2 \oplus V_3$ such that $\langle \rho_{31}(v) \rangle = E_2$. Since $v \in E_2 \oplus V_3 \leq V_1 \oplus V_3$, also $\epsilon_1 \sigma_{14}(v) \in V_1 \oplus V_3$. Now

$$\epsilon_1 \sigma_{14}(v) = \rho_{21} \rho_{31}(v)$$

and so we conclude that ρ_{21} has a diagram




Note that ρ_{21} either has a diagram




or $p_1(\rho_{21}(E_2)) = E_1$, since $\dim E_1 = 1$.


If $\rho_{21}(E_2)$ projects surjectively onto E_1 , applying $\rho_{31}\rho_{21} = \epsilon_1\sigma_{14}$ to E_2 yields a diagram for ρ_{31} of the form



and, after conjugating by σ_{23} (see Example 3.11), ρ_{21} has a diagram




Requiring $\rho_{21}^{\epsilon_1} = \rho_{21}^{-1}$ yields two possibilities for a diagram for ρ_{21} :




The first diagram is as required. The second diagram gives a required diagram after applying Lemma 3.12.

We still have to consider the case of a diagram




for ρ_{21} . Applying $\epsilon_1\sigma_{14} = \rho_{21}\rho_{31}$ to V_0 yields a diagram for ρ_{21} of the form



The second of these diagrams is as required.

Let us now focus on the first of the above diagrams. Note that, by Example 3.11, this is also a diagram for λ_{21} , and that a diagram for ρ_{12}^{-1} is as follows:




Let v_1 be a generator of E_1 . Apply $\epsilon_2\sigma_{12} = \rho_{21}\rho_{12}^{-1}\lambda_{21}$ to v_1 and observe that $\epsilon_2\sigma_{12}(v_1) = v_2$, a generator of E_2 . Now let x be the E_1 component of $\lambda_{21}(v_1)$. Note that $\rho_{12}^{-1}\lambda_{21}(v_1)$ has a non-trivial E_1 component if and only if x is not zero. But such a non-trivial component yields a non-zero component in $E_1 \oplus V_0$ of $\rho_{21}\rho_{12}^{-1}\lambda_{21}(v_1)$. This is impossible, since $\epsilon_2\sigma_{12}(v_1) = v_2$ has no such components. Thus $x = 0$, $\lambda_{21}(v_1)$ lies in V_0 , and

$$\rho_{12}^{-1}|_U : U \rightarrow E_2$$


is an isomorphism, where $U = \langle \lambda_{21}(v_1) \rangle$. Hence $\rho_{12}|_{E_2} : E_2 \rightarrow U$ is an isomorphism as well.

We claim that $\rho_{ij}^{\pm 1}|_{E_j}, \lambda_{ij}^{\pm 1}|_{E_j} : E_j \rightarrow U$ are all isomorphisms. We have established this for λ_{21} and ρ_{12} . Conjugating by ϵ_1 and ϵ_2 establishes the claim also for $\rho_{21}^{-1}, \rho_{21}, \lambda_{21}^{-1}, \rho_{12}^{-1}, \lambda_{12}$ and λ_{12}^{-1} . Using the fact that $\epsilon_1\sigma_{14} = \rho_{31}\rho_{21}$ preserves $V_1 \oplus V_3$ we immediately conclude that the claim also holds for ρ_{31}^{-1} , and hence in particular also for ρ_{13} (repeating the argument above). Now the relation $\epsilon_3\sigma_{34} = \rho_{13}\rho_{23}$ establishes the claim for ρ_{23} , and the claim follows.

Our calculations enable us to deduce that diagrams for ρ_{21} and λ_{23} respectively are as follows



But ρ_{21} and λ_{23} commute, and this together with the fact that $\rho_{21}(E_1) = U = \lambda_{23}^{-1}(E_3)$ yields a diagram for ρ_{21} of the form



In particular Example 3.10 implies that $\dim V_0 \neq 0$.

Now let us define $A_2 = \rho_{21}(E_3) \leq V_3$. Note that $\dim A_2 = 1$. Since ρ_{21} commutes with λ_{23} , examining the respective diagrams yields $\lambda_{23}(A_2) = E_1$.

Now, observing that each ϵ_i preserves each subspace of E_I , we see that in fact for all i

$$A_2 = \rho_{2i}^{\pm 1}(E_j) = \lambda_{2i}^{\pm 1}(E_j)$$

where j satisfies $\{i, j\} = \{1, 3\}$. We can define A_1 and A_3 similarly.

The relations $[\rho_{21}, \rho_{31}] = [\rho_{23}, \rho_{13}] = 1$, together with the structure of our diagrams, tell us that $A_2 \cap (A_1 + A_3) = \{0\}$. The relation $[\rho_{32}, \rho_{12}] = 1$ informs us that $A_1 \cap A_3 = \{0\}$ and so that $\dim(A_1 \oplus A_2 \oplus A_3) = 3$. This is a contradiction, since $V_0 \neq \{0\}$ and so $\dim V_3 = 2$.


We have thus shown that ρ_{21} has a diagram as claimed. Observe that, since the subgroup $W_3 < \text{Out}(F_3)$ preserves each V_i by construction, having a diagram for ρ_{21} of the form described in the statement of this lemma immediately implies that at least one of $V_0 \oplus V_2$ and $V_1 \oplus V_3$ is preserved by $\text{Out}(F_3) = \langle W_3, \rho_{21} \rangle$. \square

Lemma 3.14. *Let V be a \mathbb{K} -linear, six-dimensional representation of $\text{Out}(F_3)$, where \mathbb{K} is a field of characteristic other than 2 or 3. Suppose that, with notation of Definition 3.3, $\dim V_1 \oplus V_2 = 6$. Then $V = V_1 \oplus V_2$ as an $\text{Out}(F_3)$ -module.*


Proof. If $\dim V_1 = 6$ or $\dim V_2 = 6$ then the result is trivial.

Suppose that $\dim V_1 = \dim V_2 = 3$ and so $V = V_1 \oplus V_2$ as a vector space. We know (using Maschke's Theorem and our assumption on $\text{char}(\mathbb{K})$) that each V_i (for $i = 1, 2$) is either a sum of standard and trivial or a sum of signed standard and determinant representations of S_3 ; we can therefore pick vectors $v_i \in E_i, w_i \in E_{\{1,2,3\} \setminus \{i\}}$ so that each $v_i - v_j$ and $w_i - w_j$ is an eigenvector of an element of $S_3 \setminus \{1\}$.


We have a diagram for ρ_{21} of the form




and analogously one for ρ_{31} of the form




Since S_4 commutes with Δ , its action has to preserve the $(+1)$ -eigenspace of Δ (which is equal to V_1 in our case) as well as the (-1) -eigenspace (which equals V_2 in this case). We also have $[\epsilon_1, \Delta] = 1$, and so $\epsilon_1 \sigma_{14} = \rho_{31} \rho_{21}$ preserves V_2 . Hence, evaluating $\rho_{31} \rho_{21}$ on $E_{1,2}$ (an observing that $\dim E_I \leq 1$ for all I) gives us either a diagram for ρ_{21} of the form



or a diagram for ρ_{31} of the form




Suppose (for a contradiction) that we are in the latter case. Evaluating $\rho_{21}\rho_{31}$ on E_1 (and observing that the diagrams for ρ_{31} and ρ_{21} are related by conjugation by σ_{23}) yields diagrams for ρ_{21} and ρ_{31} respectively of the form




Now $\rho_{21}\rho_{31}(E_1) = E_3$ and $\rho_{31}\rho_{21}(E_1) = E_2$. But ρ_{31} commutes with ρ_{21} , which yields a contradiction.


We can repeat the argument after evaluating $\rho_{31}\rho_{21}$ on E_3 and conclude that we have a diagram for ρ_{21} of the form




Two diagram chases, starting at E_3 and $E_{1,2}$, show $\rho_{21}^{-1} = \rho_{21}^{\epsilon_1}$ requires ρ_{21} to have a diagram of the form



Suppose we are in the third case. We have diagrams for ρ_{21} and ρ_{31} respectively




Evaluating $\epsilon_1\sigma_{14} = \rho_{31}\rho_{21}$ on E_1 (and observing that $\epsilon_1\sigma_{14}(V_2) = V_2$) yields a diagram for ρ_{21} of the form



The first case is impossible, since we could have

$$E_1 = 1(E_1) = \epsilon_1\rho_{21}\epsilon_1\rho_{21}(E_1) \leq E_2 \oplus E_{1,2}.$$


After repeating the argument for $E_{2,3}$ we conclude that we have diagrams for ρ_{21} and ρ_{31} respectively as follows



Suppose that σ_{14} preserves each E_i . Then so does $\sigma_{24} = \sigma_{14}^{\sigma_{12}}$. But $\sigma_{24} = \sigma_{12}^{\sigma_{14}}$, and $\sigma_{12}(E_1) = E_2$. This is a contradiction. We can apply an analogous argument to the σ_{14} -action on the subspaces $E_{i,j}$. Now we easily deduce from $\epsilon_1\sigma_{14} = \rho_{31}\rho_{21}$ that


$$\rho_{21}(E_1) \not\subseteq E_1 \text{ and } \rho_{21}(E_{2,3}) \not\subseteq E_{2,3}.$$

We can now evaluate $\epsilon_1\sigma_{14} = \rho_{31}\rho_{21}$ and $\rho_{21}^{-1} = \rho_{21}^{\epsilon_1}$ on E_1 and $E_{2,3}$ and conclude that we have a diagram for ρ_{21} of the form




which shows that both V_1 and V_2 are $\text{Out}(F_n)$ -invariant.


Suppose now that we are in one of the first two cases, namely that there is a diagram for ρ_{21} of the form




Verifying that $\rho_{31}\rho_{21}$ keeps V_1 and V_2 invariant immediately tells us that in fact we have a diagram for ρ_{21} of the form



The element ρ_{31} keeps E_2 and $E_{1,3}$ invariant, and so, observing that $\epsilon_1\sigma_{14} = \rho_{21}\rho_{31}$ preserves $V_1 \oplus V_3$, we actually have diagrams for ρ_{21} and ρ_{31} respectively



But, in order for $\rho_{21}\rho_{31}$ to keep V_1 and V_2 invariant, we need to have



as a diagram for ρ_{21} . This finishes the proof. \square


Now let us investigate 5-dimensional representations of $\text{Out}(F_3)$ – we hope to be able to say more in this case!

Proposition 3.15. *Let V be a 5-dimensional, \mathbb{K} -linear representation of $\text{Out}(F_3)$, where \mathbb{K} is a field of characteristic other than 2 or 3. Suppose that, with the notation of Definition 3.3, $V \neq V_0 \oplus V_3$. Then $V = V_0 \oplus V_1 \oplus V_2 \oplus V_3$ is a decomposition of $\text{Out}(F_3)$ -modules, and, as S_4 -modules, V_0 is a sum of trivial, V_1 of standard, V_2 of signed standard, and V_3 of determinant representations.*

Proof. Since $\dim V = 5$, we have $V_1 = \{0\}$ or $V_2 = \{0\}$. Let us suppose that we have the latter, the other case being entirely similar.

Step 0: We first claim that V_0 is a sum of trivial S_4 -modules.

Lemma 3.13 gives us two possibilities for a diagram for ρ_{21} , namely



The same lemma also tells us that $V/(V_1 \oplus V_3)$ is a representation of $\text{Out}(F_3)$. Its dimension is at most 2 and therefore Lemma 3.1 tells us that it is a direct sum of two trivial representations of $\text{Out}(F_3)$ (since we know how ϵ_1 acts), and so the same statement holds for $V/(V_1 \oplus V_3)$ as an S_4 -module. Hence it also holds for V_0 , since V_0 is an S_4 -module isomorphic to $V/(V_1 \oplus V_3)$.

Note that an identical argument shows that V_3 is a sum of determinant S_4 -modules in the case when $V_1 = \{0\}$.

Step 1: We now claim that $V_0 \oplus V_1$ is $\text{Out}(F_3)$ -invariant. Suppose for a contradiction that it is not the case.

Let U be the projection of $\rho_{21}(E_3)$ onto V_3 . Note that $\dim U = 1$ since we have assumed $V_0 \oplus V_1$ not to be $\text{Out}(F_3)$ -invariant. Our aim now is to show that U is $\text{Out}(F_3)$ -invariant.

If V_3 is $\text{Out}(F_3)$ -invariant, then it is an $\text{Out}(F_3)$ -module of dimension at most two, and hence we can use Lemma 3.5 to conclude that it is in fact a sum of determinant representations. Hence, in particular, U is $\text{Out}(F_3)$ -invariant.

Now suppose that V_3 is not $\text{Out}(F_3)$ -invariant. Checking that $\rho_{31}\rho_{21}(E_3) = \rho_{21}\rho_{31}(E_3)$ on both of our diagrams for ρ_{21} yields that $\rho_{31}(U) = U$. Note that U is the unique non-trivial invariant subspace of V_3 for both ρ_{21} and ρ_{31} , as otherwise V_3 would be invariant under the action of

$$\langle S_3, \rho_{21} \rangle = \langle S_3, \rho_{31} \rangle = \text{Out}(F_3).$$

Hence U is σ_{23} -invariant. But V_3 is a 2-dimensional S_3 -module, and if it were irreducible, then the trace of each transposition would be zero. Hence V_3 is a sum of two 1-dimensional modules of S_3 , and therefore S_3 preserves U . From this we conclude that $\text{Out}(F_3)$ preserves U .

Lemma 3.5 informs us that U is a determinant representation of $\text{Out}(F_3)$. Since $\rho_{21}^{-1} = \rho_{21}^{\epsilon_1}$, we must have


$$\forall v \in E_3 : \rho_{21}(v) \in v + U.$$

Using similar relations we establish that, when restricted to $E_3 \oplus U$, λ_{21} acts as ρ_{21} , and ρ_{12} acts as $\rho_{21}^{\pm 1}$. Hence, taking $v \in E_3$,

$$v + (2 \mp 1)u = \rho_{21}^{2 \mp 1}(v) = \rho_{21}\rho_{12}^{-1}\lambda_{21}(v) = \epsilon_1\sigma_{12}(v) \in E_3$$

where $u = \rho_{21}(v) - v \in U$. This shows that $u = 0$, and hence $V_0 \oplus V_1$ is $\text{Out}(F_3)$ -invariant, which is the desired contradiction.


We have thus shown that there is a diagram for ρ_{21} of the form



Step 2: We claim that V_1 is a standard S_4 -module.

As an S_3 -module, both V_1 and V_2 are sums of one standard and one either trivial or determinant representation. The branching rule tells us that therefore, as S_4 -modules, each of the subspaces can be either a standard or a signed standard representation, or the one corresponding to partition $(2, 2)$. The last case is ruled out by Lemma 3.1, since $(V_0 \oplus V_1)/V_0$ is clearly not a sum of trivial and determinant $\text{Out}(F_3)$ -modules.

Focusing only on V_1 , we have a diagram for ρ_{21} of the form



Note that in both cases these are the minimal diagrams for ρ_{21} when restricted to V_1 , since otherwise σ_{12} could not permute E_1 and E_2 .

Let us pick vectors $v_i \in E_i$ in such a way that each $v_i - v_j$ is an eigenvector of σ_{ij} . Let us also set $v = v_1 + v_2 + v_3$. The way in which S_4 acts on V_1 in our case is determined by one parameter; we can calculate it by finding $\mu \in \mathbb{C}$ such that $v_1 + \mu v$ is an eigenvector of σ_{14} . The eigenvalue of this eigenvector will also determine the way in which S_4 acts. Let us note that we can also find this parameter μ by computing $\sigma_{14}(v_2 - v_1) = \mu v + v_2$.

In the case of the first diagram for ρ_{21} , we immediately see that

$$\sigma_{14}(v_2 - v_1) = \epsilon_1\rho_{21}\rho_{31}(v_2 - v_1) \in E_1 \oplus E_2,$$

and hence $\mu = 0$. Now both ρ_{23} and ρ_{31} preserve E_1 , and so observing that $\rho_{21}^{-1} = [\rho_{23}^{-1}, \rho_{31}^{-1}]$ yields that ρ_{21} acts trivially on E_1 . By an analogous argument so does ρ_{31} . Hence $\sigma_{14}(v_1) = \epsilon_1(v_1)$. In our case this shows that we are dealing with a standard representation; if however V_1 is trivial, ϵ_1 acts as plus one on the appropriate vector, and we see that V_2 is a signed standard S_4 -module.

In the case of the second diagram we immediately see two eigenspaces of σ_{14} , namely E_2 and E_3 . These spaces are interchanged by the action of σ_{23} which


commutes with σ_{14} , and hence must have the same eigenvalue. In a standard or a signed standard representation of S_4 each σ_{ij} has always exactly two repeated eigenvalues, and it is this eigenvalue that determines the representation. It is enough for us then to find a third eigenvector of σ_{14} and compute its eigenvalue. The vector must have a non-trivial E_1 -component, and our diagram tells us that it is enough to check how σ_{14} acts on E_1 . By an argument similar to the one above we show that $\epsilon_1\sigma_{14}(v_1) \in v_1 + E_2 \oplus E_3$, and the claim follows.

Step 3: We now claim that V_3 is a sum of determinant S_4 -representations.


As an S_3 -module, $V_1 \oplus V_3$ is a sum of one standard, one trivial and some number of determinant representations (depending on the dimension of V_3). We have already found one standard representation of S_4 , and the branching rule tells us that there can only be determinant representations of S_4 left. If at least one of them does not lie entirely in V_3 , then it would appear in $(V_1 \oplus V_3)/V_3$ by Schur's Lemma. This is not possible, since $(V_1 \oplus V_3)/V_3$ is a standard representation of S_4 . Hence all the other irreducible S_4 -modules lie within V_3 .

Step 4: Our last claim is that each V_i is $\text{Out}(F_3)$ -invariant.

We have already shown this for $V_1 \oplus V_0$. We have just shown that V_3 is S_4 invariant, and so, $\rho_{21}\rho_{31} = \epsilon_1\sigma_{14}$ keeping V_3 invariant yields a diagram for ρ_{21} of the form



We have already shown in step 2 that in both cases $\rho_{21}(v) \in v + E_2 \oplus V_0$ for each $v \in E_1$. Also, $(V_0 \oplus V_1)/V_1$ is an $\text{Out}(F_3)$ -module of dimension at most 2, and hence is described by Lemma 3.5. In particular $\rho_{12}(w) = w + E_1$ for all $w \in V_0$. Analogous statements hold for ρ_{31} and so observing that σ_{23} acts as ± 1 on $E_1 \oplus V_0$ and that $\rho_{21} = \rho_{31}^{\sigma_{23}}$ yields that $\epsilon_1\sigma_{12} = \rho_{31}\rho_{21}(V_0)$ has a non-trivial V_1 -component if and only if $\rho_{21}(V_0)$ does, and similarly that $\epsilon_1\sigma_{12} = \rho_{31}\rho_{21}(V_1)$ has a non-trivial V_0 -component if and only if $\rho_{21}(V_1)$ does. Hence we have a diagram



for ρ_{21} , which was what we claimed. \square

Lemma 3.16. *Suppose V is a \mathbb{K} -linear representation of $\text{Out}(F_3)$, such that, using notation of Definition 3.3, $V_0 \oplus V_2$ and $V_1 \oplus V_3$ are $\text{Out}(F_3)$ -invariant.*

Then the representation factors through the natural projection

$$\pi : \text{Out}(F_3) \rightarrow \text{GL}_3(\mathbb{Z}).$$

Proof. Note that $\phi(\Delta)$ lies in the product

$$Z(\text{GL}(V_0 \oplus V_2)) \times Z(\text{GL}(V_1 \oplus V_3))$$

of the centres of the general linear groups of the components $V_0 \oplus V_2$ and $V_1 \oplus V_3$. Therefore we have $\phi(\rho_{ij}) = \phi(\rho_{ij})^{\phi(\Delta)} = \phi(\lambda_{ij})$ for each $i \neq j$, and so ϕ factors as

$$\begin{array}{ccc} \text{Out}(F_3) & \xrightarrow{\phi} & \text{GL}(V) \\ \downarrow & & \uparrow \\ \text{Out}(F_3) / \langle\langle \{\rho_{ij} \lambda_{ij}^{-1} \mid i \neq j\} \rangle\rangle & \xrightarrow{\cong} & \text{GL}_3(\mathbb{Z}) \end{array}$$

This finishes the proof. \square

Observe an immediate consequence of the above.

Lemma 3.17. *Suppose V is a \mathbb{K} -linear representation of $\text{Out}(F_3)$ of dimension at most 5, where the characteristic of \mathbb{K} is not 2 or 3. Then the representation factors through the natural projection $\pi : \text{Out}(F_3) \rightarrow \text{GL}_3(\mathbb{Z})$.*

Proof. Using the notation of Definition 3.3, we have $V = V_0 \oplus V_1 \oplus V_2 \oplus V_3$ as a vector space. Suppose first that $V_1 \oplus V_2$ is trivial. Then Lemma 3.6 tells us that $V = V_0 \oplus V_3$ as an $\text{Out}(F_3)$ -module.

Supposing that $V_1 \oplus V_2 \neq \{0\}$ allows us to use Proposition 3.15, and conclude that each V_i is $\text{Out}(F_3)$ -invariant. We can now use Lemma 3.16 and finish the proof. \square

Before proceeding further we need to recall a standard fact of representations theory.

Proposition 3.18. *Let A be the kernel of the map $\text{SL}_3(\mathbb{Z}) \rightarrow \text{SL}_3(\mathbb{Z}_2)$ induced by the surjection $\mathbb{Z} \rightarrow \mathbb{Z}_2$. Let V be the standard, 3-dimensional \mathbb{K} -linear representation of $\text{GL}_3(\mathbb{Z})$. Suppose further that \mathbb{K} is a field of characteristic 0 or at least 3. Then $U = \text{Sym}^2(V^*)$, the second symmetric power of the dual module of V , is irreducible as an A -module.*

Proof. Let $U \leq V$ be an irreducible A -submodule of V , and let $\{v_1, v_2, v_3\}$ be the standard basis of V . Suppose $v \in U \setminus \{0\}$. Then

$$v = \sum_{i \leq j} \mu_{ij} v_i \otimes v_j$$

for some collection of scalars μ_{ij} .

We are going to abuse notation by using the symbols ϵ_i and ρ_{ij} to denote the images of respective elements under $\pi : \text{Out}(F_3) \rightarrow \text{GL}_3(\mathbb{Z})$. Note that $\epsilon_i \epsilon_j \in A$ and $\rho_{ij}^2 \in A$ for each appropriate $i \neq j$. Now

$$\epsilon_1 \epsilon_2(v) - v = -2 \mu_{23} v_2 \otimes v_3 - 2 \mu_{13} v_1 \otimes v_3$$

and hence

$$\epsilon_1 \epsilon_3 (\epsilon_1 \epsilon_2 (v) - v) - v = 4 \mu_{13} v_1 \otimes v_3.$$

Hence, if $\mu_{ij} \neq 0$ for some $i \neq j$, then $v_i \otimes v_j \in U$.

Furthermore

$$\rho_{13}^2(v_1 \otimes v_3) - v_1 \otimes v_3 = -2 v_1 \otimes v_1$$

and

$$\rho_{23}^2(v_1 \otimes v_3) - v_1 \otimes v_3 = -2 v_1 \otimes v_2$$

and therefore if $\mu_{ij} \neq 0$ for some $i \neq j$, then $U = V$.

Suppose that

$$v = \sum_i \mu_{ii} v_i \otimes v_i$$

Without loss of generality let us assume that $\mu_{11} \neq 0$. Then

$$\rho_{21}^2(v) - v = -\mu_{11} (2 v_1 \otimes v_2 - 4 v_2 \otimes v_2) = v' \in U$$

We can now apply our argument to v' and conclude that $U = V$. \square

We are now ready for the main result of this section.

Theorem 3.19. *Suppose V is a \mathbb{K} -linear representation of $\text{Out}(F_3)$ of dimension at most 6, where the characteristic of \mathbb{K} is not 2 or 3. Then the representation factors through the natural projection $\pi : \text{Out}(F_3) \rightarrow \text{GL}_3(\mathbb{Z})$.*

Proof. Let $\phi : \text{Out}(F_3) \rightarrow \text{GL}(V)$ be our representation. Using the notation of Definition 3.3, we have $V = V_0 \oplus V_1 \oplus V_2 \oplus V_3$ as a vector space. We need to consider a number of cases.

Suppose first that $V_1 \oplus V_2$ is trivial. Then Lemma 3.6 tells us that $V = V_0 \oplus V_3$ as an $\text{Out}(F_3)$ -module. Suppose now that $V_0 \oplus V_3$ is trivial. Lemma 3.14 tells us that $V = V_1 \oplus V_2$ as an $\text{Out}(F_3)$ -module. In both situations we can apply Lemma 3.16.

We are left with the most general case: suppose that $\dim V_1 \oplus V_2 = 3$. We are going to assume that in fact $V_2 = \{0\}$, the other case being analogous. Applying Lemma 3.13 gives us two $\text{Out}(F_3)$ -representations $r : \text{Out}(F_3) \rightarrow V/(V_1 \oplus V_3)$ and $s : \text{Out}(F_3) \rightarrow V/V_0$, where at least one of them occurs as a submodule of V . Also, r and s factor through π by Lemma 3.16. If any of these representations has dimension 0, then we are done. In what follows we shall suppose that the dimension of both r and s is non-zero, and thus that V is reducible as an $\text{Out}(F_3)$ -module. We can choose a basis for V so that the matrices in $\phi(\text{Out}(F_3))$ are all in a block-upper-triangular form, with diagonal blocks corresponding to representations r and s .

Let $\overline{\text{IA}}_3 = \ker \pi$ be the Torelli subgroup. Our aim is to show that $\overline{\text{IA}}_3$ lies in the kernel of ϕ .

Elements in $\overline{\text{IA}}_3$ map to matrices with identities on the diagonal, and all non-zero entries located in the block in the top-right corner. Hence $\overline{\text{IA}}_3$ maps to an abelian group isomorphic to \mathbb{K}^m , where $m \in \{5, 8, 9\}$ depends on the dimension of r .

Note that all products $\epsilon_i \epsilon_j$ lie in the kernel of r , and hence so do all elements

$$\rho_{kj}^2 = (\rho_{kj}^{\epsilon_i \epsilon_j} \rho_{kj}^{-1})^{-1},$$

where we took $k \neq i$. The work of Mennicke [14] now shows that in fact r factors through a finite group: when restricted to $\text{SOut}(F_3) = \pi^{-1}(\text{SL}_3(\mathbb{Z}))$, it factors through

$$\text{SOut}(F_3) \rightarrow \text{SL}_3(\mathbb{Z}) \rightarrow \text{SL}_2(\mathbb{Z}_2).$$

Let A denote the kernel of this map. Note that $\overline{\text{IA}}_3 < A$.

We have shown above that $r|_A$ is trivial, and so A maps to the identity matrix in the block corresponding to r . Note that $\phi(A)$ acts by conjugation on the abelian group of matrices with identity blocks on the diagonal, and a trivial block in the bottom-left corner. As remarked above, this group is isomorphic to \mathbb{K}^m . Each row or column (depending on which diagonal block corresponds to r) in the top-right corner corresponds to an A -submodule, and so the group \mathbb{K}^m splits as an A -module into

$$\mathbb{K}^5, 2.\mathbb{K}^4, 3.\mathbb{K}^3, 4.\mathbb{K}^2 \text{ or } 5.\mathbb{K},$$

depending on the dimension of r , where the multiplicative notation indicates the number of direct summands.

Let $T = \overline{\text{IA}}_3 / [\overline{\text{IA}}_3, \overline{\text{IA}}_3]$ denote the abelianisation of the Torelli group seen as an $\text{Out}(F_3) / \overline{\text{IA}}_3 = \text{GL}_3(\mathbb{Z})$ -module, where the action is the one induced by the conjugation action $\text{Out}(F_3) \curvearrowright \overline{\text{IA}}_3$. The structure of this module is known (see Kawazumi [10]) – it is the second symmetric power of the dual of the standard $\text{GL}_3(\mathbb{Z})$ -module. After tensoring T with \mathbb{K} , we can apply Proposition 3.18, and conclude that $T \otimes_{\mathbb{Z}} \mathbb{K}$ is an irreducible A -module of dimension 6. By Schur's Lemma, if we have an A -equivariant quotient of T , it is either isomorphic to T or equal to $\{0\}$.

Now consider the action of $\phi(A)$ on $\phi(\overline{\text{IA}}_3) \otimes_{\mathbb{Z}} \mathbb{K}$ by conjugation. It is at the same time an equivariant quotient of an irreducible 6-dimensional module and a submodule of

$$\mathbb{K}^5, 2.\mathbb{K}^4, 3.\mathbb{K}^3, 4.\mathbb{K}^2 \text{ or } 5.\mathbb{K}.$$

This implies that the image of $\overline{\text{IA}}_3$ under ϕ is trivial. This finishes the proof. \square

4 Small graphs with transitive automorphism groups

In this section we will establish some lemmata concerning graphs of rank at most 5 with groups W_3 and G_3 acting on them.

Definition 4.1 (Admissible graphs). Let X be a connected graph with no vertices of valence 2, and suppose we have a group G acting on it. We say that X is *G-admissible* if and only if there is no G -invariant non-trivial (i.e. with at least one edge) forest in X . We also say that X is *admissible* if and only if it is $\text{Aut}(X)$ -admissible.

The following theorem is due to Marc Culler [7], Dmitri Khramtsov [11] and Bruno Zimmermann [17] (each independently).

Theorem 4.2 (Culler [7]; Khramtsov [11]; Zimmermann [17]). *Let $n \geq 2$. Suppose*

$$G \hookrightarrow \text{Out}(F_n)$$

is an embedding of a finite group G . Then there exists a finite G -admissible graph X of rank n (with a fixed isomorphism $\pi_1 X \cong F_n$), so that the composition

$$G \rightarrow \text{Aut}(X) \rightarrow \text{Out}(F_n)$$

is the given embedding.

The reason for requiring $n \geq 2$ in the above theorem is that there are no admissible graphs of rank 1 (i.e. with the fundamental group \mathbb{Z}). Of course any finite subgroup of $\text{Out}(F_1) \cong \mathbb{Z}/2\mathbb{Z}$ can be realised as an action on a graph with one vertex and one edge.

Since we will be dealing with homology of finite graphs quite frequently in this section, let us observe the following.

Lemma 4.3. *Let X be a finite, oriented graph. Recall that Definition 2.1 gives us two maps $\iota, \tau : E(X) \rightarrow V(X)$. We have the following identification*

$$H_1(X, \mathbb{C}) \cong \left\{ f : E(X) \rightarrow \mathbb{C} \mid \forall v \in V(X) : \sum_{\iota(e)=v} f(e) = \sum_{\tau(e)=v} f(e) \right\}.$$

We will often refer to each such function f as a *choice of weights* of edges in X .

Before proceeding any further, we need to introduce a concept of *collapsing maps* of graphs.

Definition 4.4 (Collapsing map). Let $\pi : X \rightarrow X'$ be a surjective morphism of graphs X and X' . We say that π is a *collapsing map* if and only if the preimages of points in X' under π are path-connected.

Remark 4.5. Let us observe three facts:

1. for a given graph X , giving a subset of $E(X)$ which will be collapsed specifies a collapsing map π ;
2. any collapsing map $\pi : X \rightarrow X'$ induces a surjective map on homology by pushing forward weights of edges which it does not collapse. We will often abuse notation and refer to this induced map as π ;
3. if $\pi : X \rightarrow X'$ is a G -equivariant collapsing map (where G is a group acting on X and X'), then the induced map on homology is also G -equivariant.

It is now time to focus on the main area of our interest here.

Definition 4.6. Let G be a group acting on a graph X , and let e be an edge of X . We define X_e to be the graph obtained from X by collapsing all edges *not contained* in the G -orbit of e .

Note that the action of G on such an X_e is edge-transitive.

Lemma 4.7. *Suppose X is a G -admissible graph of rank at most 5, where G is a group, and e is any edge of X . Then X_e has no vertices of valence 1 or greater than 10 and satisfies*

$$8 \geq 2v_2 + v_3 + 2v_4 + 3v_5 + 4v_6 + 5v_7 + 6v_8 + 7v_9 + 8v_{10} \quad (*)$$

where v_i is the number of vertices of valence i in X_e .

Proof. First note that there are no vertices of valence 1 in X_e , since they could only occur if there were separating edges in X . But X is admissible, and so there are no such edges.

A simple Euler characteristic count yields

$$2(\text{rank}(X_e) - 1) \geq \sum_{i=3}^{\infty} (i-2)v_i$$

and hence in particular $v_i = 0$ for all $i > 10$, as X_e has rank at most 5.

Since X is admissible, each vertex of X_e of valence two comes from collapsing a subgraph of X which is not a tree, hence

$$\text{rank}(X_e) \leq 5 - v_2$$

and the result follows. \square

We will now consider graphs satisfying (*) with a transitive action of $W_3 \cong G_3$ yielding particular representations on the \mathbb{C} -homology of the graph.

Proposition 4.8. *Let X be a graph of rank 5 on which $G \in \{W_3, G_3\}$ acts so that the representation of G on $V = H_1(X, \mathbb{C})$ induced by the action decomposes as $V = V_0 \oplus V_1 \oplus V_2 \oplus V_3$, where*

- *if $G = W_3$ then the decomposition is the one described in Definition 3.3, and, as S_3 -modules, V_1 is a permutation, V_2 signed permutation, V_0 a sum of trivial and V_3 a sum of determinant representations;*
- *if $G = G_3$ then Δ acts as identity on $V_0 \oplus V_2$ and as minus the identity on $V_1 \oplus V_3$, and as S_4 -modules, V_1 is a standard, V_2 signed standard, V_0 a sum of trivial and V_3 a sum of determinant representations.*

Then, there is a subgraph $Y \leq X$ isomorphic to a 3-rose, on which G acts in such a way that, as an S_3 -module (where $S_3 < W_3 \cap G_3$), $H_1(Y, \mathbb{C})$ contains the standard representation.

Proof. Let $v \in V$ be a vector belonging to a standard representation of $S_3 < G$. It is represented by a choice of weights on edges of X . Let e be an edge with a non-zero weight. Then the image of v in $H_1(X_e, \mathbb{C})$ is non-trivial, and hence Schur's Lemma informs us that $H_1(X_e, \mathbb{C})$ contains a standard S_3 -module.

Let $Z = X_e$. Lemma 4.7 tells us that Z satisfies (*). Also, since G acts transitively on edges of Z , there are at most two vertex-orbits of this action, and hence in particular at most two values v_i can be non-zero. Let us list all possible values of v_i , noting that $iv_i = jv_j$ if there are vertices of valence i and j in Z , and that v_i must be even if i is odd and there are only vertices of valence i in Z . All possible cases are summarised in Figure 4.1.

Now, in order to have a standard representation of S_3 , we need at least 3 edges in Z , and the rank of Z has to be at least 2. We can therefore immediately rule out cases (1), (3), (5), (6) and (15). Also, since the action of G on the edges of Z is transitive, their number has to divide $|G| = 48$. Hence we can additionally rule out cases (8), (16) and (20). We are left with the cases listed in Figure 4.2.

We will need to deal with these cases one by one:

Case (2): Here we have three vertices of valence two, on which S_3 has to act transitively. Each of these comes from collapsing a graph of non-zero rank

Figure 4.1: Case table

Case number	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_{10}	edges	rank
(1)	4									4	1
(2)	3	2								6	2
(3)	3									3	1
(4)	2		1							4	2
(5)	2									2	1
(6)	1									1	1
(7)		8								12	5
(8)		6								9	4
(9)		4								6	3
(10)		2			1					6	4
(11)		2								3	2
(12)			4							8	5
(13)			3							6	4
(14)			2							4	3
(15)			1							2	2
(16)				2						5	4
(17)					2					6	5
(18)					1					3	3
(19)							1			4	4
(20)									1	5	5

Figure 4.2: Reduced case table

Case number	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_{10}	edges	rank
(2)	3	2								6	2
(4)	2		1							4	2
(7)		8								12	5
(9)		4								6	3
(10)		2			1					6	4
(11)		2								3	2
(12)			4							8	5
(13)			3							6	4
(14)			2							4	3
(17)					2					6	5
(18)					1					3	3
(19)							1			4	4

in X , hence the sum of homologies of these graphs contains another standard representation of S_3 . This contradicts our assumptions.

Case (4): Here Z is a subdivided 2-rose, so we cannot get a standard representation of S_3 on the homology of this graph.

Case (7): There are four graphs with an edge-transitive group action with at most 8 vertices each of valence 3, namely a 3-cage, the 1-skeleton of a tetrahedron, the complete bipartite graph $K(3,3)$, and the 1-skeleton of a cube. Clearly, only the last one has the required number of vertices. An S_3 -action yielding a standard representation has to be the one given by fixing two vertices and permuting 3 edges incident at one of them in a natural way. This however yields two copies of the standard representation when acting on homology, which contradicts our assumptions.

Case (9): In this case we are dealing with an edge-transitive G -action on 1-skeleton of a tetrahedron. Such an action has to also be vertex-transitive, but S_3 cannot act transitively on 4 points.

Case (10): In this case Z is obtained by taking a wedge of two 3-cage graphs, C_1 and C_2 , say. Since the action of G is edge-transitive, for any edge e in Z we have

$$|\text{Stab}_G(e)| = 8$$

and hence each 3-cycle acts freely. Since a 3-cycle cannot swap C_1 and C_2 , it must act in the natural way on edges of both. Hence there have to be two copies of the standard representation of S_3 in $H_1(Z, \mathbb{C})$, which is not the case.

Case (11): In this case Z is a 3-cage. If $G = G_3$, then $S_4 < G_3$ cannot act on Z yielding the desired standard or signed standard representation. Suppose now that $G = W_3$. If ϵ_1 preserves exactly one edge, then so do ϵ_2 and ϵ_3 ; these edges are distinct, as otherwise we would have some ϵ_i and ϵ_j acting in the same way where $i \neq j$, and so $H_1(Z, \mathbb{C}) \leq V_0 \oplus V_3$, where S_3 cannot have a standard representation. Since the edges are distinct, ϵ_1 and ϵ_2 cannot commute. This shows that each ϵ_i preserves all edges of Z , and hence $H_1(Z, \mathbb{C}) \leq V_0 \oplus V_3$, which is a contradiction.

Case (12): The graph Z is a bipartite graph with 4 vertices and exactly zero or two edges connecting each pair of vertices. Hence Z admits a G -equivariant quotient map to a square (i.e. a single cycle made of 4 edges) which is a 2-to-1 map on edges. Each 3-cycle acts trivially on the square; moreover it cannot act non-trivially on the preimages of edges of the square. We conclude that each 3-cycle acts trivially, which is a contradiction.

Case (13): We easily check that the graph Z consists of three vertices, each of which has exactly two edges connecting it to each of the other two. Since the 3-cycle in S_3 acts non-trivially, it has to act transitively on vertices, and so either each vertex in Z comes from collapsing a subgraph which was not a tree in X , or none of them does. Neither of these two cases is possible, since the rank of X is 5.

Case (14): In this case Z is a 4-cage. Each edge in Z has a corresponding edge in X , and the fact that X is G -admissible implies that these edges do not form a forest. Hence they can form either a single simple loop, a pair of simple loops, or a 4-cage in X . The first two cases are impossible, since they would yield a trivial action of the 3-cycle of S_3 on Z . Hence X contains Z as a subgraph.

Our assumption on the representations of G tells us that either Δ or each transposition in S_3 has to flip Z , and so X is a 4-cage with a loop of length one

attached to each vertex. Let a and b be two vectors in $H_1(X, \mathbb{C})$, each given by putting a weight 1 on exactly one of the loops.

If Δ flips the graph, then $a+b$ and $a-b$ span two one-dimensional eigenspaces of Δ , one with eigenvalue $+1$, and one with eigenvalue -1 . Hence transpositions in S_3 have to map one of these vectors to itself, and the other to minus itself; this is only possible if they flip the graph, which contradicts our assumptions.

A similar argument works if the transpositions in S_3 flip X .

Case (17): In this case $Z = X$ is a 6-cage. As before we have

$$|\text{Stab}_G(e)| = 8$$

for any edge e in Z . Hence each 3-cycle in S_3 acts freely and so we have two copies of the standard S_3 -representation, which is a contradiction.

Case (18): In this case Z is a 3-rose. If Z is actually a subgraph of X , then we are done. Suppose it is not.

As Z only has one vertex, there is a connected subgraph X' of X that we collapsed when constructing Z . Since X' is of rank 2, after erasing vertices of valence 2 (in X'), we are left with two cases: a 2-rose or a 3-cage. Since Z is not a subgraph of X , and the preimages of edges of Z in X cannot form a forest, they either form a simple loop (of length three), or a disjoint union of three loops (each of length one). In any event, we have three vertices on which the 3-cycle in S_3 acts transitively. Hence X' has to be a 3-cage, with the three vertices lying on the three edges of the cage. But then we get two standard representations of S_3 inside the homology of X , which is a contradiction.

Case (19): In this case Z is a 4-rose. The 3-cycle in S_3 acts by permuting three petals, and fixing one; let us call this fixed edge f . We easily check that f is preserved setwise by S_3 , and hence also by Δ , since Δ commutes with S_3 .

If $G = W_3$ then the one-dimensional subspace in $H_1(Z, \mathbb{C})$ spanned by a vector corresponding to f is contained either in V_0 or in V_3 , and hence f has to be preserved by all elements in G . This contradicts transitivity of the action of G on Z .

Suppose $G = G_3$. Note that there is only one way (up to isomorphism) in which S_4 can act on a set of four elements transitively. Therefore, as Δ commutes with S_4 , Δ acts as plus or minus the identity on $H_1(Z, \mathbb{C})$. Now $H_1(Z, \mathbb{C})$ as an S_4 -representation is a sum of standard and trivial or signed standard and determinant representations. In particular, our hypothesis tells us that Δ cannot act as either plus or minus the identity. This is a contradiction. \square

Lemma 4.9. *Suppose $\phi : \text{Out}(F_3) \rightarrow \text{Out}(F_5)$ is a homomorphism. Let*

$$\psi : \text{Out}(F_5) \rightarrow \text{GL}(V) \cong \text{GL}_5(\mathbb{C})$$

be the natural map, where $V = H_1(F_5, \mathbb{C})$. Note that $\psi \circ \phi$ gives a representation of $\text{Out}(F_3)$ on V . If, as a W_n -module, V splits as $V_0 \oplus V_3$ (with the notation of Definition 3.3) then the image of ϕ is finite.

Proof. The fact that $V = V_0 \oplus V_3$ as a W_n -module implies that $\psi \circ \phi(\epsilon_i \Delta) = 1$ for each i . Now a result of Baumslag–Taylor [2] tells us that the kernel of ψ is torsion-free, and so $\phi(\epsilon_i \Delta) = 1$. But this means that we have the following

commutative diagram:

$$\begin{array}{ccc}
\text{Out}(F_3) & \xrightarrow{\phi} & \text{Out}(F_5) \\
\downarrow & \nearrow & \uparrow \phi' \\
\text{Out}(F_3)/\langle\langle \{\rho_{ij} = \lambda_{ij} : i \neq j\} \rangle\rangle & \xrightarrow{\cong} & \text{GL}_3(\mathbb{Z})
\end{array}$$

This allows us to use a result of Bridson and Farb [4], who have shown that such a ϕ' necessarily has finite image. Therefore the image of ϕ is finite. \square

Theorem 4.10. *Suppose $\phi : \text{Out}(F_3) \rightarrow \text{Out}(F_5)$ is a homomorphism. Then the image of ϕ is finite.*

Proof. Consider the natural map $\psi : \text{Out}(F_5) \rightarrow \text{GL}_5(\mathbb{C}) \cong \text{GL}(V)$ as above. Again as above, the composition $\eta = \phi \circ \psi$ gives us a 5-dimensional complex linear representation $\eta : \text{Out}(F_3) \rightarrow \text{GL}_5(\mathbb{C})$.

Suppose first that, with the notation of Definition 3.3, V satisfies

$$V = V_0 \oplus V_3$$

Then Lemma 4.9 yields the result.

Now suppose that $V = V_0 \oplus V_1 \oplus V_2 \oplus V_3$ where $V_1 \oplus V_2 \neq \{0\}$. We apply Proposition 3.15 to η .

We will now apply Theorem 4.2 to two finite subgroups of $\text{Out}(F_5)$, namely $\phi(G_3)$ and $\phi(W_3)$, to obtain two graphs X and Y respectively, on which the groups G_3 and W_3 act. Note that $H_1(X, \mathbb{C}) \cong H_1(Y, \mathbb{C}) \cong V$, and the representations of G_3 and W_3 induced by the actions of the groups on homology of the respective graphs are isomorphic to the ones given by restricting η . Hence the conclusions of Proposition 3.15 apply to these representations, and so we can apply Proposition 4.8 to the actions $G_3 \curvearrowright X$ and $W_3 \curvearrowright Y$.

We conclude that both X and Y have a subgraph, preserved by the action of the respective group, isomorphic to a 3-rose. We also know that we can label the petals as e_1, e_2, e_3 , so that S_3 acts on this rose by permuting petals in the natural way, with the transpositions potentially also flipping all petals.

Knowing that $V_1 \oplus V_2 \neq \{0\}$ implies that in the W_3 case, either each ϵ_i flips e_i and leaves the other petals fixed, or each ϵ_i fixes e_i and flips the other petals. In the G_3 case, we see that there is only one way in which S_4 can act on the 3-rose inducing a standard or a signed standard representation. Each σ_{i4} has to interchange the two petals with labels different than e_i and preserve the third one; additionally, it either flips e_i and keeps some orientation of the other two fixed, or it flips the other two and fixes e_i . These two cases depend on the action of σ_{ij} for $i, j \leq 3$.

In any case, we have

$$\phi(\sigma_{14}) = \phi(\sigma_{23}\epsilon_2\epsilon_3)$$

and so

$$\phi(\lambda_{21}) = \phi(\lambda_{21}^{\sigma_{14}\epsilon_1}) = \phi(\lambda_{21})^{\phi(\Delta\sigma_{23})} = \phi(\rho_{31})$$

Therefore

$$1 = \phi([\rho_{23}^{-1}, \lambda_{21}^{-1}]) = \phi([\rho_{23}^{-1}, \rho_{31}^{-1}]) = \phi(\rho_{21})^{-1}$$

It follows that all Nielsen moves (which generate an index 2 subgroup of $\text{Out}(F_3)$) lie in the kernel of ϕ , and so the image of ϕ is of size at most 2, determined by $\phi(\epsilon_1)$. \square

References

- [1] J. Aramayona, C. J. Leininger, and J. Souto. Injections of mapping class groups. *Geom. Topol.*, 13(5):2523–2541, 2009.
- [2] G. Baumslag and T. Taylor. The centre of groups with one defining relator. *Math. Ann.*, 175:315–319, 1968.
- [3] O. Bogopol’skiĭ and D. Puga. On embeddings of $\text{Out}(F_n)$, the outer automorphism group of the free group of rank n , into $\text{Out}(F_m)$ for $m > n$. *Algebra and Logic*, 41(2):69–73, 2002.
- [4] M. R. Bridson and B. Farb. A remark about actions of lattices on free groups. *Topology Appl.*, 110(1):21–24, 2001.
- [5] M. R. Bridson and K. Vogtmann. Homomorphisms from automorphism groups of free groups. *Bull. London Math. Soc.*, 35(6):785–792, 2003.
- [6] M. R. Bridson and K. Vogtmann. Abelian covers of graphs and maps between outer automorphism groups of free groups. *Mathematische Annalen*, pages 1–34, 2011.
- [7] M. Culler. Finite groups of outer automorphisms of a free group. In *Contributions to group theory*, volume 33 of *Contemp. Math.*, pages 197–207. Amer. Math. Soc., Providence, RI, 1984.
- [8] S. M. Gersten. A presentation for the special automorphism group of a free group. *J. Pure Appl. Algebra*, 33(3):269–279, 1984.
- [9] F. Grunewald and A. Lubotzky. Linear representations of the automorphism group of a free group. *Geometric and Functional Analysis*, 18(5):1564–1608, June 2006.
- [10] N. Kawazumi. Cohomological aspects of magnus expansions. *arXiv:0505497*.
- [11] D. G. Khramtsov. Finite groups of automorphisms of free groups. *Mat. Zametki*, 38(3):386–392, 476, 1985.
- [12] D. G. Khramtsov. Outer automorphisms of free groups. In *Group-theoretic investigations (Russian)*, pages 95–127. Akad. Nauk SSSR Ural. Otdel., Sverdlovsk, 1990.
- [13] D. Kielak. Outer automorphism groups of free groups: linear and free representations. *J. London Math. Soc.*, 87(3):917–942, 2013.
- [14] J. L. Mennicke. Finite factor groups of the unimodular group. *Ann. of Math. (2)*, 81:31–37, 1965.
- [15] A. Potapchik and A. Rapinchuk. Low-dimensional linear representations of $\text{Aut } F_n$, $n \geq 3$. *Trans. Amer. Math. Soc.*, 352(3):1437–1451, 2000.
- [16] V. Turchin and T. Willwacher. Hochschild–Pirashvili homology on suspensions and representations of $\text{Out}(F_n)$. *In preparation*.
- [17] B. Zimmermann. Finite groups of outer automorphisms of free groups. *Glasgow Math. J.*, 38(3):275–282, 1996.