

# ZERO-DIMENSIONAL SYMPLECTIC ISOLATED COMPLETE INTERSECTION SINGULARITIES

WOJCIECH DOMITRZ

ABSTRACT. We study the local symplectic algebra of the 0-dimensional isolated complete intersection singularities. We use the method of algebraic restrictions to classify these symplectic singularities. We show that there are non-trivial symplectic invariants in this classification.

## 1. INTRODUCTION

The problem of symplectic classification of singular varieties was introduced by V. I. Arnold in [A1]. Arnold showed that the  $A_{2k}$  singularity of a planar curve (the orbit with respect to the standard  $\mathcal{A}$ -equivalence of parameterized curves) split into exactly  $2k + 1$  symplectic singularities (orbits with respect to the symplectic equivalence of parameterized curves). He distinguished different symplectic singularities by different orders of tangency of the parameterized curve with the *nearest* smooth Lagrangian submanifold. Arnold posed a problem of expressing these new symplectic invariants in terms of the local algebra's interaction with the symplectic structure and he proposed to call this interaction the **local symplectic algebra**. This problem was studied by many authors mainly in the case of singular curves.

In [IJ1] G. Ishikawa and S. Janeczko classified symplectic singularities of curves in the 2-dimensional symplectic space. A symplectic form on a 2-dimensional manifold is a special case of a volume form on a smooth manifold. The generalization of results in [IJ1] to volume-preserving classification of singular varieties and maps in arbitrary dimensions was obtained in [DR]. The orbit of the action of all diffeomorphism-germs agrees with the volume-preserving orbit in the  $\mathbb{C}$ -analytic category for germs which satisfy a special weak form of quasi-homogeneity e.g. the weak quasi-homogeneity of varieties is a quasi-homogeneity with non-negative weights  $\lambda_i \geq 0$  and  $\sum_i \lambda_i > 0$ .

P. A. Kolgushkin classified stably simple symplectic singularities of parameterized curves in the  $\mathbb{C}$ -analytic category ([K]).

In [DJZ2] the local symplectic algebra of singular quasi-homogeneous subsets of a symplectic space was explained by the algebraic restrictions of the symplectic form to these subsets. The generalization of the Darboux-Givental theorem ([AG]) to germs of arbitrary subsets of the symplectic space obtained in [DJZ2] reduces the problem of symplectic classification of germs of quasi-homogeneous subsets to the

---

1991 *Mathematics Subject Classification*. Primary 53D05. Secondary 58K40, 58K50, 58A10, 14H20.

*Key words and phrases*. symplectic manifold, local symplectic algebra, algebraic restrictions, relative Darboux theorem, isolated complete intersection singularities.

The work of W. Domitrz was supported by Polish MNiSW grant no. N N201 397237.

problem of classification of algebraic restrictions of symplectic forms to these subsets. For non-quasi-homogeneous subsets there is one more cohomological invariant apart of the algebraic restriction ([DJZ2], [DJZ1]). The method of algebraic restrictions is a very powerful tool to study the local symplectic algebra of 1-dimensional singular analytic varieties since the space of algebraic restrictions of closed 2-forms to a 1-dimensional singular analytic variety is finite-dimensional ([D]). By this method complete symplectic classifications of the  $A - D - E$  singularities of planar curves and the  $S_5$  singularity were obtained in [DJZ2]. These results were generalized to other 1-dimensional isolated complete intersection singularities: the  $S_\mu$  symplectic singularities for  $\mu > 5$  in [DT1], the  $T_7 - T_8$  symplectic singularities in [DT2] and the  $W_8 - W_9$  symplectic singularities in [T].

In this paper we show that some non-trivial symplectic invariants appear not only in the case of singular curves but also in the case of multiple points. We consider the symplectic classification of the 0-dimensional isolated complete intersection singularities (ICISs) in the symplectic space  $(\mathbb{C}^{2n}, \omega)$ . We need to introduce a symplectic  $V$ -equivalence to study this problem since we consider the ideals of function-germs that have not got the property of zeros.

We recall that  $\omega$  is a  $\mathbb{C}$ -analytic symplectic form on  $\mathbb{C}^{2n}$  if  $\omega$  is a  $\mathbb{C}$ -analytic nondegenerate closed 2-form, and  $\Phi : \mathbb{C}^{2n} \rightarrow \mathbb{C}^{2n}$  is a symplectomorphism if  $\Phi$  is a  $\mathbb{C}$ -analytic diffeomorphism and  $\Phi^*\omega = \omega$ .

**Definition 1.1.** Let  $f, g : (\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}^k, 0)$  be  $\mathbb{C}$ -analytic map-germs on the symplectic space  $(\mathbb{C}^{2n}, \omega)$ .  $f, g$  are **symplectically  $V$ -equivalent** if there exist a symplectomorphism-germ  $\Phi : (\mathbb{C}^{2n}, 0, \omega) \rightarrow (\mathbb{C}^{2n}, 0, \omega)$  and a  $\mathbb{C}$ -analytic map-germ  $M : (\mathbb{C}^{2n}, 0) \rightarrow GL(k, \mathbb{C})$  such that  $f \circ \Phi = M \cdot g$ .

If  $\Phi : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$  is a  $\mathbb{C}$ -analytic map-germ then for an ideal  $I$  in the ring of  $\mathbb{C}$ -analytic function-germs on  $\mathbb{C}^n$  we denote by  $\Phi^*I$  the following ideal  $\{f \circ \Phi : f \in I\}$  in the ring of  $\mathbb{C}$ -analytic function-germs on  $\mathbb{C}^n$ . The (symplectic)  $V$ -equivalence of map-germs  $f = (f_1, \dots, f_k), g = (g_1, \dots, g_k) : (\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}^k, 0)$  corresponds to the following (symplectic) equivalence of finitely-generated ideals  $\langle f_1, \dots, f_k \rangle$  and  $\langle g_1, \dots, g_k \rangle$  (see [AVG]).

**Definition 1.2.** Ideals  $\langle f_1, \dots, f_k \rangle$  and  $\langle g_1, \dots, g_k \rangle$  of  $\mathbb{C}$ -analytic function-germs at 0 on the symplectic space  $(\mathbb{C}^{2n}, \omega)$  are **symplectically equivalent** if there exists a symplectomorphism-germ  $\Phi : (\mathbb{C}^{2n}, 0, \omega) \rightarrow (\mathbb{C}^{2n}, 0, \omega)$  such that  $\Phi^* \langle f_1, \dots, f_k \rangle = \langle g_1, \dots, g_k \rangle$ .

In this paper we present the complete symplectic classification of the  $I_{a,b}, I_{2a+1}, I_{2a+4}, I_{a+5}, I_{10}^*$  singularities. For  $n = 1$  all  $V$ -orbits coincide with symplectic  $V$ -orbits. The situation for  $n \geq 2$  is different: the  $I_{a,b}$  singularities split into two symplectic  $V$ -orbits, the  $I_{2a+1}, I_{2a+4}, I_{a+5}$  singularities split into three symplectic orbits and finally  $I_{10}^*$  singularity splits into four symplectic  $V$ -orbits. The symplectic  $V$ -orbits of a map  $f = (f_1, \dots, f_{2n})$  are distinguished by the order of vanishing of a pullback of the germ of the symplectic form to a  $\mathbb{C}$ -analytic non-singular submanifold  $M$  of the minimal dimension such that the ideal of  $\mathbb{C}$ -analytic function-germs vanishing  $M$  is contained in the ideal  $\langle f_1, \dots, f_{2n} \rangle$  (see Definition 3.2).

To obtain these results we need some reformulation and modification of the method of algebraic restrictions. We present it in Section 2. In Section 3 we give the definitions of discrete symplectic invariants which completely distinguish symplectic  $V$ -singularities considered in this paper. We recall basic facts on the

classification of  $V$ -simple maps in Section 4. In Section 5 we prove the symplectic  $V$ -classification theorem for 0-dimensional ICISs (Theorem 5.1).

## 2. THE METHOD OF ALGEBRAIC RESTRICTIONS FOR THE SYMPLECTIC $V$ -EQUIVALENCE.

In this section we present basic facts on the method of algebraic restrictions adapted to the case of the symplectic  $V$ -equivalence. The proofs of all results are small modifications of the proofs of analogous results in [DJZ2].

Given a germ at 0 of a non-singular  $\mathbb{C}$ -analytic submanifold  $M$  of  $\mathbb{C}^m$  denote by  $\Lambda^p(M)$  the space of all germs at 0 of  $\mathbb{C}$ -analytic differential  $p$ -forms on  $M$ . By  $\mathcal{O}(M)$  denote the ring of  $\mathbb{C}$ -analytic function-germs on  $M$  at 0. Given an ideal  $I$  in  $\mathcal{O}(M)$  introduce the following subspace of  $\Lambda^p(M)$ :

$$\mathcal{A}_0^p(I, M) = \{\alpha + d\beta : \alpha \in I\Lambda^p(M), \beta \in I\Lambda^{p-1}(M)\}.$$

The relation  $\omega \in I\Lambda^p(M)$  means that  $\omega = \sum_{i=1}^k f_i \alpha_i$ , where  $\alpha_i \in \Lambda^p(M)$  and  $f_i \in I$  for  $i = 1, \dots, k$ .

**Definition 2.1.** Let  $I$  be an ideal of  $\mathcal{O}(M)$  and let  $\omega \in \Lambda^p(M)$ . The **algebraic restriction** of  $\omega$  to  $I$  is the equivalence class of  $\omega$  in  $\Lambda^p(M)$ , where the equivalence is as follows:  $\omega$  is equivalent to  $\tilde{\omega}$  if  $\omega - \tilde{\omega} \in \mathcal{A}_0^p(I, M)$ .

**Notation.** The algebraic restriction of the germ of a  $p$ -form  $\omega$  on  $M$  to the ideal  $I$  in  $\mathcal{O}(M)$  will be denoted by  $[\omega]_I$ . Writing  $[\omega]_I = 0$  (or saying that  $\omega$  has zero algebraic restriction to  $I$ ) we mean that  $[\omega]_I = [0]_I$ , i.e.  $\omega \in \mathcal{A}_0^p(I, M)$ .

**Definition 2.2.** Two algebraic restrictions  $[\omega]_I$  and  $[\tilde{\omega}]_{\tilde{I}}$  are called **diffeomorphic** if there exists the germ of a diffeomorphism  $\Phi : M \rightarrow \tilde{M}$  such that  $\Phi^*(\tilde{I}) = I$  and  $[\Phi^*\tilde{\omega}]_I = [\omega]_I$ .

**Definition 2.3.** The germ of a function, a differential  $k$ -form, or a vector field  $\alpha$  on  $(\mathbb{C}^m, 0)$  is **quasi-homogeneous** in a coordinate system  $(x_1, \dots, x_m)$  on  $(\mathbb{C}^m, 0)$  with positive integer weights  $(\lambda_1, \dots, \lambda_m)$  if  $\mathcal{L}_E \alpha = \delta \alpha$ , where  $E = \sum_{i=1}^m \lambda_i x_i \frac{\partial}{\partial x_i}$  is the germ of the **Euler vector field** on  $(\mathbb{C}^m, 0)$  and the integer  $\delta$  is called the quasi-degree.

It is easy to show that  $\alpha$  is quasi-homogeneous in a coordinate system  $(x_1, \dots, x_m)$  with weights  $(\lambda_1, \dots, \lambda_m)$  if and only if  $F_t^* \alpha = t^\delta \alpha$ , where

$$(2.1) \quad F_t(x_1, \dots, x_m) = (t^{\lambda_1} x_1, \dots, t^{\lambda_m} x_m).$$

**Definition 2.4.** A finitely generated ideal  $I$  of  $\mathcal{O}(\mathbb{C}^m)$  is **quasi-homogeneous** if there exist generators of  $I$  which are quasi-homogeneous in the same coordinate system  $(x_1, \dots, x_m)$  on  $\mathbb{C}^m$  with the same positive integer weights  $(\lambda_1, \dots, \lambda_m)$ .

A map-germ  $f = (f_1, \dots, f_k) : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^k, 0)$  is **quasi-homogeneous** if function-germs  $f_1, \dots, f_k$  are quasi-homogeneous in the same coordinate system  $(x_1, \dots, x_m)$  on  $\mathbb{C}^m$  with the same positive integer weights  $(\lambda_1, \dots, \lambda_m)$ .

To prove the generalization of Darboux-Givental theorem suitable for the symplectic  $V$ -equivalence of maps or the symplectic equivalence of ideals of function-germs we need the following version of the Relative Poincaré Lemma.

**Lemma 2.5.** *Let  $I$  be a finitely generated quasi-homogeneous ideal in  $\mathcal{O}(\mathbb{C}^m)$ . If  $\omega \in I\Lambda^p(\mathbb{C}^m)$  is closed then there exists  $\alpha \in I\Lambda^{p-1}(\mathbb{C}^m)$  such that  $\omega = d\alpha$ .*

*Proof.* We use the method described in [DJZ1]. We can find a coordinate system  $(x_1, \dots, x_m)$  on  $(\mathbb{C}^m, 0)$  and positive integer weights  $(\lambda_1, \dots, \lambda_m)$  and quasi-homogeneous function-germs  $f_1, \dots, f_k \in \mathcal{O}(\mathbb{C}^m)$  (in this coordinate systems with these weights) such that  $I = \langle f_1, \dots, f_k \rangle$ . Let  $\delta_i$  be a quasi-degree of  $f_i$  for  $i = 1, \dots, k$ .

Let  $F_t$  be a map defined in (2.1) and let  $V_t$  be a vector field along  $F_t$  for  $t \in [0; 1]$  such that  $V_t \circ F_t = F_t'$ .

Then we have  $F_0^* \omega = 0$  and it implies that

$$\omega = F_1^* \omega - F_0^* \omega = \int_0^1 (F_t^* \omega)' dt = \int_0^1 F_t^* d(V_t \lrcorner \omega) dt = d \left( \int_0^1 F_t^* (V_t \lrcorner \omega) dt \right).$$

Let  $\alpha = \int_0^1 F_t^* (V_t \lrcorner \omega) dt$ , then  $\omega = d\alpha$ . But  $\omega$  belongs to  $I\Lambda^p(\mathbb{C}^m)$ . It implies that there exist germs of  $p$ -forms  $\beta_i$  in  $\Lambda^p(\mathbb{C}^m)$  for  $i = 1, \dots, k$  such that  $\omega = \sum_{i=1}^k f_i \beta_i$ . So we have that

$$\alpha = \int_0^1 F_t^* (V_t \lrcorner \sum_{i=1}^k f_i \beta_i) dt = \sum_{i=1}^k f_i \int_0^1 t^{\delta_i} F_t^* (V_t \lrcorner \beta_i) dt.$$

Thus  $\alpha$  belongs to  $I\Lambda^{p-1}(\mathbb{C}^m)$ .  $\square$

The method of algebraic restrictions applied to finitely-generated quasi-homogeneous ideals is based on the following theorem.

**Theorem 2.6** (a modification of Theorem A in [DJZ2]). *Let  $I$  be a finitely generated quasi-homogeneous ideal in  $\mathcal{O}(\mathbb{C}^{2n})$ .*

- (1) *If  $\omega_0, \omega_1$  are germs at 0 of symplectic forms on  $\mathbb{C}^{2n}$  with the same algebraic restriction to  $I$  then there exists a  $\mathbb{C}$ -analytic diffeomorphism-germ  $\Phi$  of  $\mathbb{C}^{2n}$  at 0 of the form  $\Phi(x) = (x_1 + \phi_1(x), \dots, x_{2n} + \phi_{2n}(x))$ , where  $\phi_i \in I$  for  $i = 1, \dots, 2n$ , such that  $\Phi^* \omega_1 = \omega_0$ .*
- (2)  *$\mathbb{C}$ -analytic quasi-homogeneous map-germs  $f = (f_1, \dots, f_k), g = (g_1, \dots, g_k) : (\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}^k, 0)$  on the symplectic space  $(\mathbb{C}^{2n}, \omega)$  are symplectically  $V$ -equivalent if and only if algebraic restrictions  $[\omega]_{\langle f_1, \dots, f_k \rangle}$  and  $[\omega]_{\langle g_1, \dots, g_k \rangle}$  are diffeomorphic.*

**Remark 2.7.** It is obvious that if  $\Phi(x) = (x_1 + \phi_1(x), \dots, x_{2n} + \phi_{2n}(x))$  where  $\phi_i \in I$  for  $i = 1, \dots, 2n$  then  $\Phi^* I = I$

A proof of Theorem 2.6 can be obtain by a small modification of the proof of Theorem A in [DJZ2]. One only needs Lemma 2.5 and the following fact.

**Lemma 2.8.** *Let  $I$  be a finitely generated ideal in  $\mathcal{O}(\mathbb{C}^m)$ . Let  $X_t = \sum_{i=1}^m f_{i,t} \frac{\partial}{\partial x_i}$  for  $t \in [0; 1]$  be a family of germs of  $\mathbb{C}$ -analytic vector fields on  $\mathbb{C}^m$  such that  $f_{i,t} \in I$  for  $i = 1, \dots, m$ .*

*If  $\Phi_t$  for  $t \in [0, 1]$  is a family of diffeomorphism-germs of  $(\mathbb{C}^m, 0)$  such that*

$$(2.2) \quad \frac{d}{dt} \Phi_t = X_t \circ \Phi_t$$

*then*

$$(2.3) \quad \Phi_t(x) = (x_1 + \phi_{1,t}(x), \dots, x_{2n} + \phi_{2n,t}(x)),$$

*where  $\phi_{i,t} \in I$  for  $i = 1, \dots, 2n$ .*

*A sketch of the proof.* The map  $t \mapsto \Phi_t(x)$  is a solution of ODE  $\frac{dy}{dt} = X_t(y)$  with the initial condition  $y(0) = x$ . So  $\Phi_t(x)$  can be obtained as a limit  $\lim_{n \rightarrow \infty} T^n \Psi$  where  $\Psi(t, x) \equiv x$  and  $(T\Psi)(t, x) = x + \int_0^t X_s(\Psi(s, x)) ds$  is the Picard's operator. It is easy to see that if  $\Psi$  has the form (2.3) then  $T\Psi$  has the form (2.3) too. The ideal  $I$  is finitely generated. Thus  $\Phi_t$  has also this form.  $\square$

Theorem 2.6 reduces the problem of symplectic classification of quasi-homogeneous ideals to the problem of classification of the algebraic restrictions of the germ of the symplectic form to quasi-homogeneous ideals.

The meaning of the zero algebraic restriction is explained by the following theorem.

**Theorem 2.9** (a modification of Theorem B in [DJZ2]). *A finitely generated quasi-homogeneous ideal  $I$  of  $\mathcal{O}(\mathbb{C}^{2n})$  contains the ideal of  $\mathbb{C}$ -analytic function-germs vanishing on the germ of a non-singular Lagrangian submanifold of the symplectic space  $(\mathbb{C}^{2n}, \omega)$  if and only if the symplectic form  $\omega$  has zero algebraic restriction to  $I$ .*

We now formulate the modifications of basic properties of algebraic restrictions ([DJZ2]). First we can reduce the dimension of the manifold due to the following propositions.

If the ideal  $I$  in  $\mathcal{O}(\mathbb{C}^m)$  contains an ideal  $I(M)$  of function-germs vanishing on a non-singular submanifold  $M \subset \mathbb{C}^m$  then the classification of the algebraic restrictions to  $I$  of  $p$ -forms on  $\mathbb{C}^m$  reduces to the classification of the algebraic restrictions to  $I|_M = \{f|_M : f \in I\}$  of  $p$ -forms on  $M$ . At first note that the algebraic restrictions  $[\omega]_I$  and  $[\omega|_{TM}]_{I|_M}$  can be identified:

**Proposition 2.10.** *Let  $I$  be an ideal in  $\mathcal{O}(\mathbb{C}^m)$  which contains an ideal of function-germs vanishing on a non-singular submanifold  $M \subset \mathbb{C}^m$  and let  $\omega_1, \omega_2$  be germs of  $p$ -forms on  $\mathbb{C}^m$ . Then  $[\omega_1]_I = [\omega_2]_I$  if and only if  $[\omega_1|_{TM}]_{I|_M} = [\omega_2|_{TM}]_{I|_M}$ .*

The following, less obvious statement, means that the *orbits* of the algebraic restrictions  $[\omega]_I$  and  $[\omega|_{TM}]_{I|_M}$  also can be identified.

**Proposition 2.11.** *Let  $I_1, I_2$  be ideals in the ring  $\mathcal{O}(\mathbb{C}^m)$ , which contain  $I(M_1)$  and  $I(M_2)$  respectively, where  $M_1, M_2$  are equal-dimensional non-singular submanifolds. Let  $\omega_1, \omega_2$  be two germs of  $p$ -forms. The algebraic restrictions  $[\omega_1]_{I_1}$  and  $[\omega_2]_{I_2}$  are diffeomorphic if and only if the algebraic restrictions  $[\omega_1|_{TM_1}]_{I_1|_{M_1}}$  and  $[\omega_2|_{TM_2}]_{I_2|_{M_2}}$  are diffeomorphic.*

To calculate the space of algebraic restrictions of germs of 2-forms we will use the following obvious properties.

**Proposition 2.12.** *If  $\omega \in \mathcal{A}_0^k(I, \mathbb{C}^{2n})$  then  $d\omega \in \mathcal{A}_0^{k+1}(I, \mathbb{C}^{2n})$  and  $\omega \wedge \alpha \in \mathcal{A}_0^{k+p}(I, \mathbb{C}^{2n})$  for any germ of  $\mathbb{C}$ -analytic  $p$ -form  $\alpha$  on  $\mathbb{C}^{2n}$ .*

Then we need to determine which algebraic restrictions of closed 2-forms are realizable by symplectic forms. This is possible due to the following fact.

**Proposition 2.13.** *Let  $I$  be an ideal of  $\mathcal{O}(\mathbb{C}^{2n})$ . Let  $r$  be the minimal dimension of non-singular submanifolds  $M$  of  $\mathbb{C}^{2n}$  such that  $I$  contains the ideal  $I(M)$ . The algebraic restriction  $[\theta]_I$  of the germ of a closed 2-form  $\theta$  is realizable by the germ of a symplectic form on  $\mathbb{C}^{2n}$  if and only if  $\text{rank}(\theta|_{T_0M}) \geq 2r - 2n$ .*

## 3. DISCRETE SYMPLECTIC INVARIANTS.

We use discrete symplectic invariants to distinguish symplectic singularity classes. We modify definitions of these invariants introduced in [DJZ2] for the symplectic  $V$ -equivalence.

The first invariant is a symplectic multiplicity ([DJZ2]) introduced in [IJ1] as a symplectic defect of a curve.

Let  $f : (\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}^k, 0)$  be the germ of a  $\mathbb{C}$ -analytic map on the symplectic space  $(\mathbb{C}^{2n}, \omega)$ .

**Definition 3.1.** The **symplectic multiplicity**  $\mu_{\text{sympl}}(f)$  of  $f$  is the codimension of the symplectic  $V$ -orbit of  $f$  in the  $V$ -orbit of  $f$ .

The second invariant is the index of isotropy [DJZ2].

**Definition 3.2.** The **index of isotropy**  $\iota(f)$  of  $f = (f_1, \dots, f_k)$  is the maximal order of vanishing of the 2-forms  $\omega|_{TM}$  over all smooth submanifolds  $M$  such that the ideal  $\langle f_1, \dots, f_k \rangle$  contains  $I(M)$ .

These invariants can be described in terms of algebraic restrictions.

**Proposition 3.3** ([DJZ2]). *The symplectic multiplicity of the germ of a quasi-homogeneous map  $f = (f_1, \dots, f_k)$  on the symplectic space  $(\mathbb{C}^{2n}, \omega)$  is equal to the codimension of the orbit of the algebraic restriction  $[\omega]_{\langle f_1, \dots, f_k \rangle}$  with respect to the group of diffeomorphism-germs preserving the ideal  $\langle f_1, \dots, f_k \rangle$  in the space of the algebraic restrictions of closed 2-forms to  $\langle f_1, \dots, f_k \rangle$ .*

**Proposition 3.4** ([DJZ2]). *The index of isotropy of the germ of a quasi-homogeneous map  $f = (f_1, \dots, f_k)$  on the symplectic space  $(\mathbb{C}^{2n}, \omega)$  is equal to the maximal order of vanishing of closed 2-forms representing the algebraic restriction  $[\omega]_{\langle f_1, \dots, f_k \rangle}$ .*

We will use these invariants to distinguish symplectic singularities.

4.  $V$ -SIMPLE MAPS

We recall some results on classification of  $V$ -simple germs (for details see [AVG]).

**Definition 4.1.** The germ  $f : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^n, 0)$  is said be  **$V$ -simple** if its  $k$ -jet, for any  $k$ , has a neighborhood in the small jet space  $J_{0,0}^k(\mathbb{C}^m, \mathbb{C}^n)$  that intersects only a finite number of  $V$ -equivalence classes (bounded by a constant independent of  $k$ ).

**Definition 4.2.** The  **$p$ -parameter suspension of the map-germ**  $f : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^n, 0)$  is the map germ

$$F : (\mathbb{C}^m \times \mathbb{C}^p, 0) \ni (y, z) \mapsto (f(y), z) \in (\mathbb{C}^n \times \mathbb{C}^p, 0).$$

**Theorem 4.3** (see [AVG]). *The  $V$ -simple map-germs  $(\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^n, 0)$  with  $m \geq n$  belong, up to  $V$ -equivalence and suspension, to one of the three lists: the  $A-D-E$  singularities of map-germs  $\mathbb{C}^m \rightarrow \mathbb{C}$  (hypersurfaces with an isolated singularity),  $S-T-U-W-Z$  singularities of map-germs  $\mathbb{C}^3 \rightarrow \mathbb{C}^2$  (1-dimensional ICISs) and singularities of map-germs  $\mathbb{C}^2 \rightarrow \mathbb{C}^2$  (0-dimensional ICISs) presented in Table 1.*

The normal forms in Table 1 were obtained in [G] by M. Giusti.

Notation	Normal form	Restrictions
$I_{a,b}$	$(yz, y^a + z^b)$	$a \geq b \geq 2$
$I_{2a+1}$	$(y^2 + z^3, z^a)$	$a \geq 3$
$I_{2a+4}$	$(y^2 + z^3, yz^a)$	$a \geq 2$
$I_{a+5}$	$(y^2 + z^a, yz^2)$	$a \geq 4$
$I_{10}^*$	$(y^2, z^4)$	-

TABLE 1.  $V$ -simple map-germs  $\mathbb{C}^2 \rightarrow \mathbb{C}^2$ .

## 5. SYMPLECTIC 0-DIMENSIONAL ICISS

We use the method of algebraic restrictions to obtain a complete classification of singularities presented in Table 1.

**Theorem 5.1.** *Any map-germ  $(\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}^{2n}, 0)$  from the symplectic space  $(\mathbb{C}^{2n}, \sum_{i=1}^n dp_i \wedge dq_i)$  which is  $V$ -equivalent (up to a suitable suspension) to one of the normal forms in Table 1 is symplectically  $V$ -equivalent to one and only one of the following normal forms presented in Table 2*

Symplectic class	Normal forms	$cod$	$\mu_{sympl}$	$i$
$I_{a,b}^0, (n \geq 1)$	$(p_1q_1, p_1^a + q_1^b, p_2, q_2, \dots, p_n, q_n)$	0	0	0
$I_{a,b}^1, (n \geq 2)$	$(p_1p_2, p_1^a + p_2^b, q_1, q_2, p_3, q_3, \dots, p_n, q_n)$	1	1	$\infty$
$I_{2a+1}^0, (n \geq 1)$	$(p_1^2 + q_1^3, q_1^a, p_2, q_2, \dots, p_n, q_n)$	0	0	0
$I_{2a+1}^1, (n \geq 2)$	$(p_1^2 + p_2^3, p_2^a, q_1, q_2 + p_1p_2, p_3, q_3, \dots, p_n, q_n)$	1	1	1
$I_{2a+1}^2, (n \geq 2)$	$(p_1^2 + p_2^3, p_2^a, q_1, q_2, p_3, q_3, \dots, p_n, q_n)$	2	2	$\infty$
$I_{2a+4}^0, (n \geq 1)$	$(p_1^2 + q_1^3, p_1q_1^a, p_2, q_2, \dots, p_n, q_n)$	0	0	0
$I_{2a+4}^1, (n \geq 2)$	$(p_1^2 + p_2^3, p_1p_2^a, q_1, q_2 + p_1p_2, p_3, q_3, \dots, p_n, q_n)$	1	1	1
$I_{2a+4}^2, (n \geq 2)$	$(p_1^2 + p_2^3, p_1p_2^a, q_1, q_2, p_3, q_3, \dots, p_n, q_n)$	2	2	$\infty$
$I_{a+5}^0, (n \geq 1)$	$(p_1^2 + q_1^a, p_1q_1^2, p_2, q_2, \dots, p_n, q_n)$	0	0	0
$I_{a+5}^1, (n \geq 2)$	$(p_1^2 + p_2^a, p_1p_2^2, q_1, q_2 + p_1p_2, p_3, q_3, \dots, p_n, q_n)$	1	1	1
$I_{a+5}^2, (n \geq 2)$	$(p_1^2 + p_2^a, p_1p_2^2, q_1, q_2, p_3, q_3, \dots, p_n, q_n)$	2	2	$\infty$
$I_{10}^{*0}, (n \geq 1)$	$(p_1^2, q_1^4, p_2, q_2, \dots, p_n, q_n)$	0	0	0
$I_{10}^{*1}, (n \geq 2)$	$(p_1^2, p_2^4, q_1, q_2 + p_1p_2, p_3, q_3, \dots, p_n, q_n)$	1	1	1
$I_{10}^{*2}, (n \geq 2)$	$(p_1^2, p_2^4, q_1, q_2 + p_1p_2^2, p_3, q_3, \dots, p_n, q_n)$	2	2	2
$I_{10}^{*3}, (n \geq 2)$	$(p_1^2, p_2^4, q_1, q_2, p_3, q_3, \dots, p_n, q_n)$	3	3	$\infty$

TABLE 2. Classification of symplectic 0-dimensional isolated complete intersection singularities,  $cod$  – codimension of the classes;  $\mu_{sympl}$  – symplectic multiplicity;  $i$  – index of isotropy.

*Proof.* In the case  $n = 1$  the proof follows from results in [DR] where it was proved that for quasi-homogeneous singularities in the  $\mathbb{C}$ -analytic category  $V$ -orbits coincide with volume-preserving  $V$ -orbits. For general  $n$  we present the proof in the case of the  $I_{10}^*$  singularity where there are 4 different symplectic singularity classes, and in the case of the  $I_{a+5}$  singularity. The proofs in other cases are very similar.

For the  $I_{10}^*$  singularity we calculate the space of algebraic restrictions of 2-forms to the ideal  $I = \langle y^2, z^4, x_1, \dots, x_{2n-2} \rangle$ . The ideal generated by  $x_1, \dots, x_{2n-2}$  is contained in  $I$ . So by Proposition 2.10 we may consider the following ideal  $J = I|_{\{x_1 = \dots = x_{2n-2} = 0\}} = \langle y^2, z^4 \rangle$  in the ring  $\mathcal{O}(\mathbb{C}^2)$ . By Proposition 2.12 germs

of 1-forms  $d(1/2y^2) = ydy$ ,  $d(1/4z^4) = z^3dz$  and germs of 2-forms  $ydy \wedge dz$ ,  $z^3dy \wedge dz$  have zero algebraic restriction to  $J$ . So any algebraic restriction of the germ of a closed 2-forms to  $J$  can be presented in the following form  $[\omega]_J = A[dy \wedge dz]_J + B[zdy \wedge dz]_J + C[z^2dy \wedge dz]_J$ , where  $A, B, C \in \mathbb{C}$ .

If  $A \neq 0$  then we obtain  $\Phi^*[\omega]_J = [dy \wedge dz]_J$  by the diffeomorphism-germ of the form  $\Phi(y, z) = (y, z(A + 1/2Bz + 1/3Cz^2))$ . If  $A = 0$  and  $B \neq 0$  then we obtain  $\Phi^*[\omega]_J = [zdy \wedge dz]_J$  by the diffeomorphism-germ of the form  $\Phi(y, z) = (y, z\phi(z))$ , where  $\phi^2(z) = B + 2/3Cz$ . If  $A = B = 0$  and  $C \neq 0$  then we obtain  $\Phi^*[\omega]_J = [z^2dy \wedge dz]_J$  by the diffeomorphism-germ of the form  $\Phi(y, z) = (Cy, z)$ .

Since the minimal dimension  $r$  of the germ of a non-singular submanifold  $M$  such that  $I(M) \subset I$  is 2 then by Proposition 2.13 for  $n = 1$  only the algebraic restriction  $[dy \wedge dz]_I$  is realizable by the germ of a symplectic form.

For  $n > 1$  all algebraic restrictions are realizable by the following symplectic forms:

$$(5.1) \quad dy \wedge dz + \sum_{i=1}^{n-1} dx_{2i-1} \wedge dx_{2i},$$

$$(5.2) \quad zdy \wedge dz + dy \wedge dx_1 + dz \wedge dx_2 + \sum_{i=2}^{n-1} dx_{2i-1} \wedge dx_{2i},$$

$$(5.3) \quad z^2dy \wedge dz + dy \wedge dx_1 + dz \wedge dx_2 + \sum_{i=2}^{n-1} dx_{2i-1} \wedge dx_{2i},$$

$$(5.4) \quad dy \wedge dx_1 + dz \wedge dx_2 + \sum_{i=2}^{n-1} dx_{2i-1} \wedge dx_{2i}.$$

By a simple change of coordinates we obtain the normal forms in Table 2.

For the  $I_{a+5}$  singularity the space algebraic restrictions of germs of closed 2-forms to the ideal  $I = \langle y^2 + z^a, yz^2, x_1, \dots, x_{2n-2} \rangle$  can be calculated in the same way. We obtain that any algebraic restriction of the germs of a closed 2-forms on  $\mathbb{C}^2 = \{x_1 = \dots = x_{2n-2} = 0\}$  to  $J = I|_{\{x_1 = \dots = x_{2n-2} = 0\}} = \langle y^2 + z^a, yz^2 \rangle$  can be presented in the following form

$$(5.5) \quad [\omega]_J = A[dy \wedge dz]_J + B[zdy \wedge dz]_J,$$

where  $A, B \in \mathbb{C}$ .

First assume that  $A \neq 0$ . Let  $E$  denote the germ of the Euler vector field  $ay \frac{\partial}{\partial y} + 2z \frac{\partial}{\partial z}$ . Then it is easy to check that a flow  $\Phi_t$  of the germ of a vector field  $X = \frac{B}{(a+4)A} zE$  preserves  $J$ ,  $\mathcal{L}_X(Ady \wedge dz) = Bzdy \wedge dz$ ,  $[\mathcal{L}_X(Bzdy \wedge dz)]_J = 0$ . Therefore  $\Phi_t^*[Ady \wedge dz + tBzdy \wedge dz]_J = [Ady \wedge dz]_J$  for  $t \in [0, 1]$  (see [D]). Finally by a linear change of coordinates of the form  $(y, z) \mapsto (Cy, Dz)$ , where for  $C, D \in \mathbb{C}$  such that  $C^2 = D^a$  and  $CD = A$  we show that if  $A \neq 0$  then the algebraic restriction (5.5) is diffeomorphic to  $[dy \wedge dz]_J$ . By a similar change of coordinates preserving  $J$  we show that if  $A = 0$  and  $B \neq 0$  then the algebraic restriction (5.5) is diffeomorphic to  $[zdy \wedge dz]_J$ . As in the previous case, for  $n = 1$  only  $[dy \wedge dz]_I$  can be realizable by the germ of a symplectic form. For  $n \geq 2$  algebraic restrictions are realizable by (5.1), (5.2) and (5.4). Normal forms in Table 2 are obtained by an obvious change of coordinates.  $\square$



## REFERENCES

- [A1] V. I. Arnold, *First step of local symplectic algebra*, Differential topology, infinite-dimensional Lie algebras, and applications. D. B. Fuchs' 60th anniversary collection. Providence, RI: American Mathematical Society. Transl., Ser. 2, Am. Math. Soc. 194(44), 1999,1-8.
- [AG] V. I. Arnold, A. B. Givental *Symplectic geometry*, in Dynamical systems, IV, 1-138, Encyclopedia of Mathematical Sciences, vol. 4, Springer, Berlin, 2001.
- [AVG] V. I. Arnold, S. M. Gusein-Zade, A. N. Varchenko, *Singularities of Differentiable Maps*, Vol. 1, Birhauser, Boston, 1985.
- [D] W. Domitrz, *Local symplectic algebra of quasi-homogeneous curves*, Fundamentae Mathematicae 204 (2009), 57-86.
- [DJZ1] W. Domitrz, S. Janeczko, M. Zhitomirskii, *Relative Poincaré lemma, contractibility, quasi-homogeneity and vector fields tangent to a singular variety*, Ill. J. Math. 48, No.3 (2004), 803-835.
- [DJZ2] W. Domitrz, S. Janeczko, M. Zhitomirskii, *Symplectic singularities of varieties: the method of algebraic restrictions*, J. reine und angewandte Math. 618 (2008), 197-235.
- [DR] W. Domitrz, J. H. Rieger, *Volume preserving subgroups of  $\mathcal{A}$  and  $\mathcal{K}$  and singularities in unimodular geometry*, Mathematische Annalen 345 (2009), 783-817.
- [DT1] W. Domitrz, Ż. Trębska, *Symplectic  $S_\mu$  singularities*, Real and Complex Singularities, Contemporary Mathematics, 569, Amer. Math. Soc., Providence, RI, 2012, 45-65.
- [DT2] W. Domitrz, Ż. Trębska, *Symplectic  $T_7$ ,  $T_8$  singularities and Lagrangian tangency orders*, to appear in Proceedings of the Edinburgh Mathematical Society.
- [G] M. Giusti, *Classification des singularités isolées d'intersections complètes simples*, C. R. Acad. Sci., Paris, Sr. A 284 (1977), 167-170 .
- [IJ1] G. Ishikawa, S. Janeczko, *Symplectic bifurcations of plane curves and isotropic liftings*, Q. J. Math. **54**, No.1 (2003), 73-102.
- [IJ2] G. Ishikawa, S. Janeczko, *Symplectic singularities of isotropic mappings*, Geometric singularity theory, Banach Center Publications **65** (2004), 85-106.
- [K] P. A. Kolgushkin, *Classification of simple multigerms of curves in a space endowed with a symplectic structure*, St. Petersburg Math. J. **15** (2004), no. 1, 103-126.
- [L] E. J. M. Looijenga *Isolated Singular Points on Complete Intersections*, London Mathematical Society Lecture Note Series 77, Cambridge University Press 1984.
- [T] Ż. Trębska, *Symplectic  $W_8$  and  $W_9$  singularities*, to appear in Journal of Singularities.
- [Z] M. Zhitomirskii, *Relative Darboux theorem for singular manifolds and local contact algebra*, Can. J. Math. **57**, No.6 (2005), 1314-1340.

WARSAW UNIVERSITY OF TECHNOLOGY, FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, PLAC POLITECHNIKI 1, 00-661 WARSAW, POLAND,  
*E-mail address:* domitrz@mini.pw.edu.pl