

# Nonlocal Gravitational Models and Exact Solutions

Sergey Yu. Vernov\*

Instituto de Ciencias del Espacio, Institut d'Estudis Espacials de Catalunya

Campus UAB, Facultat de Ciències, Torre C5-Parell-2a planta,

E-08193, Bellaterra (Barcelona), Spain

Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University,

Leninskie Gory 1, 119991, Moscow, Russia

## Abstract

A nonlocal gravity model with a function  $f(\square^{-1}R)$ , where  $\square$  is the d'Alembert operator, is considered. The algorithm, allowing to reconstruct  $f(\square^{-1}R)$ , corresponding to the given Hubble parameter and the state parameter of the matter, is proposed. Using this algorithm, we find the functions  $f(\square^{-1}R)$ , corresponding to de Sitter solutions.

## 1 Nonlocal gravitational models

In this paper we consider nonlocal gravity models, which are describing by the action

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} \left[ R \left( 1 + f(\square^{-1}R) \right) - 2\Lambda \right] + \mathcal{L}_{\text{matter}} \right\}, \quad (1)$$

where  $\kappa^2 \equiv 8\pi/M_{\text{Pl}}^2$ , the Planck mass being  $M_{\text{Pl}} = 1.2 \times 10^{19}$  GeV. We use the signature  $(-, +, +, +)$ ,  $g$  is the determinant of the metric tensor  $g_{\mu\nu}$ ,  $\Lambda$  is the cosmological constant,  $f$  is a differentiable function, and  $\mathcal{L}_{\text{matter}}$  is the matter Lagrangian. Note that the modified gravity action (1) does not include a new dimensional parameter. This nonlocal model has a local scalar-tensor formulation. Introducing two scalar fields,  $\eta$  and  $\xi$ , we can rewrite action (1) in the following local form:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} \left[ R \left( 1 + f(\eta) - \xi \right) + \xi \square \eta - 2\Lambda \right] + \mathcal{L}_{\text{matter}} \right\}. \quad (2)$$

By varying the action (2) over  $\xi$ , we get  $\square \eta = R$ . Substituting  $\eta = \square^{-1}R$  into action (2), one reobtains action (1). Varying action (2) with respect to the metric tensor  $g_{\mu\nu}$ , one gets

$$\begin{aligned} & \frac{1}{2} g_{\mu\nu} \left[ R \left( 1 + f(\eta) - \xi \right) - \partial_\rho \xi \partial^\rho \eta - 2\Lambda \right] - R_{\mu\nu} \left( 1 + f(\eta) - \xi \right) + \\ & + \frac{1}{2} \left( \partial_\mu \xi \partial_\nu \eta + \partial_\mu \eta \partial_\nu \xi \right) - \left( g_{\mu\nu} \square - \nabla_\mu \partial_\nu \right) \left( f(\eta) - \xi \right) + \kappa^2 T_{\text{matter} \mu\nu} = 0, \end{aligned} \quad (3)$$

---

\*E-mail: vernov@ieec.uab.es, svernov@theory.sinp.msu.ru

where  $\nabla_\mu$  is the covariant derivative and  $T_{\text{matter}\mu\nu}$  the energy–momentum tensor of matter.

Variation of action (2) with respect to  $\eta$  yields  $\square\xi + f'(\eta)R = 0$ , where the prime denotes derivative with respect to  $\eta$ . If the scalar fields  $\eta$  and  $\xi$  depend on time only, then in the spatially flat Friedmann–Lemaître–Robertson–Walker metric with the interval

$$ds^2 = -dt^2 + a^2(t) (dx_1^2 + dx_2^2 + dx_3^2), \quad (4)$$

equations (3) are equivalent to the following ones:

$$-3H^2(1 + \Psi) + \frac{1}{2}\dot{\xi}\dot{\eta} - 3H\dot{\Psi} + \Lambda + \kappa^2\rho_m = 0, \quad (5)$$

$$\left(2\dot{H} + 3H^2\right)(1 + \Psi) + \frac{1}{2}\dot{\xi}\dot{\eta} + \ddot{\Psi} + 2H\dot{\Psi} - \Lambda + \kappa^2 P_m = 0, \quad (6)$$

where  $\Psi(t) = f(\eta(t)) - \xi(t)$ ,  $H = \dot{a}/a$  is the Hubble parameter, differentiation with respect to time  $t$  is denoted by a dot. For a perfect matter fluid, we have  $T_{\text{matter}00} = \rho_m(t)$  and  $T_{\text{matter}ij} = P_m(t)g_{ij}$ . The equation of state (EoS) is

$$\dot{\rho}_m = -3H(P_m + \rho_m). \quad (7)$$

The equations of motion for the scalar fields  $\eta$  and  $\xi$  are as follows

$$\ddot{\eta} + 3H\dot{\eta} = -6\left(\dot{H} + 2H^2\right), \quad (8)$$

$$\ddot{\xi} + 3H\dot{\xi} = 6\left(\dot{H} + 2H^2\right)f'(\eta). \quad (9)$$

Note that the considered system of equations does not include the function  $\eta$ , but only  $f(\eta)$ ,  $f'(\eta)$  and time derivatives of  $\eta$ . Also, one can add a constant to  $f(\eta)$  and the same constant to  $\xi$ , without changing of equations. So,  $f(\eta)$  can be determined up to a constant.

For the model with action (2), contained a perfect fluid with a constant state parameter  $w_m$ , a reconstruction procedure has been made [1] in terms of functions of the scale factor  $a$ . For the model, describing by the initial nonlocal action (1), a technique for choosing the distortion function so as to fit an arbitrary expansion history has been derived in [2].

Our goal is to demonstrate how one can reconstruct  $f(\eta)$  and get a model with exact solutions for the given Hubble parameter  $H(t)$  and the state parameter  $w_m(t) = P_m(t)/\rho_m(t)$ . We show that to do this it is enough to solve only linear differential equations.

The algorithm is as follows:

- Assume the explicit form of the functions  $H(t)$  and  $w_m(t)$ .
- Solve (7) and get  $\rho_m(t)$ .
- Solve (8) and get  $\eta(t)$ .
- Subtracting equation (5) from equation (6), get a linear differential equation

$$\ddot{\Psi} + 5H\dot{\Psi} + \left(2\dot{H} + 6H^2\right)(1 + \Psi) - 2\Lambda + \kappa^2(w_m - 1)\rho_m = 0. \quad (10)$$

- Using the known  $H(t)$ ,  $w_m(t)$ , and  $\rho_m(t)$ , solve (10) and get  $\Psi(t)$ .

- Substituting  $\xi(t) = f(\eta(t)) - \Psi(t)$  into equation (9), we get a linear differential equation for  $f(\eta)$

$$f''(\eta)\dot{\eta}^2 - 12\left(\dot{H} + 2H^2\right)f'(\eta) = \ddot{\Psi} + 3H\dot{\Psi}. \quad (11)$$

To get (11) we also use the inverse function  $t(\eta)$ . Note that equation (11) is a necessary condition that the model has solutions in the given form.

- Solve (11) and get the sought-for function  $f(\eta)$ .
- Substitute the obtained function  $f(\eta)$  to equation (5) and equation (6) to check the existence of the solutions in the given form.

## 2 Nonlocal models with de Sitter solutions

To demonstrate how the algorithm work we seek a such  $f(\eta)$  that the model has a de Sitter solution, in other words, the Hubble parameter is a nonzero constant:  $H = H_0$ . In this case, equation (8) has the following general solution:

$$\eta(t) = -4H_0(t - t_0) - \eta_0 e^{-3H_0(t-t_0)}, \quad (12)$$

with integration constants  $t_0$  and  $\eta_0$ . All equations are homogeneous. If a solution exists at  $t_0 = 0$ , then it exists at an arbitrary  $t_0$ . So, without loss of generality we can set  $t_0 = 0$ .

Note that equation (11) has been obtained without any restrictions on solutions and the perfect matter fluid. To demonstrate how one can get  $f(\eta)$ , which admits the existence of de Sitter solutions, in the explicit form, we restrict ourself to the case  $\eta_0 = 0$ . In this case, equation (11) has the following form:

$$16H_0^2 f''(\eta) - 24H_0^2 f'(\eta) = \Phi(\eta), \quad (13)$$

where  $\Phi(\eta) = \Phi(-4H_0 t) \equiv \ddot{\Psi} + 3H_0 \dot{\Psi}$ . We get the following solution

$$f(\eta) = \frac{1}{16H_0^2} \int \left\{ \int_{\tilde{\zeta}}^{\eta} \Phi(\tilde{\zeta}) e^{-3\tilde{\zeta}/2} d\tilde{\zeta} + 16C_3 H_0^2 \right\} e^{3\zeta/2} d\zeta + C_4, \quad (14)$$

where  $C_3$  and  $C_4$  are arbitrary constants. We can fix  $C_4$  without loss of generality.

Following [3], we consider the matter with the state parameter  $w_m \equiv P_m/\rho_m$  to be a constant, not equal to  $-1$ . Thus, equation (7) has the following general solution

$$\rho_m = \rho_0 e^{-3(1+w_m)H_0 t}, \quad (15)$$

where  $\rho_0$  is an arbitrary constant. Equation (10) has the following general solution:

At  $w_m \neq 0$  and  $w_m \neq -1/3$ ,

$$\Psi_1(t) = C_1 e^{-3H_0 t} + C_2 e^{-2H_0 t} - 1 + \frac{\Lambda}{3H_0^2} - \frac{\kappa^2 \rho_0 (w_m - 1)}{3H_0^2 w_m (1 + 3w_m)} e^{-3H_0 (w_m + 1)t},$$

At  $w_m = -\frac{1}{3}$ ,  $\Psi_2(t) = C_1 e^{-3H_0 t} + C_2 e^{-2H_0 t} - 1 + \frac{\Lambda}{3H_0^2} + \frac{4\kappa^2 \rho_0}{3H_0} e^{-2H_0 t} t,$

$$\text{At } w_m = 0, \quad \Psi_3(t) = C_1 e^{-3H_0 t} + C_2 e^{-2H_0 t} - 1 + \frac{\Lambda}{3H_0^2} - \frac{\kappa^2 \rho_0}{H_0} e^{-3H_0 t} t,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

Substituting the explicit form of  $\Psi(t)$ , we get

$$f_1(\eta) = \frac{C_2}{4} e^{\eta/2} + C_3 e^{3\eta/2} + C_4 - \frac{\kappa^2 \rho_0}{3(1+3w_m)H_0^2} e^{3(w_m+1)\eta/4}, \quad \text{at } w_m \neq -\frac{1}{3}, \quad (16)$$

$$\tilde{f}_1(\eta) = \frac{C_2}{4} e^{\eta/2} + C_3 e^{3\eta/2} + C_4 + \frac{\kappa^2 \rho_0}{4H_0^2} \left(1 - \frac{1}{3}\eta\right) e^{\eta/2}, \quad \text{at } w_m = -\frac{1}{3}, \quad (17)$$

where  $C_3$  and  $C_4$  are arbitrary constants. Note that  $C_2$  is an arbitrary constant as well.

One can see that the key ingredient of all functions  $f_i(\eta)$  is an exponential function. For the models with  $f(\eta)$  equal to an exponential function or a sum of exponential functions, particular de Sitter solutions have been found in [3, 4]. De Sitter solutions in the case of the exponential function  $f$  have been generalized and those stability have been analysed in [5].

### 3 Conclusion

Exact solutions play an important role in modern cosmological models, in particular, in nonlocal cosmological models [3, 4, 5, 6, 7, 8, 9]. The main result of this paper is the algorithm, using which one can reconstruct  $f(\square^{-1}R)$ , corresponding to the given Hubble parameter and the state parameter of the matter. We have found that the function  $f$  corresponding to de Sitter solutions is an exponential function or a sum of exponential functions<sup>1</sup>. In the case of the exponential function  $f$ , expanding universe solutions  $a \sim t^n$  have been found in [4, 8]. We plan to analyse possible forms of the corresponding function  $f$  in future investigations.

The author is grateful to the organizers of the Dubna International Workshop "Supersymmetries and Quantum Symmetries" (SQS'2011) for hospitality and financial support. The author wishes to express his thanks to Emilio Elizalde, Sergei D. Odintsov, Ekaterina O. Pozdeeva, Richard P. Woodard, and Ying-li Zhang for useful and stimulating discussions. The work is supported in part by the RFBR grant 11-01-00894, by the Russian Ministry of Education and Science under grants NSh-4142.2010.2 and NSh-3920.2012.2, and by contract CPAN10-PD12 (ICE, Barcelona, Spain).

### References

- [1] Koivisto T., *Dynamics of Nonlocal Cosmology*, Phys. Rev. D **77** (2008) 123513, arXiv:0803.3399
- [2] Deffayet C., Woodard R.P., *Reconstructing the Distortion Function for Nonlocal Cosmology*, JCAP **0908** (2009) 023, arXiv:0904.0961
- [3] Bamba K., Nojiri Sh., Odintsov S.D., Sasaki M., *Screening of cosmological constant for De Sitter Universe in non-local gravity, phantom-divide crossing and finite-time future singularities*, Gen. Rel. Grav. **44** (2012) 1321–1356, arXiv:1104.2692

---

<sup>1</sup>If the model includes the perfect fluid with  $w_m = -1/3$  the form of  $f$  is more complicated (formula (17)).

- [4] Nojiri Sh., Odintsov S.D., *Modified non-local- $F(R)$  gravity as the key for the inflation and dark energy*, Phys. Lett. B **659** (2008) 821, arXiv:0708.0924
- [5] Elizalde E., Pozdeeva E.O., Vernov S.Yu., *De Sitter universe in nonlocal gravity*, Phys. Rev. D **85** (2008) 044002, arXiv:1110.5806
- [6] Biswas T., Mazumdar A., Siegel W., *Bouncing Universes in String-inspired Gravity*, JCAP **0603** (2006) 009, arXiv:hep-th/0508194;  
 Biswas T., Koivisto T., Mazumdar A., *Towards a resolution of the cosmological singularity in non-local higher derivative theories of gravity*, JCAP **1011** (2010) 008, arXiv:1005.0590;  
 Koshelev A.S., Vernov S.Yu., *On bouncing solutions in non-local gravity*, arXiv:1202.1289
- [7] Aref'eva I.Ya., Joukovskaya L.V., Vernov S.Yu., *Dynamics in nonlocal linear models in the Friedmann–Robertson–Walker metric*, J. Phys. A: Math. Theor. **41** (2008) 304003, arXiv:0711.1364;  
 Vernov S.Yu., *Localization of the SFT inspired Nonlocal Linear Models and Exact Solutions*, Phys. Part. Nucl. Lett. **8** (2011) 310–320, arXiv:1005.0372
- [8] Zhang Y.l., Sasaki M., *Screening of cosmological constant in non-local cosmology*, Int. J. Mod. Phys. D **21** (2012) 1250006, arXiv:1108.2112
- [9] Biswas T., Cembranos J.A.R., Kapusta J.I., *Finite Temperature Solitons in Non-Local Field Theories from  $p$ -Adic Strings*, Phys. Rev. D **82** (2010) 085028, arXiv:1006.4098