Reply to "Comment on 'Vortex-assisted photon counts and their magnetic field dependence in single-photon superconducting detectors"

L. N. Bulaevskii, Matthias Graf¹ and V.G. Kogan²

¹Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA and

² Ames National Laboratory, Ames, Iowa 50011, USA

(Dated: October 25, 2018)

1. In the Comment by Gurevich and Vinokur¹ and in Refs. 2–4, the vortex crossing rate was calculated in the framework of the London theory of superconductivity treating vortex as a particle. It allows one to derive the energy of a single vortex in thin-film superconducting strips as presented in Eq. (1) of the Comment. However, one drawback of the London theory is that one needs to introduce a cutoff ξ_{cut} , due to the size of the vortex core, of the order of the coherence length $\xi(T)$ at the temperature T. Since the London theory treats vortices as pointlike objects with a phase, Eq. (1) in the Comment is not valid at distances smaller than $\xi(T)$ from the strip edges. The cutoff, an inherent short-coming of the model, cannot be fixed within the London theory. On the other hand, the vortex crossing rate, $R_v(I,T)$, is very sensitive to the value of ξ_{cut} :

$$R_v(I,T) \approx \Omega(I,T,\nu) \left(\frac{e\pi I}{2I_0}\right)^{\nu+1} \left(\frac{\xi_{cut}}{w}\right)^{\nu+1}, \quad (1)$$

where $\Omega(I,T,\nu) \approx 4\pi T c^2 R_{\rm eff}/(e^2\Phi_0^2)(\nu/2\pi)^{1/2}$ is the attempt frequency. The characteristic current is $I_0=c\Phi_0/(8\pi\Lambda)$, w is the film's width, $R_{\rm eff}$ is the effective resistance and $e\simeq 2.718$. The exponent $\nu\approx 110$ was obtained in Ref. 3 from measurements of $R_v(I,T)$ on NbN.⁵ Thus a change in ξ_{cut} by 10% results in a large change of $R_v(I,T)$ by the factor 3×10^4 . In Refs. 2 and 4 different values for ξ_{cut} were used, thus resulting in very different crossing rates. We stress that without employing a microscopic theory neither the London, nor the Ginzburg-Landau theories are sufficient to obtain ξ_{cut} accurately at temperatures well below T_c , where experiments of interest are performed.

In this situation, we can only estimate the cutoff parameter ξ_{cut} from the data by Bartolf $et~al.^5$ for $R_v(I,T)$. We obtain $\xi_{cut}(5.5{\rm K})\approx 3.9\,{\rm nm}$ for sample $1\,(T_c=12.73\,{\rm K})$. This value should be compared with $\xi(5.5{\rm K})\approx 3.2\,{\rm nm}$ estimated from H_{c2} measurements, see Fig. 3 of Ref. 5. For sample 2 the results are similar. On the other hand, according to Gurevich and Vinokur, $\xi_{cut}(5.5{\rm K})\approx 3.9\,{\rm nm}/(2\times0.34)=5.7\,{\rm nm}$. We do not think that "our" cutoff is any better justified than "theirs". As argued above, the discussion about the proper definition of ξ_{cut} cannot be settled within the framework of the London theory.

- 2. The concept of "vortex as a particle" is a rather crude approximation to describe the vortex energy near the film edges. It is questionable for the process of vortex entry and exit, because such processes occur on the length scale of $\xi(T)$. In our view, only a microscopic theory can provide the correct description. We do not think that the "vortex as a particle" model can be improved significantly irrespective of the boundary conditions used. Moreover, we are not aware of a convincing argument in favor of the periodic boundary conditions preferred by the authors of the Comment. Hence, we do not think that the corrections, presented in the Comment, improve this situation. To put this all in perspective, one should keep in mind that the uncertainty in the factor $(\xi_{cut}/w)^{110}$ is much bigger than any corrections to the attempt frequency Ω , due to a particular choice of boundary conditions.
- 3. In Refs. 2 and 4 the single vortex crossing rate was derived for uncorrelated crossings along the strip length L. Vortices in the strip are noninteracting (uncorrelated) at distances bigger than w. That is the reason why we evaluated the number of uncorrelated crossings of single vortices as L/w.

¹ A. Gurevich and V. M. Vinokur, arXiv:1201.5347, January 2012.

² L. N. Bulaevskii, M. J. Graf, and V. G. Kogan, Phys. Rev. B 85, 014505 (2012).

³ L. N. Bulaevskii, M. J. Graf, C. D. Batista, and V. G. Kogan, Phys. Rev. B 83, 144526 (2011).

⁴ A. Gurevich and V. M. Vinokur, Phys. Rev. Lett. **100**, 227007 (2008).

⁵ H. Bartolf, A. Engel, A. Schilling, K. Il'in, M. Siegel, H.-W. Hübers and A. Semenov, Phys. Rev. B **81**, 024502 (2010).