

EKHARA — new gamma gamma generator

Henryk Czyż^a, Sergiy Ivashyn^{b,*}

^aA. Chelkowski Institute of Physics, University of Silesia, Katowice PL-40007, Poland

^bITP, NSC “Kharkov Institute of Physics and Technology”, Kharkov UA-61108, Ukraine

Abstract

The Monte Carlo generator EKHARA allows to simulate the process $e^+e^- \rightarrow e^+e^-\pi^0$ at energies of meson factories. We review the motivation, describe the current status of the generator, and discuss features, which will appear in the forthcoming releases: the two-photon production of η and η' mesons.

Keywords: Monte Carlo generator, Pion transition form factor, Pion pair production, Two-photon processes, e^+e^- annihilation

1. Introduction

The two-photon transition form factors of the light pseudoscalar mesons ($\mathcal{P} = \pi^0, \eta, \eta'$) are being intensively discussed during the last years. Recently the BaBar experiment has provided us with new information in the region where the virtuality of the first photon is high (Q^2) and that of the second one is small (and both photons are spacelike) [1, 2]. The process, which gives an access to the given kinematic region, is $e^+e^- \rightarrow e^+e^-\mathcal{P}$ in the “single-tag” mode, see Fig. 1 (left). Within the range of the previous experiments (CELLO [3], CLEO [4]) there was a fair agreement with the theory. The results of BaBar on the pion transition form factor [1] demonstrated an unexpected deviation from the established theoretical predictions and has triggered the reconsideration of the theoretical description of these form factors [5, 6, 7, 8, 9, 10, 11, 12, 13, 14]. Over the last year several theoretical advances appeared, which were able (or at least claimed to be able) to reconcile theory and experiment (e.g., [6, 15]) for the π^0, η and η' simultaneously. Nevertheless, there are also indications that the data of Ref. [1] on π^0 cannot be easily accommodated by theory (see [7, 10, 11, 12, 16, 17]). Therefore, there are still open questions and further research is foreseen both on the theory and the experiment side.

The new experiments, e.g., KLOE-2 [18, 19], and BES-III [20] will soon start measuring the two-photon form factors of the light pseudoscalars and related observables. In case of a double-tag experiment, see Fig. 1 (right), the form factor as a function of both photon virtualities is potentially of interest. Currently, the

double-tag experiments are the one performed with the KEDR detector at Novosibirsk [21] and the one performed by KLOE-2 Collaboration at Frascati [18, 19]. Unfortunately, in the above experiments the shape of the form factor as a function of both photon virtualities is not accessible. Notice that the form factor unknown dependence on the second photon virtuality has possible impact on the precision of the experimental extraction of the quoted values of the form factor even in case of a single-tag experiment, as has been stressed in [22] and studied in [23]. In any case, a realistic simulation has to account for both photon virtualities.

The transition form factor description is not only a milestone in our understanding of the $U(1)$ axial anomaly [9], general principles of the strong interaction dynamics [11] and $SU(3)$ flavor symmetry breaking [24], but it is also very important in the context of the Standard Model tests at low energy [5], mainly in the problem of the anomalous magnetic moment of the muon a_μ . The transition form factors (though in a different kinematic region) are required for the calculation of the hadronic light-by-light (hLbyL) scattering part of the a_μ usually denoted as a_μ^{LbyL} [25, 26, 27, 28, 29, 30]. In view of the new $g-2$ experiments to be carried out at Fermilab [31] and JPARC [32], it is crucial to improve the accuracy of a_μ^{LbyL} ¹. In context of hLbyL the improvement of the models for the two-photon form factors is important, which requires the high precision of the form factor measurements [19]. In turn, precision measurements require precision simulation.

¹ This subject has been a topic of the “INT Workshop on the Hadronic Light-by-Light Contribution to the Muon Anomaly” (Seattle, February 28 – March 4, 2011) <http://www.int.washington.edu/PROGRAMS/11-47w/>

*Corresponding author

Email address: s.ivashyn@gmail.com (Sergiy Ivashyn)




Figure 1: The concept of single-tag (left) and double-tag (right) $e^+e^- \rightarrow e^+e^-\mathcal{P}$ experiment.

Summing up, the state of the art is that both theory and experiment need a computer program for the process $e^+e^- \rightarrow e^+e^-\mathcal{P}$ – the Monte Carlo generator, which would allow to simulate the events under the conditions similar to the experimental ones. Our objective is to develop and to maintain a free and public tool for $e^+e^- \rightarrow e^+e^-\mathcal{P}$ simulation (π^0, η, η') to be used by experimentalists and theorists. The guiding principles are the following:

- to account for light meson phenomenology in detail via precise modeling of the form factors, in agreement with available data;
- full description of both virtual photons;
- high generator efficiency: accounting for the peaking behaviour of the cross section.

This approach is being realized in the Monte Carlo generator EKHARA [33, 34]². In the following Section we review the current status of this program.

2. EKHARA Monte Carlo generator

The Monte Carlo generator EKHARA is a computer program written in FORTRAN using the quadruple precision numbers. The principal description of the program is given in the dedicated paper [33]. The generator can be used a stand-alone code or be interfaced to another (e.g., an experiment's) simulation program.

The first version of EKHARA [35, 34] allows to simulate the reaction $e^+e^- \rightarrow \pi^+\pi^-e^+e^-$ with the focus on a specific event selection, namely that of the radiative return method for cross section $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$ measurement. It was used for the background [36] simulation by KLOE Collaboration [37, 38]. This channel

can be important for the radiative return program of the BES-III experiment [39, 20]. The $e^+e^- \rightarrow \pi^+\pi^-e^+e^-$ mode in EKHARA has the following features, which a user should be aware of: it generates weighted events and the applied phenomenology of the $\gamma^*\gamma^* \rightarrow \pi^+\pi^-$ subprocess is *not* tuned to reflect the line shapes known from the dedicated $\gamma^*\gamma^* \rightarrow \pi^+\pi^-$ experiments. An improvement of the above issue is on the agenda.

The second version of EKHARA [33] features a new channel, $e^+e^- \rightarrow e^+e^-\mathcal{P}$. The implementation of this channel and the applied phenomenology is oriented on the $\gamma^*\gamma^*$ physics. The phase space construction and the matrix element calculation is based on the exact formulae and the exact kinematics (in contrast to the Equivalent Photon Approximation, which was used in many generators in this field).

It is important to stress that the (user-provided) form factors in EKHARA have to be meaningful both in the time-like and space-like regions of the photon, because both s - and t -channel amplitudes and their interference can be switched on in the simulation, see Fig. 2. In general, the provided form factors have to be functions of *two* photon virtualities. The above criteria considerably reduces the user's choice, because the majority of the published formulae for the form factors within different theoretical approaches hold only for the case with one photon being real and the other — space-like and virtual. Strictly speaking, from the point of view of the numerical simulation, such a case is very specific and is *never* realized for the t -channel amplitude because the leptons are not massless.

In order to support a user, we have recently [23] supplied EKHARA with the transition form factors for π^0, η and η' mesons. Inclusion of η and η' meson production in EKHARA is a new feature with respect to [33]. The form factor modeling [23] was performed in the framework of the effective field theory, namely the chiral theory with resonances [40, 41, 42] was used. The

² EKHARA web page: <http://prac.us.edu.pl/~ekhara>




Figure 2: The s - and t -channel amplitudes in $e^+e^- \rightarrow e^+e^-\mathcal{P}$.

quality of the model is good: by fitting only one free parameter to the data for π^0 , η and η' form factors (simultaneously) we achieved $\chi^2/d.o.f. \approx 1.2$. The existing data from single-tag experiments in space-like region (CELLO [3], CLEO [4] and BaBar [1, 2]) were used not only as a reference for model improvement, but also for the generator verification. The verification is done by comparison of the simulated cross sections $\frac{d\sigma}{dQ^2}$ for $e^+e^- \rightarrow e^+e^-\mathcal{P}$ under the event selection similar to that of the single-tag experiments. The pattern of data/simulation agreement is the same as for the form factors and this is an important consistency check.

Recently, EKHARA was used by KLOE-2 Collaboration [19] for the feasibility studies of $e^+e^- \rightarrow e^+e^-\pi^0$. On the basis of simulation, a per cent level of precision is foreseen for the measurement of $\Gamma_{\pi^0 \rightarrow \gamma\gamma}$ by the KLOE-2 experiment at Frascati. KLOE-2 experiment can also perform the first measurement of pion transition form factor $F_{\pi^0\gamma^*\gamma}(Q^2)$ in the space-like region in the vicinity of the origin: $0.01 < Q^2 < 0.1 \text{ GeV}^2$. The impact of the proposed measurements on the evaluation of the Standard Model prediction for the anomalous magnetic moment of the muon, a_μ , was estimated [19].

3. Summary

A brief review of the Monte Carlo event generator EKHARA is presented with the focus on the physics motivation and several new features. It can be used for the simulation of the processes $e^+e^- \rightarrow e^+e^-\mathcal{P}$ and $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$. The channels $e^+e^- \rightarrow e^+e^-\pi^0$, η , η' are important for $\gamma^*\gamma^*$ physics programs of the modern experiments (e.g., KLOE-2 and BES III) and can be used for the simulation of the two-photon form factor measurements at meson factories. The program is supplied with the form factors for π^0 , η and η' mesons in agreement with the modern data on the transition form factors and corresponding differential cross sections $\frac{d\sigma}{dQ^2}$.

The future plans of EKHARA development include the electromagnetic radiative corrections for the channel $e^+e^- \rightarrow e^+e^-\mathcal{P}$. After the inclusion of the radiative corrections we plan to perform a detailed comparison of our generator with GGRESRC [43], which was used by the BaBar experiment [1, 2]. The major update of the $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$ channel in context of possible $\gamma^*\gamma^*$ physics motivation is also foreseen.

Acknowledgments

This research was partly supported by Polish Ministry of Science and High Education from budget for science for years 2010-2013: grant number N N202 102638, by Sonderforschungsbereich SFB1044 of the Deutsche Forschungsgemeinschaft and by National Academy of Science of Ukraine contract 50/53—2011. This work is a part of the activity of the “Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies”³.

References

- [1] B. Aubert, et al., Measurement of the $\gamma\gamma^* \rightarrow \pi^0$ transition form factor, Phys. Rev. D80 (2009) 052002. arXiv:0905.4778, doi:10.1103/PhysRevD.80.052002.
- [2] P. del Amo Sanchez, et al., Measurement of the $\gamma\gamma^* \rightarrow \eta$ and $\gamma\gamma^* \rightarrow \eta'$ transition form factors, Phys.Rev. D84 (2011) 052001. arXiv:1101.1142, doi:10.1103/PhysRevD.84.052001.
- [3] H. J. Behrend, et al., A Measurement of the π^0 , η and η' electromagnetic form-factors, Z. Phys. C49 (1991) 401–410. doi:10.1007/BF01549692.
- [4] J. Gronberg, et al., Measurements of the meson photon transition form factors of light pseudoscalar mesons at large momentum transfer, Phys. Rev. D57 (1998) 33–54. arXiv:hep-ex/9707031, doi:10.1103/PhysRevD.57.33.
- [5] A. E. Dorokhov, How the recent BABAR data for $P \rightarrow \gamma\gamma^*$ affect the Standard Model predictions for the rare decays $P \rightarrow l^+l^-$, JETP Lett. 91 (2010) 163–169. arXiv:0912.5278, doi:10.1134/S0021364010040016.
- [6] P. Kroll, The form factors for the photon to pseudoscalar meson transitions - an update, Eur.Phys.J. C71 (2011) 1623. arXiv:1012.3542, doi:10.1140/epjc/s10052-011-1623-4.
- [7] S. J. Brodsky, F.-G. Cao, G. F. de Teramond, Meson Transition Form Factors in Light-Front Holographic QCD, Phys.Rev. D84 (2011) 075012. arXiv:1105.3999, doi:10.1103/PhysRevD.84.075012.
- [8] S. J. Brodsky, F.-G. Cao, G. F. de Teramond, Evolved QCD predictions for the meson-photon transition form factors, Phys. Rev. D84 (2011) 033001. arXiv:1104.3364, doi:10.1103/PhysRevD.84.033001.
- [9] Y. N. Klopot, A. G. Oganesian, O. V. Teryaev, Axial anomaly and mixing: from real to highly virtual photons, Phys.Rev. D84 (2011) 051901. arXiv:1106.3855, doi:10.1103/PhysRevD.84.051901.

³<http://www.inf.infn.it/wg/sighad/>

- [10] I. Balakireva, W. Lucha, D. Melikhov, Pion elastic and (π^0, η, η') $\rightarrow \gamma\gamma^*$ transition form factors in a broad range of momentum transfers, [arXiv:1110.6904](#).
- [11] A. Bakulev, S. Mikhailov, A. Pimikov, N. Stefanis, Pion-photon transition: The New QCD frontier, *Phys.Rev. D84* (2011) 034014. [arXiv:1105.2753](#), [doi:10.1103/PhysRevD.84.034014](#).
- [12] N. Stefanis, A. P. Bakulev, S. Mikhailov, A. Pimikov, Pion-Photon Transition Form Factor and Pion Distribution Amplitude in QCD: Facing the Enigmatic Behavior of the BaBar Data, [arXiv:1111.7137](#).
- [13] A. Dorokhov, Pseudoscalar meson form factors and decays, [arXiv:1109.3754](#).
- [14] Y. Bystritskiy, V. Bytev, E. Kuraev, A. Ilyichev, A model of a transition neutral pion form factor measured in annihilation and scattering channels at high momentum transfer, *Phys.Part.Nucl.Lett.* 8 (2011) 73–77. [arXiv:0912.3668](#), [doi:10.1134/S154747711102004X](#).
- [15] S. Noguera, S. Scopetta, The eta-photon transition form factor, [arXiv:1110.6402](#).
- [16] A. Bakulev, S. Mikhailov, A. Pimikov, N. Stefanis, Pion-photon transition form factor using light-cone sum rules: theoretical results, expectations, and a global-data fit, [arXiv:1108.4344](#).
- [17] W. Lucha, D. Melikhov, The puzzle of the $\pi \rightarrow \gamma\gamma^*$ transition form factor, [arXiv:1110.2080](#).
- [18] G. Amelino-Camelia, et al., Physics with the KLOE-2 experiment at the upgraded DAΦNE, *Eur. Phys. J. C68* (2010) 619–681. [arXiv:1003.3868](#), [doi:10.1140/epjc/s10052-010-1351-1](#).
- [19] D. Babusci, H. Czyz, F. Gonnella, S. Ivashyn, M. Mascolo, et al., On the possibility to measure the $\pi^0 \rightarrow \gamma\gamma$ decay width and the $\gamma^*\gamma \rightarrow \pi^0$ transition form factor with the KLOE-2 experiment, [arXiv:1109.2461](#).
- [20] D. Asner, T. Barnes, J. Bian, I. Bigi, N. Brambilla, et al., Physics at BES-III, *Int.J.Mod.Phys. A24* (2009) S1–794. [arXiv:0809.1869](#).
- [21] V. Zhilich, KEDR tagger, Talk at PhiPsi11 Workshop.
- [22] P. Lichard, Vector meson dominance and the π^0 transition form factor, *Phys. Rev. D83* (2011) 037503. [arXiv:1012.5634](#), [doi:10.1103/PhysRevD.83.037503](#).
- [23] H. Czyż, S. Ivashyn, A. Korchin, O. Shekhovtsova, Two-photon form factors of the π^0 , η and η' mesons in the chiral theory with resonances, *IFIC/11-63, FTUV/2011-1110*. [arXiv:1202.1171](#).
- [24] T. Feldmann, P. Kroll, Flavor symmetry breaking and mixing effects in the $\eta\gamma$ and $\eta'\gamma$ transition form factors, *Eur. Phys. J. C5* (1998) 327–335. [arXiv:hep-ph/9711231](#), [doi:10.1007/s100520050276](#).
- [25] K. Melnikov, A. Vainshtein, Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment revisited, *Phys. Rev. D70* (2004) 113006. [arXiv:hep-ph/0312226](#), [doi:10.1103/PhysRevD.70.113006](#).
- [26] A. E. Dorokhov, W. Broniowski, Pion pole light-by-light contribution to $g - 2$ of the muon in a nonlocal chiral quark model, *Phys. Rev. D78* (2008) 073011. [arXiv:0805.0760](#), [doi:10.1103/PhysRevD.78.073011](#).
- [27] J. Prades, E. de Rafael, A. Vainshtein, Hadronic Light-by-Light Scattering Contribution to the Muon Anomalous Magnetic Moment (2009) 303–317 In Roberts, Lee B., Marciano, William J. (eds.): *Lepton dipole moments (Advanced series on directions in high energy physics. 20)*. [arXiv:0901.0306](#).
- [28] A. Nyffeler, Hadronic light-by-light scattering in the muon $g - 2$: a new short-distance constraint on pion-exchange, *Phys. Rev. D79* (2009) 073012. [arXiv:0901.1172](#), [doi:10.1103/PhysRevD.79.073012](#).
- [29] F. Jegerlehner, A. Nyffeler, The Muon $g - 2$, *Phys. Rept.* 477 (2009) 1–110. [arXiv:0902.3360](#), [doi:10.1016/j.physrep.2009.04.003](#).
- [30] A. E. Dorokhov, A. E. Radzhabov, A. S. Zhevlakov, The pseudoscalar hadronic channel contribution of the light-by-light process to the muon ($g - 2$) within the nonlocal chiral quark model, *Eur. Phys. J. C71* (2011) 1702. [arXiv:1103.2042](#), [doi:10.1140/epjc/s10052-011-1702-6](#).
- [31] R. M. Carey, et al., The New ($g - 2$) Experiment: A proposal to measure the muon anomalous magnetic moment to ± 0.14 ppm precision, FERMILAB-PROPOSAL-0989.
- [32] T. Mibe, J-PARC $g - 2$ collaboration, New $g - 2$ experiment at J-PARC, *Chinese Physics C 34* (6) (2010) 745.
- [33] H. Czyż, S. Ivashyn, EKHARA: A Monte Carlo generator for $e^+e^- \rightarrow e^+e^-\pi^0$ and $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$ processes, *Comput. Phys. Commun.* 182 (2011) 1338–1349. [arXiv:1009.1881](#), [doi:10.1016/j.cpc.2011.01.029](#).
- [34] H. Czyż, E. Nowak-Kubat, The reaction $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$ and the pion form factor measurements via the radiative return method, *Phys. Lett. B634* (2006) 493–497. [arXiv:hep-ph/0601169](#), [doi:10.1016/j.physletb.2006.02.024](#).
- [35] H. Czyż, E. Nowak-Kubat, Radiative return via electron pair production: Monte Carlo simulation of the process $e^+e^- \rightarrow \pi^+\pi^-e^+e^-$, *Acta Phys. Polon. B36* (2005) 3425–3434. [arXiv:hep-ph/0510287](#).
- [36] H. Czyż, E. Nowak, $e^+e^- \rightarrow \pi^+\pi^-e^+e^-$: A potential background for $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$ measurement via radiative return method, *Acta Phys. Polon. B34* (2003) 5231–5238. [arXiv:hep-ph/0310335](#).
- [37] A. Aloisio, et al., Measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma)$ and extraction of $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$ below 1 GeV with the KLOE detector, *Phys. Lett. B606* (2005) 12–24. [arXiv:hep-ex/0407048](#), [doi:10.1016/j.physletb.2004.11.068](#).
- [38] F. Ambrosino, et al., Measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ and the dipion contribution to the muon anomaly with the KLOE detector, *Phys. Lett. B670* (2009) 285–291. [arXiv:0809.3950](#), [doi:10.1016/j.physletb.2008.10.060](#).
- [39] S. Actis, et al., Quest for precision in hadronic cross sections at low energy: Monte Carlo tools vs. experimental data, *Eur. Phys. J. C66* (2010) 585–686. [arXiv:0912.0749](#), [doi:10.1140/epjc/s10052-010-1251-4](#).
- [40] G. Ecker, J. Gasser, H. Leutwyler, A. Pich, E. de Rafael, Chiral Lagrangians for Massive Spin 1 Fields, *Phys. Lett. B223* (1989) 425. [doi:10.1016/0370-2693\(89\)91627-4](#).
- [41] G. Ecker, J. Gasser, A. Pich, E. de Rafael, The Role of Resonances in Chiral Perturbation Theory, *Nucl. Phys. B321* (1989) 311. [doi:10.1016/0550-3213\(89\)90346-5](#).
- [42] J. Prades, Massive spin 1 field chiral Lagrangian from an extended Nambu-Jona-Lasinio model of QCD, *Z. Phys. C63* (1994) 491–506. [arXiv:hep-ph/9302246](#), [doi:10.1007/BF01580330](#).
- [43] V. P. Druzhinin, L. A. Kardapoltsev, V. A. Tayursky, The event generator for the two-photon process $e^+e^- \rightarrow e^+e^-R$ ($J^{PC} = 0^{++}$) in the single-tag mode, [arXiv:1010.5969](#).