

Partial spectral multipliers and partial Riesz transforms for degenerate operators

A.F.M. ter Elst¹ and E.M. Ouhabaz²

Abstract

We consider degenerate differential operators $A = \sum_{k,j=1}^d \partial_k(a_{kj}\partial_j)$ on $L^2(\mathbb{R}^d)$ with real symmetric bounded measurable coefficients. Given a function $\chi \in C_b^\infty(\mathbb{R}^d)$ (respectively, Ω a bounded Lipschitz domain) and suppose that $(a_{kj}) \geq \mu > 0$ a.e. on $\text{supp } \chi$ (resp., a.e. on Ω). We prove a spectral multiplier type result: if $F: [0, \infty) \rightarrow \mathbb{C}$ is such that $\sup_{t>0} \|\varphi(\cdot)F(t)\|_{C^s} < \infty$ for some non-trivial function $\varphi \in C_c^\infty(0, \infty)$ and some $s > d/2$ then $M_\chi F(I + A)M_\chi$ is weak type $(1, 1)$ (resp. $P_\Omega F(I + A)P_\Omega$ is weak type $(1, 1)$). We also prove boundedness on L^p for all $p \in (1, 2]$ of the partial Riesz transforms $M_\chi \nabla(I + A)^{-1/2} M_\chi$. The proofs are based on a criterion for a singular integral operator to be weak type $(1, 1)$.

January 2012

AMS Subject Classification: 42B15, 45F05.

Keywords: Spectral multipliers, Riesz transforms, singular integral operators, degenerate operators, Gaussian bounds.

Home institutions:

- | | |
|--|---|
| 1. Department of Mathematics
University of Auckland
Private bag 92019
Auckland
New Zealand | 2. Univ. Bordeaux,
IMB, CNRS UMR 5251,
351, Cours de la Libération
33405 Talence
France |
|--|---|

1. INTRODUCTION

Let A be a non-negative self-adjoint uniformly elliptic operator in divergence form. More precisely, let $a_{kj} = a_{jk}: \mathbb{R}^d \rightarrow \mathbb{R}$ be a bounded measurable function for all $j, k \in \{1, \dots, d\}$ and assume that there exists a $\mu > 0$ such that

$$(1) \quad \sum_{k,j=1}^d a_{kj}(x) \xi_k \xi_j \geq \mu |\xi|^2 \text{ for all } \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d \text{ and } x \in \mathbb{R}^d$$

The operator $A = - \sum_{k,j=1}^d \partial_k (a_{kj} \partial_j)$, defined by quadratic form techniques, is self-adjoint on $L^2(\mathbb{R}^d)$. It is a standard fact that $-A$ is the generator of a strongly continuous semigroup $(e^{-tA})_{t>0}$ on $L^2(\mathbb{R}^d)$. The well known Aronson estimates assert that e^{-tA} is given an integral kernel p_t (called the heat kernel of A) which satisfies the Gaussian upper bound:

$$(2) \quad |p_t(x, y)| \leq C t^{-d/2} e^{-c \frac{|x-y|^2}{t}} \text{ for all } t > 0 \text{ and } x, y \in \mathbb{R}^d.$$

Here C and c are positive constants.

In recent years, harmonic analysis of operator of type A has attracted a lot of attention and substantial progress have been made in which upper bounds for the heat kernel play a fundamental role. We mention for example the theory of Hardy and BMO spaces associated with such operators (see for example [DY] and [HLM MY]), spectral multipliers [DOS] and Riesz transforms (see [DM1], [Aus], [Ouh], [She] and the references therein). Concerning spectral multipliers, it is known that if $F: [0, \infty) \rightarrow \mathbb{C}$ is a bounded measurable function then the operator $F(A)$, which is well defined on L^2 by spectral theory, extends to a bounded operator on L^p for all $1 < p < \infty$ provided F satisfies the condition

$$(3) \quad \sup_{t>0} \|\varphi(\cdot) F(t)\|_{C^s} < \infty$$

for some $s > d/2$ and some non-trivial auxiliary function $\varphi \in C_c^\infty(0, \infty)$. See Duong–Ouhabaz–Sikora [DOS], where a more general result is proved. Note that condition (3) is satisfied if F has $[d/2] + 1$ derivatives such that

$$\sup_{\lambda>0} \lambda^k |F^{(k)}(\lambda)| < \infty \text{ for all } k \in \{0, 1, \dots, [d/2] + 1\}.$$

As an example, one obtains polynomial estimates on L^p for imaginary powers of type $\|A^{is}\|_{p \rightarrow p} \leq C(1+|s|)^{\beta_p}$ for all $\beta_p > d|\frac{1}{2} - \frac{1}{p}|$. Taking $F(\lambda) := (1 - \frac{\lambda}{R})_+^\alpha$ one obtains Bochner–Riesz summability for all $\alpha > d/2$.

Concerning Riesz transforms $\mathcal{R}_k := \partial_k A^{-1/2}$, it is an obvious consequence of the ellipticity assumption (1) that \mathcal{R}_k is bounded on $L^2(\mathbb{R}^d)$ for all $k \in \{1, \dots, d\}$. As for multiplier results, the Gaussian bound (2) combined with recent developments on singular integral operators allow to prove that \mathcal{R}_k is bounded on $L^p(\mathbb{R}^d)$ for all $p \in (1, 2)$ with the sole assumption (1) and bounded measurable coefficients (see Duong–McIntosh [DM1], Auscher [Aus], Ouhabaz [Ouh]). Under weak regularity assumption on the coefficients one obtains boundedness of \mathcal{R}_k on $L^p(\mathbb{R}^d)$ for all $p \in (2, \infty)$ (cf. Auscher [Aus], Shen [She]).

In the present paper we wish to study similar problems for degenerate operators. Instead of (1) we merely assume that

$$(4) \quad \sum_{k,j=1}^d a_{kj}(x) \xi_k \xi_j \geq 0 \text{ for all } \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d \text{ and } x \in \mathbb{R}^d.$$

In this case, we define the form

$$(5) \quad \mathfrak{a}_0(u, v) = \sum_{k,j=1}^d \int_{\mathbb{R}^d} a_{kj} (\partial_j u) (\partial_k v)$$

with form domain $D(\mathfrak{a}_0) = C_c^\infty(\mathbb{R}^d)$. If this form is closable, then A will be the self-adjoint operator associated with its closure. If not, we take the regular part and consider A as the operator associated with the closure of this regular part (see [Sim] and [AE]).

Proving results like the previous ones for these operators seems unattainable because Gaussian (or Poisson) upper bounds are not true in general. Even L^1 - L^∞ estimates of e^{-tA} are not valid in general. What we will do is to restrict the operators to parts where the matrix (a_{kj}) is elliptic. It is proved by ter Elst and Ouhabaz [EO1] that if $\chi \in C_b^\infty(\mathbb{R}^d)$ and $\mu > 0$ are such that $(a_{kj}(x)) \geq \mu I$ for a.e. $x \in \text{supp } \chi$, then $M_\chi e^{-tA} M_\chi$ has a Hölder continuous kernel K_t which satisfies the Gaussian bound

$$(6) \quad |K_t(x, y)| \leq C t^{-d/2} e^{-c \frac{|x-y|^2}{t}} (1+t)^{d/2} \text{ for all } t > 0 \text{ and } x, y \in \mathbb{R}^d.$$

Here M_χ is the multiplication operator by χ . The same result holds for $P_\Omega e^{-tA} P_\Omega$ if Ω is a bounded Lipschitz domain such that $(a_{kj}(x)) \geq \mu I$ for

a.e. $x \in \Omega$ for some $\mu > 0$. Here P_Ω is the multiplication operator by the indicator function $\mathbf{1}_\Omega$ of Ω .

Note that in general one cannot get rid of the extra term $(1+t)^{d/2}$ in the right hand side of (6). For example, if $a_{kj} = \delta_{kj}$ on a smooth bounded domain Ω , then A is the Neumann Laplacian on $L^2(\Omega)$ and 0 on $L^2(\mathbb{R}^d \setminus \Omega)$. It is then clear that the Gaussian bound is not valid without the additional term $(1+t)^{d/2}$. Because of that additional term in (6), we shall consider in the sequel $I + A$ instead of A (of course, one can take $\varepsilon I + A$ for any $\varepsilon > 0$ to absorb the factor $(1+t)^{d/2}$).

For spectral multipliers and Riesz transforms we will prove the following results. Given $\chi \in C_b^\infty(\mathbb{R}^d)$ (resp., Ω a bounded Lipschitz domain) such that $(a_{kj}(x)) \geq \mu I$ for a.e. $x \in \text{supp } \chi$ (resp., for a.e. $x \in \Omega$) for some constant $\mu > 0$. The main theorems of this paper are the following.

Theorem 1.1. *Let $F: [0, \infty) \rightarrow \mathbb{C}$ be a bounded function such that*

$$\sup_{t>0} \|\varphi(\cdot)F(t)\|_{C^s} < \infty$$

for some $s > d/2$ and some non-trivial function $\varphi \in C_c^\infty(0, \infty)$. Then $M_\chi F(I + A)M_\chi$ (resp., $P_\Omega F(I + A)P_\Omega$) is bounded on $L^p(\mathbb{R}^d)$ for all $1 < p < \infty$.

Theorem 1.2. *For all $k \in \{1, \dots, d\}$ the Riesz transform type operator $M_\chi \partial_k (I + A)^{-1/2} M_\chi$ is bounded on $L^p(\mathbb{R}^d)$ for all $1 < p \leq 2$.*

Now we discuss how we prove these results. In the elliptic case, besides the Gaussian bound (2), the proof of boundedness of spectral multipliers or Riesz transforms rely on a criterion proved by Duong and McIntosh [DM2] for singular integral operators to be weak type $(1, 1)$. This criterion says that if T is bounded on L^2 with a (singular) kernel K such that there exists a family of operators $(A_t)_{t>0}$ given by integral kernels a_t which satisfy Gaussian (or Poisson) bounds, TA_t is also given by a (singular) kernel K_t and there are $C, \delta > 0$ such that

$$(7) \quad \int_{|x-y| \geq \delta\sqrt{t}} |K(x, y) - K_t(x, y)| dx \leq C$$

for all $t > 0$ and a.e. y , then T is weak type $(1, 1)$. In applications to spectral multipliers of elliptic operators we start with $T = F(A)$ and one takes $A_t = e^{-tA}$. Therefore, K_t is the kernel of the operator $F(A)e^{-tA}$ which

can be seen as a regularization of $F(A)$. In the degenerate case and because of (6), it is tempting to choose $A_t = M_\chi e^{-t(I+A)} M_\chi$. Then,

$$TA_t = M_\chi F(I+A) M_\chi M_\chi e^{-t(I+A)} M_\chi = M_\chi F(I+A) M_\chi^2 e^{-t(I+A)} M_\chi.$$

Now, the presence of M_χ^2 does not allow to regularize $F(I+A)$ by $e^{-t(I+A)}$. The simple fact that we do not have $F(I+A)$ next to $e^{-t(I+A)}$ in the expression of TA_t destroys this strategy. The same problem occurs for the Riesz transform $M_\chi \partial_k (I+A)^{-1/2} M_\chi$. To overcome this difficulty we prove a version of the Duong–McIntosh criterion that is suitable for our purpose. It reads as follows (see Theorems 2.1 and 2.3 together with Remark 2.2 for precise and quantitative statements).

Theorem 1.3. *Let T be a bounded linear operator on L^2 and $(A_t)_{t>0}$ a family of linear operators which satisfy L^1 - L^2 off-diagonal estimates. Suppose that there exists a bounded linear operator S on L^2 and $\delta, W > 0$ such that*

$$(8) \quad \int_{|x-y| \geq (1+\delta)t} |(T - SA_t)u(y)| dy \leq W \|u\|_1$$

for all $x \in \mathbb{R}^d$, $t > 0$ and $u \in L^1 \cap L^\infty$ supported in the ball $B(x, t)$. Then T is weak type $(1, 1)$.

Note that the estimate (8) is satisfied if T and SA_t are given by (singular) kernels K and K_t and there are $C, \delta > 0$ such that

$$\int_{|x-y| \geq \delta\sqrt{t}} |K(x, y) - K_t(x, y)| dx \leq C$$

for all $t > 0$ and a.e. $y \in \mathbb{R}^d$.

Theorem 1.3 gives more freedom by choosing any appropriate operator S and not necessarily $S = T$. Coming back to spectral multipliers for degenerate operators A , we had $T = M_\chi F(I+A) M_\chi$ and we choose now $S = M_\chi F(I+A)$ and $A_t = e^{-t(I+A)} M_\chi$. Then $TA_t = M_\chi F(I+A) e^{-t(I+A)} M_\chi$ for which we can prove the estimate in Theorem 1.3. Similarly for the Riesz transforms where $T = M_\chi \partial_k (I+A)^{-1} M_\chi$, we take $S = M_\chi \partial_k (I+A)^{-1}$ which turns to be bounded on L^2 and $A_t = e^{-t(I+A)} M_\chi$. We emphasize that $A_t = e^{-t(I+A)} M_\chi$ satisfies L^1 - L^2 off-diagonal estimates but it is not known whether it satisfies Gaussian upper bounds in general¹. We believe that our

¹ under the additional assumption that $a_{kj} \in W^{1,\infty}(\mathbb{R}^d)$, we proved recently in [EO2] that $e^{-t(I+A)} M_\chi$ has a kernel which satisfies a Gaussian bound.

version of the Duong–McIntosh criterion can be used in other circumstances in which a products of several operators come into play. Also, as in [DM2], our version holds for operators on domains of spaces of homogeneous type.

Notation. We fix some notation which we will use throughout this paper. If (X, ρ, μ) is a metric measured space, $x \in X$, $r > 0$ and $j \in \mathbb{N}$, then we denote by $B(x, r) := \{y \in X : \rho(x, y) < r\}$ the open ball of X with centre x and radius r , the annulus $C_j(x, r) = B(x, 2^{j+1}r) \setminus B(x, 2^j r)$ if $j \geq 2$ and $C_1(x, r) = B(x, 4r)$. Let $v(x, r) = \mu(B(x, r))$ be the volume of the ball $B(x, r)$. Next, $\|T\|_{p \rightarrow q}$ is the norm of T as an operator from L^p to L^q . If E is a measurable set, then P_E denotes the multiplication operator by the indicator function $\mathbb{1}_E$ of E . If $s \in (0, \infty) \setminus \mathbb{N}$ then we denote by C^s the space of all Lipschitz functions on $[0, \infty)$ of order s (i.e., functions which are continuously differentiable up to $[s]$ and the derivative of order $[s]$ is Hölder continuous of order $s - [s]$). By $W^{r,p}$ we denote the classical Sobolev spaces on \mathbb{R}^d .

All our operators are linear operators.

We emphasize that we shall use C, C', c, \dots for all inessential constants. A constant C may differ from line to line, even within one line.

2. SINGULAR INTEGRAL OPERATORS

Let (X, μ, ρ) be a metric measured space. We shall assume that $0 < v(x, r) < \infty$ for all $x \in X$ and $r > 0$ and that X is a space of homogeneous type. This means that it satisfies the following doubling condition

$$(9) \quad v(x, 2r) \leq C_0 v(x, r)$$

for some $C_0 > 0$, uniformly for all $x \in X$ and $r > 0$. If (9) is satisfied then there exist positive constants C_1 and d such that

$$(10) \quad v(x, \lambda r) \leq C_1 \lambda^d v(x, r)$$

for all $x \in X$ and $r \geq 1$. Let Ω be an open subset of X . It is endowed with ρ and μ but (Ω, μ, ρ) is not necessarily a space of homogeneous type. Let T be a bounded linear operator on $L^{p_0}(\Omega) := L^{p_0}(\Omega, \mu)$ for some $p_0 \in [1, \infty)$. We say that T is given by a kernel $K: \Omega \times \Omega \rightarrow \mathbb{C}$ if K is measurable and

$$(11) \quad Tu(x) = \int_{\Omega} K(x, y) u(y) d\mu(y)$$

for all $u \in L^{p_0}(\Omega)$ with bounded support and a.e. x outside the support of u . We also say that K is the associated kernel of T . A classical problem in harmonic analysis is to find conditions on the kernel K which allow to extend the operator T from $L^{p_0}(\Omega)$ to other $L^p(\Omega)$ -spaces. Several results are known in this direction. We refer the reader to [Ste], [DM2], [BK], [Aus] and the references therein.

The main result in [DM2] states that if there exists a family of bounded operators A_t , with $t > 0$, which are given by integral kernels a_t satisfying a Gaussian or Poisson estimate and if the associated kernel of $T - TA_t$ does not oscillate too much in a certain sense then T is weak type $(1, 1)$. Here we prove by the same method that if there exists a bounded operator S on $L^{q_0}(X)$ for some $q_0 \in (1, \infty)$ such that the associated kernel of $T - SA_t$ does not oscillate too much then T is weak type $(1, 1)$. As explained in the introduction, this new version gives more freedom by choosing any appropriate S which may not coincide with T . This extension turns out to be powerful to prove spectral multiplier type results as well as Riesz transforms for degenerate operators, whereas it is not clear how to apply the condition from [DM2]. Note also that, following ideas from [BK] and [Aus] we can weaken the assumption on the kernel of A_t . Instead of assuming a Gaussian or Poisson bound, we merely assume an L^1 - L^{q_0} off-diagonal estimate (see (12) below). This difference is again illustrated in our application to degenerate operators. In addition it is possible to formulate the result in [DM2] without reference to the kernels (see also the remark immediately after the next theorem).

We first state and prove the result in the case $\Omega = X$.

Theorem 2.1. *Let T be a non-zero bounded linear operator on $L^{p_0}(X)$ for some $p_0 \in (1, \infty)$. Suppose there exists a bounded linear operator S on $L^{q_0}(X)$ for some $q_0 \in (1, \infty)$, a family of bounded linear operators $(A_t)_{t>0}$ on $L^{q_0}(X)$ and a sequence $(g(j))_{j \in \mathbb{N}}$ in \mathbb{R} such that*

$$(12) \quad \left(\frac{1}{v(x, 2^{j+1}t)} \int_{C_j(x,t)} |A_t f|^{q_0} \right)^{1/q_0} \leq g(j) \frac{1}{v(x, t)} \int_{B(x,t)} |f|$$

for all $x \in X$, $t > 0$, $j \in \mathbb{N}$ and $f \in L^{q_0}(B(x, t))$, and $\sum_{j=1}^{\infty} 2^{jd} g(j) < \infty$. Finally, suppose there exist $\delta, W > 0$ such that

$$(13) \quad \int_{X \setminus B(x, (1+\delta)t)} |(T - SA_t)u| \leq W \|u\|_1$$

for all $x \in X$, $t > 0$ and $u \in L^1(X) \cap L^\infty(X)$ supported in the ball $B(x, t)$. Then T is a weak type $(1, 1)$ operator with

$$(14) \quad \|T\|_{L^1 \rightarrow L^{1,w}} \leq C(1 + \delta)^d (W + \|T\|_{p_0 \rightarrow p_0} + \|S\|_{q_0 \rightarrow q_0}^{q_0} \|T\|_{p_0 \rightarrow p_0}^{1-q_0}).$$

Here C is a constant depending only on the constants in (10). In particular, T extends to a bounded operator on $L^p(X)$ for all $p \in (1, p_0)$.

Remark 2.2. Let $p_0, q_0 \in (1, \infty)$, $T \in \mathcal{L}(L^{p_0}(X))$ and for all $t > 0$ let $S, A_t \in \mathcal{L}(L^{q_0}(X))$. Suppose that T and SA_t have a kernel K and K_t , respectively. Let $\delta, W > 0$ and assume that

$$(15) \quad \int_{\rho(x,y) \geq \delta t} |K(x, y) - K_t(x, y)| d\mu(x) \leq W < \infty,$$

for all $t > 0$ and $y \in X$. Fix now $x \in X$, $t > 0$ and $u \in L^1(X) \cap L^\infty(X)$ supported in the ball $B(x, t)$. Then

$$\begin{aligned} & \int_{X \setminus B(x, (1+\delta)t)} |(T - SA_t)u(y)| d\mu(y) \\ &= \int_{X \setminus B(x, (1+\delta)t)} \left| \int_{B(x, t)} (K(y, z) - K_t(y, z)) u(z) d\mu(z) \right| d\mu(y) \\ &\leq \int_X \int_{\rho(y, z) \geq \delta t} |K(y, z) - K_t(y, z)| d\mu(y) |u(z)| d\mu(z) \\ &\leq W \|u\|_1. \end{aligned}$$

Thus, (13) is satisfied. The condition (15) is the direct analogue of the condition in Duong–McIntosh [DM2].

We also observe that one does not need kernels for both operators T and TA_t but a kernel $H_t(x, y)$ for the difference $T - SA_t$. We may then replace $K(x, y) - K_t(x, y)$ in (15) by $H_t(x, y)$. On the other hand the local estimate (13) which does not appeal to kernels may have advantage of avoiding measurability questions with respect to x and y of the expected singular kernels.

Proof. As mentioned before, the arguments are similar to the arguments used in [DM2] and [Aus]. We give the details for convenience. Recall we denote by C all inessential constants.

We begin by the classical Calderón–Zygmund decomposition. There exist $c, N > 0$ such that the following is valid. Fix $f \in L^1(X) \cap L^\infty(X)$ and

$\alpha > \frac{\|f\|_1}{\mu(X)}$. There exist $g, b_1, b_2, \dots \in L^1(X) \cap L^\infty(X)$ such that

$$f = g + b = g + \sum_i b_i$$

and

- (i) $|g(x)| \leq c\alpha$ for a.e. $x \in X$,
- (ii) each b_i is supported in a ball $B_i = B(x_i, r_i)$ and $\frac{\|b_i\|_1}{v(x_i, r_i)} \leq c\alpha$,
- (iii) $\sum_i v(x_i, r_i) \leq c \frac{\|f\|_1}{\alpha}$, and,
- (iv) there exists a constant N such that $\sum_i \mathbb{1}_{B_i}(x) \leq N$ for a.e. $x \in X$.

See Section III.2 in [CW].

We proceed in several steps.

Step 1 Using the boundedness of T on L^{p_0} we have

$$\begin{aligned} \mu(\{x \in X : |(Tg)(x)| > \alpha\}) &\leq \frac{\|T\|_{p_0 \rightarrow p_0}^{p_0} \|g\|_{p_0}^{p_0}}{\alpha^{p_0}} \\ &\leq C \alpha^{p_0-1} \frac{\|T\|_{p_0 \rightarrow p_0}^{p_0} \|g\|_1}{\alpha^{p_0}} \\ &\leq C \frac{\|T\|_{p_0 \rightarrow p_0}^{p_0} \|g\|_1}{\alpha}. \end{aligned}$$

It follows from (ii) and (iii) above that $\|b\|_1 \leq c\|f\|_1$ and hence $\|g\|_1 \leq (1+c)\|f\|_1$. Therefore,

$$(16) \quad \mu(\{x \in X : |(Tg)(x)| > \alpha\}) \leq C \frac{\|T\|_{p_0 \rightarrow p_0}^{p_0} \|f\|_1}{\alpha}.$$

Step 2 We shall prove that

$$(17) \quad \left\| \sum_i A_{r_i} b_i \right\|_{q_0} \leq C \alpha^{1-\frac{1}{q_0}} \|f\|_1^{1/q_0}.$$

We follow similar arguments as in [Aus]. Fix $u \in L^{q'_0}$ with $\|u\|_{q'_0} = 1$, where q'_0 is the dual exponent of q_0 . Let $i, j \in \mathbb{N}$ and set $C_{i,j} := C_j(x_i, r_i)$. Then

$$\begin{aligned} \int_{C_{i,j}} |A_{r_i} b_i| |u| &\leq \left(\int_{C_{i,j}} |A_{r_i} b_i|^{q_0} \right)^{1/q_0} \left(\int_{C_{i,j}} |u|^{q'_0} \right)^{1/q'_0} \\ &\leq g(j) \frac{v(x_i, 2^{j+1}r_i)^{1/q_0}}{v(x_i, r_i)} \left(\int |b_i| \right) \left(\int_{C_{i,j}} |u|^{q'_0} \right)^{1/q'_0} \\ &\leq c\alpha g(j) v(x_i, 2^{j+1}r_i) \left(\frac{1}{v(x_i, 2^{j+1}r_i)} \int_{C_{i,j}} |u|^{q'_0} \right)^{1/q'_0}, \end{aligned}$$

where we have used (12) and property (ii) in the Calderón–Zygmund decomposition. Denote by \mathcal{M} the Hardy–Littlewood maximal operator. Then

$$\frac{1}{v(x_i, 2^{j+1}r_i)} \int_{C_{i,j}} |u|^{q'_0} \leq \mathcal{M}(|u|^{q'_0})(y)$$

for all $y \in B_i$. Combining the previous inequalities and using the doubling condition (10) one estimates

$$\int_{C_{i,j}} |A_{r_i} b_i| |u| \leq C\alpha 2^{jd} g(j) v(x_i, r_i) \left(\mathcal{M}(|u|^{q'_0})(y) \right)^{1/q'_0}.$$

Taking the integral over $y \in B_i$ gives

$$\int_{C_{i,j}} |A_{r_i} b_i| |u| \leq C\alpha 2^{jd} g(j) \int_{B_i} \left(\mathcal{M}(|u|^{q'_0})(y) \right)^{1/q'_0} d\mu(y).$$

We sum over j, i and use $\sum_j 2^{jd} g(j) < \infty$ together with property (iv) in the Calderón–Zygmund decomposition to obtain

$$\begin{aligned} \int_X \left| \sum_i A_{r_i} b_i \right| |u| &\leq C\alpha \int_X \mathbf{1}_{\cup_i B_i}(y) \left(\mathcal{M}(|u|^{q'_0})(y) \right)^{1/q'_0} d\mu(y) \\ &\leq C\alpha \|\mathbf{1}_{\cup_i B_i}\|_{q_0} \left\| \left(\mathcal{M}(|u|^{q'_0}) \right)^{1/q'_0} \right\|_{q'_0, w} \\ &\leq C\alpha \left(\sum_i v(x_i, r_i) \right)^{1/q_0} \| |u|^{q'_0} \|_1^{1/q'_0}. \end{aligned}$$

Note that we have used the fact that \mathcal{M} is weak type $(1, 1)$ to obtain the last inequality. Using now (iii) of the Calderón–Zygmund decomposition and $\|u\|_{q'_0} = 1$, we obtain (17).

By assumption, S is bounded on L^{q_0} . Hence

$$\mu(\{x \in X : |(S \sum_i A_{r_i} b_i)(x)| > \alpha\}) \leq \frac{1}{\alpha^{q_0}} \|S\|_{q_0 \rightarrow q_0}^{q_0} \sum_i \|A_{r_i} b_i\|_{q_0}^{q_0}.$$

Now we use (17) to obtain

$$(18) \quad \mu(\{x \in X : |(S \sum_i A_{r_i} b_i)(x)| > \alpha\}) \leq \frac{C}{\alpha} \|S\|_{q_0 \rightarrow q_0}^{q_0} \|f\|_1.$$

Step 3 Let δ be as in (13) and for all $i \in \mathbb{N}$ set $Q_i := B(x_i, (1 + \delta)r_i)$, the ball of centre x_i and radius $(1 + \delta)r_i$. Then

$$\begin{aligned}
& \mu(\{x \in X : |\sum_i (T - SA_{r_i})b_i(x)| > \alpha\}) \\
& \leq \sum_i \mu(Q_i) + \mu(\{x \in X \setminus \bigcup_j Q_j : |\sum_i ((T - SA_{r_i})b_i)(x)| > \alpha\}) \\
& \leq C(1 + \delta)^d \sum_i v(x_i, r_i) + \frac{1}{\alpha} \int_{X \setminus \bigcup_j Q_j} |\sum_i ((T - SA_{r_i})b_i)(x)| d\mu(x) \\
& \leq \frac{C(1 + \delta)^d}{\alpha} \|f\|_1 + \frac{1}{\alpha} \sum_i \int_{X \setminus Q_i} |((T - SA_{r_i})b_i)(x)| d\mu(x) \\
& \leq \frac{C(1 + \delta)^d}{\alpha} \|f\|_1 + \frac{W}{\alpha} \sum_i \int |b_i(y)| d\mu(y) \\
& \leq \frac{C(1 + \delta)^d(1 + W)}{\alpha} \|f\|_1.
\end{aligned}$$

Note that the penultimate inequality follows from assumption (13) and the last one from properties (ii) and (iii) in the Calderón–Zygmund decomposition. Hence

$$(19) \quad \mu(\{x \in X : |\sum_i ((T - SA_{r_i})b_i)(x)| > \alpha\}) \leq \frac{C(1 + \delta)^d(1 + W)}{\alpha} \|f\|_1.$$

Step 4 It follows from (16) that

$$\begin{aligned}
& \mu(\{x \in X : |(Tf)(x)| > \alpha\}) \\
& \leq \mu(\{x \in X : |(Tg)(x)| > \frac{\alpha}{2}\}) + \mu(\{x \in X : |(Tb)(x)| > \frac{\alpha}{2}\}) \\
& \leq C \frac{\|T\|_{p_0 \rightarrow p_0}^{p_0}}{\alpha} \|f\|_1 + \mu(\{x \in X : |(Tb)(x)| > \frac{\alpha}{2}\}).
\end{aligned}$$

For the second term we use (18) and (19) to estimate

$$\begin{aligned}
& \mu(\{x \in X : |(Tb)(x)| > \frac{\alpha}{2}\}) \\
& = \mu(\{x \in X : |\sum_i (SA_{r_i}b_i)(x) + \sum_i ((T - SA_{r_i})b_i)(x)| > \frac{\alpha}{2}\})
\end{aligned}$$

$$\begin{aligned}
&\leq \mu(\{x \in X : |(S \sum_i A_{r_i} b_i)(x)| > \frac{\alpha}{4}\}) \\
&\quad + \mu(\{x \in X : |\sum_i ((T - SA_{r_i}) b_i)(x)| > \frac{\alpha}{4}\}) \\
&\leq \frac{C(1 + \delta)^d}{\alpha} (\|S\|_{q_0 \rightarrow q_0}^{q_0} + (1 + W)) \|f\|_1.
\end{aligned}$$

We then conclude that T is of weak type $(1, 1)$ with a weak type estimate

$$(20) \quad \|T\|_{L^1 \rightarrow L^{1,w}} \leq C(1 + \delta)^d (1 + W + \|T\|_{p_0 \rightarrow p_0}^{p_0} + \|S\|_{q_0 \rightarrow q_0}^{q_0}).$$

If we replace T and S by $\|T\|_{p_0 \rightarrow p_0}^{-1} T$ and $\|T\|_{p_0 \rightarrow p_0}^{-1} S$ we obtain (13) with $\|T\|_{p_0 \rightarrow p_0}^{-1} W$ instead of W . Thus applying (20) to $\|T\|_{p_0 \rightarrow p_0}^{-1} T$, $\|T\|_{p_0 \rightarrow p_0}^{-1} S$ and $\|T\|_{p_0 \rightarrow p_0}^{-1} W$ yields (14).

Finally, by Marcinkiewicz interpolation theorem the operator T extends to a bounded operator from $L^p(X) \cap L^{p_0}(X)$ to $L^p(X)$ for all $p \in (1, p_0)$. \square

Following again an idea in [DM2] we can prove a version of the previous theorem on arbitrary domains. Let Ω be an open subset of X and assume that T is bounded on $L^{p_0}(\Omega)$ and S and A_t are bounded on $L^{q_0}(\Omega)$. We define $\tilde{T}: L^{p_0}(X) \rightarrow L^{p_0}(X)$ by

$$\tilde{T}f = \mathbb{1}_\Omega T(\mathbb{1}_\Omega f)$$

and similarly for \tilde{S} and \tilde{A}_t . If A_t satisfies (21) below then \tilde{A}_t satisfies (12). The operator T is weak type $(1, 1)$ if and only if \tilde{T} is weak type $(1, 1)$. Applying the previous theorem to \tilde{T} , \tilde{S} and \tilde{A}_t gives the following result.

Theorem 2.3. *Let T be a non-zero bounded linear operator on $L^{p_0}(\Omega)$ for some $p_0 \in (1, \infty)$. Suppose there exists a bounded operator S on $L^{q_0}(\Omega)$ for some $q_0 \in (1, \infty)$, a family of bounded operators $(A_t)_{t>0}$ on $L^{q_0}(\Omega)$ and a sequence $(g(j))_{j \in \mathbb{N}}$ in \mathbb{R} such that*

$$(21) \quad \left(\frac{1}{v(x, 2^{j+1}t)} \int_{C_j(x,t) \cap \Omega} |A_t f|^{q_0} \right)^{1/q_0} \leq g(j) \frac{1}{v(x, t)} \int_{B(x,t) \cap \Omega} |f|$$

for all ball $x \in \Omega$, $t > 0$, $j \in \mathbb{N}$ and $f \in L^{q_0}(B(x, t) \cap \Omega)$, and $\sum_{j=1}^{\infty} 2^{jd} g(j) < \infty$. Finally, suppose there exist $\delta, W > 0$ such that

$$(22) \quad \int_{\Omega \setminus B(x, (1+\delta)t)} |((T - SA_t)u)(y)| d\mu(y) \leq W \|u\|_1$$

for all $x \in X$, $t > 0$ and $u \in L^1(\Omega) \cap L^\infty(\Omega)$ supported in the ball $B(x, t) \cap \Omega$. Then T is a weak type $(1, 1)$ operator with

$$(23) \quad \|T\|_{L^1(\Omega) \rightarrow L^{1,w}(\Omega)} \leq C(1 + \delta)^d (W + \|T\|_{p_0 \rightarrow p_0} + \|S\|_{q_0 \rightarrow q_0}^{q_0} \|T\|_{p_0 \rightarrow p_0}^{1-q_0}).$$

Here C is a constant depending only on the constants in (10). In particular, T extends to a bounded operator on $L^p(\Omega)$ for all $p \in (1, p_0)$.

As in Remark 2.2 the condition (22) follows if the operators T and SA_t are given by kernels K and K_t (in the sense of (11)) and there are $\delta, W > 0$ such that

$$(24) \quad \int_{\rho(x,y) \geq \delta t} |K(x, y) - K_t(x, y)| d\mu(x) \leq W < \infty,$$

for all $t > 0$ and $y \in \Omega$. It suffices to note that the associated kernel of \widetilde{T} is the extension by 0 outside $\Omega \times \Omega$ of the kernel of T where $\widetilde{T}f = \mathbb{1}_\Omega T(\mathbb{1}_\Omega f)$ as above. Similarly for the kernel of \widetilde{SA}_t .

We may replace in the previous theorems the annulus $C_j(x, r)$ by the annulus $A(x, j, r) := B(x, (j+1)r) \setminus B(x, jr)$. In that case, $v(x, 2^{j+1}r)$ has to be replaced by $v(x, (j+1)r)$ and the condition on g becomes $\sum_j j^d g(j) < \infty$.

Following [BK] it is proved in [Aus] that a bounded operator T on $L^2(X)$ is of weak type (r, r) if

$$(25) \quad \left(\frac{1}{v(x, 2^{j+1}t)} \int_{C_j(x,t)} |T(I - A_t)f|^2 \right)^{1/2} \leq g(j) \left(\frac{1}{v(x, t)} \int_{B(x,t)} |f|^r \right)^{1/r}$$

and

$$(26) \quad \left(\frac{1}{v(x, 2^{j+1}t)} \int_{C_j(x,t)} |A_t f|^2 \right)^{1/2} \leq g(j) \left(\frac{1}{v(x, t)} \int_{B(x,t)} |f|^r \right)^{1/r}$$

for all $x \in X$, $t > 0$, $j \in \mathbb{N}$ and $f \in L^2$ supported in $B(x, t)$, and $\sum g(j)2^{dj} < \infty$. One can prove a version of this result in which $T - TA_t$ in (25) is replaced by $T - SA_t$ as in Theorem 2.1. We do not write the details here since we have no concrete application. Theorem 2.1 is suitable for our purpose.

Finally, let us mention that a Gaussian upper bound implies assumption (12). Indeed, assume that A_t is given by a kernel a_t such that

$$|a_t(x, y)| \leq \frac{C}{v(y, t^{1/m})} \exp\left\{-c \frac{\rho(x, y)^{m/(m-1)}}{t^{1/(m-1)}}\right\}$$

for all $t > 0$ and $x, y \in X$. Here $m \geq 2$ and C, c are two positive constants. Fix $x_0 \in X$. Note that the operator $\mathbb{1}_{C_j(x_0, t)} A_t^m \mathbb{1}_{B(x_0, t)}$ has the

kernel $(x, y) \mapsto \mathbb{1}_{C_j(x_0, t)}(x) a_{tm}(x, y) \mathbb{1}_{B(x_0, t)}(y)$. But

$$\begin{aligned}
& |\mathbb{1}_{C_j(x_0, t)}(x) a_{tm}(x, y) \mathbb{1}_{B(x_0, t)}(y)| \\
& \leq \frac{C}{v(y, t)} \mathbb{1}_{C_j(x_0, t)}(x) \mathbb{1}_{B(x_0, t)}(y) \exp\left\{-c \frac{\rho(x, y)^{m/(m-1)}}{t^{m/(m-1)}}\right\} \\
& \leq \frac{C}{v(y, t)} e^{-c 2^{jm/(m-1)}} \mathbb{1}_{B(x_0, t)}(y) \\
& \leq \frac{C}{v(x_0, t)} e^{-c 2^{jm/(m-1)}},
\end{aligned}$$

where we used the doubling property in the last step. This an L^1 - L^∞ estimate. By interpolation, it implies an L^1 - L^{q_0} estimate which gives (12) for every $q_0 \in (1, \infty)$.

3. A PARTIAL MULTIPLIER THEOREM FOR DEGENERATE OPERATORS

Let the coefficients a_{kj} , form \mathbf{a}_0 and the self-adjoint operator A associated with \mathbf{a}_0 be as in the introduction. For every bounded measurable function $F: [0, \infty) \rightarrow \mathbb{C}$, the operator $F(A)$ is well defined by spectral theory and is bounded on $L^2(\mathbb{R}^d)$. As mentioned in the introduction, if A is uniformly elliptic then $F(A)$ extends to a bounded operator on $L^p(\mathbb{R}^d)$ for all $p \in (1, \infty)$ provided F has a finite number of derivatives on $[0, \infty)$ which have good decay. We address here the same problem for degenerate operators. This is a difficult problem because no global Gaussian upper bounds are available for A in general. We prove a partial result by projecting on the part where the matrix (a_{kj}) is uniformly elliptic. There are two versions.

Theorem 3.1. *Let $\Omega \subset \mathbb{R}^d$ be an open bounded set with Lipschitz boundary. Suppose there exists a $\mu > 0$ such that $(a_{kj}(x)) \geq \mu I$ for almost every $x \in \Omega$ and denote by P_Ω the projection from $L^2(\mathbb{R}^d)$ onto $L^2(\Omega)$. Set $H = A + I$. Let $F: [0, \infty) \rightarrow \mathbb{C}$ be a bounded function such that*

$$(27) \quad \sup_{t>0} \|\varphi(\cdot) F(t)\|_{C^s} < \infty$$

for some $s > d/2$ and some non-trivial function $\varphi \in C_c^\infty(0, \infty)$. Then $P_\Omega F(H) P_\Omega$ is of weak type $(1, 1)$ and extends to a bounded operator on $L^p(\mathbb{R}^d)$ for all $p \in (1, \infty)$.

Theorem 3.2. *Let $\chi \in C_b^\infty(\mathbb{R}^d)$, $\mu > 0$ and suppose that $(a_{kj}(x)) \geq \mu I$ for almost every $x \in \text{supp } \chi$. Set $H = A + I$. Let $F: [0, \infty) \rightarrow \mathbb{C}$ be a bounded function such that*

$$(28) \quad \sup_{t>0} \|\varphi(\cdot)F(t)\|_{C^s} < \infty$$

for some $s > d/2$ and some non-trivial function $\varphi \in C_c^\infty(0, \infty)$. Then $M_\chi F(H)M_\chi$ is of weak type $(1, 1)$ and extends to a bounded operator on $L^p(\mathbb{R}^d)$ for all $p \in (1, \infty)$.

The proof of both theorems is almost the same. It relies mainly on weighted estimates for the associated kernel of $M_\chi F(H)M_\chi$ (or the kernel of $P_\Omega F(H)P_\Omega$) together with Theorem 2.1. The proof of weighted estimates for the kernel of $M_\chi F(H)M_\chi$ (or of $P_\Omega F(H)P_\Omega$) is based on partial Gaussian bounds proved in [EO1] and a similar strategy as in [DOS] and [Ouh].

In the sequel of this section we assume that there exists a constant $\mu > 0$ such that $(a_{kj}(x)) \geq \mu I$ for a.e. $x \in \Omega$ respectively for a.e. $x \in \text{supp } \chi \cup \text{supp } \tilde{\chi}$. In the first case Ω is a bounded Lipschitz domain of \mathbb{R}^d and in the second case $\chi, \tilde{\chi} \in C_b^\infty(\mathbb{R}^d)$. We denote by $S_t := e^{-tA}$ the holomorphic semigroup generated by $-A$ on $L^2(\mathbb{R}^d)$. We recall the following result from [EO1].

Theorem 3.3. *There are $C, c > 0$ such that for all $t > 0$ the operator $M_{\tilde{\chi}} S_t M_\chi$ (respectively $P_\Omega S_t P_\Omega$) is given by a kernel p_t which satisfies*

$$|p_t(x, y)| \leq Ct^{-d/2} e^{-c\frac{|x-y|^2}{t}} (1+t)^{d/2} \text{ for all } t > 0 \text{ and } x, y \in \mathbb{R}^d.$$

The theorem is stated in [EO1] with $\chi = \tilde{\chi}$ but the arguments work with different χ and $\tilde{\chi}$. It is also proved there that

$$(29) \quad \|M_\chi S_t\|_{2 \rightarrow \infty} \leq Ct^{-d/4} (1+t)^{d/4} \quad \text{respectively} \quad \|P_\Omega S_t\|_{2 \rightarrow \infty} \leq Ct^{-d/4} (1+t)^{d/4}.$$

If $z = t + is \in \mathbb{C}$ with $t = \text{Re } z > 0$, then

$$\begin{aligned} \|M_\chi S_z M_\chi\|_{1 \rightarrow \infty} &= \|M_\chi S_{t/2} S_{is} S_{t/2} M_\chi\|_{1 \rightarrow \infty} \\ &\leq Ct^{-d/4} (1+t)^{d/4} \|S_{is} S_{t/2} M_\chi\|_{1 \rightarrow 2} \\ &\leq Ct^{-d/4} (1+t)^{d/4} \|S_{t/2} M_\chi\|_{2 \rightarrow \infty} \\ &\leq Ct^{-d/2} (1+t)^{d/2}. \end{aligned}$$

Similarly,

$$(30) \quad \|P_\Omega S_z P_\Omega\|_{1 \rightarrow \infty} \leq C(\operatorname{Re} z)^{-d/2} (1 + \operatorname{Re} z)^{d/2}$$

for all $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$. Using the Gaussian bounds of Theorem 3.3 for real t together with the uniform bounds (30) for complex z it follows as in Theorem 3.4.8 in [Dav] or Theorem 7.2 in [Ouh] that for all $\varepsilon > 0$ the kernel $p_z^{(0)}$ of $M_\chi S_z M_\chi$ respectively $P_\Omega S_z P_\Omega$ satisfies the bound

$$(31) \quad |p_z^{(0)}(x, y)e^{-\varepsilon z}| \leq C_\varepsilon (\operatorname{Re} z)^{-d/2} \exp\left\{-c \frac{|x-y|^2}{|z|} \cos(\arg z)\right\}$$

for all $x, y \in \mathbb{R}^d$ and $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$.

Let $H = A + I$ and define $p_z(x, y) = p_z^{(0)}(x, y)e^{-z}$. Then p_z is the kernel of $M_\chi e^{-zH} M_\chi$. We shall formulate the results below for $M_\chi F(H) M_\chi$ only, but all statements are also valid for $P_\Omega F(H) P_\Omega$. In the following lemmas, we shall always assume that $(a_{kj}(x)) \geq \mu I$ for almost every $x \in \operatorname{supp} \chi$. Since associated kernels with several operators are involved in the sequel we shall denote by K_T the kernel associated to a given operator T whenever it exists.

Lemma 3.4. *For all $s > 0$ and $\varepsilon > 0$ there exists a $C > 0$ such that*

$$\int_{\mathbb{R}^d} |K_{M_\chi F(H) M_\chi}(x, y)|^2 (1 + \sqrt{r}|x-y|)^s dx \leq Cr^{d/2} \|\delta_r F\|_{C^{s/2+\varepsilon}}^2$$

for all $r > 0$, $y \in \mathbb{R}^d$ and $F \in C^{s/2+\varepsilon}$ supported in $[0, r]$. Here $(\delta_r F)(\lambda) := F(r\lambda)$.

Proof. The arguments are very similar to those of Lemma 4.3 in [DOS]. Fix $r > 0$ and assume first that F is supported in $[0, 1]$. Set $g(\lambda) := F(\lambda)e^\lambda$ and $H_r := \frac{1}{r}H$. By (31), the kernel $p_{z/r}$ of $M_\chi e^{-zH_r} M_\chi$ satisfies

$$(32) \quad |p_{z/r}(x, y)| \leq Cr^{d/2} (\operatorname{Re} z)^{-d/2} \exp\left\{-cr \frac{|x-y|^2}{|z|} \cos(\arg z)\right\}$$

for all $x, y \in \mathbb{R}^d$ and $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$, with constants C, c independent of r . We write $g(\lambda) = \int_{\mathbb{R}} \hat{g}(\xi) e^{i\lambda\xi} d\xi$, where \hat{g} is the Fourier transform of g . Then

$$F(H_r) = \int_{\mathbb{R}} \hat{g}(\xi) e^{-(1-i\xi)H_r} d\xi,$$

from which one obtains

$$(33) \quad K_{M_\chi F(H_r) M_\chi}(x, y) = \int_{\mathbb{R}} \hat{g}(\xi) p_{(1-i\xi)/r}(x, y) d\xi.$$

Let $y \in \mathbb{R}^d$. Using the estimate (32) with $z = 1 - i\xi$ gives

$$\begin{aligned}
& \int_{\mathbb{R}^d} |p_{(1-i\xi)/r}(x, y)|^2 (1 + \sqrt{r}|x - y|)^s dx \\
& \leq Cr^d \int \exp\left\{-2cr \frac{|x - y|^2}{1 + \xi^2}\right\} (1 + \sqrt{r}|x - y|)^s dx \\
& \leq Cr^d (1 + \xi^2)^{s/2} \int \exp\left\{-cr \frac{|x - y|^2}{1 + \xi^2}\right\} dx \\
& \leq Cr^d (1 + \xi^2)^{s/2} \left(\frac{1 + \xi^2}{r}\right)^{d/2} \\
& = Cr^{d/2} (1 + \xi^2)^{(d+s)/2}.
\end{aligned}$$

It follows from (33), the continuous version of the Minkowski inequality and the previous estimate that

$$\begin{aligned}
& \left(\int_{\mathbb{R}^d} |K_{M_\chi F(H_r)M_\chi}(x, y)|^2 (1 + \sqrt{r}|x - y|)^s dx \right)^{1/2} \\
& \leq \int_{\mathbb{R}} |\hat{g}(\xi)| \left(\int_{\mathbb{R}^d} |p_{(1-i\xi)/r}(x, y)|^2 (1 + \sqrt{r}|x - y|)^s dx \right)^{1/2} d\xi \\
& \leq Cr^{d/4} \int_{\mathbb{R}} |\hat{g}(\xi)| (1 + \xi^2)^{(d+s)/4} d\xi \\
& \leq Cr^{d/4} \|g\|_{W^{(d+s+2)/2, 2}} \\
(34) \quad & \leq Cr^{d/4} \|F\|_{W^{s/2+\alpha, 2}}.
\end{aligned}$$

Here $\alpha = (d + 2)/2$ and the constants are independent of r and y . On the other hand $M_\chi F(H_r)M_\chi = M_\chi g(H_r)e^{-H_r}M_\chi$. It follows from (29) that

$$\|e^{-H_r}M_\chi\|_{1 \rightarrow 2} \leq Cr^{d/4} \left(1 + \frac{1}{r}\right)^{d/4} e^{-1/r} \leq Cr^{d/4}$$

for all $r > 0$. Moreover, $\|M_\chi g(H_r)\|_{2 \rightarrow 2} \leq e\|\chi\|_\infty \|F\|_\infty$. Therefore

$$(35) \quad \int_{\mathbb{R}^d} |K_{M_\chi F(H_r)M_\chi}(x, y)|^2 dx \leq \|M_\chi F(H_r)M_\chi\|_{1 \rightarrow 2}^2 \leq Cr^{d/2} \|F\|_\infty^2.$$

This is valid for all F with support in $[0, 1]$ and for all $s > 0$. The estimates (34) and (35) together with an interpolation argument (see [MM], p. 151 and [DOS], p. 455) give then that for all $s > 0$ there exists a $C > 0$ such

that

$$(36) \quad \int_{\mathbb{R}^d} |K_{M_\chi F(H_r)M_\chi}(x, y)|^2 (1 + \sqrt{r}|x - y|)^s dx \leq Cr^{d/2} \|F\|_{C^{s/2+\varepsilon}}^2.$$

Finally, if F has support in $[0, r]$ we use the last estimate with $\delta_r F$ and obtain the lemma. \square

Lemma 3.5. *The operators $A_t := e^{-t^2 H} M_\chi$ satisfy (12).*

Proof. Let $\psi \in W^{1,\infty}(\mathbb{R}^d, \mathbb{R})$ be such that $|\nabla\psi| \leq 1$. For all $\rho \in \mathbb{R}$ define $U_\rho = M_{e^{\rho\psi}}$ and set $S_t^\rho := U_\rho e^{-tH} U_{-\rho}$. It follows from [EO1] Proposition 3.6 by duality and a limit $n \rightarrow \infty$ that there exist $C, \omega > 0$, independent of t, ρ and ψ , such that

$$(37) \quad \|S_t^\rho M_\chi\|_{1 \rightarrow 2} \leq Ct^{-d/4} e^{\omega\rho^2 t}.$$

Now fix two bounded open non-empty sets E and F of \mathbb{R}^d and choose $\psi(x) := d(x, E) \wedge N$, where $N = \sup\{|x - y| : x \in E, y \in F\} + 1$. For all $h \in L^2(E)$ and $\rho \geq 0$ one has

$$M_\chi e^{-tH} h = M_\chi U_{-\rho} S_t^\rho h.$$

Therefore

$$\|M_\chi e^{-tH} h\|_{L^\infty(F)} \leq e^{-\rho d(E,F)} \|M_\chi S_t^\rho h\|_\infty \leq Ct^{-d/4} e^{-\rho d(E,F)} e^{\omega\rho^2 t} \|h\|_2.$$

Choosing $\rho = \frac{d(E,F)}{2\omega t}$ yields the Davies–Gaffney type estimate

$$(38) \quad \|P_F(M_\chi e^{-tH}) P_E\|_{2 \rightarrow \infty} \leq Ct^{-d/4} e^{-\frac{d(E,F)^2}{4\omega t}}.$$

In particular,

$$\|P_{C_j(x,t)} e^{-t^2 H} M_\chi P_{B(x,t)}\|_{1 \rightarrow 2} \leq Ct^{-d/2} e^{-c4^j}$$

for all $x, y \in \mathbb{R}^d$ and $j \in \mathbb{N}$. This shows the lemma. \square

Proof of Theorems 3.1 and 3.2. As mentioned above, the proof of both theorems is almost the same. We consider $M_\chi F(H)M_\chi$ only. The proof is based on Theorem 2.1 and the previous lemmas. It is in the same spirit as in the elliptic case where a Gaussian bound holds (cf. [DOS], [Ouh]). Let $\varphi \in C_c^\infty(0, \infty)$ be such that $\text{supp } \varphi \subset [1/4, 1]$ and

$$\sum_{n=-\infty}^{\infty} \varphi(2^{-n}\lambda) = 1$$

for all $\lambda > 0$. Then

$$F(\lambda) = \sum_{n=-\infty}^{\infty} \varphi(2^{-n}\lambda)F(\lambda) =: \sum_{n=-\infty}^{\infty} F_n(\lambda).$$

We apply Theorem 2.1 to $M_\chi F_n(H)M_\chi$ for each fixed $n \in \mathbb{Z}$. We choose

$$S := M_\chi F_n(H) \quad \text{and} \quad A_t := e^{-t^2 H} M_\chi.$$

By Lemma 3.5, the operators A_t satisfy (12). It remains to prove (15). For this we have to estimate for all $y \in \mathbb{R}^d$ the integral

$$I_{n,t} := \int_{|x-y| \geq t} |K_{M_\chi G_{n,t}(H)M_\chi}(x, y)| dx,$$

where $G_{n,t}(\lambda) = F_n(\lambda) - F_n(\lambda)e^{-t^2\lambda} = \varphi(2^{-n}\lambda)F(\lambda)(1 - e^{-t^2\lambda})$. First, by the Cauchy–Schwarz inequality we have

$$(39) \quad I_{n,t} \leq \left(\int_{\mathbb{R}^d} |K_{M_\chi G_{n,t}(H)M_\chi}(x, y)|^2 (1 + 2^{n/2}|x - y|)^{2s} dx \right)^{1/2} \times \left(\int_{|x-y| \geq t} (1 + 2^{n/2}|x - y|)^{-2s} dx \right)^{1/2}$$

We apply Lemma 3.4 with $r = 2^n$ and obtain

$$(40) \quad \int_{\mathbb{R}^d} |K_{M_\chi G_{n,t}(H)M_\chi}(x, y)|^2 (1 + 2^{n/2}|x - y|)^{2s} dx \leq C2^{nd/2} \|\delta_{2^n} G_{n,t}\|_{C^{s+\varepsilon}}^2.$$

Simple computations show that there exists a $C > 0$, independent of n and t , such that

$$(41) \quad \begin{aligned} \|\delta_{2^n} G_{n,t}\|_{C^{s+\varepsilon}} &= \|\varphi(\cdot)F(2^n \cdot)(1 - e^{-t^2 2^n \cdot})\|_{C^{s+\varepsilon}} \\ &\leq C \sup_{t' > 0} \|\varphi(\cdot)F(t' \cdot)\|_{C^{s+\varepsilon}} \min(1, t^2 2^n). \end{aligned}$$

On the other hand (see [DOS] or (7.46) in [Ouh]) one estimates

$$(42) \quad \int_{|x-y| \geq t} (1 + 2^{n/2}|x - y|)^{-2s} dx \leq C2^{-nd/2} \min(1, (t2^{n/2})^{d-2s}).$$

Using (39), (40), (41) and (42) we obtain

$$I_{n,t} \leq C \min(1, t^2 2^n) \min(1, (t2^{n/2})^{\frac{d}{2}-s}) \sup_{t' > 0} \|\varphi(\cdot)F(t' \cdot)\|_{C^{s+\varepsilon}}.$$

Hence

$$\sum_{n=-\infty}^{\infty} I_{n,t} \leq C \left(\sum_{n \in \mathbb{Z}, t^2 2^n \leq 1} t^2 2^n + \sum_{n \in \mathbb{Z}, t2^{n/2} > 1} (t2^{n/2})^{\frac{d}{2}-s} \right) \sup_{t' > 0} \|\varphi(\cdot)F(t' \cdot)\|_{C^{s+\varepsilon}}$$

and the RHS is bounded by a constant independent t since $s > \frac{d}{2}$. This proves Theorem 3.1. \square

As explained in the introduction, the reason why we consider $H = A + I$ instead of A in the previous results comes from the fact the Gaussian upper bound in Theorem 3.3 is valid with the extra factor $(1+t)^{d/2}$. If one considers the case where $a_{kj} = \delta_{kj}$ on a smooth bounded domain Ω , then A is the Neumann Laplacian on $L^2(\Omega)$ and 0 on $L^2(\mathbb{R}^d \setminus \Omega)$. It is then easy to see that L^2 - L^∞ estimates (respectively, Gaussian bounds) for $M_\chi e^{-tA}$ or $P_\Omega e^{-tA}$ (respectively, $M_\chi e^{-tA} M_\chi$ or $P_\Omega e^{-tA} P_\Omega$) cannot hold without an extra factor $(1+t)^{d/4}$ (respectively, $(1+t)^{d/2}$). On the other hand we can replace in the previous theorems $H = A + I$ by $H = A + \varepsilon I$ for any $\varepsilon > 0$.

It may be possible that if $(a_{kj}) \geq \mu I$ on a connected subset F of \mathbb{R}^d which is ‘large enough’ (in some sense) one obtains Theorem 3.3 without the extra factor $(1+t)^{d/2}$ in the Gaussian bound. This remains to be proved. We mention that if such bound holds, we obtain by the same proof Theorems 3.1 and 3.2 for $F(A)$ rather than $F(H)$.

We emphasize also that we consider here general degenerate operators with non-smooth coefficients. One may obtain global results for some specific operators which are degenerate at every point and the coefficients are not continuous at every point. For example, one might take a pure second-order subelliptic operator in divergence form with real measurable coefficients on a Lie group with polynomial growth. Then global Gaussian bounds are valid by [SS], Théorème 1 together with regularization argument (see, for example Section 2.1 in [ER]). Therefore a global spectral multiplier result for such operators follows directly from [DOS]. Note however that the order of smoothness required on the function F is larger than half of Euclidean dimension. On the other hand, the operators that we consider in this paper are allowed to vanish on big sets.

Examples 3.6. We give some examples which are direct applications of the previous theorems.

Imaginary powers. Set $F(\lambda) = \lambda^{is}$ where $s \in \mathbb{R}$. Then Theorems 3.1 and 3.2 together with the Riesz–Thorin interpolation theorem imply that for all $\varepsilon > 0$ and $p \in (1, \infty)$ there exists a $C > 0$ such that

$$\|M_\chi H^{is} M_\chi\|_{p \rightarrow p} \leq C_\varepsilon (1 + |s|)^{(d+\varepsilon)|\frac{1}{2} - \frac{1}{p}|}$$

and

$$\|P_\Omega H^{is} P_\Omega\|_{\mathcal{L}(L^p)} \leq C_\varepsilon (1 + |s|)^{(d+\varepsilon)|\frac{1}{2}-\frac{1}{p}|}$$

for all $s \in \mathbb{R}$.

The Schrödinger group. Set $F(\lambda) = (1 + \lambda)^{-\alpha} e^{it\lambda}$ with $t \in \mathbb{R}$ and $\alpha > d/2$. The operators $M_\chi(I + H)^{-\alpha} e^{itH} M_\chi$ and $P_\Omega(I + H)^{-\alpha} e^{itH} P_\Omega$ are bounded on $L^p(\mathbb{R}^d)$ for all $p \in (1, \infty)$. Their L^p -norms are estimated by $C(1 + |t|)^\alpha$. By interpolation, we obtain boundedness on L^p for all $\alpha > d|\frac{1}{2} - \frac{1}{p}|$ and $t \in \mathbb{R}$.

Remark. Using the same proof as in [CCO], these results can be obtained directly from the Gaussian upper bound of Theorem 3.3 without appealing to Theorems 3.1 and 3.2.

Wave operators. Set $F(\lambda) = (1 + \lambda)^{-\alpha/2} e^{it\sqrt{\lambda}}$ with $t \in \mathbb{R}$ and $\alpha > d/2$. The operators $M_\chi(I + H)^{-\alpha} e^{it\sqrt{H}} M_\chi$ and $P_\Omega(I + H)^{-\alpha} e^{it\sqrt{H}} P_\Omega$ are bounded on $L^p(\mathbb{R}^d)$ for all $p \in (1, \infty)$.

4. RIESZ TRANSFORMS

The aim in this section is to prove boundedness on $L^p(\mathbb{R}^d)$ of a type of Riesz transform operator $M_\chi \nabla(I + A)^{-1/2} M_\chi$. We keep the same notation as in the previous section. The main result of this section is the next theorem.

Theorem 4.1. *Let $\chi \in C_b^\infty(\mathbb{R}^d)$, $\mu > 0$ and suppose that $(a_{kj}(x)) \geq \mu I$ for almost every $x \in \text{supp } \chi$. Set $H = A + I$. Then for every $k \in \{1, \dots, d\}$, the operator $M_\chi \partial_k H^{-1/2} M_\chi$ is of weak type $(1, 1)$ and is bounded on $L^p(\mathbb{R}^d)$ for all $p \in (1, 2]$.*

Here ∂_k denotes the distributional derivative. The proof is based on Theorem 2.1 and uses some ideas from [CD], [DM1] and Chapter 7 in [Ouh] in the uniformly elliptic case. We start with the following lemma. Let \mathfrak{a} be the closure of the regular part of the form \mathfrak{a}_0 defined in (5).

Lemma 4.2. *Let $\chi \in C_b^\infty(\mathbb{R}^d)$, $\mu > 0$ and suppose that $(a_{kj}(x)) \geq \mu I$ for almost every $x \in \text{supp } \chi$. Then $\chi u \in W^{1,2}(\mathbb{R}^d)$ and*

$$\|\chi \partial_k u\|_2^2 \leq \frac{\|\chi\|_\infty^2}{\mu} \|H^{1/2} u\|_2^2$$

for all $u \in D(\mathfrak{a}) = D(H^{1/2})$ and $k \in \{1, \dots, d\}$.

Proof. Let $u \in D(\mathbf{a})$. Then there exists a sequence $(u_n)_{n \in \mathbb{N}}$ in $D(\mathbf{a}_0) = C_c^\infty(\mathbb{R}^d)$ such that $\lim u_n = u$ in $L^2(\mathbb{R}^d)$ and $\mathbf{a}(u) = \lim \mathbf{a}_0(u_n)$. By the ellipticity assumption on the support of χ one deduces

$$(43) \quad \mu \int_{\mathbb{R}^d} \chi^2 |\nabla u_n|^2 \leq \sum_{k,j=1}^d \int_{\mathbb{R}^d} a_{kj} (\partial_k u_n) (\partial_j u_n) \chi^2 \leq \|\chi\|_\infty^2 \mathbf{a}_0(u_n)$$

for all $n \in \mathbb{N}$. Therefore $(\chi u_n)_{n \in \mathbb{N}}$ is bounded in $W^{1,2}(\mathbb{R}^d)$. Hence it has a weakly convergent subsequence in $W^{1,2}(\mathbb{R}^d)$. Since $\lim \chi u_n = \chi u$ in $L^2(\mathbb{R}^d)$ it follows that $\chi u \in W^{1,2}(\mathbb{R}^d)$. Then taking the limit $n \rightarrow \infty$ in (43) one estimates

$$\mu \int_{\mathbb{R}^d} \chi^2 |\partial_k u|^2 \leq \|\chi\|_\infty^2 \mathbf{a}(u) \leq \|\chi\|_\infty^2 \|H^{1/2} u\|_2^2$$

for all $k \in \{1, \dots, d\}$. \square

Lemma 4.3. *Let $\chi, \tilde{\chi} \in C_b^\infty(\mathbb{R}^d)$, $\mu > 0$ and assume that $(a_{kj}(x)) \geq \mu I$ for almost every $x \in \text{supp } \chi \cup \text{supp } \tilde{\chi}$. Then for all $\beta > 0$ small enough we have*

$$\int_{\mathbb{R}^d} |(M_{\tilde{\chi}} e^{-sH} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} dy \leq C s^{-d/2} e^{2\beta t^2/s} e^{-s} \|u\|_1^2$$

for all $t > 0$, $s > 0$, $x \in \mathbb{R}^d$ and $u \in L^2(\mathbb{R}^d)$ with $\text{supp } u \subset B(x, t)$.

Proof. By Theorem 3.3 one estimates

$$\begin{aligned} & |(M_{\tilde{\chi}} e^{-sH} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} \\ &= \left| \int_{B(x,t)} e^{-s} p_s(y, z) u(z) dz \right|^2 e^{\beta \frac{|x-y|^2}{s}} \\ &\leq C e^{-s} \left(\int_{B(x,t)} s^{-d/2} e^{-c \frac{|y-z|^2}{s}} e^{\beta \frac{|x-y|^2}{2s}} |u(z)| dz \right)^2 \\ &\leq C e^{-s} \left(\int_{\mathbb{R}^d} s^{-d/2} e^{-(c-\beta) \frac{|y-z|^2}{s}} |u(z)| dz \right)^2 e^{2\beta t^2/s} \\ &\leq C e^{-s} s^{-d/2} \|u\|_1 \int_{\mathbb{R}^d} s^{-d/2} e^{-(c-\beta) \frac{|y-z|^2}{s}} |u(z)| dz e^{2\beta t^2/s}. \end{aligned}$$

Taking $\beta < \frac{1}{2}c$ and integrating over y yields the lemma. \square

Since $e^{-sH} L^2(\mathbb{R}^d) \subset D(\mathbf{a})$ for all $s > 0$ we obtain from Lemma 4.2 the inclusion $M_\chi \nabla e^{-sH} M_\chi (L^2(\mathbb{R}^d)) \subset W^{1,2}(\mathbb{R}^d)$ for all $s > 0$. The following weighted L^2 -estimate is in the same spirit as weighted gradient estimates for heat kernels (see [CD], [Gri] and Theorem 6.19 in [Ouh]).

Lemma 4.4. *For all $\beta > 0$ small enough we have*

$$\int_{\mathbb{R}^d} |(M_\chi \nabla e^{-sH} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} dy \leq C s^{-d/2-1} e^{6\beta t^2/s} \|u\|_1^2$$

for all $t > 0$, $s > 0$, $x \in \mathbb{R}^d$ and $u \in L^2(\mathbb{R}^d)$ with $\text{supp } u \subset B(x, t)$.

Proof. In order to avoid domain problems of forms we shall proceed by approximation. First, we prove the lemma for uniformly elliptic coefficients with constants β and C depending only on $\mu > 0$ such that $(a_{kj}(x)) \geq \mu I$ a.e. $x \in \text{supp } \chi$.

Assume that there exists a $\mu_0 > 0$ such that $(a_{kj}(x)) \geq \mu_0 I$ for a.e. $x \in \mathbb{R}^d$. In this case the form \mathbf{a} has domain $W^{1,2}(\mathbb{R}^d)$. We use similar ideas as in the proof of Theorem 6.19 in [Ouh] but we want to prove that the constants in the estimates are independent of μ_0 . Let $\psi \in C_c^\infty(\mathbb{R}^d)$ be such that $\psi(x) = 1$ for all $x \in B(0, 1)$ and $0 \leq \psi \leq 1$. For all $n \in \mathbb{N}$ define $\psi_n \in C_c^\infty(\mathbb{R}^d)$ by $\psi_n(x) = \psi(\frac{1}{n}x)$. Set

$$I_n := \int_{\mathbb{R}^d} |\chi(y) (\nabla e^{-sH} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} \psi_n(y) dy$$

and define $f := e^{-sH} M_\chi u$. Then,

$$\begin{aligned} I_n &\leq \frac{1}{\mu} \sum_{k,j} \int_{\mathbb{R}^d} a_{kj}(y) (\partial_k f)(y) (\partial_j f)(y) e^{\beta \frac{|x-y|^2}{s}} \chi(y)^2 \psi_n(y) dy \\ &= \frac{1}{\mu} \sum_{k,j} \int_{\mathbb{R}^d} a_{kj} (\partial_k f) \partial_j \left(f e^{\beta \frac{|x-y|^2}{s}} \chi^2 \psi_n \right) \\ &\quad + \frac{1}{\mu} \sum_{k,j} \int_{\mathbb{R}^d} a_{kj}(y) (\partial_k f)(y) f(y) \frac{2\beta(x_j - y_j)}{s} e^{\beta \frac{|x-y|^2}{s}} \chi(y)^2 \psi_n(y) dy \\ &\quad - \frac{2}{\mu} \sum_{k,j} \int_{\mathbb{R}^d} a_{kj}(y) (\partial_k f)(y) f(y) e^{\beta \frac{|x-y|^2}{s}} (\partial_j \chi)(y) \chi(y) \psi_n(y) dy \\ &\quad - \frac{1}{n\mu} \sum_{k,j} \int_{\mathbb{R}^d} a_{kj}(y) (\partial_k f)(y) f(y) e^{\beta \frac{|x-y|^2}{s}} \chi(y)^2 (\partial_j \psi)\left(\frac{1}{n}y\right) dy \\ &=: J_{1,n} + J_{2,n} + J_{3,n} + J_{4,n}. \end{aligned}$$

Since $y \mapsto f(y)e^{\beta\frac{|x-y|^2}{s}}\chi(y)^2\psi_n(y)$ is an element of $W^{1,2}(\mathbb{R}^d)$ we have

$$\begin{aligned} J_{1,n} &= \frac{1}{\mu} \mathbf{a}(f, f e^{\beta\frac{|x-\cdot|^2}{s}} \chi^2 \psi_n) \\ &= \frac{1}{\mu} \int_{\mathbb{R}^d} (Ae^{-sH} M_\chi u) (e^{-sH} M_\chi u) e^{\beta\frac{|x-\cdot|^2}{s}} \chi^2 \psi_n \\ &\leq \frac{\|\chi\|_\infty}{\mu} \|He^{-sH} M_\chi u\|_2 \|e^{\beta\frac{|x-\cdot|^2}{s}} M_\chi e^{-sH} M_\chi u\|_2. \end{aligned}$$

The standard estimate $\|He^{-sH}\|_{2 \rightarrow 2} \leq s^{-1}$ and Lemma 4.3 give

$$(44) \quad J_{1,n} \leq C s^{-d/2-1} e^{2\beta t^2/s} \|u\|_1^2$$

if β is small enough. Using the obvious inequality $\frac{|x_j - y_j|}{s} \leq \frac{1}{\sqrt{\varepsilon} s} e^\varepsilon \frac{|x-y|^2}{s}$ we have

$$\begin{aligned} |J_{2,n}| &\leq \frac{C}{\sqrt{s}} \sum_k \int_{\mathbb{R}^d} |\partial_k e^{-sH} M_\chi u| \chi^2 |e^{-sH} M_\chi u| e^{2\beta\frac{|x-\cdot|^2}{s}} \\ &\leq \frac{C}{\sqrt{s}} \sqrt{I_n} \left(\int_{\mathbb{R}^d} |(M_\chi e^{-sH} M_\chi u)(y)|^2 e^{3\beta\frac{|x-y|^2}{s}} dy \right)^{1/2}. \end{aligned}$$

Therefore Lemma 4.3 implies

$$(45) \quad |J_{2,n}| \leq C \sqrt{I_n} s^{-\frac{d}{4}-\frac{1}{2}} e^{-s/2} e^{3\beta t^2/s} \|u\|_1.$$

We estimate the third term in a similar way.

$$\begin{aligned} |J_{3,n}| &\leq C \sum_k \int_{\mathbb{R}^d} |\partial_k e^{-sH} M_\chi u| |\chi \partial_j \chi| |e^{-sH} M_\chi u| e^{\beta\frac{|x-\cdot|^2}{s}} \psi_n \\ &\leq C \sqrt{I_n} \left(\int_{\mathbb{R}^d} |M_{\partial_j \chi} e^{-sH} M_\chi u|^2 e^{\beta\frac{|x-\cdot|^2}{s}} \right)^{1/2} \\ (46) \quad &\leq C \sqrt{I_n} s^{-\frac{d}{4}-\frac{1}{2}} e^{-s/3} e^{\beta t^2/s} \|u\|_1. \end{aligned}$$

Finally,

$$\begin{aligned}
|J_{4,n}| &\leq \frac{C}{n} \sum_{k,j} \int_{\mathbb{R}^d} |(\chi \partial_k e^{-sH} M_\chi u)(y)| |(M_\chi e^{-sH} M_\chi u)(y)| e^{\beta \frac{|x-y|^2}{s}} \times \\
&\hspace{20em} \times |(\partial_j \psi)(\frac{1}{n} y)| dy \\
&\leq \frac{C}{n} \|M_\chi \nabla e^{-sH} M_\chi u\|_2 \left(\int_{\mathbb{R}^d} |(M_\chi e^{-sH} M_\chi u)(y)|^2 e^{2\beta \frac{|x-y|^2}{s}} dy \right)^{1/2} \\
(47) \quad &\leq \frac{C}{n} \|M_\chi \nabla e^{-sH} M_\chi u\|_2 s^{-d/4} e^{-s/2} e^{2\beta t^2/s} \|u\|_1.
\end{aligned}$$

Therefore, we obtain from (44), (45), (46) and (47) that

$$I_n \leq C s^{-d/2-1} e^{6\beta t^2/s} \|u\|_1^2 + \frac{C}{n} \|M_\chi \nabla e^{-sH} M_\chi u\|_2 s^{-d/4} e^{2\beta t^2/s} \|u\|_1.$$

Letting $n \rightarrow \infty$ and then use Fatou's lemma yields

$$(48) \quad \int_{\mathbb{R}^d} |(M_\chi \nabla e^{-sH} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} dy \leq C s^{-d/2-1} e^{6\beta t^2/s} \|u\|_1^2.$$

The constants C and β are independent of μ_0 .

Now we prove the lemma for degenerate operators. For all $n \in \mathbb{N}$ set $a_{kj}^{(n)} = a_{kj} + \frac{1}{n} \delta_{kj}$. Then $(a_{kj}^{(n)}(x)) \geq \frac{1}{n} I$ for a.e. $x \in \mathbb{R}^d$ and $(a_{kj}^{(n)}(x)) \geq \mu I$ for a.e. $x \in \text{supp } \chi$. Moreover, $\|a_{kj}^{(n)}\|_\infty \leq 1 + \|a_{kj}\|_\infty$. We denote by A_n the elliptic operator with the coefficients $a_{kj}^{(n)}$ and $H_n = I + A_n$. We apply (48) to H_n and obtain

$$(49) \quad \int_{\mathbb{R}^d} |(M_\chi \nabla e^{-sH_n} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} dy \leq C s^{-d/2-1} e^{6\beta t^2/s} \|u\|_1^2.$$

for some constants C and $\beta > 0$ which are independent of n . Let $k \in \{1, \dots, d\}$. Then

$$|(e^{-sH_n} M_\chi u, \partial_k M_\chi (\varphi e^{\beta \frac{|x-\cdot|^2}{2s}}))| \leq C s^{-\frac{d}{4}-\frac{1}{2}} e^{3\beta t^2/s} \|u\|_1 \|\varphi\|_2$$

for all $\varphi \in C_c^\infty(\mathbb{R}^d)$. On the other hand, e^{-tH_n} converges strongly in $L^2(\mathbb{R}^d)$ to e^{-tH} (see [AE] Corollary 3.9). It follows then that

$$|(e^{-sH} M_\chi u, \partial_k M_\chi (\varphi e^{\beta \frac{|x-\cdot|^2}{2s}}))| \leq C s^{-\frac{d}{4}-\frac{1}{2}} e^{3\beta t^2/s} \|u\|_1 \|\varphi\|_2.$$

Since this is true for all $\varphi \in C_c^\infty(\mathbb{R}^d)$ we have by density

$$\|(M_\chi \partial_k e^{-sH} M_\chi u) \cdot e^{\beta \frac{|x-\cdot|^2}{2s}}\|_2 \leq C s^{-\frac{d}{4}-\frac{1}{2}} e^{3\beta t^2/s} \|u\|_1.$$

This proves the lemma. \square

Proof of Theorem 4.1. It follows from Lemma 4.2 that the Riesz transform $M_\chi \partial_k A^{-1/2}$ is bounded on $L^2(\mathbb{R}^d)$.

In order to prove a weak type estimate for $T = M_\chi \partial_k H^{-1/2} M_\chi$ we apply Theorem 2.1 with $S = M_\chi \partial_k H^{-1/2}$ and $A_t = e^{-t^2 H} M_\chi$. These operators are bounded on $L^2(\mathbb{R}^d)$ and by Lemma 3.5 the operators A_t satisfy assumption (12). It remains then to check (13). By the formula

$$H^{-1/2} = \frac{1}{2\sqrt{\pi}} \int_0^\infty e^{-sH} \frac{ds}{\sqrt{s}}$$

we have

$$H^{-1/2} e^{-t^2 H} = \frac{1}{2\sqrt{\pi}} \int_0^\infty e^{-(s+t^2)H} \frac{ds}{\sqrt{s}} = \frac{1}{2\sqrt{\pi}} \int_0^\infty e^{-sH} \mathbf{1}_{\{s>t^2\}} \frac{ds}{\sqrt{s-t^2}}.$$

Let $\beta > 0$ be as in Lemma 4.4 and let $\delta > 0$. Fix $x \in \mathbb{R}^d$, $t > 0$ and let $u \in L^2(\mathbb{R}^d)$ with $\text{supp } u \subset B(x, t)$. Set $\nu(s, t) = \left| \mathbf{1}_{\{s>t^2\}} \frac{1}{\sqrt{s-t^2}} - \frac{1}{\sqrt{s}} \right|$. Then

$$\begin{aligned} & 2\sqrt{\pi} \int_{\mathbb{R}^d \setminus B(x, (1+\delta)t)} |((T - SA_t)u)(y)| dy \\ & \leq \int_0^\infty \int_{\mathbb{R}^d \setminus B(x, (1+\delta)t)} |(M_\chi \partial_k e^{-sH} M_\chi u)(y)| dy \nu(s, t) ds \\ & \leq \int_0^\infty \nu(s, t) \left(\int_{\mathbb{R}^d} |(M_\chi \partial_k e^{-sH} M_\chi u)(y)|^2 e^{\beta \frac{|x-y|^2}{s}} dy \right)^{1/2} \times \\ & \quad \times \left(\int_{\mathbb{R}^d \setminus B(x, (1+\delta)t)} e^{-\beta \frac{|x-y|^2}{s}} dy \right)^{1/2} ds \\ & \leq C \int_0^\infty \nu(s, t) s^{-\frac{d}{4}-\frac{1}{2}} e^{3\beta t^2/s} \|u\|_1 \left(\int_{\mathbb{R}^d \setminus B(x, (1+\delta)t)} e^{-\beta \frac{|x-y|^2}{s}} dy \right)^{1/2} ds. \end{aligned}$$

Note that we have used Lemma 4.4 in the last inequality. Now

$$\int_{\mathbb{R}^d \setminus B(x, (1+\delta)t)} e^{-\beta \frac{|x-y|^2}{s}} dy \leq e^{-\beta \frac{(1+\delta)^2 t^2}{2s}} \int_{\mathbb{R}^d} e^{-\beta \frac{|x-y|^2}{2s}} dy \leq C s^{d/2} e^{-\beta \frac{(1+\delta)^2 t^2}{2s}}.$$

Choosing $\delta \geq 4$ we obtain a positive constant γ such that

$$\int_{\mathbb{R}^d \setminus B(x, (1+\delta)t)} |(T - SA_t)u(y)| dy \leq C \int_0^\infty \nu(s, t) s^{-1/2} e^{-\gamma t^2/s} ds.$$

The latter integral is bounded by some constant M independent of t . This proves (13) and hence $T = M_\chi \partial_k H^{-1/2} M_\chi$ is weak type $(1, 1)$. By interpolation, it is bounded on $L^p(\mathbb{R}^d)$ for all $1 < p \leq 2$. \square

As discussed at the end of the previous section, we note that if one proves a version of Theorem 3.3 without the extra factor $(1+t)^{d/2}$ if $(a_{kj}(x)) \geq \mu I$ for a.e. x in a ‘big’ domain, then Theorem 4.1 holds with A in place of H . That is $M_\chi \partial_k A^{-1/2} M_\chi$ is weak type $(1, 1)$ and bounded on $L^p(\mathbb{R}^d)$ for all $1 < p \leq 2$. In [EO2], we prove by a different method that if the coefficients $a_{kj} \in W^{1,\infty}(\mathbb{R}^d)$, then $M_\chi \partial_k (I + A)^{-1/2}$ and $M_\chi \partial_k \partial_j (I + A)^{-1}$ are bounded on $L^p(\mathbb{R}^d)$ for all $p \in (1, \infty)$. Moreover, if $a_{kj} \in W^{\nu,\infty}(\mathbb{R}^d, \mathbb{C})$ then we show that $M_\chi \partial_k (I + A)^{-1/2} M_\chi$ is bounded on L^p for all $p \in (1, \infty)$.

Acknowledgements Most of this work was carried out whilst the second named author was visiting the Department of Mathematics at the University of Auckland. He wishes to thank both the University of Auckland and the CNRS for financial support. Part of this work is supported by the Marsden Fund Council from Government funding, administered by the Royal Society of New Zealand.

REFERENCES

- [AE] ARENDT, W. and ELST, A. F. M. TER, Sectorial forms and degenerate differential operators. *J. Operator Theory* **67** (2012), 101–140.
- [Aus] AUSCHER, P., On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates. *Memoirs Amer. Math. Soc.* **186**, No. 871 (2007).
- [BK] BLUNCK, S. and KUNSTMANN, P. C., Calderón–Zygmund theory for non-integral operators and the H^∞ -functional calculus. *Rev. Mat. Iberoamericana* **19** (2003), 919–942.
- [CCO] CARRON, G., COULHON, T. and OUHABAZ, E.-M., Gaussian estimates and L_p -boundedness of Riesz means. *J. Evol. Equ.* **2** (2002), 299–317.
- [CW] COIFMAN, R. R. and WEISS, G., *Analyse harmonique non-commutative sur certains espaces homogènes*. Lect. Notes in Math. 242. Springer-Verlag, Berlin etc., 1971.
- [CD] COULHON, T. and DUONG, X. T., Riesz transforms for $1 \leq p \leq 2$. *Trans. Amer. Math. Soc.* (1999), 1151–1169.
- [Dav] DAVIES, E. B., *Heat kernels and spectral theory*. Cambridge Tracts in Mathematics 92. Cambridge University Press, Cambridge etc., 1989.
- [DM1] DUONG, X. T. and MCINTOSH, A., The L^p boundedness of Riesz transforms associated with divergence form operators. In CRANNY, T. and THOMPSON, B., eds., *Joint Australian-Taiwanese workshop on analysis and applications (Brisbane, 1997)*, vol. 37 of Proc. Centre Math. Anal. and its Appl., 1999, 15–25.

- [DM2] DUONG, X. T. and MCINTOSH, A., Singular integral operators with non-smooth kernels on irregular domains. *Rev. Mat. Iberoamericana* **15** (1999), 233–265.
- [DOS] DUONG, X. T., OUHABAZ, E.-M. and SIKORA, A., Plancherel-type estimates and sharp spectral multipliers. *J. Funct. Anal.* **196** (2002), 443–485.
- [DY] DUONG, X. T. and YAN, L., Duality of Hardy and BMO spaces associated with operators with heat kernel bounds. *J. Amer. Math. Soc.* **18** (1975), 943–973.
- [EO1] ELST, A. F. M. TER and OUHABAZ, E.-M., Partial Gaussian bounds for degenerate differential operators. *Potential Analysis* **35** (2011), 175–199.
- [EO2] ———, Partial Gaussian bounds for degenerate differential operators II, 2011.
- [ER] ELST, A. F. M. TER and ROBINSON, D. W., Second-order subelliptic operators on Lie groups I: complex uniformly continuous principal coefficients. *Acta Appl. Math.* **59** (1999), 299–331.
- [Gri] GRIGOR'YAN, A., Upper bounds of derivatives of the heat kernel on an arbitrary complete manifold. *J. Funct. Anal.* **127** (1995), 363–389.
- [HLMMY] HOFMANN, S., LU, G., MITREA, D., MITREA, M. and YAN, L., Hardy spaces associated to non-negative self-adjoint operators satisfying Davies–Gaffney estimates. *Memoirs Amer. Math. Soc.* **214**, No. 1007 (2011).
- [MM] MAUCERI, M. and MEDA, S., Vector-valued multipliers on stratified groups. *Rev. Mat. Iberoamericana* **6** (1990), 141–154.
- [Ouh] OUHABAZ, E.M., *Analysis of heat equations on domains*, vol. 31 of London Mathematical Society Monographs Series. Princeton University Press, Princeton, NJ, 2005.
- [SS] SALOFF-COSTE, L. and STROOCK, D. W., Opérateurs uniformément sous-elliptiques sur les groupes de Lie. *J. Funct. Anal.* **98** (1991), 97–121.
- [She] SHEN, Z., Bounds of Riesz transforms on L^p spaces for second order elliptic operators. *Ann. Inst. Fourier (Grenoble)* **55** (2005), 173–197.
- [Sim] SIMON, B., A canonical decomposition for quadratic forms with applications to monotone convergence theorems. *J. Funct. Anal.* **28** (1978), 377–385.
- [Ste] STEIN, E. M., *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton University Press, Princeton, 1993.