

The HARPS-TERRA project I. Description of the algorithms, performance and new measurements on a few remarkable stars observed by HARPS.

Guillem Anglada-Escudé^{1,2} R. Paul Butler¹

anglada@dtm.ciw.edu,

ABSTRACT

Doppler spectroscopy has uncovered or confirmed all the known planets orbiting nearby stars. Two main techniques are used to obtain precision Doppler measurements at optical wavelengths. The first approach is the gas cell method, which consists on the least-squares matching of the spectrum of Iodine imprinted on the spectrum of the star. The second method relies on the construction of a stabilized spectrograph externally calibrated in wavelength. The most precise stabilized spectrometer in operation is HARPS, operated by ESO in La Silla Observatory/Chile. The Doppler measurements obtained with HARPS are typically obtained using the Cross-Correlation Function technique (CCF). It consists of multiplying the stellar spectrum with a weighted binary mask and finding the minimum of such product as a function of the Doppler shift. It is known that CCF is suboptimal in exploiting the Doppler information in the stellar spectrum. Here, we describe an algorithm to obtain precision RV measurements using least-squares matching of each observed spectrum to a high signal-to-noise ratio template derived from the same observations. Such algorithm is implemented in our software called HARPS-TERRA (Template Enhanced Radial velocity Re-analysis Application). New radial velocity measurements on a representative sample of stars observed by HARPS is used to illustrate the benefits of the proposed method. We show that, compared to CCF, template matching provides a significant improvement in accuracy, specially when applied to M dwarfs.

Subject headings: Methods: data analysis – Techniques: radial velocities – Stars: planetary systems – Stars: individual : Tau Ceti, HD 85512, Barnard star, Proxima, Kapteyn star, GJ 676A, ϵ Eridani, HD 69830

¹Carnegie Institution of Washington, Department of Terrestrial Magnetism, 5241 Broad Branch Rd. NW, Washington D.C., 20015, USA

²Universität Göttingen, Institut für Astrophysik, Friedrich-Hund-Platz 1,37077 Göttingen, Germany

1. Introduction

The Doppler technique has been the most successful method to detect and confirm the presence of extrasolar planets around nearby stars (Mayor & Queloz 1995; Marcy & Butler 1996). The stability of the spectrographs and the data analysis techniques used to obtain precision radial velocity (RV) measurements have been steadily improving during the last 16 years of exoplanet discoveries. In particular, the vast majority of candidates detected via Doppler spectroscopy have been obtained using two approaches : the gas cell method and the stabilized spectrograph approach.

Because an absorption cell can be installed at low cost on any general purpose echelle spectrometer, it is the most broadly used technique. In this approach, a glass cell filled with Iodine gas at low pressure is inserted just before the entrance slit of the spectrometer. Because the stellar spectrum is imprinted with Iodine prior to entering the spectrograph, the cell spectrum tracks the same instrumental distortions suffered by the stellar lines. Thanks to this, the wavelength solution and the instrumental profile can be fitted simultaneously to the Doppler shift of the star allowing nominal RV precisions at the level of 1-2 m s⁻¹. The data analysis method required to extract such precise measurements consists in forward-modeling the spectrum of Iodine multiplied to a high SNR template of the star, and convolving the product with a parameterized model of the instrumental profile. This method was pioneered by Butler et al. (1996) and has been used to infer the presence of more than 300+ exoplanets. Due to the large number of parameters involved, this technique is computationally very intensive. Also, it requires a high SNR realization of the real stellar spectrum at higher resolution than the actual observations. Because obtaining a high resolution perfectly calibrated spectrum of the star is usually not possible, obtaining a realistic template from the deconvolution of observations without Iodine is a key element of the method and is thought to be one of its major limitations (e.g., see supplementary material in Howard et al. 2010).

The other leading method to obtain precision radial velocity measurements consists of building a fiber fed, very stable spectrograph which is then wavelength calibrated using an external source such as a Th/Ar emission lamp. This technique has been developed and refined by the group lead by M. Mayor (from now on the Geneva group) over the past 20 years. This approach provided the first clear detection of an extrasolar planet around a solar-like star (Mayor & Queloz 1995) using the ELODIE spectrograph at Observatoire de Haute-Provence/France. The interested reader can find the basic elements in the construction of a fiber-fed stabilized spectrograph in Baranne et al. (1996). While the gas cell technique requires the simultaneous adjustment of the wavelength scale, instrumental profile and Doppler offset of the star; the stabilized spectrograph approach allows tackling each problem separately. First, the instrumental profile is constant by design thanks to the use of

an image scrambling system coupled with the optical fibers. In the design by Baranne et al. (1996), the wavelength calibration is obtained using two optical fibers closely packed together at the entrance of the spectrometer. Therefore, they follow almost identical optical paths within the spectrometer optics tracking almost the same optical distortions. The first fiber (science fiber) is illuminated with a wavelength calibration source (e.g., Th/Ar lamp) at the beginning of the night and the nominal absolute wavelength solution relative to that source is obtained (Pepe et al. 2002). The second fiber is fed with the same calibration source during the calibration of the science fiber and during the science observations. Although there might be a significant RV offset between fibers, they are similarly affected by changes in the instrument and, as a consequence, they share similar intra-night wavelength drifts. This second fiber is only used to monitor intra-night changes in the instrument and the information it provides is an RV drift to be added to the measured RVs. Finally, the RV measurement on a fully calibrated spectrum is obtained by the so-called Cross-Correlation Function method (or CCF). CCF is based on multiplying the observed spectrum by a weighted binary mask. The binary mask is different from zero on the nominal positions of the stellar lines, and each non-zero chunk is weighted according to the relative depth of the stellar line against the local continuum. This binary mask is then Doppler shifted and the CCF is evaluated again. The minimum of the CCF as a function of the Doppler offset is the desired RV measurement. To achieve higher precision, the shape of the CCF is centroided using a Gaussian profile. The CCF method and some details on how the binary masks are obtained is outlined in Queloz (1995) and Pepe et al. (2002).

The flagship stabilized spectrometer build by the Geneva group is HARPS (High Accuracy Radial velocity Planet Searcher) installed in the 3.6 m Telescope at the European Southern Observatory (ESO) site in La Silla/Chile (Pepe et al. 2003). Thanks to a careful design and construction (vacuum sealed tank, high mechanical stability and accurate temperature control), the wavelength solution and instrumental profile are very stable. For example, intra-night Doppler drifts as measured by the calibration fiber are typically smaller than 0.5 m s^{-1} . Long exposures ($t > 200 \text{ sec}$) using the calibration fiber cause leaks of the Th/Ar spectrum on the science spectrum, so the calibration fiber is only used when observing very bright targets and when extreme RV precision is required. The list of planets detected by HARPS is long and varied as can be seen in the 34 papers of the series *The HARPS search for southern extra-solar planets*. Instead of citing all of them, we refer the interested reader to the latest HARPS results presented in Pepe et al. (2011); Mayor et al. (2011). HARPS has demonstrated an RV stability at the level of 1 m s^{-1} on time-scales of several years.

Despite these impressive results, it is known that the CCF method implemented in the HARPS Data Reduction Software is suboptimal in the sense that it does not exploit the full Doppler information on the stellar spectrum (e.g. see Queloz 1995; Pepe et al. 2002). We

asked ourselves if a least squares approach where the observed stellar spectrum is matched to a high SNR template could be used to extract higher RV precision on HARPS observations. Given that a number of stabilized spectrographs are under construction (e.g. HARPS-North, ESPRESSO/ESO, Carmones/CaHa), and that there is significant investment in hardware development to achieve higher RV stability, it is important that the used data analysis methods are as optimal as possible. Also, precision RVs are difficult to reproduce and, given that the stakes are ambitious (detection of potentially habitable worlds), the use of different RV measurement improves the detection confidence of low amplitude signals. In this work we derive from first principles the algorithms of the template matching technique to be applied on stabilized spectrographs and implement it to public HARPS observations. HARPS is an ESO instrument and, as such, all the data obtained from it becomes publicly available after a proprietary period of a few months (or years). Since January 2011, the data products derived from the HARPS-ESO Data Reduction Software (wavelength calibrated spectra, but also CCF measured RVs among others) are publicly available through a dedicated webpage in the ESO website ¹. All the HARPS data used in this work has been obtained from there.

The core of our project is our software tool called HARPS-TERRA, where TERRA stands for *Template Enhanced Radial velocity Re-analysis Application*. HARPS-TERRA handles the full process of unpacking the HARPS-ESO archive files, generation of a high SNR template and the obtention of the final RV measurement in a single command line call. This software is custom made, fully coded in Java, and can run on any machine supporting a Java Run-time Environment 1.6 or newer.

In Section 2, we derive and describe the basic algorithms used to obtain RV measurements from HARPS reduced spectra. In Section 3, we investigate the optimal RV extraction parameters and define the *standard setup* to obtain optimal RV precision with minimum human intervention on G, K and M dwarfs. Section 4 shows the performance of HARPS-TERRA on a few representative data sets; demonstrating a significant improvement in precision, especially on M dwarfs. In the same section, we use new RV measurements obtained with HARPS-TERRA to discuss planetary systems proposed on a few representative stars observed by HARPS.

HARPS-TERRA is still in development but can be distributed for particular applications upon request. Given that other HARPS programs could benefit from using it (e.g. asteroseismology programs, binary stars), we plan a public release of the tool in the near future.

¹<http://archive.eso.org/wdb/wdb/eso/repro/form>

2. Description of the algorithms

The proposed algorithm is based on minimizing the differences of the observed spectrum against a parameterized template. In our particular implementation, we first use the higher SNR observation as a preliminary template. In a second iteration, a very high SNR template is obtained by coadding all the observations and the RVs are measured again. Note that the template will be already convolved with the instrumental profile so this method (as the CCF technique does) relies on the long term stability of the instrumental profile and wavelength calibration strategy.

HARPS is a cross-dispersed echelle spectrograph. Therefore, the stellar spectrum is split in diffraction orders over the detector (also called *echelle apertures*, or *apertures*). Each aperture has to be extracted very carefully in a complex process from the raw CCD images. The wavelength calibration is usually made on a nightly basis using a standard set of calibration frames (i.e., flat, darks, Th/Ar lamps) that are taken once at the beginning of the night. Thankfully, all this extraction and the corresponding wavelength solution is efficiently implemented by the HARPS-ESO Data Reduction Software (hereafter HARPS DRS) developed by the HARPS-ESO team², and will not be discussed here. The wavelength calibrated spectra are provided in conveniently formatted 'fits' files described in the HARPS DRS manual. As of November 2011, HARPS-TERRA is designed to work with the output files of the HARPS DRS v3.5, but should be easily adapted to future updates of the HARPS DRS and/or other stabilized spectrographs. An overview of the HARPS spectral format is given in Table 1. As described in the introduction, the secondary calibration fiber is typically only used on bright stars to achieve maximum precision. If the observer used this secondary fiber, such drift is also provided by the HARPS DRS and will be added to the final RV measurement of each echelle aperture. In all that follows we treat each echelle aperture as an independent spectrum and the final RV measurement will be a weighted mean of the RVs measured across all such apertures.

Let us assume that we have a very high SNR, wavelength calibrated spectrum of the star (template). Given a wavelength calibrated observation and a stable instrumental profile, the observed spectrum differs from the template only by a Doppler shift (due to Earth motion around the Sun and/or the presence of companions) and a flux normalization function across each aperture. We observed that this flux normalization is time dependent at the few % level most likely due to observational and instrumental effects, such as atmospheric differential refraction, differential absorption or telescope tracking errors. Let us define the difference R

²<http://www.eso.org/sci/facilities/lasilla/instruments/harps/doc/index.html>

between the template F and the observed flux f at each wavelength λ as

$$R[\lambda; \hat{\alpha}] = F[\alpha_v \lambda] - f[\lambda] \sum_{m=0}^M \alpha_m (\lambda - \lambda_c)^m \quad (1)$$

where λ is the wavelength of the observation transformed to the Solar System barycenter reference frame. The set of free parameters is represented by $\hat{\alpha} = [\alpha_v, \alpha_0, \dots, \alpha_M]$. The first term on the right side is the template evaluated at $\alpha_v \lambda_i$, where α_v is the Doppler factor on which we are interested. To derive a differential RV measurement from this Doppler factor, we use the very simple expression $\alpha_v = 1 - v_r/c$ (Stumpff 1979), where c is the speed of light and v_r is the relative radial velocity between the observer and the star on which we are interested. The second term is the observed flux f multiplied by an M-degree polynomial accounting for the flux normalization across each aperture (also called blaze function correction), and λ_c is the central wavelength of a given aperture. Note that we could apply this flux normalization to the template F instead to the observed flux f . However, this would couple the flux normalization coefficients α_m to the Doppler factor α_v in a non-linear fashion which, from a numerical point of view, is an undesirable complication.

Now this difference R can be Taylor expanded around some nominal values for the parameters $\hat{\alpha}_{(0)}$ in powers of the parameter increments $\delta \hat{\alpha}$ as

$$R[\lambda; \hat{\alpha}] \simeq R[\lambda; \hat{\alpha}_{(0)}] + \left. \frac{\partial R}{\partial \alpha_v} \right|_{\lambda; \hat{\alpha}_{(0)}} \delta \alpha_v + \sum_{m=0}^M \left. \frac{\partial R}{\partial \alpha_m} \right|_{\lambda; \hat{\alpha}_{(0)}} \delta \alpha_m \quad (2)$$

$$\left. \frac{\partial R}{\partial \alpha_v} \right|_{\lambda; \hat{\alpha}_{(0)}} = \lambda \left. \frac{dF}{d(\alpha_v \lambda)} \right|_{\alpha_v(0) \lambda} \quad (3)$$

$$\left. \frac{\partial R}{\partial \alpha_m} \right|_{\lambda; \hat{\alpha}_{(0)}} = -f[\lambda] (\lambda - \lambda_c)^m. \quad (4)$$

The weighted sum of these R over all the observed wavelengths λ_i (or pixels) is defined as

$$\chi^2 = \sum_{i=1}^{N_{pix}} \omega_i R[\lambda_i; \hat{\alpha}]^2. \quad (5)$$

and is the quantity to be minimized. The partial derivative of χ^2 with respect to each increment $\delta \alpha_v, \delta \alpha_0, \delta \alpha_1, \dots, \delta \alpha_M$, equated to 0 generates the a system of equations for such increments (the so-called normal equations) which can then can be solved using standard matrix techniques. The quantities ω_i are the weights assigned to each λ_i and their precise value will be discussed later. The resulting system of equations reads

$$\sum_i^{N_{pix}} \omega_i \left. \frac{\partial R}{\partial \alpha_v} \frac{\partial R}{\partial \alpha_v} \right|_{\lambda_i; \hat{\alpha}_{(0)}} \delta \alpha_v + \dots + \sum_i^{N_{pix}} \omega_i \left. \frac{\partial R}{\partial \alpha_v} \frac{\partial R}{\partial \alpha_M} \right|_{\lambda_i; \hat{\alpha}_{(0)}} \delta \alpha_M = - \sum_i^{N_{pix}} \omega_i R \left. \frac{\partial R}{\partial \alpha_v} \right|_{\lambda_i; \hat{\alpha}_{(0)}} \quad (6)$$

$$\begin{aligned} & \dots \\ & \dots \\ \sum_i^{N_{pix}} \omega_i \frac{\partial R}{\partial \alpha_M} \frac{\partial R}{\partial \alpha_v} \Big|_{\lambda_i; \hat{\alpha}_{(0)}} \delta \alpha_v + \dots + \sum_i^{N_{pix}} \omega_i \frac{\partial R}{\partial \alpha_M} \frac{\partial R}{\partial \alpha_M} \Big|_{\lambda_i; \hat{\alpha}_{(0)}} \delta \alpha_M &= - \sum_i^{N_{pix}} \omega_i R \frac{\partial R}{\partial \alpha_M} \Big|_{\lambda_i; \hat{\alpha}_{(0)}} \end{aligned} \quad (7)$$

where the values of R are obtained using Equation 1 and the partial derivatives are computed using Equations 3 and 4. Because the Doppler factor α_v is nonlinear, this system of equations has to be solved iteratively. Each iteration consists in 1) compute the normal equations using the current values of the parameters $\hat{\alpha}_{(0)}$, 2) solve for the parameter increments $\delta \hat{\alpha}$, 3) update the parameter values $\hat{\alpha}_{new} = \hat{\alpha}_0 + \delta \hat{\alpha}$. These equations can be written in matrix form as

$$A_{lk} \delta \alpha_k = b_l, \quad (8)$$

where A_{lk} is the $M+1 \times M+1$ matrix of coefficients multiplying the parameter increments $\delta \alpha_k$ in the normal equations 6–7. A_{lk} is equivalent to the Hessian matrix of the χ^2 and sometimes is called curvature matrix. By straight-forward error propagation (Press et al. 1992) one finds that, when the solution converges to the χ^2 minimum, the formal uncertainties in each free parameter can be derived from the inverse of the \mathbf{A} matrix as

$$\sigma_{\alpha_k} = \sqrt{(\mathbf{A}^{-1})_{kk}}. \quad (9)$$

where \mathbf{A}^{-1} is usually called the *covariance matrix*. Even though this prescription ignores correlation between parameters, it still provides a good estimate of the formal precision of the measurement if the weights ω_i are properly estimated. The formal uncertainty in the Doppler coefficient $\delta \alpha_v$ multiplied by the speed of light ($\sigma_v = c \delta \alpha_v$) is used to compute the initial weight of each echelle aperture in the computation of the mean epoch RV. Therefore one needs to be sure that these formal uncertainties are as realistic as possible (see sub-section 2.1.2)

The radial velocity measurement obtained on each order, its formal uncertainty and other information about the fit (flux normalization coefficients, number of iterations, number of masked pixels, RMS of the fit, etc.) are stored in a file for further processing (see Section 2.2). At this point, no assumption is done on the nature of the star and all the apertures are processed irrespective of their SNR or overall quality.




Fig. 1.— Result of the fit to a small chunk of the spectrum of Tau Ceti (G8.5V). The average SNR of the observation is 180. Such SNR would correspond to an RMS of 0.55% if only photon noise were involved. The actual RMS is 0.65%. Even though it is a very good fit, this slight excess of RMS illustrates that assuming Poisson statistics is usually too optimistic to obtain the weight of each pixel and derive a realistic formal RV precision. A cubic flux normalization polynomial has been applied to match the blaze function of the observations to the blaze function of the template. The nominal RV uncertainty for this aperture (aperture 54) is 5.4 m s^{-1} .

2.1. Algorithm implementation details

Even though the basic algorithm has already be outlined, a number of technical details must be carefully addressed to reach maximal RV precision in an efficient numerical fashion. Here we describe those technical aspects that require special attention and provide the practical solutions we have adopted.

2.1.1. Template interpolation

Until now, we have assumed a theoretical continuous and differentiable function for the template. In reality, we need to build the template from the same observations. Since the observed wavelengths do not necessarily coincide with the wavelengths where the template is sampled, F and its derivatives must be obtained through interpolation. In our implementation, the interpolation of the discretely sampled template is obtained using a cubic spline (Press et al. 1992). Cubic splines have the property of producing a continuous version of the derivative, which is critical for the convergence properties of the non-linear least-squares




Fig. 2.— Results of the template matching process to a small chunk of the spectrum of GJ 699 (M4V). The average SNR of the observation is 83 which should correspond to an RMS of 1.22% compared to the actual 1.23% obtained. Even though the SNR is almost a third compared to the SNR of the G dwarf in Figure 1, the much more abundant Doppler information in the spectral features of the M dwarf gives a nominal uncertainty of 2.68 m s⁻¹ for this observation.

algorithm. The details on how such a template is generated are given in Section 2.3.

2.1.2. Pixel weighting

Each weight ω_i should be a function of the uncertainty in R_i . Assuming a very high SNR template and Poisson statistics, the uncertainty in the flux of each pixel is just $\sigma_i = \sqrt{f_i}$. However, we found that assuming such uncertainty produced suboptimal results because (1) it tends to overweight pixels with high flux and little real Doppler information (continuum), and (2) Poisson statistics never applies in realistic situations (e.g., pixel non-linearities, imperfect flat fielding, unmasked telluric features, imperfect template). Instead, we re-scale the Poisson uncertainty of each pixel using an empirical approach. In the first iteration of equations 6–7, Poisson statistics are assumed on each pixel so $\omega_{i(0)} = \sqrt{f_i}$. After that, we compute the RMS of the R_i differences over all the pixels and we call it $\sigma_{(R)}$. If Poisson statistics applied, $\sigma_{(R)}$ should be equal to the square root of the mean pixel flux ($\sqrt{\langle f \rangle}$). However, if there are additional sources of noise $\sigma_{(R)}/\sqrt{\langle f \rangle} = \kappa$, where $\kappa > 1$. The weights to be used in the next iteration are therefore re-scaled with this quality factor κ as $\omega_{i(new)} = (\kappa^2 f_i)^{-1}$. The κ value of the last least-squares iteration is also stored in a file and provides useful information to assess the level of systematic noise on each echelle order.

2.1.3. Telluric masking and outlier filtering

The weights are also used to mask those pixels with undesirable properties. This is, any ω_i on wavelengths coincident with telluric absorption features deeper than 1% is set to 0. A synthetic spectrum of the atmosphere was used to identify such telluric lines and generate a list of wavelengths (or telluric mask) to be avoided. The same telluric mask is applied to all the observations.

Because of the Barycentric motion of the Earth ($\sim \pm 15 \text{ km s}^{-1}$), the stellar spectrum at the border of each echelle order is not always present in all the epochs. This going in-and-out of the borders can generate small systematic RV offsets correlated with the Earth barycentric motion. Even though a more sophisticated solution could be adopted and for the sake of simplicity, the weights of pixels within 0.8 \AA of each aperture border are also set to 0. Although some Doppler information is lost in the process, the flux at the borders is significantly lower than at the center of the apertures and some extra random noise is preferred over correlated noise with a 1 year periodicity. We tested less restrictive border masking obtaining almost identical results.

After the first least squares iteration is obtained, the weights of possible outliers are also set to 0. A pixel is considered an outlier if it has a residual 4 times larger than the empirically determined $\sigma_{\langle R \rangle}$ (i.e., $4\text{-}\sigma$ clipping). This threshold is arbitrary but we found it does a good job removing pixel outliers (e.g., cosmic ray hits), while preserving most of the well behaved ones.

2.1.4. Parameter initialization and barycentric correction

The described algorithm is a non-linear least squares solver and, as such, it requires an initial guess for the values of the free parameters. The initial value for the flux polynomial coefficient α_0 is set to the mean pixel flux $\langle f \rangle$ and all the other α_m are set to 0. Note that the real flux normalization function (or blaze function) of the template is unknown. However, we are only interested in correcting relative changes compared to the template, so a low order polynomial correction should be sufficient given a stabilized instrument such as HARPS. Section 3.1 investigates the optimal degree required for such polynomial.

All the analysis described above is done in barycentric wavelengths. The correction from observed to barycentric wavelengths is also implemented using the aforementioned simplistic recipe for the Doppler factor using the projected radial velocity of the observer as computed by Stumpff (1980) (also provided in header of the HARPS DRS files). Therefore, as a first guess, α_v is initialized to 1. Before the least-squares adjustment begins, α_v is initialized

by finding the approximate χ^2 minimum using 5 in a bisection algorithm with α_v as the independent variable. The obtained RV from this bisection is typically within 10 m s^{-1} of the final RVs, minimizing the required number of computationally expensive least-squares iterations.

Note that the simplistic definition of the Doppler factor we are using is only accurate to first order on v/c . Since we are only interested in differential RV measurements, one can show that the second order terms ($\sim v^2/c^2$ but also gravitational terms proportional to G) are mostly constant and that this expression for α_v is good at the few tens of cm s^{-1} . While higher order terms should be certainly taken into account, such refinement is useless unless a full revision of the Barycentric correction model is implemented. Critical parts in this model that need to be revised are: 1) use of an up-to-date model for Earth rotation, 2) use of the most recent version of the ephemeris for Earth (e.g., DE405 or newer, Standish 1998) and 3) inclusion of all the relativistic corrections to the first post-Newtonian order (gravitational and kinematic). Even though such model might improve some of the RVs presented here, further work is required to produce a functional and reliable code. A tool to provide Barycentric Doppler corrections to 1 cm s^{-1} level is in development and we plan to present it in a future publication.

2.1.5. Convergence criteria

Since the Doppler factor is the only strongly non-linear parameter, the convergence criteria is established as the iteration requiring a $\delta\alpha_v \times c$ smaller than 1 cm s^{-1} . Thanks to the simplicity of the model, only 3 to 6 iterations are typically required to reach convergence.

The typical nominal uncertainty of an echelle aperture with a SNR of 100 is between 2 to 15 m s^{-1} , strongly depending on spectral type and aperture under consideration. Examples of template matched observations for a G8.5V star (Tau Ceti) and an M4 dwarf (GJ 699) are shown in Figures 1 and 2.

2.2. Obtaining the final RV measurements

A final refinement is useful to produce the highest quality RV. Because the template has uncertainties, there can be ambiguous radial velocity zero-point offsets associated to each aperture. Also, some apertures might have extra uncertainty due to instrumental effects not accounted in the formal uncertainties (extraction issues, problems with the wavelength

solution, etc.). With the purpose of computing a more realistic weighted mean to each epoch, the zero–point and nominal uncertainty of each order can be empirically reassessed as follows.

As a first step, the weighted mean μ_e of each epoch e is computed as

$$\mu_e = \frac{1}{Z} \sum_{a=0}^{N_a} \frac{v_{e,a}}{\sigma_{e,a}^2}, \quad (10)$$

$$Z = \sum_{i=0}^{N_a} \frac{1}{\sigma_{e,a}^2}. \quad (11)$$

where the subindex a runs over all the N_a apertures being analyzed, $v_{e,a}$ is the RV measurement of the a -th aperture at epoch e , and $w_{e,a}$ is the corresponding formal uncertainty in the RV (σ_{α_v} as obtained from equation 9 multiplied by the speed of light). Z is the sum of all the weights and normalizes the weighted mean.

Let us now concentrate in one aperture, say aperture 15. On each epoch e , we compute the difference between the epoch weighted mean μ_e and $v_{e,15}$. The weighted average of this difference over all the epochs is the zero–point of order 15, which we call $v_{15}^{(0)}$. The weighted standard deviation of these differences over all the epochs is the new nominal uncertainty $\sigma'_{e,15}$. This process is then repeated for all the orders to obtain the corresponding $v_i^{(0)}$ and $\sigma'_{e,i}$. After all the weights and zero-points are obtained, the final radial velocity measurement and its precision on a given epoch is given by

$$RV_e = \frac{1}{Z'} \sum_{i=0}^{N_o} \frac{v_{e,i} - v_i^{(0)}}{\sigma_{e,i}'^2}, \quad (12)$$

$$\sigma_e^2 = \frac{1}{Z'(N_o - 1)} \sum_{i=0}^{N_o} \frac{(v_{e,i} - v_i^{(0)})^2}{\sigma_{e,i}'^2}, \quad (13)$$

$$Z' = \sum_{i=0}^{N_o} \frac{1}{\sigma_{e,i}'^2}. \quad (14)$$

Note that, because $v_i^{(0)}$ and $\sigma'_{e,i}$ are computed with respect to the initial guess of the epoch mean value, the zero–point correction does not remove offsets shared by all the apertures (i.e. Keplerian signals are not removed). In case the user is interested in the RV measurements without this re-normalization of the weights, the original μ_e and the individual RV measurements of each order are also provided as a data product by HARPS-TERRA.

As mentioned in the introduction, these algorithms and all the code necessary to extract the HARPS spectra from the ESO reduced data-products are included in the HARPS-TERRA software. HARPS-TERRA has been designed to achieve the highest precision but while being computationally efficient. It takes between 1 and 5 seconds to process one full spectrum on a Linux operating system running on a 2.0 GHz processor. The construction of a high SNR template (see next Section) takes around 5 minutes when using 200 spectra. A data set consisting on 100 spectra can be fully processed in less than 10 minutes.

2.3. Construction of the template

In order to obtain maximal precision, one would like a template with the highest possible SNR. This can be achieved by carefully coadding all the available spectra. Still, some precautions must be taken in the construction of such template. This is an outline of the process we use to generate them.

In a first pass over all the spectra, the RV and flux normalization coefficients are obtained with respect to a preliminary template (highest SNR observation). This preliminary RV measurement and the heliocentric motion of the observer are used to obtain the barycentric wavelengths of each observed spectrum at each epoch.

Spectra obtained in different epochs are not sampled at the same barycentric wavelengths and cannot be co-added without some kind of interpolation. To solve this, the preliminary template is used as a reference to generate a grid of regularly spaced reference wavelengths. The number of re-sampled reference wavelengths is 4 times the number of original pixels in each aperture of the preliminary template (this is $4096 \times 4 = 16384$). Each observed spectrum can then be interpolated on such reference wavelengths using a cubic spline. Finally, the flux normalization is applied and the template value at each re-sampled wavelength is computed as a $3\text{-}\sigma$ clipped mean over all the epochs.

The flux normalization polynomial is only applied if the average SNR of a given aperture is higher than 5. SNR lower than 5 are not rare on the bluer apertures of M dwarfs and the corresponding normalization polynomials are very unreliable. Because the blaze variability is of the order of 1%, it is safer to simply apply a scale factor and match the average flux of the observation to the average flux of the preliminary template.

Telluric features must also be removed from the coadding. As in the RV measurement, the flux measurements coincident with telluric features deeper than 1% are masked out. Let us note that, if a star has been observed in different phases of the Earth motion around the Sun, this process allows to generate a telluric free spectrum on regions with mild telluric

contamination. When a significant number of spectra is available ($N > 20$), the result of this coadding can be very spectacular, specially on M dwarfs whose spectrum has almost no continuum and it is very hard to distinguish pure noise from real features (see Figure 2 as an example).

3. Performance

Even though the described algorithms are relatively simple, the extraction of precise radial velocities still depends on a number of parameters that have to be tuned by hand. The two parameters we investigate here are; (1) the optimal degree for the flux normalization polynomial, and (2) bluest echelle aperture to be used (e.g., M dwarfs can have 10 to 100 times more flux on the red than on the blue and low SNR spectra are typically more affected by systematic noise). HARPS-TERRA has been designed to be very flexible on all such parameters but a general procedure to produce optimal results with minimal human intervention is still desirable. To illustrate the effect of these two parameters we use observations of Tau Ceti (GJ 71, HD 10700, a very stable G8.5V dwarf), HD 85512 (GJ 370, a quiet K6V dwarf with a very low amplitude candidate planet) and Barnard star (GJ 699, a relatively quiet halo M4 dwarf). These three stars have abundant data in the ESO archive and bracket the highest priority targets for the search of very low mass companions (G, K and M dwarfs). The direct comparison of the final RVs obtained with CCF and HARPS-TERRA on a larger sample is given in Section 4. All the stars we discuss here are nearby and show a significant linear trend due to perspective acceleration (Zechmeister et al. 2009). In all that follows, such perspective acceleration has been subtracted from the measured RVs.

Tau Ceti. For this experiment we use 84 spectra of the 4000+ available in the HARPS-ESO archive. Since it is a very quiet and stable star, Tau Ceti had been used as a standard star by several HARPS programs. Not all such programs aim to achieve the highest RV precision. This results in a very heterogeneous sample of public observations with varying exposure times and a wide range of SNR. The SNR of some of the available spectra is so high (> 500) that saturation or extraction problems seem to be seriously affecting the reddest orders. For a consistent analysis, we prepare a sub-sample from the *HARPS high precision survey for exoplanets in the southern hemisphere* (e.g. Lovis et al. 2006; Mayor et al. 2009). This subset also belongs to the sample presented in (Pepe et al. 2011) showing an RMS of 0.92 m s^{-1} over a time-span of 5 years. Pepe et al. (2011) noted that, to achieve such precision, several spectra in a given night had to be averaged to mitigate the effects of stellar pulsation and granulation. Because this star is one of the 10 higher priority targets in the *HARPS-GTO survey for Earth-like planets*, only the observations taken before 2009

Table 1. HARPS spectral parameters relevant to this work

Parameter	Value
Spectral resolution $\lambda/\delta\lambda$ at 5500Å	120 000
Number of diffraction orders (or echelle apertures)	72
Pixels in each aperture	4096
Wavelength range (Blue CCD)	3780 – 5300 Å
Wavelength range (Red CCD)	5330 – 6910 Å
Sampling per resolution element	4.1 pix

Table 2. Relevant parameters to this work of each star. A more detailed description can be found in the SIMBAD database and references therein. All quantities are given to the last significant digit.

Parameter	GJ 676A	Tau Ceti	HD 85512	Barnard’s	Kapteyn’s	Proxima	ε Eri	HD 69830
$\mu_{R.A.}^*$ [mas yr ⁻¹] ^a	-260	-1721	461	-798	6505	-3776	-975	279
μ_{Dec} [mas yr ⁻¹] ^a	-184	854	-472	10328	-5731	766	19	-987
Parallax [mas] ^a	61	274	90	548	256	772	311	80
V ^b	9.58	3.5	7.651	9.51	8.85	11.05	3.73	5.95
K ^b	5.82	1.79	–	4.52	5.05	4.38	1.78	4.16
Sp. type ^b	M0V	G8.5V	K6V	M4V	M1V	M6V	K3V	G8V
Mass [M _⊙] ^c	0.71	0.78	0.69	0.16	0.27	0.12	0.82	0.86

^aProper motions and parallaxes from HIPPARCOS (van Leeuwen 2007) are required to subtract the perspective acceleration effect (see text). $\mu_{R.A.}^*$ corresponds to $\mu_{R.A.} \cos \delta$ and is the proper motion in the direction of increasing R.A. in a local tangent plane defined on the star’s nominal coordinates at the catalog reference epoch. $\mu_{R.A.}$ is the obsolete coordinate dependent definition of the secular change in R.A., which is singular at the celestial poles and has not direct physical interpretation.

^bUnless noted in the manuscript, V and K photometry and nominal spectral type are obtained from the SIMBAD database.

^cStellar masses for M dwarfs have been derived from Delfosse et al. (2000) using absolute K magnitudes. These masses should be accurate at the 5-10% level. Masses of the K and G dwarfs are obtained from various references (see section on each star).

are publicly available. In any case, we only need a statistically significant sample with consistently measured RVs using the CCF method to be used as a reference. Still, to work with a more manageable sample, we only use the first observation of each night. The final sample contains exposure times between 30 and 150 seconds and a typical SNR in the range between 100 to 250 at 6000 Å. The CCF RVs for this sub-sample show an RMS of 1.53 m s⁻¹. The RMS from HARPS-TERRA using all the apertures is 1.52 m s⁻¹. Since we are only using one spectra per night; this sub-sample shows, as expected, a larger RMS than the nightly averages used in Pepe et al. (2011).

HD 85512 (GJ 370), is a quiet K6V dwarf with a very low amplitude candidate planet (Pepe et al. 2011) that could support liquid water if it were significantly covered by clouds (Kaltenegger et al. 2011). It is also a target of the current HARPS-GTO program to search for low mass companions, so only the first 5 months of observations (Dec 2008 to March 2009) are publicly available. The typical integration times are between 400 and 600 seconds and the SNR at 6000 Å ranges from 100 to 250. For our purpose here, it is only relevant that the CCF measurements show an RMS as low as 1.10 m s⁻¹ using a K5 binary mask (a very stable star indeed). The HARPS-TERRA RVs from the same 122 observations show an RMS of ~ 1.0 m s⁻¹, which already represents a significant improvement. For comparison, the proposed planet candidate has an RV semi-amplitude of 0.8 m s⁻¹ and the detection is based on 250+ measurements in Pepe et al. (2011). A quadratic trend (also present in the Ca H+K S-index activity index) had to be removed to cleanly detect the candidate in Pepe et al. (2011). No trend nor planet signals were subtracted to the RVs discussed here.

Barnard’s Star (GJ 699) is the star with the highest proper motion and the second closest star system to the sun. It is an M4 dwarf with halo kinematics (total velocity with respect to the Sun is ~ 150 km s⁻¹) and it is known to be slightly metal poor. Even though it is a low mass star and relatively faint in absolute terms, its proximity to the Sun (1.82 pc) allows to obtain a typical SNR between 50 and 80 at 6000 Å in 900 sec integrations. 22 spectra are available in the HARPS-ESO archive over a time-span of one year (April 2007 to May 2008). Even though it is classified as active (V2500 Oph), only occasional flares have been reported on it. Barnard star was observed within the *ESO-UVES search of low mass companions around M dwarfs* (Zechmeister et al. 2009). Those measurements demonstrated its radial velocity stability down to 2.5 m s⁻¹ using the Iodine cell technique. They also found that a possible RV wobble with a 40 days period was significantly correlated with the strength of its H α emission. Therefore, we *a priori* expected that the RV measurements obtained with HARPS would also show some activity induced jitter. The CCF RVs as extracted from the archive are obtained using an M2 binary mask and show an RMS of 1.54 m s⁻¹. The RVs from the HARPS-TERRA RV measurements using the full spectrum have an RMS of 1.23 m s⁻¹, which again represents a significant improvement. A quick-look analysis of the RVs

does not show evidence of the previously reported periodicity at 40 days, but the number of observations is too low and the time-sampling is too sparse to rule out signals in this period domain.

Two more stable M dwarfs. In addition to Barnard’s star, we will briefly use HARPS RV measurements of two other M dwarfs : Proxima (GJ 551, M6Ve) and Kapteyn’s star (GJ 191, M1V subdwarf). The data set on Proxima consists on 27 radial velocities taken between May 2005 and February 2009. The CCF RMS is 2.29 m s^{-1} and recurrent flaring events can be detected in more than one epoch. The data set on GJ 191 contains 30 spectra taken between December 2003 and February 2009. The CCF RMS is 2.45 m s^{-1} . These two stars will be used to illustrate some features that seem to be common on M dwarfs other than Barnard’s.

3.1. Flux normalization polynomial

Because RV measurements rely on the slopes of strong spectral features, the flux normalization correction is a key element to achieve the highest possible RV precision. For example, much better precision is obtained from deep-sharp lines than broad-shallow ones. In a perfectly calibrated instrument, the blaze function should be constant over time. However, several instrumental/observational effects can cause variability of the effective blaze function. For example, due to atmospheric differential refraction, the photocenter of a star at the entrance fiber will depend on wavelength causing wavelength dependent flux losses as a function of the airmass. Also, bluer wavelengths are more efficiently dispersed by the atmosphere adding additional airmass dependent variability. At the end of the day, the combination of several effects causes time variable blaze function shapes at the level of a few % which (as shown below) can severely perturb the RV measurements.

Because we are interested in relative measurements, we only need to correct for the differential variations of the blaze as compared to the template. In figure 3, we show the RV measurements of our three test stars when using different flux normalization polynomials. When only a constant normalization is applied (0 degree polynomial), the scatter in the radial velocities of the G and the K dwarfs is very large and has a clearly systematic behavior. The RV precision dramatically improves when using a linear flux correction (first degree polynomial) on both stars. No significant improvement in precision is obtained by using polynomial degrees higher than 3. On the other hand, the RV measurements on the M dwarf are much less sensitive to the degree of the polynomial used. The improvement on the precision is modest but still significant with a linear correction, and optimal results are also obtained when a cubic polynomial is applied.




Fig. 3.— Radial velocity measurements as a function of time obtained on the three test stars using different flux normalization polynomials. Unless corrected, the flux variability strongly perturbs the RV measurements on the G star (left) as can be seen in the obtained RMS when only an overall normalization factor is applied (order 0 in black). The effect of the flux variability is still significant for the K dwarf and has very little effect on the M dwarf. An order 1 polynomial (blue) provides a significant improvement on both the G and the K dwarfs and an order 3 polynomial (red) seems to reach a good compromise between computing efficiency and precision on all spectral types. RV offset has been added to the measurements of the 0 and 1 degree polynomials to improve visualization.

The sensitivity to the flux normalization correction as a function of the spectral type was expected. The spectra of typical G and K dwarfs consist on well isolated sharp lines against a smooth continuum. If not properly corrected, the changes in the slopes induced by the blaze variability on the continuum contain spurious Doppler information that strongly perturbs the Doppler measurement. On the other hand, the spectrum of an M dwarf is dominated by heavily blended molecular absorption bands which strongly dominate over the shifts induced by blaze variability. Given that high degree flux corrections are computationally expensive, we set the nominal blaze correction to a cubic polynomial.

3.2. Most useful echelle apertures

The final uncertainty in the radial velocity measurement depends on many instrumental effects in addition to the formal statistical uncertainties and the stellar spectrum. K and, specially, M dwarfs have significantly less flux in the blue than in the red. If noise in bluer wavelengths were purely random, these apertures would be properly down-weighted through the formal uncertainties and the final measurement would be unaffected. However low SNR spectra can be more sensitive to instrumental systematic effects, so using arbitrarily low SNR

data can be counterproductive. As an example, the M2 binary masks used by the HARPS DRS, only uses apertures redder than the 22nd one ($\lambda > 4400\text{\AA}$). Also, given that all the stars are active at some level and that activity should affect the measured RV differently at different wavelengths (Reiners et al. 2010), one could expect some apertures to provide more reliable measurements than others. As an example, star spots are known to have higher contrast at bluer wavelengths, so one would expect stronger RV stellar jitter on the blue. The wavelength dependence of the stellar jitter has been exploited in the past to rule out possible companions around young stars using complementary RV measurements in the near infrared, e.g., see the RV measurements obtained in the H band by Huélamo et al. (2008); Figueira et al. (2010) on TW Hya using the CRIRES spectrograph at the ESO/VLT. This section is devoted to develop a strategy to determine the wavelength dependence of the RV precision and define the bluest aperture to be used for each spectral type.

Given that the typical formal uncertainty of a given aperture is never better than $2\text{--}3 \text{ m s}^{-1}$, it is hard to distinguish random noise from systematic effects (intrinsic to the star or instrumental) when looking at the RV measurements of single echelle apertures. Instead, we use the following procedure to assess the dependence of the precision with the bluest aperture used. First, we extract the radial velocities on all the apertures. Then we measure RVs on all the epochs as described in Section 2.2 but only using the apertures between 71 (reddest available one) to some bluer one (say aperture 68), and obtain the RMS of these time series. We then repeat this process adding one aperture at a time and plot the obtained RMS as a function of the bluest aperture used. This is illustrated on Figure 4 on our three test stars. On the right hand-side of each panel, the RMS is expected to be higher because less apertures are used. However, if the noise were purely random, we would expect a reduced RMS as more apertures are added to the blue. The K6V dwarf nicely follows this behavior.

However, the M4V star behaves quite differently. The RMS reaches a minimum (81 cm s^{-1}) at aperture 58 when only the 14 reddest apertures are used. Then, it starts increasing as more apertures are included towards the blue ($\sim 1.2 \text{ m s}^{-1}$ at HARPS aperture 0). The typical SNR is still very significant on apertures bluer than 58 ($\text{SNR} > 30$) so instrumental systematic effects are not expected to be important at this level. This combined with the fact that the K6V star does not show such an excess, suggests that a significant fraction of this extra noise is not instrumental but intrinsic to the star. For this star in particular, the 58-th aperture ($\lambda \sim 6000\text{\AA}$) seems to be the middle point where the activity induced variability and the photon+instrumental noise equally contribute to the error budget. This behavior is also present in other *stable* M dwarfs. For example, we show the same diagrams for Proxima (M6V) and Kapteyn’s star (M1.5V, GJ 191) in Figure 5. Because both stars are fainter than Barnard’s, one has to use bluer apertures to reach the point where the instrumental noise meets this wavelength dependent jitter. In an ideal situation the bluest aperture should be




Fig. 4.— Radial velocity RMS as a function of the bluer aperture used. The RMS derived from the public CCF measurements is indicated as a horizontal line. The bluer aperture used in the CCF analysis is marked as a vertical thick line. While the RMS of the K6V dwarf decreases as bluer apertures are added, neither the M dwarf nor the G dwarf show the same behavior. The origin of this RMS excess in the blue is still unknown. The fact that it only shows on some stars very stable stars, indicates that is likely due to stellar induced jitter rather than an instrumental effect.

selected for each star, but if this effect is really related to stellar activity, this sweet spot is difficult to predict a priori. This behavior is under investigation and hints towards new ways of assessing if a signal could be activity induced. Also, the wavelength dependence of the RMS illustrates the importance of moving to redder wavelengths to efficiently search for low mass companions around cool stars (Reiners et al. 2010). Figures 4 and 5 also show the RMS obtained from the CCF measurements (horizontal line) illustrating that the template matching technique works better even without *cherry picking* the bluer aperture to be used. As mentioned before, the CCF M2 binary masks used by the HARPS DRS do not consider apertures bluer than the 22nd. Given that aperture 22 also seems a reasonable choice for HARPS-TERRA, we will also use it as the default bluer aperture to be used when extracting RVs from M dwarfs.

Tau Ceti is known to be one of the most RV stable G dwarfs. A bit unexpectedly, we find that the RMS as a function of the bluer aperture also shows a minimum of 1.36 m s^{-1} at aperture 40, and then it starts increasing again towards the blue (1.52 m s^{-1} at aperture 0). Given that the SNR is still high (~ 40 at order 0), again this should not happen in a perfectly stable star. As for the M dwarf, this indicates that some scatter is due to stellar activity and has a chromatic component (e.g. stellar pulsation should be achromatic). The RMS obtained at the sweet-spot (order 40th) can be used to estimate the magnitude of this chromatic noise as $1.36/\sqrt{2} \sim 0.96 \text{ m s}^{-1}$, which is quite significant. For comparison, an Earth-mass in the habitable zone on Tau Ceti would have an RV amplitude of 10–20 cm s^{-1} .




Fig. 5.— RMS as a function of the bluer aperture used on Proxima and Kapteyn’s star (GJ 191) showing a similar behavior as Barnard star. This is, the RMS starts decreasing as bluer wavelengths are added until it starts increasing again (GJ 191) or does not change significantly (Proxima). For Proxima, the SNR below aperture 10 is so low ($\text{SNR} < 5$) that adding such apertures to the analysis only adds systematic noise to the RV measurement. As a general rule, the cut at aperture 22 used by the HARPS DRS seems a reasonable assumption.

Given that it is one of the most quiet Sun-like stars being surveyed for rocky planets, this issue requires a more detailed analysis which is beyond the scope of this paper. Given the increased precision achievable by selecting the sweet spot aperture, a reprocessing of the full data set in Pepe et al. (2011) with HARPS-TERRA could lead to a significant increase in the overall precision and enhanced sensitivity to lower mass companions.

In conclusion to this section, we find that a cubic polynomial is sufficient to obtain an optimal flux normalization correction. The maximal precision on quiet stars (such as HD 85512) is obtained using the whole spectrum. However, in several stars a moderate improvement in the overall precision could be achieved by finding the sweet spot between random noise (instrumental+photon) and the wavelength dependent jitter. The determination of this sweet-spot requires a better understanding of its physical origin and will not be further discussed here. To make a fair and consistent comparison, all the apertures will be used when analyzing spectra of G and K dwarfs. For M dwarfs, only apertures redder than aperture 22 ($\sim 4400 \text{ \AA}$) will be used. This bluest aperture used together with the cubic flux normalization polynomial discussed in Section 3.1 define the *standard setup* of HARPS-TERRA when analyzing spectra from G, K and M dwarf stars.

3.3. Ca II H+K activity indicator

Pseudo-radial velocity variations can be caused by stellar activity. As mentioned before, it is suspected that some apparent RV offsets are closely related to the magnetic activity of the star and related surface features such as spots. Also, enhanced magnetic activity can cause local or global changes in the convection patterns of the stellar surfaces. When convection is enhanced, hotter and bluer material emerging from the convection cells (e.g. stellar granulation) causes apparent blue shifts to the integrated stellar spectrum. For one reason or another, one could expect apparent radial velocity jitter whenever the magnetic field of the star experiences changes. A detailed discussion on the topic can be found in Lovis et al. (2011) and references therein. Lovis et al. (2011) also provides the recipes to compute one of the most commonly used activity indicators, the Mount Wilson S-index, using HARPS data. This index measures the relative flux of the Ca II H and K lines in emission ($\lambda_H = 3933.664\text{\AA}$ and $\lambda_K = 3968.470$) compared to a local continuum. These lines in emission are formed in the hot plasma of the chromospheres of stars and, as a consequence, their intensity varies with the strength of the stellar magnetic field. We incorporated the automatic measurement of the S-index as one of the outputs of HARPS-TERRA. To obtain the correct flux estimates, it is necessary to know the absolute heliocentric radial velocity of the star at few hundred m s^{-1} accuracy. Since HARPS-TERRA cannot provide this information, we use the median of the heliocentric CCF RV measurements to estimate the barycentric wavelengths of the lines on all the epochs. Because the H and K lines and the continuum bands defined in Lovis et al. (2011) appear in different echelle apertures (aperture 5 and 6), we use the re-sampled, blaze corrected full spectrum also provided by the HARPS DRS. The full spectrum is then interpolated using a cubic spline and the indices are computed on a regularly sampled grid of 0.01\AA conveniently Doppler shifted to match the heliocentric radial velocity of the star.

To validate this procedure, we obtained time-series of the S-index on published HARPS stars obtaining perfect agreement in all cases. The S-index measurements on Tau Ceti and HD 85512 are illustrated in Figure 6 and show the same behavior reported by Pepe et al. (2011). No coherent variability is observed on Tau Ceti (relative scatter is 0.7%), and long term variability is observed on HD 85521. The S-index as a function of time is also plotted for Barnard’s star. Barnard’s (M4V) shows a similar amount of relative variability as the K dwarf except for a mild flaring event that doubles the Ca II H+K flux in one of the epochs. In case a tentative signal is found, the time series of the S-index will be used to investigate possible correlation of the RVs with magnetic activity. An example of this is illustrated in section 4.4, where we find that a promising long period quadratic trend in the RV is correlated with an almost identic trend in the S-index. A popular form of the same activity indicator is the so-called $R'_H K$ index. $R'_H K$ represents the flux in emission of the Ca H+K lines over the total stellar spectrum and requires some assumptions on the spectral




Fig. 6.— S-Index time series on Tau Ceti, HD 88512 and Barnard’s star. The mean value $\langle S \rangle$ and the relative variability of the index $\sigma / \langle S \rangle$ are given for each star. One of the epochs for Barnard star shows an S value two times higher than usual which is indicative of a flaring event occurring during the observation. That point was excluded from the computation of the mean S value and its relative variability. The error bars represent the photon noise in the measurement of S.

energy distribution of the star. Still, $R'_H K$ is obtained as a linear relation from S and, as a consequence, it will experience the same time variability. Since the computation of $R'_H K$ requires additional information about the star (e.g. effective temperature and metallicity Lovis et al. 2011) and does not provide extra information about the time variability, $R'_H K$ is not provided by HARPS-TERRA.

The HARPS DRS also provides three other activity indicators derived from the shape of the Cross-Correlation Function. The first one is the so-called Bisector Span of the CCF (BIS). BIS is a measure of the asymmetry of the average spectral line and should correlate with the RV if the observed variability is caused by spots or plages rotating with the star (Queloz et al. 2001). The second index is the full-width-half-maximum (FWHM) of the CCF and is a measure of the width on the mean spectral line. The variability of the FWHM is thought to be a direct consequence of changes in the convective patterns on the surface of a star; effectively changing the shapes of the integrated line profiles but could also depend on other physical processes related to the stellar magnetic field. The third index is the contrast of the CCF (CONTRAST), which is sensitive to the changes in the depth of the average spectral line profiles. These three numbers provide important diagnostics to distinguish genuine Doppler signals from activity induced periodicities. Since they are already computed by the HARPS DRS, they will not be further discussed here. Equivalent activity indicators optimally designed for the template matching technique are in development and will be given in a future publication.

4. Comparison : CCF vs template matching

Comparing actual radial velocities is more informative than discussing which method (CCF or template matching) works better on theoretical grounds. This section is devoted to illustrate the performance of the template matching approach compared to the pipelined HARPS CCF measured RVs. We discuss a few remarkable systems in terms of reported planet abundance but also their reported RV stability. We show that, while the performance on G and K stars is similar using both techniques, template matching works significantly better on M dwarfs and moderately active stars. Given the relative youth of the HARPS-TERRA code compared to the many years of refinement of the HARPS-CCF algorithms, we can only expect further improved precision in the future. In addition to alternative RV measurements, the template matching approach should allow us to perform a new set of diagnostics such as the determination of the bluest aperture to be used for an optimal RV extraction already discussed in Section 3.2.

Whenever signals are present and orbital fits are required, we use the SYSTEMIC interface (Meschiari & Laughlin 2010) as provided in August 2010³ to obtain the orbital parameters and their uncertainties. SYSTEMIC allows the interactive adjustment of multi-planetary systems and is able to generate a large variety of data products and figures. When periodograms are required to illustrate the detection of a signal, we use a custom made version of a least-squares periodogram. A sinusoidal signal is adjusted to a list of 10^4 test periods between 1.1 and 10 000 days. The F-ratio statistic of the least-squares solution is then plotted against the period so the higher peak represents the most likely periodicity in the data. These periodograms are based on the definitions given in Cumming (2004), and they are formally equivalent to the so-called generalized Lomb-Scargle periodograms (GLS) described by Zechmeister & Kürster (2009). The CCF and HARPS-TERRA RV measurements used in this section are given Appendix A.

4.1. Proxima, a flaring M4V

Proxima Centauri is the nearest stellar neighbor to the Sun and, therefore, has a special interest among the planet hunting community and the public in general. So far, there have not been any serious claim of a companion around this star. It is the common proper motion pair to the α Centauri binary, two stars with masses similar to the sun in a long period orbit. Proxima Cen has been intensively monitored by the ESO/UVES planet search program at

³<http://oklo.org>




Fig. 7.— Measured radial velocities on Proxima using HARPS-Terra (black) compared to the ones obtained with the CCF method (red)

a median precision of 3 m s^{-1} using the Iodine cell technique and no significant periodicity was detected (Endl & Kürster 2008). Between 2004 and 2009, it has been observed 27 times with HARPS on different programs.

Visual inspection of Proxima’s spectrum strongly suggests it is an active star as several activity indicators in the HARPS wavelength range are in strong emission (H_{α} , NaD, Mg, Ca, etc). Perhaps for this reason and because it is relatively faint, the star has not been as intensively monitored as other earlier type nearby M dwarfs such as GJ 581 (Forveille et al. 2011b) or GJ 876 (Rivera et al. 2010). The HARPS-TERRA RVs have been obtained using the standard setup for M dwarfs. Figure 7 shows the CCF and HARPS-TERRA measurements as a function of time. In July 16 2004, Proxima experienced an energetic event (probably a flare) causing all the activity indicators to go to strong emission. The flare event happened on the second epoch and the emission lines (such as H_{α}) show intensities 2 to 10 times stronger than their quiescent state (S-index goes from a median value of 8.7 to 58). Still, neither the CCF nor the HARPS-TERRA measurement show a significant offset on that particular epoch. Instead, the largest CCF offset (7.5 m s^{-1}) occurred 3 days later with no apparent counter part in the activity indices. For comparison, the HARPS-TERRA measurement on this same is only 3.2 m s^{-1} away the mean RV. When excluding the RV outlier, the RMS of the CCF goes from 2.37 to 1.98 m s^{-1} and TERRA goes from 2.05 to 1.88 m s^{-1} . The extra noise to be added in quadrature to the TERRA RMS to match the RMS of the CCF is 1.2 m s^{-1} considering all the measurements, or 0.6 m s^{-1} if the outlying event is excluded. An overview of the data sets on Proxima Cen and the other two M dwarfs discussed in previous sections (Barnard’s and Kapteyn’s) is given in Table 3.

Concerning possible signals, the HARPS-TERRA RVs periodogram show a very marginal peak at 5.6 days that is also barely visible in the CCF values when the outlying RV measurement is removed. So, unfortunately, no promising signals are yet detected on Proxima




Fig. 8.— Measured radial velocities using CCF left and HARPS-TERRA on the right. Only the 34 epochs in the archive coincident with the published RVs are used for this fit. This figure clearly shows that the observed reduction in the RMS of M dwarfs is not a spurious effect of the data analysis techniques we apply.

Cen.

4.2. GJ 676A, an M1V with an eccentric gas giant

In the previous sections, we have illustrated the improvement on the precision thanks to the use of the template matching technique on stars with no planets. To illustrate that the increased precision is not an artifact of the data reduction process, we now analyze the spectra of a planet hosting M dwarf with a large amplitude signal. GJ 676A is orbited by a gas giant candidate with a period of 1060 days and a RV semi-amplitude of $\sim 120 \text{ m s}^{-1}$. GJ 676Ab was announced by Forveille et al. (2011a) and is one of the few gas giants detected around low mass stars. The RV measurements provided by the HARPS DRS contain two epochs where the CCF failed to converge giving spurious offsets of 38 km s^{-1} and 76 km s^{-1} respectively. The HARPS-TERRA measurements in these two epochs look perfectly reasonable. For comparison purposes, these two points were removed from the orbital analysis. As seen in figure 8, the HARPS-TERRA RVs show essentially the same RV signal as the CCFs, demonstrating that our fitting procedure is not knocking out real RV offsets and that our RMS estimates on stable stars are realistic.

After subtracting the best fit solution for one planet, the CCF RVs shows an RMS of 6.31 m s^{-1} , while the HARPS-Terra ones have an RMS of 6.08 m s^{-1} . Given that the typical

photon noise is of the order of 1-2 m s⁻¹, it is obvious that something else is happening on this star. As suggested by Forveille et al. (2011a), we added a linear trend to the fit and adjusted all the free parameters again. The improvement on both fits is quite significant (HARPS-Terra RMS is 3.13 m s⁻¹ and the RMS from the CCF values is 3.77 m s⁻¹, see Table 4). As already noted by (Forveille et al. 2011a), the fact that the detected trend is significantly larger than the maximal expected acceleration due to GJ 676B strongly suggests the presence of an additional companion to GJ 676A with a period of several thousand days at least. Table 4 contains the best fit to the CCF and TERRA data sets. Only those epochs (34 of them) that are present in both data sets are included in both fits. The uncertainties correspond to the 68% confidence intervals as obtained using the Monte Carlo Markov Chain algorithm included in SYSTEMIC. The MCMC jump lengths β of each parameter were tuned so the acceptance rate of all parameters was between 15 and 30% (Ford 2005) and 5 $\cdot 10^6$ MCMC iterations were used to generate the desired distributions and confidence levels in Table 4. The same data analysis technique will be used to characterize the uncertainties in the orbital parameters given in the forthcoming sections.

As we have seen on the other *stable* M stars, the post-fit RMS from TERRA is significantly lower than the one obtained with CCF, giving further proof that, for M dwarfs, template matching significantly outperforms CCF. In this case, the uncertainty that has to be added in quadrature to the HARPS-TERRA measurements is 2.1 m s⁻¹, which is very significant. As noted in the discovery paper of GJ 676A, further low mass planets could be present around GJ 676A in shorter periods. Given the significantly better accuracy, it is likely that such signals could already be detected applying HARPS-TERRA to the complete data set. However, even though these results were already published, only 38 spectra are available through the data base compared to the 69 used in Forveille et al. (2011a), so the full reanalysis cannot be done here. Also, we found that several RV measurements are missing in Forveille et al. (2011a). We checked that these missing measurements coincide most of the time with the outliers found in the archive. By comparison, no spectrum failed to provide a useful RV measurement using HARPS-TERRA demonstrating the robustness of the algorithms.

4.3. ϵ Eridani, active K3V with a gas giant companion?

ϵ Eridani is a K3V star of approximately 0.82 M_⊙ (Butler et al. 2006) and a close neighbor to the Sun (3.2 pc). ϵ Eridani was first reported to be RV variable by (Campbell et al. 1988) and (Walker et al. 1995) explicitly reported a possible variability with a period between 5 and 10 years. A 6.9 years periodicity was first reported in Cumming et al. (1999)

but, at that time, it was considered suspicious given the high chromospheric activity on the star. Using additional RV measurements, Hatzes et al. (2000) proposed that the observed 6.9 years variability was most likely caused by the presence of a planet. The proposed planet would have a significantly eccentric orbit ($e \sim 0.7$) and had a minimum mass of $0.86 M_{jup}$. The reanalysis of several RV data sets combined with Fine Guidance Sensor/Hubble Space Telescope astrometry indicated that the companion was an actual planet with a true mass of $\sim 1.5 M_{jup}$ (Benedict et al. 2006). We want to remark that the RMS of those early RV measurements was around $10\text{--}20 \text{ m s}^{-1}$ while the amplitude of the claimed candidate is about 18 m s^{-1} . Also, the astrometric amplitude reported in (Benedict et al. 2006) is very close to the epoch-to-epoch systematic errors of FGS/HST. Because of the bias towards large amplitudes affecting astrometric measurements (Pourbaix 2001), such astrometric measurement should be understood as an upper limit. Evidence of gaps in the debris around ϵ Eridani suggest an additional very long period companion (not yet confirmed) which should have a negligible RV signature Backman et al. (e.g. 2009). ϵ Eridani is significantly more active than the Sun and has a strong stellar magnetic field (Ruedi et al. 1997). Magnetic related activity (flaring, bright and dark spots, etc), is the supposed source of the observed excess in RV variability in time-scales of weeks. Stellar global magnetic cycles could also be the responsible for part of the observed long term variability.

The star was regularly observed by HARPS between 2004 and 2007. The archive contains 113 spectra of ϵ Eri. Ten spectra showed anomalous shapes giving poor mismatches at the order of 20% in several echelle apertures. A closer look did show severe extraction problems on all of them. Three of them are from Nov 6, 2003; and three more are from Feb 4, 2004, all showing an anomalous flux deficit between 5500 and 6000 Å. The other four are from Sep 13, 2004 and Nov 02, 2004 and all show very bumpy blaze function shapes probably related to extraction issues (according to the headers, these spectra were obtained on engineering time). After excluding these 10 measurements, the final sample we use contains 103 HARPS observations on 23 different nights.

The CCF measurements as extracted from the archive have an RMS of 9.1 m s^{-1} , while the photon noise of each observation is of the order of $\sim 1 \text{ m s}^{-1}$. Such variability shows no temporal coherence (no peaks in the periodogram) and therefore these observations seem to confirm that the stellar activity is contributing significantly to the observed scatter in the RVs. The spectra were processed using the standard HARPS-TERRA setup for K dwarfs. The resulting RV measurements show a remarkably lower RMS (6.8 m s^{-1}) than the CCF RVs. In Figure 9, we present the nightly averages (23 equivalent epochs) overlapped to the nominal solution of planet b as given by Benedict et al. (2006). In the case of the CCF, when the proposed planet is subtracted, the RMS goes from 9.3 to 11.7 and the HARPS-Terra RMS increases from 6.8 to 10.0 m s^{-1} . Based on this it appears that the planet is not real,




Fig. 9.— 23 nightly averages of the 103 spectra used to measure the RVs. Red symbols represent the CCF measurements and the HARPS-TERRA ones are shown as solid black dots. No coherent RV variability coincident with previous orbital solutions is observed in either data set.

or a major revision of its orbital solution needs to be obtained.

Because of its proximity to the Sun and its brightness, ϵ Eridani has been observed by several programs with different instruments over the years. The six data sets available to the date are provided in the current distribution of SYSTEMIC and were extracted from Benedict et al. (2006) and Butler et al. (2006). Let us note that, because the data used in (Butler et al. 2006) was restricted to fewer measurements, the orbital solution presented there (only RV) was already quite different from the one reported by Hatzes et al. (2000) and Benedict et al. (2006). In order to check if there is still an orbital solution compatible with the new RV data, we add the new HARPS-TERRA measurements to a Keplerian fit with the other 6 data sets. In addition to the 5 Keplerian parameters (period, minimum mass, eccentricity, initial mean anomaly and argument of the node), the model must include 7 constants to account for the zero-point of each instrument. We make a first tentative fit using the nominal reported uncertainties. The obtained solution is significantly different to the one reported by Hatzes et al. (2000), Benedict et al. (2006) or Butler et al. (2006). In particular, the new radial velocity semi-amplitude is only $\sim 11 \text{ m s}^{-1}$ compared to the $\sim 18.0 \text{ m s}^{-1}$ given on all the previous studies and the orbital period changes from 6.9 to 7.25 yr. This changes are driven by the lack coherent variability observed in the new HARPS-TERRA RVs. The obtained eccentricity is slightly smaller and the argument of the node is also forced to very different values (see table 5). Especially in the most recent datasets with smaller formal uncertainties, it is obvious that the RV scatter is dominated by stellar noise rather than photon noise. Since the HARPS-TERRA measurements show the smaller RMS, we use them to estimate the amount of stellar jitter that has to be added in quadrature to the nominal uncertainties in order to recover the obtained scatter. We




Fig. 10.— Best orbital solution satisfying all the radial velocities publicly available to the date on ϵ Eridani. This solution is significantly different to the one presented in previous works. The new HARPS-TERRA measurements are the black dots on the right. The naming of the data sets follows the prescriptions given in Benedict et al. (2006).

find that this jitter amounts to $\sim 6.6 \text{ m s}^{-1}$. By adding this 6.6 m s^{-1} in quadrature to all the reported uncertainties, we derive a more realistic orbital solution and corresponding uncertainties. The definitive orbital fit is quite similar to the one obtained without adding the jitter but has a χ^2 per degree of freedom of 1.15, indicating that 6.6 m s^{-1} is a reasonable estimate for the stellar jitter. Figure 10 and Table 5 present the best fit to the 7 RV datasets with the jitter included in the error bars. Even if this orbit seems a good fit to all the observations, we need to remark that the orbital solution is significantly different than the previously reported ones (e.g. the period P is at $4.2\text{--}\sigma$ from previous estimates). We find this difference very suspicious and seem to indicate that the long term RV variability of ϵ Eri is due to stellar activity cycles (non-strictly periodic) rather a putative planet. Let us note that, even if the planet is real, the astrometric measurement of the orbital inclination and mass reported by Benedict et al. (2006) are no longer valid due the much more weight of the RV measurements in the determination of the orbit. In the light of this result, and given that significant efforts are being devoted to attempt direct imaging of this candidate (e.g. Janson et al. 2007; Heinze et al. 2008), a reassessment of the allowed orbital parameters combining astrometry with RV measurements using modern Monte Carlo




Fig. 11.— RMS vs bluest aperture used in the RV measurement. Even though it is an active star, the RMS always decreases as bluer apertures are included indicating that the chromatic component of the RMS is not a dominant source of RV jitter. The CCF measurements also use all the apertures.

techniques (e.g. Reffert & Quirrenbach 2011; Anglada-Escudé et al. 2011) is mostly needed. Also, a few additional HARPS observations fully covering the putative period should be sufficient to confirm that, at least, the observed variability is still present over a full orbital period. Precision radial velocities in the near infrared over a full orbital period should provide definitive confirmation/refutation of the existence of such planet.

In overview, the precision in the RV obtained using the TERRA code provides a significant reduction on the measured RMS compared to the public CCF RVs. Given that ϵ Eridani is an active star, the stellar lines used in the CCF might be peculiar when compared to quiet K dwarfs, contributing to the excess of scatter. Even though stellar activity is the most likely cause for most of the observed jitter, we do not see the chromatic jitter effect seen on M dwarfs or Tau Ceti. Figure 11 shows how the RMS always decreases as bluer apertures are included. This indicates that the leading effect causing the observed RV jitter on active stars is quite different to the one affecting more quiet dwarfs.

4.4. HD 69830, three Neptunes and a long period trend?

HD 69830 made it to the news in 2006 (Lovis et al. 2006) being the first planetary system hosting 3 Neptune mass planets. HD 69830 is a G8V star slightly less luminous and less massive than the sun ($0.6 L_{\odot}$ and $0.86 M_{\odot}$ respectively). The detection was based on 74 HARPS epochs taken between October 2003 and January 2006. The final fit to the solution had an RMS below 1.0 m s^{-1} .

The ESO archive contains 529 spectra of the star. The headers of two spectra were




Fig. 12.— 527 Radial velocity measurements obtained with HARPS-TERRA (black) compared to the 454 RVs obtained using a K5 binary mask with the CCF method (red circles). The CCF RVs are shifted 15 m s^{-1} to improve visualization.

corrupted rendering those two observations unusable (both observations are from Feb 5th, 2007). The resulting 527 spectra were taken between October 2003 and March 2008. The SNR at 6000\AA ranges from 100 to 300 and the exposure time varies between 180 to 400 seconds. The spectra were processed using the standard HARPS-TERRA setup for G dwarfs. By comparison, the CCF RV show several outliers with RV offsets of several km s^{-1} away from the average (9 spectra on 3 different nights). When removed, the RMS of the full CCF sample is still rather high compared to the TERRA one ($\sim 6.74 \text{ m s}^{-1}$). By direct inspection of the CCF RV measurements, we could see that several RVs show negative offsets of the order of 20 m s^{-1} . We found that all such measurements (67 of them) correspond to spectra processed by the HARPS DRS with a G2 binary mask instead of the K5 binary mask used for the majority of the observations (454 of them). Because the HARPS DRS is not public, we could not consistently reobtain all the CCF RVs. We show all the RV measurements obtained with HARPS-TERRA and the CCF ones obtained using the K5 binary mask in Figure 12.

The purpose of this section is to illustrate that HARPS-TERRA can detect signals at the limit of the demonstrated HARPS stability. Given the general good agreement among the two data-sets and for the sake of simplicity, only the HARPS-TERRA measurements will be discussed on all that follows.

In order to perform the orbital analysis, we consolidate these 527 observations in 176 nightly averages. The periodogram of these RVs directly shows the three signals reported by (Lovis et al. 2006). The best fit solution to these three planets compares well to the solution given in the discovery paper and has an RMS of 1.1 m s^{-1} .




Fig. 13.— **Left.** RV residuals to a fully Keplerian fit to the planets already reported by Lovis et al. (2006). A significant quadratic trend (red line) is observed in the data. As reported by (Lovis et al. 2006), the early measurements (left) have larger uncertainties due to lower SNR and other instrumental issues. **Right.** S index measurements as obtained from the original 529 spectra. A clear quadratic trend is also observed on this index pointing towards activity as the origin of the observed long term variability in the RV.

However, the periodogram to the residuals to the 3 planet fit still shows a strong peak at 490 days (see Figure 14) and an even higher power beyond 1000 days. By visual inspection of the residuals (see Figure 13, one can see that the long period power is due an apparent *parabolic* shape in the RV residuals. When we tried solving for the 490 days signal, we found that the coverage in phase of such solution was very spotty. This is characteristic of long period signals aliased with the seasonal availability of the star (see Dawson & Fabrycky 2010, for a detailed discussion of the yearly alias). Given that the power at longer periods is higher than the 490 days peak, from now on we assume that the long period trend is the most likely signal left in the data.

To assess what are the chances of this signal being generated by random noise, we compute its empirical false alarm probability as follows. We generate 10^4 synthetic data sets by keeping the observed epochs but doing random permutations of the residuals to the 3 planet fit. While these synthetic data sets will have the same distribution of random errors, the random permutations destroy the temporal coherence of any signal present. We then compute the periodogram for each synthetic data set and count the number of times we obtain a power higher than the one we find in the real data. This only happened once indicating that the FAP is low ($\sim 0.01\%$), which means that such trend is statistically significant and cannot be ignored. This method of computing FAP is described in more detail in Cumming (2004).

In order to assess what could be the cause of this trend, we examined the measurements on the S-index as described in Section 3.3. Unfortunately, the S-index shows a trend with the same shape and relative variability as the RV curve. This is characteristic of spurious offsets




Fig. 14.— Periodogram to the residuals of HD 69830 to a 3 planet fit (top) and to a 4 planet fit (bottom). Even though there is a suggestive periodicity around 490 days, this signal is most likely an alias of the long period parabolic trend seen in Figure 13. When the best circular orbit to the long period trend is removed, no additional signals are present in the data.

caused by the magnetic activity cycle of the star and has been observed in other stable stars such as Tau Ceti (Pepe et al. 2011). A tentative fit of the RV residuals provides a period of $\sim 8500 \pm 2000$ which roughly matches the expected duration of the activity cycle of a quiet star similar to our Sun. Even though the signal could be the combination of a long period planet and the activity induced offset, we cannot conclude that there is solid evidence for an additional companion to HD 69830. Once a circular orbit is subtracted to the data, the signal at 490 days completely disappears from the periodogram, confirming that both peaks in Figure 14 correspond to the same aliased trend.

The number of observations used here is significantly larger than those available at the time of discovery of the first 3 candidates. Also, only the long time-span of the observations (~ 5 years) allows to detect the quadratic trend and its correlation with the S-index variability. As shown in the bottom panel of Figure 14, no more periodicities can be inferred from the residuals to the 4 signal solution. Removing the trend with a circular orbit leaves an RMS 0.92 m s^{-1} . Note that, even with the trend, this is one of the most RV stable G dwarfs observed by HARPS. As a final note, we analyzed the CCF RV obtained with the K5 binary mask using the same procedure. We recovered the same orbital solution for the




Fig. 15.— Best fit to the 4 signals present in the HD 69830 measurements obtained with HARPS-TERRA. The first three planets were already reported by (Lovis et al. 2006). A circular orbit for planet e has been assumed to generate this plot. The RMS of the residuals to the 4 signals fit is only 0.92 m s^{-1}

three reported candidates, but the long period trend and its alias at 490 days appear both with less significance. While one would be tempted to claim that HARPS-TERRA obtains higher precision also in this case, this extreme cannot be confirmed here for three reasons : 1) the CCF data set contains fewer measurements, 2) the RMS of both measurements after removing the first 3 planets is almost identical, 3) such precision is at the limit of the long term stability of HARPS and any signal at this level (specially long period ones) has to be taken with due caution.

4.4.1. Wavelength dependence of the signals in HD 69830

We performed an additional test to assess the reality of the aforementioned trend. Given the similar spectral type of HD 69830 (G8V) to Tau Ceti (G8.5V) we asked ourselves if the chromatic jitter effect detected in the RMS of M dwarfs and Tau Ceti could be exploited to investigate the nature of the trend. To do this, we first plotted the RMS as a function of the bluer aperture used and found its minimum (see Figure 16). Let us note that, although




Fig. 16.— Radial velocity RMS as a function of the bluer aperture used on HD 69830. Even though this raw RMS contains the signal of three low amplitude companions, the contribution of the Keplerian signals and the chromatic jitter should add in quadrature, providing an optimal bluer aperture to be used.

there are three low amplitude candidates contributing to the initial RMS, Keplerian signals should be achromatic and, therefore, a minimum in the RMS should still be present when the wavelength dependent noise is added in quadrature. Figure 16 shows that such minimum in the RMS is found at aperture 37, which is similar to the optimal bluer aperture found for Tau Ceti.

We then used the RVs obtained using only apertures redder than 37 and derived the full Keplerian solution for the first three planets candidates. The obtained orbits were compatible with the previously reported orbits. However, when we computed the periodogram of the residuals to the three planet fit, we found that the quadratic trend and the corresponding 490 days alias were completely gone. Also, the RMS to the 3 planet fit was already very low (1.01 m s^{-1}), clearly indicating that the secular signal was mainly driven by apparent RV offsets at the bluest wavelengths. Even though instrumental effects might be involved, we think this result strongly suggests that, 1) this chromatic jitter is intrinsic to the star, 2) it is a significant source of the stellar RV noise observed in other relatively quiet stars, and 3) might pose a fundamental limit to the maximum RV precision achievable on G stars.




Fig. 17.— Periodogram to the residuals of HD 69830 to a 3 planet fit (top) when only apertures redder than 37 are used to produce the RV measurements. No hint of the trend or any other periodicity is seen in the data.

5. Summary and conclusions

We present a new method to obtain precision radial velocities from wavelength calibrated public HARPS spectra. Our new velocities compare well with those obtained with the CCF method and are able to detect the same signals reported by previous HARPS discoveries. For stable G and K dwarfs, the difference between CCF and HARPS-TERRA is at the level of $\sim 0.5 \text{ m s}^{-1}$ RMS. Still, the template matching approach seems to be less sensitive to offsets induced by the stellar activity (e.g. see Section 4.3 on ϵ Eridani) and requires almost no assumptions on the nature of the star. Because the template matching technique makes a more optimal use of the Doppler information in the stellar spectrum, HARPS-TERRA provides a significant increase in accuracy when applied to the heavily blended spectra of M dwarfs.

We have shown the importance of correcting for blaze function variability (flux normalization polynomial) to achieve sub m s^{-1} precision. While the CCF method needs some preprocessing of the spectra to account for such variability, least-squares template matching performs such correction in a self-consistent way.

We find that several stars show excess variability when the bluer echelle apertures are used. Even if the SNR is typically lower in the blue, this increase in the *jitter* should not happen on perfectly stable stars. We find significant evidence that this *chromatic jitter* is likely related to the star itself rather than an instrumental effect and that the wavelength dependence of the RV offsets can be exploited to confirm or rule out suspicious candidate signals. Further research is necessary to assess the nature of this excess.

We also have demonstrated that HARPS-TERRA can detect already reported signals and we have applied it to a number of interesting stars with abundant public data. While we

can reproduce well other detections, the new HARPS-TERRA measurements do not firmly confirm the planet candidate reported around ϵ Eridani. The orbital solution allowed by the new observations is significantly different compared to the previously reported ones, casting some doubts on the reality of this candidate.

We also report the detection of a quadratic trend in the residuals of HD 69830. Despite the high significance of the signal, we find that it correlates well with a similar trend in the S-index. We also find that such trend completely disappears when only the redder half of the spectrum is used to derive the RV measurements. This leads us to conclude that the most likely explanation for the observed trend is the activity cycle of HD 69830 inducing wavelength dependent RV shifts that are stronger towards the blue. Although we do not report any new planet, this example demonstrates that HARPS-TERRA is also able to robustly detect and diagnose unreported signals at the limit of the HARPS instrumental stability. Given that HD 69830 is a nearby star, a planet candidate with such a long period would lie around $0.7''$ from the central star and might be imaged with the next generation of adaptive optics systems (e.g. Lagrange et al. 2010). Even though everything points to stellar activity as the most likely explanation for the trend, we provide, just in case, an estimate of the equivalent circular orbit.

We have shown that the template matching on stabilized spectrometers requires few assumptions compared to the elaborate binary masks required by the CCF method. The capability of reproducing precision RV measurements with two different data analysis methods is also a powerful ally to double check the significance of very low amplitude signals. Given the significant increase in precision achieved on low mass stars, it is likely that HARPS-TERRA can uncover undetected low amplitude signals in already existing data sets (Anglada-Escudé et al. 2012).

Acknowledgements The research of GA has been supported by the Carnegie Fellowship Postdoctoral Program. We thank Mathias Zechmeister, Steve Vogt, Alan Boss and Alycia Weinberger for useful discussions. RPB gratefully acknowledges support from NASA OSS Grant NNX07AR40G and from the Carnegie Institution of Washington.

REFERENCES

- Anglada-Escudé, G., Arriagada, P., Vogt, S. S., & et al. 2012, accepted to ApJL, –,
Anglada-Escudé, G., Boss, A. P., Weinberger, A. J., Thompson, I. B., Butler, R. P., Vogt, S. S., & Rivera, E. J. 2011, ArXiv e-prints
Backman, D., et al. 2009, ApJ, 690, 1522

- Baranne, A., et al. 1996, *A&AS*, 119, 373
- Benedict, G. F., et al. 2006, *AJ*, 132, 2206
- Butler, R. P., Marcy, G. W., Williams, E., McCarthy, C., Dosanjuh, P., & Vogt, S. S. 1996, *PASP*, 108, 500
- Butler, R. P., et al. 2006, *ApJ*, 646, 505
- Campbell, B., Walker, G. A. H., & Yang, S. 1988, *ApJ*, 331, 902
- Cumming, A. 2004, *MNRAS*, 354, 1165
- Cumming, A., Marcy, G. W., & Butler, R. P. 1999, *ApJ*, 526, 890
- Dawson, R. I., & Fabrycky, D. C. 2010, *ApJ*, 722, 937
- Delfosse, X., Forveille, T., Ségransan, D., Beuzit, J.-L., Udry, S., Perrier, C., & Mayor, M. 2000, *A&A*, 364, 217
- Endl, M., & Kürster, M. 2008, *A&A*, 488, 1149
- Figueira, P., et al. 2010, *A&A*, 511, A55+
- Ford, E. B. 2005, *AJ*, 129, 1706
- Forveille, T., et al. 2011a, *A&A*, 526, A141+
- . 2011b, ArXiv e-prints
- Hatzes, A. P., et al. 2000, *ApJ*, 544, L145
- Heinze, A. N., Hinz, P. M., Kenworthy, M., Miller, D., & Sivanandam, S. 2008, *ApJ*, 688, 583
- Howard, A. W., et al. 2010, *Science*, 330, 653
- Huélamo, N., et al. 2008, *A&A*, 489, L9
- Janson, M., et al. 2007, *AJ*, 133, 2442
- Kaltenegger, L., Udry, S., & Pepe, F. 2011, ArXiv e-prints
- Lagrange, A.-M., et al. 2010, *Science*, 329, 57
- Lovis, C., et al. 2006, *Nature*, 441, 305

- . 2011, ArXiv e-prints
- Marcy, G. W., & Butler, R. P. 1996, *ApJ*, 464, L147
- Mayor, M., & Queloz, D. 1995, *Nature*, 378, 355
- Mayor, M., et al. 2009, *A&A*, 507, 487
- . 2011, ArXiv e-prints
- Meschiari, S., & Laughlin, G. P. 2010, *ApJ*, 718, 543
- Pepe, F., Mayor, M., Galland, & et al. 2002, *A&A*, 388, 632
- Pepe, F., Rupprecht, G., Avila, G., & el al. 2003, in *SPIE Conference Series*, Vol. 4841, 1045–1056
- Pepe, F., et al. 2011, *A&A*, 534, A58+
- Pourbaix, D. 2001, *A&A*, 369, L22
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical recipes in FORTRAN. The art of scientific computing* (Cambridge: University Press, —c1992, 2nd ed.)
- Queloz, D. 1995, in *IAU Symposium*, Vol. 167, *New Developments in Array Technology and Applications*, ed. A. G. D. Philip, K. Janes, & A. R. Upgren, 221
- Queloz, D., et al. 2001, *A&A*, 379, 279
- Reffert, S., & Quirrenbach, A. 2011, *A&A*, 527, A140
- Reiners, A., Bean, J. L., Huber, K. F., Dreizler, S., Seifahrt, A., & Czesla, S. 2010, *ApJ*, 710, 432
- Rivera, E. J., Laughlin, G., Butler, R. P., Vogt, S. S., Haghhighipour, N., & Meschiari, S. 2010, *ApJ*, 719, 890
- Rueddi, I., Solanki, S. K., Mathys, G., & Saar, S. H. 1997, *A&A*, 318, 429
- Standish, E. M. 1998, *JPL Tech Note*, IOM 312.F-98-048
- Stumpff, P. 1979, *A&A*, 78, 229
- . 1980, *A&AS*, 41, 1

van Leeuwen, F. 2007, *A&A*, 474, 653

Walker, G. A. H., Walker, A. R., Irwin, A. W., Larson, A. M., Yang, S. L. S., & Richardson, D. C. 1995, *Icarus*, 116, 359

Zechmeister, M., & Kürster, M. 2009, *A&A*, 496, 577

Zechmeister, M., Kürster, M., & Endl, M. 2009, *A&A*, 505, 859

A. Radial velocity measurements

This appendix contains tables with the most relevant time-series used through the paper. Unless stated otherwise, the first two columns are the RVs as obtained with HARPS-TERRA using the standard setup for each spectral type. The corresponding CCF values are also provided if necessary for comparison purposes. An offset (average RV) has been subtracted to all the RVs to improve readability and all RV measurements are Doppler offsets measured in the Solar System Barycentric system. The perspective acceleration effect has also been removed from all the presented RVs. No nightly averages have been applied. The tables also contain the S-index values in the Mount-Wilson system as measured by HARPS-TERRA. In a few cases, the RV measurements using the redder part of the spectrum are provided instead the S-index or the CCF measurements. Check each table caption for further information.

Table 3. Overview of the RV measurements on the three M dwarfs without planets discussed in the text.

Parameter	Proxima	Barnard's	Kapteyn's
Mean SNR at 6100 Å	39	120	85
Number of RV	27	30	22
First observation	May 2004	Dec 2003	April 2007
Last public observation	Feb 2009	Feb 2009	May 2008
RMS _{CCF} [m s ⁻¹]	2.37	2.45	1.51
RMS _{TERRA} [m s ⁻¹]	2.05	2.13	1.19
Extra noise in CCF [m s ⁻¹]	+1.18	+1.21	+0.92
Optimal bluer aperture	37	29	56
RMS at optimal aperture [m s ⁻¹]	1.94	1.95	0.82

Table 4. GJ 676A orbital solution using CCF and TERRA RVs. The numbers in parenthesis indicate the uncertainty in the last two significant digits of the parameter values. Uncertainties have been obtained using a Monte Carlo Markov Chain approach (see text).

Parameter	CCF	TERRA
P [days]	1061.7 (2.3)	1060.2 (1.8)
K [m s ⁻¹]	128.10 (66)	125.97 (58)
M ₀ [deg]	210.13 (89)	208.77 (72)
e	0.326 (92)	0.331 (78)
ω[deg]	88.8 (1.5)	89.8 (1.2)
Linear trend [m s ⁻¹ yr ⁻¹]	8.48 (66)	8.99 (56)
$M \sin i$ [M_{jup}]	4.842 (25)	4.752 (22)
a [AU]	1.81	1.81
RMS [m s ⁻¹]	3.77	3.11
σ_{O-C} [m s ⁻¹] ^a	3.51	2.97
Extra noise ^b	+2.13	-
N_{obs}	34	34
N_{par}	8	8
χ^2	113.95	114.38
$\chi^2/(N_{\text{obs}} - N_{\text{par}})$	4.38	4.39

^aWeighted RMS of the residuals as computed by Pepe et al. (2011)

^bUncertainty that has to be added in quadrature to the CCF RMS to match the RMS of the TERRA measurements.

Table 5. Orbital solution for ϵ Eridani including 7 data sets. A stellar jitter of 6.6 m s⁻¹ was added in quadrature to all nominal uncertainties. Parameter values represent the least-squares solution to the last two significant digits. The numbers in the parenthesis represent uncertainty in the last two significant digits. The uncertainties represent the 68% confidence level intervals as obtained from a Bayesian MCMC.

Parameter	CCF
P [days]	2651 (36)
K [m s ⁻¹]	11.8 (1.1)
M ₀ [deg]	09 (12)
e ^a	0.40 (11)
ω [deg]	141.4 (9.8)
$M \sin i$ [M_{jup}]	0.645 (58)
a [AU]	3.51
RMS [m s ⁻¹]	12.6
σ_{O-C} [m s ⁻¹] ^b	10.7
N_{obs}	359
N_{par}	12
χ^2	401.02
$\chi^2/(N_{\text{obs}} - N_{\text{par}})$	1.15

^aAsymmetric distribution. $0.20 < e < 0.68$ with a 99% confidence level

^bWeighted RMS of the residuals as computed by Pepe et al. (2011)

Table 6. Orbital solution for the 3 planet candidates plus trend detected on the RVs of HD 69830 when using all the echelle apertures. The proposed parameter values are obtained from the χ^2 solution. The numbers in parenthesis correspond to the uncertainty in the last two significant digits in the parameter values (68% confidence level intervals). Statistical quantities at the bottom correspond to a circular orbital fit to the quadratic trend. All the orbital elements are referred to the first epoch of observation at $JD_0=2452939.87402$ days.

	b	c	d	(activity?)
P [days]	8.6687 (12)	31.645 (28)	202.2 (1.6)	8500 ^a
K [m s ⁻¹]	3.46 (11)	2.61 (22)	1.85 (13)	12.35 ^a
M_0 [deg]	228 (10)	118 (15)	39 (17)	144 ^a
e	0.06* (05)	0.08* (05)	0.190 ⁺ (76)	0 (fixed)
ω [deg]	24 (23)	192 (40)	172 (32)	0 (fixed)
$M \sin i$ [M_{jup}]	0.0315 (11)	0.0366 (22)	0.0473 (35)	1.12 ^a
$M \sin i$ [M_{\oplus}]	10.00(32)	11.60 (69)	15.0 (1.1)	355.4 ^a
a [AU]	0.079	0.186	0.641	7.7 AU ^a
<hr/>				
RMS [m s ⁻¹]	0.92			
σ_{O-C} [m s ⁻¹] ^b	0.88			
N_{obs}	176			
N_{par}	19			
χ^2	120.35			
$\frac{\chi^2}{N_{obs}-N_{par}}$	0.76			

^aOrbital values of the equivalent circular orbit. Note that the most likely explanation to this signal is the RV offset induced by the magnetic cycle of the star.

^bWeighted RMS of the residuals as computed by Pepe et al. (2011)

*Compatible with circular orbit

⁺Slightly significant eccentricity

Table 7. Measurements on Tau Ceti.

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2453280.550009	0.42615	0.51027	0.16668	0.30930	0.13932	0.00050
2453281.551756	1.58720	0.44181	1.69523	0.28281	0.14008	0.00045
2453282.551419	0.47161	0.52024	0.92778	0.33676	0.13800	0.00053
2453283.550154	1.18747	0.54740	0.46403	0.32926	0.13992	0.00052
2453284.793218	-0.16593	0.69264	-0.05192	0.60515	0.13798	0.00089
2453309.755272	0.03438	0.28136	0.62089	0.18451	0.14151	0.00028
2453336.661904	-1.67321	0.39949	-0.85605	0.19063	0.14184	0.00029
2453337.609246	-3.42329	0.31737	-3.25516	0.17864	0.14239	0.00024
2453340.675431	0.08433	0.30174	0.90024	0.19522	0.14199	0.00029
2453760.542309	1.59136	0.46607	1.77011	0.28889	0.14087	0.00043
2453916.886879	-0.39858	0.87591	-3.22161	0.77612	0.14001	0.00111
2453930.839720	-0.31610	0.65544	-1.10805	0.32354	0.14015	0.00046
2453932.928932	1.50750	0.49047	1.74647	0.34221	0.14065	0.00045
2453933.883797	2.57307	0.45278	1.83316	0.28355	0.14048	0.00040
2453934.783019	-0.37735	0.44273	-1.43380	0.29595	0.14042	0.00043
2453936.747115	-0.89113	0.62788	-1.32457	0.34074	0.13853	0.00051
2453973.783269	0.21108	0.40290	-0.06561	0.29617	0.14152	0.00042
2453979.889092	0.93003	0.49919	0.15911	0.31815	0.14186	0.00046
2453987.661313	-1.72160	0.53284	-2.80149	0.34558	0.14182	0.00050
2454056.586134	1.24695	0.41570	1.67500	0.29186	0.14114	0.00041
2454075.526659	0.14591	0.34110	0.13332	0.24622	0.14127	0.00035
2454099.597043	4.16541	0.56336	4.59409	0.40518	0.14266	0.00058
2454100.598017	1.73427	0.67375	2.05844	0.37469	0.14187	0.00054
2454124.549301	1.66338	0.52397	2.10391	0.37496	0.14001	0.00052
2454133.563699	0.11903	0.48968	0.64645	0.31977	0.14115	0.00048
2454293.935829	-1.46651	0.41222	-0.65909	0.16860	0.14286	0.00023
2454310.880394	-1.56033	0.52724	-1.87347	0.23422	0.14279	0.00032
2454339.754289	-0.80785	0.47144	-0.24668	0.31153	0.14118	0.00043
2454350.778184	0.55266	0.51637	0.69916	0.29501	0.14207	0.00041
2454384.686692	3.14097	0.40408	3.35287	0.15295	0.14185	0.00022

Table 7—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454422.615282	0.06190	0.38518	0.31968	0.19184	0.14181	0.00027
2454423.647156	-1.30165	0.35700	-1.27200	0.15518	0.14236	0.00022
2454427.648200	1.73931	0.38704	1.93710	0.14622	0.14228	0.00019
2454428.688123	-0.51489	0.43608	0.31912	0.17355	0.14199	0.00025
2454673.890343	1.49822	0.76819	1.31597	0.47454	0.14490	0.00067
2454677.917256	-0.88101	0.51626	-0.87665	0.18420	0.14282	0.00025
2454677.928982	-0.92065	0.51027	-1.84646	0.20674	0.14251	0.00027
2454682.908129	-1.19540	0.47784	-1.12159	0.18972	0.14299	0.00025
2454700.888598	-2.23162	0.39059	-3.04735	0.17714	0.14275	0.00023
2454706.870489	1.82493	0.44845	1.02977	0.22966	0.14141	0.00032
2454707.845917	1.23706	0.36022	0.15084	0.17382	0.14265	0.00024
2454710.827579	0.82114	0.38539	0.03831	0.17969	0.14247	0.00025
2454739.718124	0.11103	0.37315	0.35369	0.17054	0.14184	0.00024
2454744.771976	-0.22687	0.40056	-0.18430	0.18562	0.14201	0.00026
2454745.675709	0.08569	0.41463	-0.10717	0.21458	0.14210	0.00030
2454746.670216	-0.13365	0.43367	0.17159	0.20121	0.14144	0.00028
2454749.586923	-0.73917	0.44729	-1.37629	0.32008	0.13981	0.00045
2454750.692281	1.13850	0.40899	0.69797	0.22472	0.14212	0.00031
2454753.573079	1.74573	0.46274	0.02863	0.37914	0.14176	0.00055
2454754.559322	-0.32681	0.53640	-0.39371	0.39162	0.14091	0.00056
2454755.679404	0.29299	0.44036	-0.01416	0.31528	0.14123	0.00043
2454758.646533	-1.95207	0.42726	-2.67768	0.28447	0.14106	0.00039
2454759.640638	0.00817	0.48013	0.27377	0.30003	0.14114	0.00040
2454760.653260	-1.16384	0.40928	-0.61169	0.20455	0.14168	0.00028
2454761.656390	-1.16540	0.41320	-1.10564	0.21845	0.14155	0.00030
2454762.644229	-1.06238	0.53653	-0.26168	0.30721	0.14097	0.00041
2454763.643444	0.30742	0.48262	-0.25132	0.28900	0.14071	0.00040
2454765.620493	1.25523	0.47284	1.64610	0.37305	0.14082	0.00051
2454766.681681	1.14891	0.40759	1.34630	0.20944	0.14154	0.00028
2454767.632523	0.53366	0.51566	0.88859	0.30774	0.14198	0.00042

Table 7—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454774.595948	-1.11180	0.48134	-0.44572	0.33713	0.14220	0.00046
2454775.604242	0.40314	0.41925	0.30552	0.34186	0.14307	0.00047
2454776.510976	-1.19330	0.59267	-1.32105	0.40581	0.14077	0.00058
2454777.518608	-0.93294	0.58974	-0.78300	0.35539	0.14292	0.00051
2454778.522152	0.38828	0.43303	1.23505	0.26072	0.14233	0.00037
2454779.518763	-2.64996	0.43771	-2.54100	0.26941	0.14110	0.00038
2454780.510881	0.30322	0.47415	0.41436	0.38485	0.14007	0.00055
2454799.640485	-1.16786	0.40783	-0.24898	0.25205	0.14160	0.00036
2454800.530029	-0.93070	0.34930	-0.54784	0.27845	0.14215	0.00039
2454801.528742	-4.88433	0.49278	-4.48218	0.24048	0.14209	0.00032
2454802.543634	-0.92240	0.39588	0.21395	0.24653	0.14236	0.00033
2454802.628512	-0.84200	0.37481	-0.09522	0.23537	0.14217	0.00033
2454802.689538	1.10213	0.55795	1.49803	0.29201	0.14081	0.00042
2454803.527323	0.12361	0.40676	1.16162	0.20761	0.14252	0.00028
2454804.514669	-1.38687	0.42670	-1.47922	0.31569	0.14111	0.00043
2454805.537775	0.23125	0.41988	1.40071	0.28830	0.14199	0.00039
2454806.509368	0.36937	0.52978	1.58339	0.26320	0.14201	0.00036
2454807.509811	0.65043	0.37536	1.47624	0.24939	0.14180	0.00035
2454849.536926	-0.71769	0.46875	-1.31384	0.21532	0.14137	0.00032
2454850.530505	1.35464	0.48487	1.13132	0.35047	0.14146	0.00053
2454852.534411	-1.67241	0.53911	-1.40068	0.26477	0.14055	0.00041
2454854.532914	-0.19676	0.46495	0.42592	0.26224	0.14022	0.00039
2455079.745110	-1.05128	0.53001	-2.17176	0.40380	0.14158	0.00056
2455080.685064	1.42864	0.85726	-0.10816	0.54625	0.14332	0.00077
2455081.847921	2.74017	0.35396	2.07335	0.27103	0.14093	0.00038
...						

Table 8. Differential RV and S-index measurements on HD 85512.

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2452990.814510	-0.88325	0.36747	0.84604	0.41236	0.30296	0.00201
2453002.833631	-1.67383	0.46082	0.26290	0.36177	0.29365	0.00160
2453051.749270	1.74874	0.70950	2.85012	0.42727	0.31230	0.00201
2453063.674457	2.03416	0.56129	2.97591	0.33228	0.29376	0.00156
2453759.772682	-1.61672	0.37845	-1.79124	0.20737	0.36436	0.00110
2454141.753872	-0.12664	0.37722	-0.04298	0.19197	0.44411	0.00111
2454422.852079	1.03102	0.31708	1.00156	0.22558	0.42707	0.00125
2454426.830209	-0.45662	0.21420	-0.06601	0.17664	0.42802	0.00097
2454521.675291	0.72385	0.38073	0.79073	0.25303	0.45941	0.00143
2454522.751630	-0.21174	0.38153	-0.72756	0.22907	0.45702	0.00137
2454523.729887	0.71248	0.42494	-0.00414	0.24571	0.45449	0.00145
2454524.719023	-0.79472	0.38560	-1.41732	0.20589	0.44699	0.00118
2454525.738981	-1.27079	0.37973	-1.35187	0.19803	0.44795	0.00117
2454526.674098	-2.49480	0.35956	-2.50058	0.18378	0.45195	0.00107
2454527.700326	-0.64342	0.31771	-0.61497	0.19459	0.45780	0.00112
2454528.663845	-1.93882	0.42210	-2.07227	0.20633	0.45447	0.00119
2454529.666263	-0.20906	0.42149	-0.42878	0.23494	0.44962	0.00132
2454530.711586	-1.55187	0.39350	-1.65076	0.18654	0.44572	0.00111
2454569.588820	1.89192	0.38085	1.88517	0.24763	0.46155	0.00143
2454571.568133	1.11252	0.41419	1.05914	0.27689	0.45361	0.00155
2454611.543521	-0.48222	0.39609	-0.77938	0.24158	0.44191	0.00140
2454616.543982	-0.45841	0.46752	-0.51327	0.29343	0.44599	0.00173
2454618.499480	1.50813	0.59271	0.46051	0.38246	0.45518	0.00223
2454620.465440	0.89537	1.27857	0.23609	0.71800	0.46258	0.00400
2454640.522808	-1.43690	0.41836	-1.12496	0.23527	0.43044	0.00143
2454643.490178	-1.85157	0.36541	-0.89792	0.23753	0.42524	0.00139
2454647.505972	-1.42305	0.37022	-0.96801	0.25259	0.41597	0.00144
2454774.860714	-0.38721	0.47769	0.05524	0.27993	0.44046	0.00155
2454775.859512	-0.30989	0.53411	0.19669	0.30566	0.42924	0.00169
2454776.864029	0.46201	0.55296	0.93342	0.32761	0.42708	0.00180

Table 8—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454777.871579	1.17839	0.47659	0.57886	0.29708	0.42361	0.00162
2454778.871212	-0.44662	0.41868	-0.31650	0.27303	0.42176	0.00148
2454779.873185	-0.34746	0.41142	-0.38435	0.25825	0.42267	0.00139
2454780.871141	0.91430	0.37413	0.94712	0.31935	0.41173	0.00173
2454799.829864	1.30904	0.35045	0.90019	0.29279	0.41535	0.00158
2454800.786091	-0.76367	0.31431	-1.20127	0.27028	0.41468	0.00147
2454800.870020	-1.07304	0.28155	-1.13122	0.23297	0.41452	0.00124
2454801.790654	-1.37285	0.27065	-1.98554	0.22682	0.41855	0.00122
2454802.794987	0.14598	0.35821	-0.78923	0.24573	0.41731	0.00132
2454802.855430	-0.62136	0.34001	-1.30106	0.23499	0.41415	0.00125
2454803.763935	0.54865	0.43427	-0.03745	0.26604	0.41851	0.00145
2454803.846162	0.93043	0.24946	0.55284	0.26660	0.41811	0.00143
2454804.767411	1.24275	0.56758	0.68269	0.35246	0.42054	0.00193
2454804.871433	0.20996	0.38910	0.49915	0.26587	0.42154	0.00143
2454805.790969	0.06369	0.42291	0.39387	0.31746	0.41935	0.00171
2454805.868196	0.39308	0.52777	-0.24704	0.32898	0.41910	0.00175
2454806.742012	0.97093	0.39321	0.67610	0.29908	0.42119	0.00163
2454806.856498	0.94230	0.37880	0.02110	0.27962	0.42065	0.00152
2454807.770293	-0.90188	0.30669	-0.66582	0.25313	0.42153	0.00137
2454807.864778	1.24293	0.51895	0.72756	0.29899	0.42501	0.00149
2454847.713287	-0.41941	0.49128	-0.34358	0.32522	0.43756	0.00175
2454847.796196	-1.06151	0.21402	-1.72131	0.24845	0.43472	0.00135
2454847.870702	-0.51850	0.42245	-0.39152	0.28663	0.43494	0.00165
2454848.721476	1.61499	0.40274	1.11845	0.23200	0.43571	0.00126
2454848.807105	-0.22149	0.29688	0.17581	0.22951	0.43800	0.00129
2454848.880361	0.42829	0.38082	0.82343	0.23811	0.43475	0.00133
2454849.703181	1.87891	0.29701	0.83659	0.26975	0.43776	0.00148
2454849.786704	1.41224	0.43852	0.90121	0.28946	0.44444	0.00163
2454849.843894	2.31995	0.35957	2.31916	0.32507	0.43362	0.00182
2454850.724483	-0.73645	0.37336	-1.05234	0.25105	0.43520	0.00137

Table 8—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454850.783409	-1.29569	0.33110	-1.40547	0.25251	0.43613	0.00136
2454850.876215	0.71488	0.36731	0.86683	0.24514	0.43550	0.00134
2454851.699231	1.28151	0.44178	0.93072	0.29420	0.44048	0.00166
2454851.786295	0.00994	0.30641	-0.14545	0.27635	0.44314	0.00156
2454851.855430	2.15976	0.43750	2.60306	0.25418	0.43608	0.00142
2454852.716538	0.22951	0.33106	0.17189	0.24752	0.43686	0.00136
2454852.795627	0.11070	0.41366	0.52717	0.28490	0.44224	0.00159
2454852.852216	0.11579	0.40216	0.74560	0.23595	0.43442	0.00131
2454853.721876	0.60190	0.38248	0.73183	0.22650	0.43522	0.00125
2454853.797713	-0.01413	0.33530	1.04335	0.26091	0.43589	0.00150
2454853.864638	1.02154	0.62632	1.59037	0.39743	0.43915	0.00233
2454854.716507	0.94744	0.46220	1.31099	0.29051	0.43959	0.00161
2454854.796094	1.47125	0.34617	1.16622	0.27374	0.43271	0.00154
2454854.855900	1.24581	0.32214	0.85167	0.24503	0.43756	0.00140
2454878.574702	-0.51082	0.29326	-0.50583	0.28400	0.42845	0.00159
2454878.710898	-1.50188	0.41384	-1.67796	0.26378	0.43332	0.00142
2454878.843806	-0.52454	0.37263	-0.52288	0.26447	0.43179	0.00149
2454879.569485	0.22277	0.52824	-0.31650	0.31389	0.43211	0.00175
2454879.710427	-0.48386	0.42351	-0.57509	0.28279	0.44457	0.00160
2454879.856217	-0.47865	0.89424	0.08908	0.52442	0.44097	0.00299
2454880.577127	-0.05848	0.62577	-0.13535	0.37976	0.43333	0.00211
2454880.732455	-2.93431	0.37924	-3.07860	0.19755	0.43588	0.00106
2454880.863766	-1.48004	0.45109	-1.37422	0.25401	0.43665	0.00148
2454881.659769	-1.02299	0.43062	-1.40499	0.26240	0.43115	0.00146
2454881.741797	-0.90042	0.55500	-0.33495	0.31296	0.42561	0.00175
2454881.844068	0.29924	0.42419	-0.57878	0.28644	0.42729	0.00164
2454882.610208	-0.52737	0.37543	-0.22098	0.23025	0.43755	0.00126
2454882.749042	-1.74323	0.41568	-1.16582	0.26158	0.43727	0.00148
2454883.565785	-0.50086	0.43124	-0.19048	0.27727	0.43553	0.00155
2454883.703150	-1.16985	0.43829	-1.14082	0.25273	0.42819	0.00138

Table 8—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454884.601155	0.30836	0.52500	-0.09374	0.33109	0.42940	0.00181
2454884.714768	0.38128	0.51397	-0.01591	0.30849	0.43218	0.00180
2454884.813798	-0.32888	0.52706	-0.32277	0.32075	0.43222	0.00179
2454885.556406	0.81803	0.36237	-0.31406	0.26520	0.43314	0.00146
2454885.728852	-0.49381	0.47021	-1.70702	0.27560	0.43337	0.00153
2454886.550638	-1.44476	0.51389	-1.17486	0.30589	0.43232	0.00171
2454886.700479	-2.59850	0.33154	-3.04046	0.21746	0.43626	0.00120
2454886.835945	-0.85837	0.44853	-0.73663	0.31310	0.43298	0.00179
2454913.493753	0.83847	0.40403	1.13385	0.24517	0.41286	0.00130
2454913.645893	-0.77191	0.43475	-0.84361	0.24561	0.41101	0.00131
2454913.711355	0.13483	0.42695	0.37127	0.26330	0.41319	0.00145
2454914.536879	0.94527	0.23518	0.88401	0.23935	0.40543	0.00128
2454914.617734	-0.28178	0.49101	-0.69604	0.28937	0.40698	0.00153
2454914.694215	0.16920	0.40655	0.45984	0.25245	0.40627	0.00138
2454915.495236	1.22463	0.47423	1.43712	0.31993	0.40492	0.00172
2454915.635339	0.46154	0.46259	1.55398	0.32859	0.41621	0.00175
2454915.740384	1.17023	0.52067	0.87881	0.33709	0.40557	0.00186
2454916.497608	0.99710	0.48945	1.43081	0.27653	0.40770	0.00148
2454916.630280	-0.24596	0.35840	-0.09291	0.24775	0.40610	0.00135
2454916.736899	-0.77722	0.44409	-0.52787	0.26355	0.40774	0.00148
2454917.519967	0.81803	0.49113	0.70067	0.29837	0.40882	0.00157
2454917.613554	0.49259	0.36575	0.31994	0.25146	0.40559	0.00136
2454917.718691	0.20512	0.39450	-0.35377	0.25412	0.40908	0.00141
2454918.531863	-0.43317	0.35222	-0.17746	0.22672	0.40899	0.00120
2454918.629211	-0.10029	0.40764	0.25174	0.25856	0.41666	0.00139
2454918.696849	0.00913	0.39490	0.13772	0.23837	0.40713	0.00129
2454919.505320	1.87824	0.38133	1.89198	0.21122	0.40799	0.00110
2454919.615249	-0.30477	0.44376	-0.12338	0.25361	0.40575	0.00132
2454919.704784	0.06071	0.48345	0.90951	0.24209	0.40894	0.00134
2454920.549308	-0.09708	0.41839	0.39410	0.21830	0.40590	0.00115

Table 8—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454920.631714	0.36906	0.49770	0.99815	0.28876	0.40549	0.00157
2454920.732487	1.00852	0.48964	1.42499	0.28499	0.40602	0.00157

Table 9. RV measurements on Barnard’s star (GJ 699). Last two columns are the RV measurements obtained using the *optimal* redmost part of the spectrum only ($6000 < \lambda < 6800 \text{ \AA}$, see Section 3.2).

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$	$\sigma_{\text{TERRA-Red}}$
2454194.893890	0.00000	0.34651	0.57194	0.32376	0.00000	0.34651
2454196.883399	0.81911	0.35716	0.84845	0.30566	0.81911	0.35716
2454197.894815	-0.78946	0.33255	-0.38601	0.31239	-0.78946	0.33255
2454198.890257	1.50797	0.45543	2.70673	0.43272	1.50797	0.45543
2454199.918201	0.44577	0.32082	0.35208	0.26366	0.44577	0.32082
2454200.920621	-0.50497	0.31501	-0.27627	0.30062	-0.50497	0.31501
2454202.918690	0.85727	0.29046	1.12713	0.30629	0.85727	0.29046
2454227.843799	1.28708	0.35269	0.93223	0.34197	1.28708	0.35269
2454291.596937	-1.26042	0.43720	-1.52277	0.48978	-1.26042	0.43720
2454298.714512	0.38788	0.48088	-0.32340	0.45487	0.38788	0.48088
2454315.661296	2.57554	0.24627	0.33993	0.67525	2.57554	0.24627
2454316.647304	0.96320	0.42271	1.40579	0.46300	0.96320	0.42271
2454319.637970	0.73630	0.26919	-0.48904	0.34144	0.73630	0.26919
2454320.650245	-0.23790	0.32574	0.42950	0.31232	-0.23790	0.32574
2454340.650587	-0.43332	0.30364	-1.05476	0.28830	-0.43332	0.30364
2454343.557195	-0.19728	0.28865	-1.01755	0.27217	-0.19728	0.28865
2454347.558141	-0.87450	0.34079	-0.25382	0.31584	-0.87450	0.34079
2454385.523878	2.06296	0.64341	0.40371	0.77195	2.06296	0.64341
2454389.501586	2.88514	0.80941	3.56973	1.08852	2.88514	0.80941
2454523.881299	-0.70082	0.50892	-1.69989	0.43730	-0.70082	0.50892
2454551.908998	-1.03745	0.41725	-2.49500	0.33980	-1.03745	0.41725
2454588.921002	-1.75406	0.54486	-3.16873	0.42538	-1.75406	0.54486

Table 10. RV measurements on Kapteyn’s star (GJ 191). Last two columns are the RV measurements obtained using the *optimal* redmost part of the spectrum only ($4670 < \lambda < 6800 \text{ \AA}$, see Section 3.2).

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]
2452985.741114	4.17510	1.03818	4.55727	0.86033	3.81838	0.83699
2452996.763214	2.60098	0.26086	2.77118	0.56030	2.70283	0.21930
2453337.800966	-1.04323	1.09789	0.12095	0.76332	-0.74766	0.90860
2453668.835375	-2.28865	0.59397	-3.79258	0.49284	-2.21699	0.54708
2453670.739478	-3.94631	0.58109	-4.59149	0.41933	-4.25875	0.56210
2454141.574447	0.03985	0.52542	0.72655	0.37960	-0.13958	0.50784
2454167.530533	-1.47217	0.50752	-2.15028	0.38433	-1.66431	0.47212
2454169.505216	-2.43451	0.47790	-2.29499	0.34512	-2.51582	0.48129
2454173.500808	-2.17285	0.48345	-2.22839	0.36009	-2.31741	0.46131
2454197.512079	1.68710	0.42683	4.28535	0.43096	1.36156	0.39197
2454229.468730	2.04523	0.76567	3.40866	0.56835	1.78758	0.68699
2454342.862585	2.03375	0.54717	2.80321	0.39053	1.98278	0.54735
2454345.879666	-0.21043	0.37798	-1.23904	0.33907	-0.05894	0.35456
2454349.834178	-0.54352	0.44040	-1.39747	0.35483	-0.21252	0.41454
2454424.731616	1.52251	0.49022	1.10564	0.37339	1.71790	0.48342
2454429.718977	4.42431	0.49714	2.61326	0.39773	4.18135	0.43460
2454449.760520	0.32913	0.58481	-0.14029	0.45151	0.20908	0.55812
2454450.647003	1.29293	0.55343	1.25197	0.40700	1.52003	0.52137
2454452.706840	0.89530	0.68933	1.28428	0.53263	0.99496	0.63206
2454459.643590	-1.27204	0.57986	0.21444	0.35396	-1.32404	0.57255
2454462.705724	1.50577	0.51209	0.50036	0.35678	1.31237	0.50473
2454463.698541	2.26304	0.44297	1.90185	0.43135	2.21079	0.44640
2454464.682434	1.08128	0.52163	0.45370	0.42748	1.09792	0.45686
2454733.897251	-1.34113	0.51035	-1.27236	0.36206	-1.21830	0.50568
2454754.838832	-3.05621	0.50697	-4.33169	0.41750	-3.20201	0.47069
2454767.758056	-0.57231	0.54277	2.08814	0.43101	-0.45009	0.49990
2454879.556051	-2.57809	0.56308	-1.40012	0.43816	-2.62889	0.53567
2454879.696855	-1.51557	0.96155	-1.03663	0.68905	-1.41452	0.84971
2454880.521651	1.20466	2.87965	-0.96773	1.28010	1.78313	2.09901
2454880.673879	-2.65391	0.88143	-3.24375	0.57402	-2.31082	0.74765

Table 11. RV measurements on Proxima Cen (GJ 551). Last two columns are the RV measurements obtained using the *optimal* redmost part of the spectrum only ($4400 < \lambda < 6800 \text{ \AA}$, see Section 3.2).

JD [days]	RV _{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV _{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	RV _{TERRA-Red} [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]
2453152.599771	0.37348	0.47936	0.79047	1.02223	0.37873	0.45254
2453202.584812	-2.97293	2.29999	-0.93952	2.25726	-2.86609	2.15906
2453203.537223	1.17250	0.68714	1.90963	1.04968	0.93396	0.61743
2453205.531679	1.16372	3.95272	7.52242	1.30372	1.87165	1.98058
2453207.505219	0.66168	1.92677	-0.91287	4.20013	0.29104	1.79977
2453207.510890	2.95883	2.08065	0.91052	3.97917	3.82038	1.82329
2453577.495602	3.73820	0.49688	1.54178	1.09223	3.60389	0.46023
2453809.893589	-2.17564	0.51088	-0.64866	0.74769	-2.11864	0.48677
2453810.783507	-1.80758	0.46309	-2.09214	0.61884	-1.91120	0.45148
2453812.782152	0.71017	0.68795	-0.99516	0.94375	0.59724	0.64054
2453816.809553	-3.81687	0.49196	-4.54173	0.77872	-3.93731	0.45792
2454173.818222	-0.46091	0.49826	-0.30837	0.73580	-0.62461	0.48062
2454293.586482	2.76021	0.59884	4.29679	0.74403	2.25631	0.52081
2454296.593448	0.00000	0.68772	-1.53705	1.32162	0.37807	0.62543
2454299.597768	5.28554	0.86045	3.38681	2.03355	5.43933	0.71328
2454300.557562	-1.96552	0.53381	-2.38955	0.73998	-1.98891	0.51131
2454646.569314	0.85745	0.58995	1.43802	1.22954	0.60146	0.56480
2454658.564359	1.43411	0.87609	0.10350	1.53635	1.72781	0.65871
2454666.543473	-0.20101	0.50861	-0.61685	0.86422	-0.16505	0.45478
2454878.863802	-0.82326	0.45059	-0.18170	0.69693	-0.98036	0.43858
2454879.830132	-0.38851	0.59892	-0.04147	1.10250	-0.58228	0.53957
2454881.873704	-1.23090	0.44044	-2.62104	0.61993	-1.00764	0.36487
2454882.853542	-2.20739	0.55524	-0.79912	0.95251	-2.02011	0.44535
2454883.861612	-0.57856	0.71330	-0.19154	1.55181	-0.49378	0.68713
2454884.835397	0.77791	0.45355	-2.41492	1.19246	0.55074	0.42495
2454885.875077	0.23612	0.54306	0.31352	0.78035	0.00000	0.50075
2454886.843595	-0.01721	0.59010	-0.98175	0.95621	-0.23690	0.53914

Table 12. Differential RV and S-index measurements on 34 epochs of GJ 676A

JD [days]	RV _{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV _{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454167.897856	143.65593	1.45050	144.59273	0.96910	1.27632	0.01023
2454169.895854	140.76651	1.38811	141.45382	0.83020	1.29254	0.00959
2454171.904445	145.84144	1.52625	143.32531	0.94088	1.23147	0.01024
2454232.818013	142.10548	1.19654	143.81091	0.78189	1.34713	0.01004
2454391.491808	-23.38607	1.40115	-18.92497	0.76311	1.39087	0.00982
2454393.489934	-27.22405	1.31049	-24.68468	0.87658	1.42889	0.01102
2454529.900847	-99.67167	1.40944	-99.46562	1.03749	1.36065	0.01155
2454547.915016	-98.53057	1.46694	-96.80962	0.79384	1.40050	0.00987
2454559.815698	-95.20497	1.91938	-86.59697	0.98344	1.65069	0.01287
2454569.903637	-103.26245	2.84609	-95.28933	1.19791	1.37069	0.01260
2454571.889460	-100.73284	1.92074	-98.96364	0.56785	1.39171	0.00782
2454582.820292	-90.95710	1.36587	-86.83138	0.82576	1.35936	0.01028
2454618.755585	-80.89806	1.79552	-80.82636	1.33777	1.43244	0.01417
2454660.661636	-61.63426	1.36018	-57.37647	0.77204	1.40788	0.01009
2454661.772229	-61.42342	0.82988	-59.37748	0.70829	1.44002	0.00977
2454662.675237	-65.33515	1.68434	-66.06418	1.00474	1.44577	0.01226
2454663.811590	-63.08516	1.49354	-58.29150	0.65510	1.38950	0.00908
2454664.790043	-57.62443	1.56774	-57.06590	1.16132	1.44093	0.01357
2454665.786377	-64.24289	1.05812	-61.16171	0.66721	1.44502	0.00944
2454666.696058	-63.89311	0.87388	-61.95570	0.59440	1.47606	0.00856
2454670.672602	-61.81325	1.46348	-59.12632	1.14012	1.48977	0.01408
2454671.603329	-60.91549	1.76724	-58.30522	0.92748	1.49472	0.01198
2454687.561959	-62.55236	1.74214	-56.57420	0.94771	1.45407	0.01196
2454721.554874	-45.94209	1.59175	-41.25367	1.10505	1.37213	0.01145
2454751.490690	-39.23628	2.74576	-32.27202	2.99088	1.42617	0.02215
2454773.502375	-20.11282	1.65477	-15.49613	0.70763	1.48276	0.01024
2454916.819805	47.72555	1.18121	45.29150	0.49907	1.54039	0.00704
2454921.892971	42.22977	1.23941	40.95217	1.17217	1.45797	0.01206
2455284.893135	174.45713	2.28031	179.61741	1.70550	1.21186	0.01423
2455355.795443	139.67896	1.15261	141.62076	0.92020	1.34806	0.01223

Table 12—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2455375.610729	113.16490	1.68466	115.74927	1.19504	1.41771	0.01569
2455387.656686	98.02398	1.50445	101.98521	1.44032	1.34647	0.01797
2455402.590925	86.01649	4.12785	88.49974	4.40646	1.46733	0.03187
2455404.645561	81.14929	2.79506	85.81422	2.54605	1.48202	0.01978

Table 13. Differential RV and S-index measurements on ϵ Eridani.

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2452949.789541	-13.27658	0.20836	-26.59166	0.24191	0.41652	0.00078
2452949.790432	-12.55701	0.13163	-26.30259	0.23060	0.41713	0.00075
2452949.791323	-13.71360	0.19637	-27.35248	0.22330	0.41628	0.00077
2453001.598269	-2.21518	0.63673	-1.45770	0.26916	0.41976	0.00090
2453039.595541	-1.14426	0.57985	-13.96471	0.25574	0.42908	0.00093
2453039.597010	-1.65446	0.53546	-14.61557	0.23938	0.43211	0.00086
2453039.598365	-1.14070	0.61844	-14.48246	0.23418	0.43038	0.00086
2453279.824466	3.86346	0.33977	4.53516	0.20579	0.40336	0.00068
2453279.826468	3.16143	0.34242	3.89520	0.21334	0.40452	0.00070
2453287.796071	-15.10212	0.50166	-14.28711	0.28492	0.39628	0.00093
2453287.796765	-14.67626	0.42027	-13.65327	0.30743	0.39432	0.00099
2453287.797460	-15.19174	0.46168	-13.94097	0.34981	0.39549	0.00114
2453287.800990	-13.94422	0.37098	-12.52541	0.36313	0.39361	0.00118
2453287.801661	-14.17952	0.47441	-13.42466	0.37418	0.39198	0.00121
2453287.802402	-15.99510	0.49595	-15.21834	0.41474	0.39653	0.00131
2453304.696457	12.78309	1.71706	13.20934	1.07309	0.40753	0.00320
2453304.697256	10.56232	1.08003	10.14149	0.73927	0.39812	0.00239
2453304.697973	11.30622	0.60135	11.18511	0.49097	0.40703	0.00159
2453304.897642	10.31486	0.51256	10.77588	0.34483	0.41251	0.00119
2453304.898418	10.01810	0.52478	10.20889	0.35370	0.41403	0.00126
2453304.899193	9.34064	0.43474	10.51016	0.32739	0.41087	0.00114
2453305.673376	4.24321	0.42609	4.76444	0.19104	0.42163	0.00064
2453305.674151	5.09995	0.43148	5.16497	0.19743	0.42104	0.00067
2453305.674915	4.83352	0.37415	5.44822	0.18181	0.42185	0.00061
2453305.798344	1.12653	0.37405	2.00091	0.17950	0.42447	0.00062
2453305.799119	1.99759	0.36819	2.80579	0.17962	0.42363	0.00062
2453305.799871	2.51138	0.37643	3.40577	0.17678	0.42439	0.00060
2453306.753420	-1.72985	0.33166	-1.33805	0.16492	0.41890	0.00053
2453306.755191	-0.92093	0.30025	-0.08969	0.15381	0.41813	0.00051
2453309.796792	-4.71858	0.31458	-4.47246	0.25026	0.41097	0.00087

Table 13—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2453309.797498	0.43306	0.46670	0.58817	0.27248	0.41146	0.00099
2453309.798204	-1.57477	0.45241	-2.07887	0.26280	0.40924	0.00091
2453332.675881	-3.05227	0.41665	-2.32218	0.22959	0.40278	0.00072
2453412.559889	0.23016	0.28596	2.07113	0.15437	0.39257	0.00052
2453412.561984	-0.91317	0.31459	1.25813	0.14285	0.39177	0.00047
2453412.564125	-0.32673	0.31609	1.56604	0.15708	0.39168	0.00053
2453430.573344	-3.85861	0.31525	-2.31239	0.15533	0.38339	0.00053
2453430.574698	-5.45693	0.29355	-3.75580	0.14803	0.38385	0.00050
2453430.576110	-6.79037	0.29477	-5.18083	0.15178	0.38353	0.00052
2453600.931282	5.81997	0.38137	8.52798	0.18076	0.37807	0.00055
2453600.932497	8.63476	0.35128	10.85005	0.18748	0.37922	0.00055
2453600.933863	10.40966	0.77078	10.83690	0.39826	0.37988	0.00107
2453604.832542	-10.13333	0.31194	-21.64840	0.13596	0.39097	0.00042
2453604.839764	-8.87536	0.32454	-20.14088	0.12622	0.39149	0.00039
2453604.846918	-10.02268	0.31671	-21.08649	0.12668	0.39045	0.00039
2453666.843509	-8.48681	0.31086	-5.87599	0.13299	0.36549	0.00041
2453666.844887	-8.97717	0.37114	-6.42853	0.13258	0.36582	0.00040
2453666.846287	-8.46288	0.33264	-6.22016	0.13320	0.36626	0.00040
2453690.797493	3.40551	0.42353	5.62341	0.24631	0.35934	0.00075
2453690.798223	2.88630	0.50651	4.83332	0.28658	0.35854	0.00091
2453690.798917	2.49051	0.52386	4.29881	0.27198	0.35833	0.00086
2453732.682986	-1.42603	0.24952	-0.04318	0.12236	0.36478	0.00039
2453732.684375	-1.33882	0.27220	0.16628	0.12599	0.36465	0.00040
2453732.686632	0.59544	0.37612	2.02145	0.17610	0.36476	0.00056
2453732.687511	-0.46478	0.34948	1.03467	0.17851	0.36417	0.00057
2453732.688368	-1.67548	0.31386	-0.11975	0.17552	0.36408	0.00055
2453933.900747	-2.66193	0.37090	-1.43334	0.24307	0.37129	0.00077
2453933.901499	-2.67314	0.36761	-1.51247	0.24013	0.37140	0.00076
2453933.902275	-1.75109	0.44490	0.06342	0.24651	0.37109	0.00078
2453933.903062	-2.72935	0.43244	-1.63089	0.26283	0.36859	0.00082

Table 13—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2453933.903814	-3.51340	0.44200	-1.88215	0.26083	0.37158	0.00082
2453933.904566	-3.03592	0.34467	-1.55349	0.25235	0.37060	0.00079
2453964.856253	3.28979	0.37050	4.01923	0.24474	0.39733	0.00079
2453964.856993	3.53272	0.43480	4.38195	0.23263	0.39782	0.00075
2453964.857780	3.92908	0.34563	4.32423	0.23184	0.39654	0.00074
2453964.858568	3.37464	0.33591	4.17304	0.22857	0.39872	0.00073
2453964.859308	4.98262	0.41784	5.03279	0.23046	0.39746	0.00074
2454056.621484	1.14378	0.43186	3.15507	0.27644	0.38066	0.00086
2454056.622248	1.75443	0.46479	3.60896	0.26876	0.38404	0.00084
2454056.623012	1.26824	0.38687	2.65103	0.26124	0.38360	0.00081
2454056.623787	2.42968	0.46621	4.22000	0.28575	0.38254	0.00089
2454056.624563	2.19753	0.43532	3.90528	0.27230	0.38247	0.00085
2454056.625338	2.93444	0.47212	4.31767	0.30988	0.38445	0.00097
2454098.597909	8.10298	0.36799	9.44468	0.28065	0.41336	0.00090
2454098.598627	9.32500	0.41250	10.90518	0.28863	0.41124	0.00093
2454098.599333	8.03716	0.44091	9.69640	0.26538	0.41137	0.00085
2454098.600073	8.88737	0.43609	10.02031	0.27559	0.41400	0.00089
2454098.600756	8.98741	0.40726	10.87771	0.28030	0.41192	0.00091
2454098.601485	7.75810	0.52626	8.70430	0.30913	0.41271	0.00100
2454125.554250	5.23617	0.39693	7.31871	0.20624	0.40760	0.00067
2454125.554956	6.88588	0.21339	7.98202	0.20767	0.40749	0.00068
2454125.555662	7.36820	0.36084	8.64821	0.21041	0.40612	0.00068
2454125.556380	6.65554	0.34632	8.10203	0.20863	0.40785	0.00068
2454125.557074	5.39622	0.36946	6.76326	0.20465	0.40881	0.00066
2454125.557792	5.79571	0.31462	7.09557	0.21234	0.40613	0.00069
2454133.609700	-6.35693	0.43111	-5.10162	0.26056	0.41900	0.00090
2454133.610406	-6.29121	0.45659	-5.08948	0.25379	0.41921	0.00087
2454133.611112	-5.99178	0.40965	-4.48487	0.25504	0.42002	0.00088
2454133.611818	-7.79234	0.47088	-5.75788	0.25976	0.41866	0.00090
2454133.612524	-5.65916	0.45851	-4.33199	0.25341	0.42149	0.00088

Table 13—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	RV_{CCF} [m s ⁻¹]	σ_{CCF} [m s ⁻¹]	S-index	σ_S
2454133.613230	-7.76059	0.42255	-5.63697	0.27075	0.42233	0.00095
2454310.940581	-0.64908	0.38286	0.69489	0.20527	0.38570	0.00063
2454310.941287	-1.31705	0.41388	0.83152	0.21414	0.38630	0.00066
2454310.941993	0.10049	0.47179	1.98065	0.21697	0.38621	0.00067
2454310.942699	0.01095	0.39670	1.71384	0.21022	0.38480	0.00064
2454310.943405	-0.20973	0.33509	1.49530	0.21543	0.38436	0.00066
2454310.944112	0.03643	0.43017	1.58340	0.21464	0.38610	0.00066
2454339.807249	6.12589	0.67453	6.79357	0.41329	0.38234	0.00132
2454339.807944	6.77749	0.49238	7.02270	0.34181	0.38261	0.00109
2454339.808673	6.45656	0.57184	6.46947	0.34501	0.38384	0.00110
2454339.809379	5.36156	0.59459	5.46278	0.31312	0.38428	0.00100
2454339.810062	6.23670	0.55792	7.44273	0.33396	0.38215	0.00106
2454339.810803	5.94256	0.51823	6.71612	0.33036	0.38294	0.00105

Table 14. Measurements on HD 69830. Instead of the CCF RVs, RV using the redder part of the spectrum (see Section 4.4.1) are provided in this case.

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2452939.874025	0.78514	0.44513	-0.44804	0.49607	0.14331	0.00049
2452943.876091	4.52766	0.54224	3.79835	0.50728	0.14377	0.00043
2452946.863070	1.66663	0.54780	-0.08756	0.62966	0.14381	0.00042
2452949.840269	5.06226	0.97553	1.66953	0.94835	0.13985	0.00070
2452949.842734	4.21729	0.92317	0.86545	0.95698	0.14056	0.00064
2452949.845246	3.97665	0.83998	2.04983	0.81015	0.13968	0.00065
2452949.849471	3.85089	12.32709	2.06592	42.48213	0.14061	0.00067
2452949.851937	5.97607	11.50930	1.61959	41.41179	0.14058	0.00066
2452949.854379	4.54371	11.17014	2.65817	41.20824	0.14083	0.00059
2452997.808952	-0.91173	0.99004	-1.11047	0.68650	0.13573	0.00099
2453000.711174	0.64741	0.54575	2.47339	0.56774	0.12771	0.00090
2453035.779711	5.30455	0.39516	3.54643	0.43669	0.13887	0.00045
2453038.699416	8.23152	0.25283	4.92453	0.27440	0.13511	0.00067
2453039.741900	3.97480	0.66940	2.22453	0.69977	0.13619	0.00056
2453039.745025	3.94953	0.63563	2.06223	0.72727	0.13703	0.00057
2453039.748231	3.49526	0.67554	1.37279	0.71206	0.13680	0.00059
2453039.760754	5.16432	9.37407	1.89493	26.71249	0.13809	0.00062
2453039.763925	5.13782	4.48621	-2.94748	9.57111	0.13792	0.00064
2453039.767073	6.33786	2.88505	-4.76363	5.81669	0.13799	0.00062
2453049.629850	0.37736	0.39724	-0.87744	0.40893	0.14024	0.00027
2453058.637539	-0.04309	0.57317	-1.74315	0.59506	0.13605	0.00050
2453064.608232	7.64635	0.64285	6.05670	0.61696	0.13736	0.00052
2453066.628043	1.77976	0.45523	0.31687	0.47809	0.13698	0.00045
2453146.503893	-3.33535	0.38280	-3.40699	0.44488	0.13948	0.00065
2453147.465287	-4.71074	0.52368	-4.79587	0.57704	0.13924	0.00077
2453150.459926	1.92931	0.20526	0.44799	0.32053	0.14244	0.00055
2453151.459791	-2.22043	0.44265	-2.38302	0.54345	0.13661	0.00089
2453154.455248	-4.17988	0.61584	-5.30788	0.58508	0.13299	0.00130
2453155.453692	0.82964	1.18025	-3.33782	0.39213	0.12019	0.00131
2453156.519401	-1.43760	0.49627	-2.47239	0.63557	0.14357	0.00064

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453158.478807	4.26279	0.45153	3.23177	0.49599	0.14042	0.00109
2453159.471884	4.99679	0.28124	4.17098	0.35774	0.14281	0.00066
2453306.865602	-1.03528	0.44712	0.53459	0.51313	0.13783	0.00038
2453306.868067	-1.26640	0.36180	-0.25589	0.43592	0.13749	0.00037
2453306.870533	-0.07581	0.39535	0.49602	0.47885	0.13762	0.00038
2453308.844196	-4.16508	0.56968	-3.42959	0.44854	0.13592	0.00064
2453308.846234	-5.32271	0.47657	-5.63243	0.50924	0.13587	0.00069
2453308.848387	-6.35650	0.50749	-4.91441	0.52385	0.13537	0.00059
2453308.850516	-6.40518	0.46166	-5.06083	0.54490	0.13619	0.00057
2453308.852635	-6.50015	0.51426	-6.02175	0.56060	0.13443	0.00058
2453309.849606	-7.23881	0.35543	-6.73419	0.46367	0.13730	0.00047
2453309.852083	-8.07103	0.41065	-7.17786	0.52661	0.13714	0.00046
2453309.854526	-6.94335	0.34825	-6.27095	0.41016	0.13754	0.00047
2453309.856991	-6.39130	0.45835	-5.19492	0.56199	0.13793	0.00047
2453310.830223	-6.28643	0.31354	-6.14452	0.43422	0.13847	0.00043
2453310.832920	-6.23299	0.27933	-6.24399	0.40112	0.13900	0.00044
2453310.835605	-6.80787	0.35359	-6.71418	0.43764	0.13865	0.00042
2453310.838302	-6.40912	0.37056	-5.77436	0.41565	0.13866	0.00042
2453315.778319	0.28864	0.29461	0.33766	0.32072	0.13381	0.00058
2453315.780646	1.60989	0.40899	2.66728	0.40787	0.13573	0.00050
2453315.783019	0.62515	0.47567	1.29782	0.44608	0.13481	0.00055
2453315.785588	0.62135	0.54968	1.12478	0.53936	0.13250	0.00058
2453315.788089	2.33039	0.35991	2.51662	0.41197	0.13662	0.00052
2453315.790473	1.94666	0.31313	2.72421	0.38453	0.13769	0.00047
2453336.872454	-8.66485	0.41742	-8.78173	0.42816	0.13599	0.00040
2453336.875938	-7.78774	0.36861	-9.45765	0.43755	0.13578	0.00039
2453336.879445	-8.68171	0.34353	-8.33832	0.42555	0.13697	0.00035
2453338.867133	-5.71559	0.37085	-6.12025	0.42995	0.13678	0.00037
2453338.869807	-5.21931	0.35225	-4.94127	0.43105	0.13661	0.00038
2453338.872469	-5.60585	0.35834	-5.28837	0.42944	0.13712	0.00037

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453338.875131	-5.12891	0.39994	-4.29192	0.43685	0.13657	0.00038
2453339.869177	-2.85457	0.34970	-2.83329	0.42413	0.13711	0.00034
2453339.872580	-2.57551	0.35140	-2.26293	0.38869	0.13674	0.00035
2453339.876156	-3.38385	0.36478	-3.33283	0.48154	0.13636	0.00035
2453340.869553	-0.78826	0.36073	-1.43562	0.39690	0.13725	0.00032
2453340.873002	-1.99708	0.33478	-2.50952	0.33537	0.13733	0.00031
2453340.876452	-3.01507	0.29598	-3.53607	0.35173	0.13746	0.00030
2453341.872961	-1.64596	0.37859	-0.84700	0.42809	0.13791	0.00031
2453341.876653	-1.40799	0.27800	-0.66491	0.28392	0.13843	0.00031
2453341.880230	-3.06807	0.36895	-1.75078	0.43886	0.13844	0.00030
2453366.798062	3.84100	0.63129	4.04858	0.61372	0.13422	0.00076
2453366.799127	2.91869	0.46893	1.73770	0.61256	0.13510	0.00071
2453366.800238	1.33860	0.59279	2.98067	0.62795	0.13612	0.00065
2453366.801291	1.53909	0.59539	3.06536	0.72532	0.13758	0.00065
2453366.802356	1.21533	0.67015	4.01545	0.66332	0.13657	0.00066
2453366.803375	2.14459	0.51185	3.19135	0.54201	0.13783	0.00068
2453366.804440	2.34518	0.63214	1.87377	0.57043	0.13700	0.00067
2453366.805481	2.41210	0.59906	2.10647	0.71665	0.13714	0.00066
2453366.806558	3.73175	0.69178	4.96263	0.61783	0.13606	0.00067
2453366.807588	2.13956	0.57380	3.67442	0.49060	0.13589	0.00066
2453400.706778	1.49934	0.37973	1.86331	0.48077	0.13569	0.00065
2453400.711686	1.85069	0.48246	1.88625	0.62451	0.13567	0.00063
2453401.697427	2.62976	0.41042	2.19572	0.46027	0.13682	0.00039
2453402.701071	2.14926	0.29702	1.34116	0.36770	0.13688	0.00037
2453402.705412	1.36853	0.30596	0.67315	0.33319	0.13826	0.00035
2453403.729703	-0.61995	0.35122	-1.61612	0.40208	0.13788	0.00030
2453403.734159	0.74778	0.32295	0.02373	0.43824	0.13804	0.00030
2453404.674421	-2.89880	0.26641	-2.47469	0.35349	0.13875	0.00032
2453404.677557	-2.58693	0.29448	-2.16310	0.41574	0.13872	0.00032
2453404.680694	-3.30718	0.40782	-2.51000	0.47496	0.13841	0.00032

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453405.777713	-3.44110	0.34339	-3.26672	0.36888	0.13678	0.00042
2453405.780641	-4.25348	0.32519	-4.40101	0.43507	0.13599	0.00043
2453405.783534	-4.08800	0.38589	-2.72189	0.44429	0.13719	0.00042
2453406.718771	-3.11293	0.33545	-2.73396	0.37865	0.13771	0.00038
2453406.721826	-3.08077	0.33568	-2.98677	0.40330	0.13725	0.00037
2453406.724974	-3.38173	0.34648	-4.05589	0.39317	0.13777	0.00036
2453407.680475	-1.63318	0.28777	-1.98054	0.30286	0.13651	0.00033
2453407.684665	-3.02023	0.34212	-2.15202	0.43389	0.13789	0.00032
2453408.642479	0.55553	0.38208	2.17886	0.42109	0.13703	0.00044
2453408.645569	0.31675	0.38109	-0.06293	0.43120	0.13580	0.00042
2453408.648740	-0.38248	0.36657	-0.09023	0.45257	0.13638	0.00046
2453409.738878	4.05788	0.53855	4.67360	0.57574	0.13716	0.00055
2453409.741991	4.57236	0.53723	4.03299	0.65031	0.13613	0.00055
2453409.745128	4.18512	0.47115	4.10425	0.52211	0.13712	0.00056
2453410.664352	5.55956	0.42933	6.85214	0.47758	0.13714	0.00051
2453410.667685	5.41082	0.34721	5.04231	0.44025	0.13621	0.00050
2453410.670775	5.52528	0.37583	5.45977	0.46299	0.13716	0.00046
2453411.654823	6.23648	0.46447	7.00862	0.49397	0.13589	0.00052
2453411.657879	7.02804	0.47792	6.22656	0.53994	0.13734	0.00050
2453411.661050	6.07873	0.51058	7.49478	0.53423	0.13685	0.00053
2453412.696056	3.09669	0.51696	2.43780	0.51878	0.13549	0.00050
2453412.699459	2.28360	0.42004	2.67686	0.40209	0.13638	0.00049
2453412.702596	2.29743	0.51836	3.36255	0.51662	0.13579	0.00054
2453413.761199	-0.18516	0.31919	-0.03160	0.38107	0.13646	0.00035
2453413.767102	-0.16032	0.36831	0.21098	0.44503	0.13679	0.00038
2453413.773294	-0.24091	0.32838	0.30477	0.36172	0.13757	0.00035
2453429.626004	0.84656	0.15878	0.87010	0.22155	0.13692	0.00032
2453429.630622	0.12921	0.26123	-0.19945	0.33139	0.13663	0.00030
2453429.635032	0.17330	0.19078	0.38698	0.26252	0.13668	0.00034
2453466.614551	-3.67489	0.28522	-3.84030	0.35807	0.13292	0.00057

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453466.616750	-5.02133	0.34516	-4.74137	0.38181	0.13138	0.00054
2453466.618995	-4.49331	0.36569	-4.93671	0.42855	0.13228	0.00057
2453466.621113	-5.10117	0.43462	-5.21700	0.46367	0.13098	0.00062
2453466.623439	-4.47501	0.30201	-5.65894	0.37143	0.13169	0.00060
2453467.520130	-4.59873	0.37068	-4.13978	0.50840	0.13295	0.00058
2453467.522340	-4.22102	0.36172	-4.46752	0.46737	0.13220	0.00058
2453467.524562	-3.21630	0.27175	-4.39147	0.40684	0.13308	0.00057
2453467.526831	-3.89848	0.60057	-4.16720	0.63279	0.13279	0.00058
2453467.529018	-4.78486	0.34442	-4.60434	0.42903	0.13196	0.00059
2453491.481929	-2.35756	0.32639	-2.25858	0.41577	0.13856	0.00039
2453491.487206	-1.03011	0.37552	-2.17801	0.39056	0.13846	0.00038
2453669.864054	5.54388	0.35388	6.36261	0.33081	0.13573	0.00035
2453669.867411	5.51263	0.38493	5.31506	0.40178	0.13551	0.00034
2453669.871011	6.15330	0.39249	6.12737	0.33784	0.13540	0.00036
2453670.819980	6.15216	0.43164	6.99326	0.47986	0.13444	0.00042
2453670.823453	6.89929	0.52437	7.12720	0.58390	0.13396	0.00046
2453670.826995	7.37551	0.40677	7.37013	0.49710	0.13516	0.00042
2453671.841491	3.48094	0.36641	2.61735	0.40319	0.13539	0.00037
2453671.845033	4.17857	0.31135	4.39141	0.36766	0.13570	0.00035
2453671.848413	3.99753	0.36193	3.12935	0.35761	0.13502	0.00038
2453672.849898	2.18991	0.37805	2.29204	0.51378	0.13531	0.00038
2453672.853417	1.90327	0.36499	2.36241	0.42131	0.13516	0.00036
2453672.856786	2.24109	0.29082	1.97685	0.37697	0.13476	0.00040
2453674.872652	-2.61646	0.31695	-2.34281	0.43045	0.13729	0.00030
2453674.876113	-1.28593	0.33521	-0.96444	0.42509	0.13666	0.00030
2453674.879539	-2.14769	0.35076	-1.64529	0.41566	0.13616	0.00031
2453675.845039	-2.29515	0.34015	-1.31071	0.44637	0.13712	0.00026
2453675.848534	-2.30702	0.22902	-1.11499	0.30967	0.13720	0.00026
2453675.851995	-2.76019	0.25803	-1.90278	0.33779	0.13698	0.00025
2453692.845470	-4.69804	0.41056	-3.95939	0.48869	0.13464	0.00042

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453692.848144	-4.23319	0.38349	-4.11985	0.47498	0.13498	0.00043
2453692.850794	-5.18519	0.40374	-4.36833	0.46324	0.13565	0.00038
2453692.853491	-4.61372	0.37279	-4.35219	0.46137	0.13619	0.00037
2453693.822865	-4.29957	0.34021	-4.47122	0.40475	0.13640	0.00030
2453693.826685	-2.88391	0.33323	-3.02631	0.39582	0.13638	0.00031
2453693.830493	-2.81502	0.34377	-2.60286	0.37831	0.13642	0.00030
2453694.836339	-0.64892	0.38707	-0.22584	0.44351	0.13603	0.00032
2453694.840263	0.26619	0.27756	0.90208	0.36565	0.13656	0.00031
2453694.844060	0.03275	0.32818	0.81068	0.38918	0.13648	0.00031
2453695.826003	3.39180	0.37351	3.68762	0.40102	0.13565	0.00035
2453695.829394	3.50801	0.33151	3.84033	0.36171	0.13599	0.00035
2453695.832971	3.47731	0.30479	4.22654	0.33654	0.13579	0.00032
2453696.810816	2.76416	0.30100	2.79594	0.37547	0.13642	0.00028
2453696.814300	2.82787	0.29056	3.07455	0.34890	0.13587	0.00029
2453696.817785	3.02973	0.33083	3.07964	0.43365	0.13604	0.00029
2453697.840089	4.01470	0.65531	5.53488	0.76016	0.12969	0.00060
2453697.843388	3.51569	0.51464	3.85310	0.52446	0.13298	0.00048
2453697.846826	3.40226	0.47967	3.79477	0.57497	0.13310	0.00050
2453698.800872	1.15261	0.48489	-0.41896	0.53564	0.13383	0.00046
2453698.804205	-0.33565	0.52095	0.40364	0.53691	0.13290	0.00049
2453698.807770	0.69819	0.50900	1.21905	0.58293	0.13177	0.00051
2453699.829807	-2.97167	0.37419	-2.26387	0.46707	0.13555	0.00033
2453699.833257	-3.56776	0.38390	-3.38822	0.48755	0.13525	0.00033
2453699.836729	-3.44800	0.37925	-3.67757	0.45236	0.13533	0.00032
2453700.861845	-2.47281	0.40332	-2.21996	0.48871	0.13588	0.00030
2453700.865074	-2.84517	0.34031	-2.24855	0.40788	0.13595	0.00028
2453700.868570	-2.01000	0.30202	-1.20080	0.37308	0.13555	0.00029
2453724.852851	-4.08797	0.34222	-4.05429	0.40500	0.13619	0.00029
2453724.856138	-3.36420	0.32372	-3.53448	0.35787	0.13564	0.00031
2453724.859379	-2.79296	0.33289	-3.26328	0.35277	0.13657	0.00031

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453725.801458	-5.62882	0.30318	-5.00371	0.38695	0.13650	0.00027
2453725.804746	-5.35079	0.34737	-4.17470	0.42981	0.13651	0.00027
2453725.807963	-5.68100	0.31899	-5.03180	0.36507	0.13628	0.00028
2453726.845139	-4.42657	0.30822	-4.99396	0.38364	0.13620	0.00032
2453726.848450	-5.50790	0.34272	-6.18978	0.37848	0.13646	0.00032
2453726.851667	-5.17599	0.34434	-6.25688	0.36673	0.13620	0.00032
2453727.761092	-4.09173	0.36566	-3.88115	0.46514	0.13724	0.00035
2453727.764322	-3.82004	0.37984	-3.70773	0.41088	0.13685	0.00034
2453727.767562	-4.62746	0.36844	-4.39414	0.45928	0.13685	0.00034
2453728.740578	-3.38973	0.32277	-3.39078	0.38166	0.13737	0.00031
2453728.743946	-3.18653	0.35360	-2.92621	0.45176	0.13656	0.00031
2453728.747106	-2.83288	0.34704	-2.77002	0.40246	0.13657	0.00030
2453729.859861	1.20141	0.41969	2.78567	0.51803	0.13615	0.00034
2453729.863032	0.77732	0.35808	0.94284	0.46667	0.13635	0.00037
2453729.866470	1.97862	0.36767	2.11902	0.41502	0.13564	0.00037
2453732.729703	2.58925	0.44791	2.78922	0.44328	0.13700	0.00042
2453732.732111	2.09890	0.33397	2.60011	0.47382	0.13693	0.00043
2453732.734564	1.96161	0.29726	2.94865	0.38741	0.13631	0.00044
2453744.812769	-5.13430	0.63333	-4.15730	0.68945	0.13711	0.00075
2453744.815246	-4.15496	0.67584	-3.27827	0.83987	0.13711	0.00065
2453744.817654	-5.67826	0.60045	-3.77693	0.61023	0.13597	0.00063
2453757.637147	-0.15337	0.42603	0.76746	0.56081	0.13615	0.00038
2453757.641024	0.97346	0.45831	0.86619	0.59199	0.13717	0.00039
2453757.644971	0.70780	0.36730	0.33700	0.51141	0.13550	0.00037
2453758.688630	1.31276	0.40488	1.29294	0.43171	0.13649	0.00036
2453758.692623	1.27766	0.33147	1.41918	0.48231	0.13652	0.00034
2453758.696396	1.15695	0.30550	2.08542	0.43302	0.13685	0.00033
2453759.644705	-1.85163	0.30115	-2.40665	0.45782	0.13610	0.00030
2453759.648479	-1.14632	0.28685	-2.11553	0.43028	0.13635	0.00031
2453759.652530	-1.21156	0.23881	-1.17221	0.36465	0.13621	0.00035

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453760.653349	-2.91288	0.36425	-3.17470	0.40547	0.13589	0.00036
2453760.657262	-2.66004	0.36360	-3.06461	0.41443	0.13624	0.00036
2453760.661127	-2.82337	0.35694	-3.30695	0.37406	0.13663	0.00035
2453761.708948	-2.51463	0.42030	-2.78156	0.55595	0.13716	0.00036
2453761.712756	-3.32690	0.37017	-3.15736	0.47879	0.13729	0.00036
2453761.716657	-3.18505	0.41857	-3.35220	0.43692	0.13644	0.00035
2453762.662114	-2.43072	0.39465	-2.25494	0.48325	0.13720	0.00035
2453762.665795	-2.43043	0.28196	-2.83444	0.35986	0.13663	0.00033
2453762.669707	-2.12679	0.36245	-2.53808	0.46394	0.13608	0.00038
2453763.698937	2.17103	0.33444	2.35671	0.43316	0.13680	0.00030
2453763.702802	1.91533	0.33692	1.43154	0.42558	0.13732	0.00030
2453763.706633	1.49771	0.35371	1.34220	0.40022	0.13658	0.00031
2453764.728616	3.67728	0.36931	4.28114	0.45642	0.13642	0.00030
2453764.732447	3.22551	0.33040	2.93520	0.42039	0.13658	0.00030
2453764.736371	3.19365	0.36708	2.87574	0.50931	0.13635	0.00032
2453765.657368	4.43218	0.33196	5.18193	0.45279	0.13680	0.00028
2453765.661211	4.40651	0.39125	4.72292	0.49458	0.13707	0.00027
2453765.665088	4.11763	0.37038	3.94850	0.49213	0.13689	0.00027
2453778.667837	-4.56145	0.37282	-4.43952	0.43250	0.13732	0.00041
2453778.670325	-4.60848	0.34830	-4.75497	0.45378	0.13707	0.00039
2453778.672767	-5.40699	0.42716	-6.09734	0.58583	0.13705	0.00040
2453782.638032	1.43300	0.19407	0.80740	0.26384	0.13665	0.00027
2453782.642904	0.76140	0.27401	0.12364	0.34314	0.13664	0.00031
2453783.654951	1.53628	0.26693	2.32227	0.29831	0.13735	0.00030
2453783.660113	1.05830	0.21507	1.31893	0.28061	0.13760	0.00033
2453784.674902	-0.86533	0.39535	-0.00028	0.45475	0.13622	0.00035
2453784.679797	-0.16788	0.34380	0.70342	0.35369	0.13590	0.00033
2453785.668983	-1.13425	0.34768	-0.84463	0.41179	0.13573	0.00035
2453785.674191	-1.01095	0.39389	-0.30691	0.48541	0.13597	0.00037
2453786.750781	-4.73138	0.32365	-4.51487	0.41579	0.13575	0.00030

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453786.755792	-4.78953	0.27751	-4.45363	0.37643	0.13661	0.00030
2453787.655545	-3.69812	0.28723	-4.29072	0.37640	0.13648	0.00027
2453787.660568	-4.08116	0.27830	-4.40042	0.31070	0.13671	0.00028
2453788.638616	-3.48309	0.33511	-3.82527	0.35611	0.13710	0.00032
2453788.643384	-3.25630	0.27196	-3.05815	0.31929	0.13703	0.00028
2453789.613804	1.49651	0.26540	1.90608	0.30363	0.13676	0.00029
2453789.618780	0.99518	0.31205	0.98293	0.38610	0.13669	0.00035
2453790.748498	4.74863	0.22779	5.23132	0.27449	0.13571	0.00034
2453790.753521	3.92697	0.20052	4.86758	0.24438	0.13665	0.00031
2453808.537626	3.90182	0.25514	3.76615	0.35204	0.13707	0.00045
2453808.540230	2.89277	0.33224	3.77691	0.45427	0.13732	0.00042
2453808.542649	2.99188	0.34280	2.46290	0.35639	0.13684	0.00046
2453809.604393	5.22656	0.16877	4.51950	0.27234	0.13662	0.00034
2453809.608317	4.26927	0.31073	3.39476	0.35890	0.13705	0.00033
2453809.612229	4.07993	0.16162	3.20713	0.26130	0.13699	0.00033
2453810.658371	-0.05851	0.17443	-0.28622	0.26078	0.13770	0.00043
2453810.662144	-0.12081	0.17455	0.63034	0.28196	0.13757	0.00039
2453810.666033	0.74261	0.15578	0.27340	0.24962	0.13639	0.00041
2453811.683044	-2.98056	0.17066	-2.38198	0.23681	0.13651	0.00035
2453811.688217	-2.85404	0.19468	-2.27285	0.27833	0.13647	0.00037
2453812.645913	-4.61105	0.18124	-4.54847	0.26531	0.13667	0.00040
2453812.649814	-4.92640	0.18006	-4.49255	0.29425	0.13689	0.00037
2453812.653598	-4.27373	0.18221	-3.19306	0.25882	0.13617	0.00038
2453813.630309	-3.19757	0.16745	-2.90646	0.27159	0.13722	0.00031
2453813.635563	-3.29842	0.16604	-3.49204	0.27488	0.13634	0.00029
2453814.661899	-4.12972	0.17099	-4.08484	0.27186	0.13608	0.00036
2453814.667061	-3.78260	0.17146	-4.33271	0.25501	0.13618	0.00033
2453815.693940	-2.33075	0.19362	-1.21793	0.31736	0.13672	0.00032
2453815.699194	-2.06644	0.17993	-1.15527	0.29776	0.13650	0.00032
2453816.638937	0.81430	0.19151	0.92143	0.31554	0.13577	0.00031

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453816.644145	0.12925	0.17988	0.77329	0.27790	0.13587	0.00031
2453817.658025	2.87955	0.19675	3.31779	0.31532	0.13621	0.00039
2453817.663175	1.98647	0.18812	2.11958	0.31398	0.13634	0.00038
2453828.534908	4.52179	0.30088	5.37743	0.39387	0.13662	0.00046
2453828.537349	5.32379	0.21992	6.31553	0.34944	0.13641	0.00044
2453828.539826	4.00680	0.25526	4.51888	0.36365	0.13701	0.00044
2453829.647509	1.45493	0.29364	1.62129	0.40690	0.13628	0.00032
2453829.652671	1.70956	0.29367	2.46572	0.37818	0.13621	0.00031
2453830.560854	0.25581	0.25232	-0.36557	0.34988	0.13730	0.00028
2453830.565900	0.30241	0.29521	0.90468	0.41969	0.13646	0.00029
2453831.596400	0.29193	0.22587	0.90937	0.34314	0.13657	0.00030
2453831.601435	0.95164	0.18922	0.85546	0.28357	0.13611	0.00034
2453832.589715	2.68455	0.26691	1.88013	0.40457	0.13545	0.00031
2453832.594750	2.50350	0.24690	1.54812	0.30242	0.13629	0.00032
2453833.585460	5.75407	0.20977	5.35009	0.29805	0.13710	0.00027
2453833.590483	5.46304	0.23538	5.62886	0.31606	0.13720	0.00028
2453834.566541	6.46400	0.21788	6.47847	0.30615	0.13730	0.00028
2453834.571668	6.69855	0.19801	6.52318	0.26755	0.13686	0.00029
2453835.612500	7.32113	0.22735	6.64022	0.30387	0.13661	0.00036
2453835.617720	6.14006	0.19927	6.58063	0.29210	0.13772	0.00034
2453836.637581	4.28250	0.23831	3.95666	0.33424	0.13689	0.00033
2453836.642604	3.86918	0.23156	3.89909	0.33420	0.13692	0.00033
2453861.572945	8.61695	0.39223	8.84252	0.54340	0.13621	0.00052
2453861.576463	7.70314	0.50845	6.99263	0.56575	0.13704	0.00055
2453861.579958	6.94769	0.33766	7.96717	0.38269	0.13565	0.00056
2453862.488945	5.57358	0.19512	6.58458	0.28957	0.13757	0.00033
2453862.492451	6.35092	0.18495	6.35603	0.29460	0.13663	0.00032
2453862.495958	4.95670	0.19257	4.86184	0.28901	0.13664	0.00033
2453863.522435	2.45371	0.29902	2.44613	0.36307	0.13694	0.00031
2453863.525941	2.96147	0.33352	3.07455	0.43428	0.13682	0.00032

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453863.529448	1.78956	0.32316	1.73863	0.37986	0.13793	0.00033
2453864.514435	1.35169	0.20031	1.86700	0.29243	0.13658	0.00042
2453864.518092	1.43072	0.20024	0.65366	0.27922	0.13615	0.00045
2453864.521668	1.13280	0.26407	1.74968	0.34313	0.13422	0.00044
2453865.492965	0.19268	0.33348	-0.86707	0.43091	0.13619	0.00032
2453865.496506	0.01718	0.20893	-1.19161	0.30685	0.13656	0.00031
2453865.500013	-1.18088	0.22671	-1.32131	0.33754	0.13695	0.00031
2453866.485116	-0.95099	0.33290	-0.14356	0.41194	0.13773	0.00028
2453866.488600	-1.41174	0.30124	-0.25903	0.38497	0.13749	0.00029
2453866.492141	-1.07988	0.23191	-0.69113	0.31023	0.13738	0.00031
2453867.521408	1.03249	0.29843	0.05133	0.38540	0.13727	0.00040
2453867.524845	1.04143	0.32170	0.22428	0.45079	0.13746	0.00036
2453867.528328	2.05492	0.30773	1.00450	0.39202	0.13679	0.00039
2453868.498052	3.70134	0.31316	3.42507	0.43858	0.13688	0.00039
2453868.501558	3.49350	0.40602	3.34016	0.44566	0.13685	0.00035
2453868.505007	3.47494	0.34737	3.85642	0.34361	0.13821	0.00038
2453869.548243	5.23166	0.23201	4.51579	0.32771	0.13787	0.00039
2453869.551761	4.14809	0.18535	3.90187	0.28355	0.13705	0.00038
2453869.555337	4.24482	0.38471	4.06305	0.49511	0.13725	0.00039
2453870.483803	2.99006	0.32238	3.12346	0.40417	0.13702	0.00032
2453870.489034	3.25602	0.25687	3.03370	0.35434	0.13636	0.00030
2453871.541390	2.12108	0.19914	1.35399	0.29712	0.13642	0.00036
2453871.545001	1.32262	0.20535	0.18534	0.31308	0.13573	0.00037
2453871.548473	-0.04792	0.23682	-0.00362	0.34152	0.13627	0.00038
2453872.569432	-4.27831	0.25573	-3.24867	0.34613	0.13608	0.00053
2453872.571897	-2.99019	0.25021	-3.34073	0.35111	0.13622	0.00052
2453872.574362	-2.61178	0.27673	-2.97922	0.33531	0.13550	0.00056
2453887.502616	4.78192	0.37816	5.47524	0.40712	0.13824	0.00063
2453887.505023	4.77403	0.59958	5.41109	0.59343	0.13675	0.00075
2453887.507569	4.33287	0.61483	4.86646	0.72567	0.13934	0.00080

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2453887.510022	4.08618	0.45585	5.29439	0.45503	0.13625	0.00065
2453887.512569	4.89602	0.37904	5.39969	0.45244	0.13634	0.00074
2454075.803672	-0.40502	0.42779	-0.46552	0.41460	0.13611	0.00040
2454075.806068	1.50431	0.39863	1.84167	0.36859	0.13517	0.00037
2454075.808487	-0.29027	0.48227	-0.01426	0.52272	0.13528	0.00037
2454075.814286	-0.50082	0.61080	0.79561	0.73358	0.13250	0.00066
2454075.815340	1.18197	0.56987	1.74401	0.60614	0.13144	0.00063
2454075.816393	-0.49125	0.70066	0.93344	0.70252	0.13311	0.00063
2454075.817458	-2.47933	0.64372	-3.36405	0.59534	0.13374	0.00064
2454075.818523	-1.06212	0.65674	0.41528	0.54008	0.13357	0.00063
2454075.819576	-0.12312	0.50990	1.74553	0.58507	0.13236	0.00060
2454075.820629	-0.93448	0.64063	0.09652	0.57924	0.13456	0.00061
2454075.821683	-0.54236	0.59284	1.42412	0.58247	0.13416	0.00058
2454075.822736	-1.11096	0.55581	0.21034	0.53026	0.13278	0.00058
2454075.823801	-2.48877	0.58117	-1.73808	0.52680	0.13302	0.00057
2454075.824854	-1.02110	0.62773	-0.13149	0.76167	0.13389	0.00056
2454075.825896	0.06052	0.48921	0.51677	0.59171	0.13297	0.00058
2454075.826961	-0.68503	0.56628	0.49149	0.55320	0.13321	0.00057
2454075.828014	0.29572	0.67012	0.89180	0.55517	0.13398	0.00058
2454075.829079	0.39239	0.50443	1.17166	0.49941	0.13395	0.00059
2454075.830121	-2.19718	0.58157	-1.87604	0.58886	0.13274	0.00058
2454075.831186	-0.01828	0.54382	1.61068	0.51508	0.13293	0.00057
2454075.832250	0.11574	0.47120	1.20656	0.46034	0.13457	0.00057
2454075.833292	-1.55500	0.57790	-0.47879	0.51895	0.13335	0.00056
2454075.834346	0.81765	0.47252	2.15214	0.59279	0.13498	0.00057
2454075.835410	-1.35892	0.51261	0.36853	0.60407	0.13426	0.00056
2454075.836464	-0.46659	0.53420	0.03961	0.64429	0.13313	0.00056
2454075.837517	-0.22228	0.54752	1.13248	0.65923	0.13335	0.00055
2454075.838582	-0.42690	0.43691	0.78817	0.52966	0.13304	0.00055
2454075.839635	0.08643	0.36252	-0.04878	0.44244	0.13490	0.00056

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2454075.840700	0.59522	0.51622	1.21554	0.40639	0.13383	0.00055
2454075.841753	-1.72468	0.46576	-0.75693	0.43396	0.13392	0.00055
2454075.842795	0.13262	0.51185	0.81648	0.58107	0.13311	0.00056
2454075.843872	0.33495	0.56241	0.02999	0.56713	0.13448	0.00054
2454075.844913	-1.41839	0.55449	0.67168	0.60732	0.13397	0.00055
2454075.846511	0.97367	0.44564	0.52098	0.42859	0.13313	0.00054
2454075.847576	-1.86750	0.49571	-2.24789	0.53002	0.13388	0.00055
2454075.848617	-2.08276	0.59974	-3.06660	0.58150	0.13356	0.00057
2454075.849671	-1.18885	0.62655	-0.50588	0.55183	0.13445	0.00055
2454075.850724	1.37196	0.58501	-0.13766	0.57698	0.13363	0.00057
2454075.851789	-1.15251	0.48630	-1.26440	0.48217	0.13226	0.00053
2454075.852831	0.12352	0.49197	0.08101	0.44527	0.13398	0.00056
2454075.853884	-0.43020	0.52295	-1.56351	0.52102	0.13339	0.00054
2454075.854949	0.13639	0.56040	-0.78612	0.56488	0.13356	0.00057
2454075.856002	-1.72402	0.52807	-1.62393	0.55841	0.13495	0.00054
2454075.857044	-0.82455	0.50620	-1.77767	0.47204	0.13225	0.00056
2454075.858097	-1.60142	0.54212	-1.25345	0.66845	0.13266	0.00054
2454075.859139	-1.28517	0.57126	0.31052	0.55655	0.13205	0.00059
2454075.860215	-0.39922	0.61575	1.50470	0.55092	0.13278	0.00059
2454078.829833	4.91606	0.36716	5.55053	0.46578	0.13575	0.00036
2454078.833294	5.35496	0.36458	6.03390	0.48105	0.13690	0.00029
2454078.837207	5.91966	0.35173	6.40174	0.42314	0.13687	0.00030
2454082.818017	0.87895	0.32950	1.11781	0.36818	0.13759	0.00034
2454082.822231	0.99663	0.41558	0.66403	0.49198	0.13683	0.00040
2454097.765399	-2.10192	0.37243	-2.28619	0.48434	0.13652	0.00039
2454097.770862	-2.99266	0.40772	-3.06580	0.53106	0.13704	0.00038
2454097.777101	-3.91556	0.32616	-3.40748	0.41512	0.13680	0.00038
2454097.781244	-3.37766	0.30950	-2.39172	0.38054	0.13751	0.00039
2454098.660233	-5.18222	0.51087	-3.80848	0.60078	0.13537	0.00056
2454098.662675	-5.00712	0.45861	-4.73114	0.53118	0.13572	0.00053

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2454098.665071	-4.58575	0.52536	-3.62798	0.55440	0.13628	0.00055
2454099.748061	-5.62188	0.43659	-4.11452	0.58974	0.13675	0.00049
2454099.750631	-5.38632	0.32017	-4.80574	0.45203	0.13447	0.00048
2454099.753050	-6.21846	0.32258	-6.35664	0.38763	0.13643	0.00045
2454114.787257	3.51793	0.44659	3.95007	0.52849	0.13793	0.00044
2454114.790683	1.66451	0.41399	2.86015	0.56535	0.13750	0.00043
2454114.793275	1.71321	0.45509	2.32482	0.58138	0.13752	0.00051
2454114.795694	1.84490	0.57601	2.96843	0.68036	0.13666	0.00053
2454114.797928	3.17423	0.30955	1.89135	0.33889	0.13762	0.00051
2454117.750890	-3.15477	0.42117	-3.69871	0.52403	0.13830	0.00039
2454117.754929	-2.33218	0.34817	-2.03636	0.42233	0.13815	0.00033
2454117.759293	-1.44703	0.30831	-1.15857	0.41670	0.13776	0.00036
2454120.749210	3.28637	0.34886	3.65548	0.40222	0.13688	0.00037
2454120.753169	0.63454	0.34052	1.50530	0.47373	0.13728	0.00035
2454120.757567	1.18980	0.31838	1.80787	0.32805	0.13779	0.00030
2454122.742851	0.88762	0.26837	0.26089	0.33678	0.13794	0.00028
2454122.748361	0.36172	0.31167	0.21773	0.34466	0.13795	0.00028
2454125.758510	-6.16717	0.40760	-5.90941	0.45161	0.13626	0.00039
2454125.760941	-7.39069	0.27547	-6.69317	0.36766	0.13633	0.00039
2454125.763360	-7.27624	0.41962	-7.53511	0.49028	0.13627	0.00040
2454132.667449	-5.37357	0.46596	-5.83357	0.52750	0.13600	0.00049
2454132.669949	-5.20446	0.46002	-5.11993	0.64998	0.13652	0.00055
2454132.672461	-5.17832	0.48068	-4.85899	0.45557	0.13742	0.00052
2454132.676917	-6.21484	0.43711	-6.52003	0.54199	0.13690	0.00044
2454132.679336	-4.73404	0.43323	-4.41567	0.44053	0.13696	0.00043
2454132.681778	-5.16766	0.37977	-5.61952	0.38389	0.13691	0.00045
2454132.691928	-6.47711	0.32261	-6.90335	0.44177	0.13615	0.00032
2454132.695875	-6.04535	0.31519	-6.22972	0.34383	0.13539	0.00032
2454132.699706	-6.79397	0.32429	-7.70438	0.39766	0.13576	0.00031
2454136.701436	-2.48902	0.37678	-1.84619	0.51179	0.13643	0.00029

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2454136.705267	-2.33357	0.29755	-1.10838	0.41958	0.13614	0.00028
2454142.677407	-2.75128	0.25751	-2.48071	0.28697	0.13646	0.00033
2454142.681308	-2.27732	0.24088	-2.74364	0.28798	0.13686	0.00032
2454142.685069	-2.84101	0.32402	-2.89733	0.37651	0.13653	0.00029
2454166.613458	-1.12035	0.22935	-1.70597	0.31335	0.13664	0.00026
2454166.619140	-1.05450	0.20261	-1.51304	0.29511	0.13582	0.00027
2454168.675542	-6.19463	0.38517	-5.76963	0.43246	0.13573	0.00032
2454168.681271	-5.33497	0.16123	-4.74094	0.23850	0.13601	0.00030
2454170.626158	-1.82166	0.22988	-1.86718	0.31584	0.13586	0.00028
2454170.631620	-2.13190	0.27144	-1.78458	0.36529	0.13626	0.00028
2454172.587035	3.20746	0.17677	3.03937	0.26125	0.13636	0.00024
2454172.592659	3.79186	0.16381	4.48580	0.24447	0.13664	0.00023
2454174.611040	4.71344	0.15733	4.66928	0.25790	0.13656	0.00026
2454174.616619	4.57324	0.16672	5.54240	0.24401	0.13672	0.00027
2454191.525052	0.65747	0.22450	-0.47619	0.33697	0.13571	0.00042
2454191.527540	1.49187	0.43920	0.81656	0.45945	0.13619	0.00048
2454191.529959	1.89339	0.20441	2.68386	0.30574	0.13800	0.00046
2454191.536127	0.74922	0.22718	0.57901	0.29643	0.13550	0.00044
2454191.538639	0.90527	0.25478	1.26925	0.37177	0.13593	0.00044
2454191.540988	2.54203	0.29026	1.85344	0.41918	0.13586	0.00042
2454194.545916	-4.49891	0.20924	-4.26572	0.30108	0.13597	0.00034
2454194.548566	-4.89952	0.25441	-4.58994	0.35891	0.13711	0.00037
2454194.551298	-5.46554	0.29277	-5.62680	0.41853	0.13627	0.00038
2454194.553809	-5.51905	0.24624	-4.90097	0.33023	0.13668	0.00041
2454197.597061	-1.97442	0.25964	-1.48502	0.39273	0.13612	0.00034
2454197.602281	-1.79455	0.21595	-1.15708	0.31176	0.13521	0.00035
2454199.542616	4.57282	0.25858	4.76014	0.33621	0.13602	0.00038
2454199.546134	3.91166	0.28303	3.21962	0.36996	0.13728	0.00035
2454199.549583	3.10526	0.27876	3.16928	0.40147	0.13656	0.00034
2454202.532565	-1.09156	0.23384	-0.28315	0.28023	0.13606	0.00036

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2454202.536130	-1.12259	0.29050	-0.32495	0.42680	0.13725	0.00036
2454202.539579	-0.72971	0.29584	-0.36586	0.41266	0.13673	0.00038
2454225.564093	2.39145	0.54862	2.21679	0.55520	0.13639	0.00055
2454225.566813	2.83234	0.49678	2.06947	0.56883	0.13675	0.00060
2454225.569510	3.92635	0.36093	3.37093	0.46116	0.13576	0.00057
2454225.572229	2.46529	0.28325	2.82865	0.34828	0.13567	0.00054
2454228.539967	-3.91008	0.38197	-4.35740	0.41459	0.13517	0.00034
2454228.545105	-4.41474	0.33846	-5.14485	0.42585	0.13457	0.00036
2454232.508224	3.10066	0.42171	2.40423	0.50818	0.13719	0.00058
2454232.512969	2.18616	0.26650	2.49974	0.34098	0.13502	0.00036
2454234.534446	5.75893	0.20196	6.00486	0.31461	0.13607	0.00039
2454234.539619	5.46665	0.32433	6.00014	0.33367	0.13608	0.00040
2454237.476763	2.68254	0.38327	2.28480	0.42313	0.13665	0.00089
2454237.479043	0.36213	0.46870	-0.23531	0.59285	0.13016	0.00086
2454237.481624	0.15950	0.35133	0.85637	0.42608	0.13275	0.00085
2454419.851821	-4.34496	0.40607	-4.04881	0.54242	0.13676	0.00028
2454419.855571	-3.96753	0.36050	-3.43975	0.45094	0.13704	0.00027
2454419.859403	-4.19767	0.35962	-3.73518	0.44202	0.13743	0.00028
2454420.821069	-2.24310	0.36632	-2.65993	0.48658	0.13692	0.00028
2454420.824900	-2.66239	0.32543	-3.09376	0.40777	0.13747	0.00029
2454420.828731	-2.06278	0.36132	-2.10228	0.43279	0.13711	0.00029
2454421.827854	0.75776	0.38545	0.51327	0.48660	0.13724	0.00027
2454421.831604	0.98427	0.38215	0.63695	0.54371	0.13738	0.00028
2454421.835482	-0.08478	0.40944	-0.36775	0.51278	0.13705	0.00028
2454422.809938	3.20943	0.44431	3.77599	0.57457	0.13616	0.00032
2454422.813699	3.15814	0.39937	3.54966	0.48905	0.13645	0.00031
2454422.817577	2.90675	0.41374	3.09528	0.49120	0.13605	0.00031
2454423.855973	4.61546	0.52499	3.41588	0.62739	0.13540	0.00043
2454423.859631	4.57756	0.47954	4.18756	0.51476	0.13678	0.00042
2454423.863497	4.89955	0.42707	4.64497	0.51368	0.13681	0.00043

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2454424.804118	6.75452	0.46645	7.06819	0.56867	0.13654	0.00030
2454424.809581	6.42086	0.40634	6.40445	0.48654	0.13731	0.00029
2454425.790494	6.91247	0.32465	6.67280	0.41677	0.13677	0.00032
2454425.796062	6.10161	0.39162	6.34050	0.51934	0.13628	0.00031
2454426.802000	4.66715	0.38287	4.61034	0.49691	0.13671	0.00025
2454426.807568	4.47686	0.37494	4.14142	0.46583	0.13673	0.00024
2454427.797763	1.50751	0.39135	0.19559	0.48838	0.13717	0.00018
2454428.838888	0.57864	0.36771	-0.66106	0.49067	0.13664	0.00019
2454429.789184	1.62763	0.38411	1.35543	0.42407	0.13666	0.00024
2454446.822355	-1.54065	0.42783	-2.07115	0.56660	0.13577	0.00049
2454446.824960	-2.26056	0.51804	-1.49126	0.60438	0.13684	0.00050
2454446.827089	-2.76374	0.51997	-3.11853	0.60962	0.13647	0.00053
2454446.829208	-2.27533	0.58886	-3.17179	0.71146	0.13621	0.00049
2454448.827313	1.57517	0.53853	0.43547	0.44517	0.13767	0.00050
2454448.829396	2.48211	0.47791	1.18330	0.58196	0.13834	0.00050
2454448.831537	2.19380	0.51251	1.29828	0.57165	0.13842	0.00052
2454448.833598	-0.28843	0.46734	0.82591	0.48180	0.13652	0.00049
2454448.835716	2.14322	0.50445	1.10211	0.53830	0.13841	0.00053
2454450.819511	10.72418	0.53350	9.04266	0.63738	0.13570	0.00050
2454450.825159	5.43642	0.55952	4.52099	0.51937	0.13578	0.00050
2454450.828064	3.81987	0.48345	3.80534	0.53510	0.13649	0.00049
2454450.830831	4.13238	0.49239	4.20373	0.59799	0.13608	0.00050
2454453.825642	0.56250	0.39649	1.57835	0.50149	0.13556	0.00039
2454453.828328	-0.33051	0.51327	0.53707	0.48145	0.13527	0.00039
2454453.831013	0.12130	0.43598	0.02916	0.51172	0.13657	0.00039
2454453.833710	0.35338	0.47824	0.56301	0.53206	0.13632	0.00038
2454479.762583	-3.57294	0.44971	-4.76022	0.49352	0.13533	0.00036
2454479.766426	-3.01337	0.40944	-4.14729	0.47906	0.13513	0.00034
2454479.770257	-2.73507	0.38540	-3.32446	0.48443	0.13604	0.00033
2454481.794820	-3.58001	0.47052	-5.10263	0.53997	0.13642	0.00031

Table 14—Continued

JD [days]	RV_{TERRA} [m s ⁻¹]	σ_{TERRA} [m s ⁻¹]	$RV_{\text{TERRA-Red}}$ [m s ⁻¹]	$\sigma_{\text{TERRA-Red}}$ [m s ⁻¹]	S-index	σ_S
2454481.798593	-3.91573	0.46729	-5.96624	0.53358	0.13529	0.00033
2454481.802471	-4.09181	0.51118	-5.96808	0.62922	0.13598	0.00033
2454483.698715	1.29746	0.48000	0.54456	0.60555	0.13636	0.00029
2454483.702476	0.36225	0.43088	-0.61381	0.58691	0.13631	0.00028
2454483.706284	0.79127	0.46459	-0.60462	0.61089	0.13612	0.00030
2454524.660927	-1.50725	0.21138	-2.41298	0.26748	0.13657	0.00028
2454524.666564	-1.09767	0.34225	-2.12258	0.41817	0.13721	0.00027
2454551.682681	0.80538	0.21556	0.20082	0.34696	0.13893	0.00068
2454551.684440	0.66815	0.21057	0.08806	0.34274	0.13615	0.00066
2454551.686199	1.67740	0.31122	1.11200	0.39440	0.13729	0.00076
2454551.687947	0.38769	0.21278	0.17045	0.32349	0.13889	0.00074
2454551.689717	0.11022	0.23703	-0.53709	0.36704	0.13859	0.00076
2454552.636480	1.20206	0.17504	0.14744	0.28928	0.13710	0.00021
2454553.586819	4.13702	0.38970	4.20592	0.54014	0.13555	0.00048
2454553.589400	3.81644	0.45388	4.04901	0.61480	0.13623	0.00049
2454553.592108	3.72233	0.25398	4.32143	0.37483	0.13629	0.00050
2454553.594793	4.38820	0.22153	4.24417	0.31672	0.13627	0.00052