

Seismic detection of acoustic sharp features in the CoRoT target HD 49933

Anwesh Mazumdar¹, Eric Michel², H. M. Antia³, and Sebastien Deheuvels^{2,4}

¹ Homi Bhabha Centre for Science Education, TIFR, V. N. Purav Marg, Mankhurd, Mumbai 400088, India.

² LESIA, Observatoire de Paris, UMR8109 CNRS, Université Pierre et Marie Curie, Université Denis Diderot, Place J. Janssen, 92195 Meudon, France.

³ Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India.

⁴ Astronomy Department, Yale University, P.O. Box 208101, New Haven, CT 065208101, USA.

ABSTRACT

The technique of determining the acoustic location of layers of sharp changes in the sound speed inside a star from the oscillatory signal in its frequencies is applied on a solar-type star, the CoRoT target, HD 49933. We are able to determine the acoustic depth of the second helium ionisation zone of HD 49933 to be 794^{+55}_{-68} s. The acoustic depth of the base of the convective zone is found to be 1855^{+173}_{-412} s where the large error bars reflect the ambiguity in the result, which is difficult to determine with present precision on the frequencies because of the intrinsically weak nature of the signal. The positions of both these layers are consistent with those in a representative stellar model of HD 49933.

Key words. Stars: individual: HD 49933 – Stars: oscillations – Stars: interiors – Stars: fundamental parameters

1. Introduction

Any localised feature in the sound speed inside a star, such as that caused by the change in the temperature gradient at the base of the convection zone, introduces an oscillatory term in frequencies as a function of radial order n , that is proportional to $\sin(2\tau_d\omega_{n,\ell} + \phi)$, (Gough 1990) where $\tau_d = \int_{r_d}^R \frac{dr}{c}$ is the acoustic depth of the localised feature, c is the speed of sound, r_d is the radial distance where the feature is located, $\omega_{n,\ell}$ is the angular frequency of a mode with radial order n and degree ℓ , and ϕ is a phase factor. This oscillatory signature has been extensively studied for the Sun in order to determine the extent of overshoot below the solar convection zone (Monteiro et al. 1994; Basu et al. 1994; Roxburgh & Vorontsov 1994). It has been proposed earlier that this may be used for distant stars also to find the position of the base of the convective envelope or the second helium ionisation zone (Monteiro et al. 2000; Mazumdar & Antia 2001; Roxburgh & Vorontsov 2003; Ballot et al. 2004; Basu et al. 2004; Mazumdar 2005). Indeed, Miglio et al. (2010) have used the modulation of the frequency separations to determine the location of the second helium ionisation zone in a red giant star.

We present here the results of applying this technique for the first time on a main sequence solar-type star: HD 49933, observed by the CoRoT mission in 2007–08 (Baglin et al. 2006).

2. The technique

The oscillatory signal in the frequencies is quite small and is embedded in the frequencies together with a smooth trend arising from the regular variation of the sound speed in the stellar interior. It can be enhanced by using the second differences $\delta^2\nu(n, \ell) = \nu(n-1, \ell) - 2\nu(n, \ell) + \nu(n+1, \ell)$, instead of the fre-

quencies themselves (see, e.g., Gough 1990; Basu et al. 1994; Mazumdar & Antia 2001; Basu et al. 2004).

The acoustic depths of the base of the convective zone (BCZ) and the second helium ionisation zone (HIZ), τ_{BCZ} and τ_{HIZ} , respectively, can be obtained from the data by fitting the second differences to a suitable function representing the oscillatory signals from these layers (Mazumdar & Antia 2001). We follow Basu et al. (2004) in choosing the following functional form:

$$\delta^2\nu = a_0 + b_0 \sin(4\pi\nu\tau_{\text{BCZ}} + \phi_{\text{BCZ}}) + (c_0 + c_2/\nu^2) \sin(4\pi\nu\tau_{\text{HIZ}} + \phi_{\text{HIZ}}), \quad (1)$$

where a_0 , b_0 , c_0 , c_2 , τ_{BCZ} , ϕ_{BCZ} , τ_{HIZ} and ϕ_{HIZ} are 8 free parameters of fitting. Since the number of data points in the observed set of frequencies for HD 49933 is quite small, we optimise the number of free parameters to strike a balance between a fair representation of the oscillatory signal and a reasonable χ^2 . We find that the smooth component is fairly constant over the range of frequencies that we use, as is the amplitude of the signal from the BCZ. However, the amplitude of the signal from the HIZ varies more sharply with frequency, and thus requires at least one frequency-dependent term. We note that Basu et al. (2004) have shown that the exact form of the amplitudes of the oscillatory signal does not affect the results significantly. To test our results, we also fit a function suggested by Houdek & Gough (2007) (simplified version of Eq. (22) in that paper, with 8 free parameters).

The fit is carried out through a nonlinear χ^2 minimisation, weighted by the errors in the data. The errors in the second differences are correlated, and this is taken care of by defining the χ^2 using a covariance matrix. The effects of the errors are considered by producing 1000 realisations of the data, where the mean values of the frequencies are skewed by random errors corresponding to a normal distribution. The successful convergence of such a non-linear fitting procedure is somewhat dependent on the choice of reasonable initial guesses. To remove the effect of

Send offprint requests to: anwesh@tifr.res.in

initial guesses affecting the final fitted parameters, we carry out the fit for multiple combinations of starting values. For each realisation, 100 random combinations of initial guesses are tried for fitting the function above and from among the resulting fits, the one which produces the minimum value of χ^2 is accepted as the fit for that particular realisation.

The median value of each parameter for 1000 realisations is taken as its fitted value. The error in the parameter is estimated from the range of values covering 68% area about the median (corresponding to 1σ error). As an example, the histogram of distribution of the parameters τ_{BCZ} and τ_{HIZ} over multiple realisations are illustrated in Fig. 1. Thus the quoted errors in these parameters reflect the width of these histograms on two sides of the median value.

The strength of the oscillatory signals are measured through the amplitudes of the respective oscillatory components in the frequencies themselves. These are denoted by A_{BCZ} and A_{HIZ} respectively, and estimated by averaging the amplitudes of the oscillatory components in Eq. 1 over the observed frequency range, and correcting for the scaling factor of $4 \sin^2(\pi\tau/\tau_0)$ arising in the second differences (see Mazumdar & Antia 2001). In the present study, we use the amplitudes only as an index of the significance of the detected depth of the acoustic glitches.

We tested the technique on the frequencies of a polytropic model, with a polytropic index of 1.5, provided by Christensen-Dalsgaard & Mullan (1994). Since there are no sharp acoustic features in such a model, we do not expect to find any oscillatory pattern in the second differences of the frequencies. Indeed, if we attempt to fit a function of the form given by Eq. 1 to the second differences of this model, we only find a signal corresponding to half of the large separation, which is the periodicity at which the successive points occur due to $\ell = 0$ or 2 and $\ell = 1$.

The range of acoustic depth that can be probed by this technique is limited. The smallest depth that can be fitted corresponds to the range in the observed frequencies, and is ~ 350 s. On the other hand, the sampling frequency of the data points determines the largest acoustic depth that can be fitted. The average large separation of HD 49933 being $85 \mu\text{Hz}$, the upper limit on the acoustic depth that can be fitted is ~ 2940 s.

3. Results

3.1. Fitting the CoRoT frequencies

We use the second differences of the frequencies obtained by Benomar et al. (2009) to extract the acoustic depths of BCZ and HIZ of HD 49933. We ignore frequencies which have errors of more than $3 \mu\text{Hz}$. This gives a total of 31 second differences to be used for $\ell = 0, 1, 2$. We consider four different subsets of the data, namely: (i) $\ell = 0$ modes only (12 data points), (ii) $\ell = 1$ modes only (16 data points), (iii) $\ell = 0, 1$ modes (28 data points), and (iv) $\ell = 0, 1, 2$ modes (31 data points). This was done to test for consistency of the results when modes of different degrees are considered, especially since the $\ell = 0$ and $\ell = 2$ modes in the data have, on the average, higher errors than the $\ell = 1$ modes. However, the number of data points in the first two cases are only marginally larger than the number of free parameters in the fitting function, and these cases are used only for testing purposes.

The fits to the second differences of the mean frequencies and the distribution of the fitted acoustic depths of BCZ and HIZ for different realisations of the data are shown in Fig 1. Although we fit 1000 realisations of the data, only those realisations with $\tau_{\text{BCZ}} \leq 2700$ s are considered due to reasons explained below.

Table 1. Results of fitting second differences of HD 49933 frequencies to Eq. 1. Median values of the acoustic depths and amplitudes and their errors are listed. The last two rows give the calculated and the fitted values of a representative stellar model of HD 49933.

ℓ	τ_{BCZ} (s)	τ_{HIZ} (s)	A_{BCZ} (μHz)	A_{HIZ} (μHz)
0	1877^{+273}_{-340}	847^{+248}_{-253}	$1.06^{+0.59}_{-0.34}$	$1.92^{+5.80}_{-0.70}$
1	1496^{+438}_{-194}	812^{+45}_{-79}	$0.75^{+0.24}_{-0.20}$	$1.47^{+0.33}_{-0.29}$
0,1	1626^{+327}_{-262}	806^{+45}_{-61}	$0.66^{+0.18}_{-0.15}$	$1.40^{+0.28}_{-0.22}$
0,1,2	1855^{+173}_{-412}	794^{+55}_{-68}	$0.71^{+0.19}_{-0.15}$	$1.32^{+0.34}_{-0.31}$
Model	1998	882		
0,1,2	1982^{+447}_{-586}	848^{+61}_{-62}	$0.47^{+0.19}_{-0.13}$	$1.37^{+0.24}_{-0.22}$

A fit to the data of subset (iv) above to the form suggested by Houdek & Gough (2007) is shown also in Fig 1. Table 1 lists the median values of the acoustic depths with estimated errors and the corresponding amplitudes of BCZ and HIZ signals with respective estimated errors for the four data subsets.

The acoustic depth of the HIZ obtained from the different sets are consistent within the errorbars. The amplitude of the HIZ oscillatory function is also significant at more than 2.5σ level in each set.

However, the BCZ oscillatory signal is not so consistent among different sets. For all values of ℓ , we get a peak in the histogram close to the value $\tau_0/2 \sim 2900$ s, which corresponds to the Nyquist frequency. The occurrence of this peak could mean that the amplitude of the oscillatory function corresponding to BCZ is close to zero, so that the fitted sine wave has periodicity equal to the natural spacing of the data points, *i.e.*, the large separation. While this seems to be the case for $\ell = 1$ modes, the second differences of the radial modes appear to have an actual oscillatory signal corresponding to this frequency! Thus this effect cannot be fully ascribed to the Nyquist effect. This is also borne out by the fact that this peak persists even when we combine data with multiple values of ℓ , which should not be the case (see, *e.g.*, Mazumdar & Antia 2001). We cannot find a satisfactory explanation as to the origin of this periodicity in the radial modes, apart from an artifact introduced due to the noise. In order to eliminate this artifact for all subsets, we discard the realisations for which the BCZ signal is fitted with an acoustic depth of around 2900 s. Few realisations yield τ_{BCZ} values higher than the acoustic midpoint of the star (~ 2900 s), and these are discarded also. Actually, we consider only those realisations for which $\tau_{\text{BCZ}} < 2700$ s, since the peak in the histogram around the acoustic midpoint has a typical spread of 200 s on either side of the peak. This results in reduction of the number of valid realisations by nearly 30%.

In all the cases, there is a peak in the histogram for τ_{BCZ} around 1950 s, which is most likely due to the base of the convective envelope in the star. For $\ell = 0$ and $\ell = 0, 1, 2$, this is the dominant peak of the histogram, while for $\ell = 1$ and $\ell = 0, 1$, this is the secondary peak. For the latter two cases, the dominant peak is centred around 1370 s. If this value is to correspond to τ_{BCZ} it would imply a very shallow convection zone in the star, which is an unlikely scenario given the expected mass and age of HD 49933. This peak cannot be associated with any artifact due to the sampling frequency, but on the other hand, we cannot assign any physical significance to it.

To understand the difference in the values of τ_{BCZ} obtained for different ℓ , we carried out the following test. For each case,




Fig. 1. Fits of Eq. 1 to the second differences of the mean CoRoT frequencies for HD 49933 and the histograms for the fitted values of τ_{HIZ} and τ_{BCZ} for different realisations of the data are shown for different subsets (corresponding to different choices of ℓ values). The second differences of the frequencies of $\ell = 0$ (blue), $\ell = 1$ (red) and $\ell = 2$ (green) modes of HD 49933 and their fit to Eq. 1 (magenta curve) are shown in the left panel. The dotted cyan line in the bottom left panel shows the fit by the Houdek & Gough (2007) form. The corresponding histograms of the fitted values of τ_{BCZ} (in red) and τ_{HIZ} (in blue) for different realisations are shown in the right panel. N denotes the total number of valid realisations after rejecting fits with $\tau_{\text{BCZ}} > 2700$ s. The solid bands at the top of the right panel indicate the range of initial guesses for the two parameters in each fit.

we held the parameter τ_{BCZ} in Eq. 1 fixed at a certain value, while allowing all other parameters to be free, and monitored the value of the reduced χ^2 obtained by the resulting fit to the mean second differences. The fixed value of τ_{BCZ} was changed continuously between 1000 s and 4800 s to find the minima in the reduced χ^2 as this parameter varies. The χ^2 is plotted as a function of τ_{BCZ} in Fig. 2. We find that the $\ell = 1$ modes show a deep minimum around 1370 s, but only a slight inflexion around 1950 s. In contrast, the location of the minimum and the inflexion are interchanged for the $\ell = 0$ modes. When these modes are combined along with the $\ell = 2$ modes, we have minima at both the values. The third minimum around 2900 s for all cases has already been discussed. The almost mirrored shapes of the graphs above τ_{BCZ} values of 2900 s merely illustrate the aliasing effect, as elaborated in Mazumdar & Antia (2001).

3.2. Fitting model frequencies

We fit Eq. 1 to the second differences of frequencies calculated from a representative stellar model for the star HD 49933 as well. The model is generated with the CESAM2k evolution




Fig. 2. The variation of χ^2 with the adopted value of τ_{BCZ} is shown for four subsets of the CoRoT data on HD 49933 corresponding to different choices of the degree, ℓ . The vertical dotted line shows the acoustic mid-point of the star.

code (Morel & Lebreton 2008) and uses standard physical inputs, including convective core overshoot, but no diffusive mixing. While we use the theoretical frequencies as the mean values




Fig. 3. Fits of Eq. 1 to the second differences of the frequencies of a theoretical model for HD 49933 (*top panel*) and the histograms for different realisations of the data for the fitted values of τ_{HIZ} and τ_{BCZ} (*bottom panel*) are shown. See Fig. 1 for explanations of colours and symbols.

in this case, we set the errors in the frequencies equal to the errors in the observed frequencies for the corresponding mode, in order to mimic the uncertainties present in the data. The results of such a fit are displayed in Fig. 3. The model values as well as the fitted median values of τ_{HIZ} and τ_{BCZ} are given in Table 1.

We find that the median values of the acoustic depths of both BCZ and HIZ are recovered from the frequencies. However, the error in the value of τ_{BCZ} is quite large due to the flat nature of the histogram in Fig. 3. In contrast, the histogram of the HIZ signal is sharply peaked and hence the estimated error in the median value of τ_{HIZ} is also smaller. We conclude that the intrinsic oscillatory signal arising from the acoustic glitch at BCZ is quite weak, and it is difficult to extract this signal even with model frequencies. It is beyond the scope of this study to investigate why this signal is so weak, and it needs to be probed whether this is a common feature of all stars in the relevant part of the HR diagram. The HIZ signal, however, is strong and can easily be used to extract the acoustic depth of the second helium ionisation zone.

4. Summary

We have applied the method of determining the acoustic depth of layers of sharp variation in sound speed in stellar interiors from the oscillatory signal in frequencies to the CoRoT primary target HD 49933.

The oscillatory signal is made more pronounced in comparison to the smooth change in sound speed by using the second differences of the frequencies. We fit a function with two oscillatory components corresponding to the two major layers of sharp variation in sound speed, *viz.*, the base of the convective envelope and the second helium ionisation zone. To prove that the technique does not produce artificial signatures of oscilla-

tory signals when there are none, we tested it on the frequencies of a polytropic model where there are no real layers of sharp variation in the sound speed. In this case the only significant oscillatory signal corresponds to the uniform spacing of the frequencies, as expected.

We find that the oscillatory signal arising due to the second helium ionisation zone is quite strong in the CoRoT data of HD 49933. We estimate the acoustic depth of this layer to be 794^{+55}_{-68} s. This is borne out by the frequencies of modes of all available degrees: $\ell = 0, 1, 2$, when considered either separately, or in combination. The acoustic depth of this layer in a representative stellar model is consistent with this value, and can also be easily extracted by our technique.

The oscillatory signal due to the base of the convective envelope is far weaker in the data, in comparison to the helium signal. Moreover, we find that the signal is not consistent between modes of different degrees. We find additional periodicities in the frequencies corresponding to different acoustic depths in modes of $\ell = 0$ and $\ell = 1$, for which we are unable to assign any physical acoustic feature inside the star. It seems likely that these are artifacts due to the errors in the data which have no physical relevance. Nevertheless, we do find an oscillatory signal corresponding to an acoustic depth of 1855^{+173}_{-412} s in the combined frequency set of different degrees, which is consistent with the position of the base of the convective envelope in a representative stellar model. This oscillatory signal is weak even in the model frequencies, and is fairly difficult to extract from the second differences. These results prove that the technique of extracting acoustic depths of sharp features inside a star can be successfully applied for real asteroseismic data.

Acknowledgements. CoRoT (Convection, Rotation and planetary Transits) space mission has been developed and is operated by the French Space agency CNES in collaboration with Austria, Belgium, Brazil, ESA's RSSd and Science Programmes, Germany and Spain. AM acknowledges support from the National Initiative on Undergraduate Science (NIUS) undertaken by the Homi Bhabha Centre for Science Education – Tata Institute of Fundamental Research (HBCSE-TIFR), Mumbai, India.

References

- Baglin, A., Auvergne, M., Barge, P., et al. 2006, ESA SP, 1306, 33
- Ballot, J., Turck-Chièze, S., & García, R. A. 2004, A&A, 423, 1051
- Basu, S., Antia, H. M., & Narasimha, D. 1994, MNRAS, 267, 209
- Basu, S., Mazumdar, A., Antia, H. M., & Demarque, P. 2004, MNRAS, 350, 277
- Benomar, O., Baudin, F., Campante, T. L., et al. 2009, A&A, 507, L13
- Christensen-Dalsgaard, J., & Mullan, D. J. 1994, MNRAS, 270, 921
- Gough, D. O. 1990, Progress of Seismology of the Sun and Stars, 367, 283
- Houdek, G., Gough, D. O. 2007, MNRAS, 375, 861
- Mazumdar, A. 2005, A&A, 441, 1079
- Mazumdar, A., & Antia, H. M. 2001, A&A, 377, 192
- Miglio, A., Montalbán, J., Carrier, F., et al. 2010, A&A, 520, L6
- Monteiro, M. J. P. F. G., Christensen-Dalsgaard, J., & Thompson, M. J. 1994, A&A, 283, 247
- Monteiro, M. J. P. F. G., Christensen-Dalsgaard, J., & Thompson, M. J. 2000, MNRAS, 316, 165
- Morel, P., & Lebreton, Y. 2008, Ap&SS, 316, 61
- Roxburgh, I. W. & Vorontsov, S. V. 1994, MNRAS, 268, 880
- Roxburgh, I. W. & Vorontsov, S. V. 2003, A&A, 411, 215