TRANSITION MATRICES FOR SYMMETRIC AND QUASISYMMETRIC HALL-LITTLEWOOD POLYNOMIALS

NICHOLAS A. LOEHR, LUIS G. SERRANO, AND GREGORY S. WARRINGTON

Abstract. We introduce explicit combinatorial interpretations for the coefficients in some of the transition matrices relating to skew Hall-Littlewood polynomials $P_{\lambda/\mu}(t)$ and Hivert's quasisymmetric Hall-Littlewood polynomials $G_{\gamma}(t)$. More specifically, we provide:

(1) the G-expansions of the Hall-Littlewood polynomials $P_{\lambda}(t)$, the monomial quasisymmetric polynomials M_{α} , the quasisymmetric Schur polynomials S_{α} , and the peak quasisymmetric functions K_{α} ;

The F-expansion of $P_{\lambda/\mu}(t)$ is facilitated by introducing starred tableaux.

1. Introduction

The ring of symmetric functions Sym and the ring of quasisymmetric functions QSym both play important roles in algebra and combinatorics. Much of the combinatorial richness arising from these rings stems from their various distinguished bases and the relationships between these bases. The goal of this paper is to present explicit, combinatorial descriptions of several such transition matrices relating to the Hall-Littlewood polynomials. Figure 1 illustrates some of the bases discussed.

In the top triangle in Figure 1 are included two classical bases for the ring of symmetric functions: the *Schur functions* s_u and the monomial symmetric functions m_{ν} . The s_u and m_{ν} are closely related to a third, one-parameter family of symmetric functions $P_{\lambda}(t)$, known as Hall-Littlewood polynomials. More specifically, $P_{\lambda}(t)$ equals s_{λ} at $t = 0$, and it equals m_{λ} at $t = 1$. The $P_{\lambda}(t)$ arose out of a problem studied by P. Hall. Hall had used his eponymous algebra (isomorphic to the algebra of symmetric functions) to encode the structure of finite abelian pgroups. However, at the time there was no known explicit basis of symmetric functions with the same structure constants as that of the natural basis for Hall's algebra. D. E. Littlewood [20] solved this problem in 1961 with his introduction of the $P_{\lambda}(t)$.

The bottom triangle of Figure 1 consists of quasisymmetric analogues of the above bases. In the context of quasisymmetric functions, the monomial quasisymmetric functions, M_{β} , are a very natural analogue of the m_{ν} . There are several possible quasisymmetric analogues of the Schur functions, including the "quasisymmetric Schur functions" S_{α} introduced in [13] and discussed later in this paper. However, for reasons described in the next paragraph, we anchor the lowerleft portion of the bottom triangle in Figure 1 by Gessel's *fundamental quasisymmetric functions*.

⁽²⁾ an expansion of $P_{\lambda/\mu}(t)$ in terms of the F_{α} 's.

Date: November 11, 2018.

²⁰¹⁰ Mathematics Subject Classification. 05E05; 05E10, 16T30.

Key words and phrases. symmetric functions, quasisymmetric functions, Hall-Littlewood polynomials, standardization, Young tableaux, noncommutative symmetric functions.

This work was partially supported by a grant from the Simons Foundation (#244398 to Nicholas Loehr).

Second author supported by a Natural Science and Engineering Research Council of Canada PDF grant.

Third author supported in part by National Security Agency grant H98230-09-1-0023 and National Science Foundation grant DMS-1201312.

This work was partially supported by a grant from the Simons Foundation (#197419 to Greg Warrington).

FIGURE 1. Prism of bases and transitions.

denoted here by F_{α} . By defining an action of the Hecke algebra on polynomials which leaves the quasisymmetric functions invariant, Hivert [15] constructed the quasisymmetric Hall-Littlewood polynomials $G_{\gamma}(t)$. (See also work of Lascoux, Novelli, and Thibon [17] for constructions of quasisymmetric and noncommutative symmetric functions with extra parameters.) Similarly to what happens in the top triangle, specialization of the $G_{\gamma}(t)$ at $t = 0$ (which corresponds to the southwest-pointing arrow in Figure 1) yields F_{γ} , while specialization at $t = 1$ yields M_{γ} .

We now motivate our choice of the F_{α} as the desired quasisymmetric analogue of the Schur functions. The Schur functions are the prototypical example of a symmetric function with combinatorial expansions in terms of both a collection of semistandard objects (i.e., semistandard Young tableaux) and of *standard objects* (i.e., standard Young tableaux). The first case is that of the classical expansion in terms of monomials weighted by the Kostka numbers. The second expansion (due to Gessel [10]) expresses the Schur functions in terms of fundamental quasisymmetric functions F_{α} . This expansion, which follows from the technique of *standardization*, is indicated by the vertical line connecting s_{μ} and F_{α} in Figure 1. Such standardizations have been used recently to give F-expansions of various symmetric functions including plethysms of Schur functions [22], the modified Macdonald polynomials [11, 12], the Lascoux-Leclerc-Thibon (LLT) polynomials [19], and (conjecturally) the image of a Schur function under the Bergeron-Garsia nabla operator [21].

Given Hivert's construction, the following question arises. Is there an expansion of the $P_{\lambda}(t)$ in terms of the $G_{\gamma}(t)$, which would interpolate between the F-expansion of the s_{μ} at $t = 0$ and the M-expansion of the m_{ν} at $t = 1$? The main purpose of this paper is to provide such an expansion, as well as other change-of-basis matrices between different bases of the Hall algebra and the algebra of quasisymmetric functions. In terms of Figure 1, we provide the middle vertical edge as well as the reverses of the two downward directed edges in the bottom face (namely, from each of F_{α} and M_{β} to $G_{\gamma}(t)$). More specifically, our principal results are:

(1) G -expansion of the P basis. In Theorem 32 we give an explicit combinatorial expansionof the Hall-Littlewood polynomials $P_{\lambda}(t)$ in terms of the Hivert quasisymmetric Hall-Littlewood polynomials $G_{\gamma}(t)$. This provides the desired t-interpolation between Gessel's F-expansion of Schur polynomials (i.e., $t = 0$) and the obvious expansion of m_{ν} 's into M_{β} 's (i.e., $t = 1$). A key step in our construction is to combine the two expansions described in the next two items.

- (2) F -expansion of the P basis. One of the main tools for our calculations is the definition of a new class of tableaux, called starred tableaux. With these, we give in Theorem 19 a combinatorial expansion of the skew Hall-Littlewood polynomials $P_{\lambda/\mu}(t)$ in terms of the fundamental quasisymmetric functions F_{α} . A minor variation to our method gives a corresponding expansion for the dual Hall-Littlewood polynomials $Q_{\lambda/\mu}$ (see Theorem 25).
- (3) G -expansions of the F basis and the M basis. In Theorems 26 and 29 we give explicitcombinatorial expansions for the F_{α} and the M_{β} in terms of the $G_{\gamma}(t)$. These are inverse matrices to those found in [15]. See Remark 1 below for the relationship to existing results in the realm of noncommutative symmetric functions.
- (4) G-expansions of the S basis and K_{α} . In Theorems 35 and 39 we give explicit combinatorial expansions for the quasisymmetric Schur functions S_β and the peak quasisymmetric functions K_{α} in terms of the $G_{\gamma}(t)$'s.

Remark 1. The algebra NSym of noncommutative symmetric functions, developed by Gelfand, Krob, Lascoux, Leclerc, Retakh, and Thibon [9], is a dual Hopf algebra to QSym. Many bases of NSym have been developed (see, for example, [13, 17, 29]). In papers such as [1, 2, 24, 30], attempts are made to construct bases that are suitable analogues in NSym of the Hall-Littlewood polynomials. Of the transition matrices we describe in §4 and §5, the only one which we know to have been constructed in the dual setting of NSym is that of $\mathcal{M}(F,G)$: Hivert introduces noncommutative Hall-Littlewood polynomials $H_{\gamma}(t)$ that are dual to the $G_{\gamma}(t)$. If we let R denote the basis for NSym of *ribbon Schur functions*, then it follows that $\mathcal{M}(F, G) = \mathcal{M}(H, R)^T$ (see [24, Theorem 6.13]). Our proof of Theorem 26 is substantially the same as Hivert's proof. We include our proof both for completeness and so as to give a derivation that does not invoke NSym.

The structure of the paper is as follows. For ease of reference, we define all bases discussed here in §2 and summarize known combinatorial transition matrices in §3. The expansions of the various polynomials in terms of the F_{α} 's and the $G_{\gamma}(t)$'s are presented in §4 and §5, respectively. §6 contains a few remarks on transition matrices for plethystically transformed Hall-Littlewood polynomials. Finally, specific examples of transition matrices discussed here are listed in Appendix 7. SAGE code for computing these transition matrices is available on the third author's website [32].

2. Review of Symmetric and Quasisymmetric Bases

This section reviews the definitions of the symmetric and quasisymmetric functions appearing in Figure 1. Logically, the precise definitions of the various bases are not needed in this paper, as the expansions found in $\S 4$ and $\S 5$ are derived from the known transition matrices of $\S 3$. However, the material of this section is included for completeness.

It will be necessary to introduce a number of functions in the variables x_1, \ldots, x_N , some of which have an extra parameter t . For brevity, we will suppress much of this notation. For example, $G_{\gamma}(x_1,\ldots,x_N;t)$, $G_{\gamma}(x;t)$, $G_{\gamma}(t)$, and G_{γ} all refer to the Hivert quasisymmetric function indexed by γ .

2.1. Compositions and Partitions. Given $n \in \mathbb{N}$, a *composition of* n is a sequence $\alpha =$ $(\alpha_1, \alpha_2, \ldots, \alpha_k)$ of positive integers (called parts) with $\alpha_1 + \cdots + \alpha_k = n$. Define the length $\ell(\alpha)$ to be the number of parts of α , and the *size* $|\alpha|$ to be the sum of its parts. For example, the 3

composition $\alpha = (2, 4, 1)$ has $\ell(\alpha) = 3$ and $|\alpha| = 7$. We may abbreviate the notation, writing α as 241, when no confusion can arise. Let Comp_n be the set of compositions of n, and let Comp_n be the set of all compositions. A composition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k) \in \text{Comp}_n$ is called a *partition* of n iff $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_k$. We write Par_n for the set of partitions of n and Par for the set of all partitions. For any composition or partition α , we define $\alpha_j = 0$ for all $j > \ell(\alpha)$.

For $n \in \mathbb{N}^+$, there are 2^{n-1} compositions of n and 2^{n-1} subsets of $[n-1] = \{1, 2, \ldots, n-1\}$. One can define natural bijections between these sets of objects as follows. Given $\alpha \in \text{Comp}_n$ as above, let

$$
sub(\alpha) = {\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3, \dots, \alpha_1 + \dots + \alpha_{k-1}} \subseteq [n-1].
$$

The inverse bijection sends any subset $T = \{t_1 < t_2 < \cdots < t_m\} \subseteq [n-1]$ to

comp(T) =
$$
(t_1, t_2 - t_1, t_3 - t_2, \dots, t_m - t_{m-1}, n - t_m) \in \text{Comp}_n
$$
.

Given $\alpha, \beta \in \text{Comp}_n$, we say β is *finer* than α , denoted $\beta \succeq \alpha$, iff $\text{sub}(\alpha) \subseteq \text{sub}(\beta)$. Informally, β is finer than α if we can chop up some of the parts of α into smaller pieces (without reordering anything) and obtain β . For example, $1111 \succeq 121 \succeq 31 \succeq 4$.

2.2. Symmetric Polynomials. Let F be a commutative ring, and let \mathfrak{S}_N denote the symmetric group on N letters. A polynomial $f \in \mathbb{F}[x_1, \ldots, x_N]$ is called *symmetric* iff

$$
w(f(x)) = f(x_{w(1)}, x_{w(2)}, \dots, x_{w(N)}) = f(x_1, x_2, \dots, x_N)
$$
 for all $w \in \mathfrak{S}_N$.

Write $\text{Sym}_N = \text{Sym}_N(\mathbb{F})$ for the ring of symmetric polynomials in N variables with coefficients in F; we usually omit the ground ring F from the notation. For each $n \geq 0$, let $Sym_N^n = Sym_N^n(\mathbb{F})$ be the subspace of Sym_N consisting of zero and the homogeneous polynomials of degree n. For $N \geq n$, bases of the vector space Sym_N^n are naturally indexed by partitions of n.

Given $\lambda \in \text{Par}_n$ of length $k \leq N$, the monomial symmetric polynomial $m_\lambda(x_1, \ldots, x_N)$ is the sum of all distinct monomials that can be obtained by permuting subscripts in x^{λ} = $x_1^{\lambda_1} x_2^{\lambda_2} \cdots x_k^{\lambda_k}$. For $N \geq n$, $\{m_\lambda(x_1, \ldots, x_N) : \lambda \in \text{Par}_n\}$ is readily seen to be a basis of Sym_N^n .

Now suppose $N \ge n$ and that $\nu \in \text{Par}_n$ is a partition with *distinct* parts. The *skew-symmetric* polynomial indexed by ν in N variables is

$$
a_{\nu}(x_1,...,x_N) = \sum_{w \in \mathfrak{S}_N} \text{sgn}(w) w(x^{\nu}) = \det ||x_i^{\nu_j}||_{1 \leq i,j \leq N}.
$$

In particular, letting $\delta_N = (N-1, N-2, \ldots, 2, 1), a_{\delta_N}(x_1, \ldots, x_N) = \prod_{1 \leq i < j \leq N} (x_i - x_j)$ is the Vandermonde determinant. Given $\lambda \in \text{Par}_n$, the Schur symmetric polynomial indexed by λ in N variables is

$$
s_{\lambda}(x_1,\ldots,x_N)=\frac{a_{\lambda+\delta_N}(x_1,\ldots,x_N)}{a_{\delta_N}(x_1,\ldots,x_N)}.
$$

For the rest of the paper, let t be an indeterminate, and let F be any field containing $\mathbb{Q}(t)$ as a subfield. Following $[23, \frac{5}{11}]$, pp. 204–7, we define the Hall-Littlewood symmetric polynomials as follows. Fix $\lambda \in \text{Par}_n$ and $N \ge n$. Define $[0]_t = 0$ and for $m \ge 1$, $[m]_t = 1 + t + t^2 + \cdots + t^{m-1}$. Also set $[m]!_t = \prod_{i=1}^m [i]_t$, and $[0]!_t = 1$. Define $v_\lambda(t) = \prod_{i \geq 0} [m_i]!_t$ where m_i denotes the number of occurrences of i as a part of λ . Then the Hall-Littlewood polynomial indexed by λ is

(1)
$$
P_{\lambda}(x;t) = \frac{1}{v_{\lambda}(t)} \sum_{w \in \mathfrak{S}_N} w\left(x^{\lambda} \prod_{i < j} \frac{x_i - tx_j}{x_i - x_j}\right).
$$

Setting $t = 0$ in P_{λ} gives s_{λ} , whereas setting $t = 1$ in P_{λ} gives m_{λ} . Thus, the Hall-Littlewood basis "interpolates" between the Schur basis and the monomial basis.

One can define Schur polynomials and Hall-Littlewood polynomials more concretely by giving combinatorial descriptions of their expansions in terms of monomial symmetric polynomials. See §3.1 and §3.5 below.

2.3. Quasisymmetric Polynomials. A polynomial $f \in \mathbb{F}[x_1, \ldots, x_N]$ is called *quasisymmetric* iff for every composition $\alpha = (\alpha_1, \ldots, \alpha_k)$ with at most N parts and every $1 \leq i_1 < i_2 < \cdots <$ $i_k \leq N$, the monomials $x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_k^{\alpha_k}$ and $x_{i_1}^{\alpha_1}$ $\frac{\alpha_1}{i_1} x_{i_2}^{\alpha_2}$ $\frac{\alpha_2}{i_2} \cdots x_{i_k}^{\alpha_k}$ $\frac{\alpha_k}{i_k}$ have the same coefficient in f. Write $\operatorname{QSym}_N = \operatorname{QSym}_N(\mathbb{F})$ for the ring of quasisymmetric polynomials in N variables with coefficients in F. For each $n \geq 0$, let $\text{QSym}_N^n = \text{QSym}_N^n(\mathbb{F})$ be the subspace of QSym_N consisting of zero and the homogeneous polynomials of degree n. For $N \geq n$, bases of the vector space QSym_N^n are naturally indexed by compositions of n. Symmetric polynomials are quasisymmetric, so Sym_N^n is a subspace of QSym_N^n .

For $\alpha \in \text{Comp}_n$ of length $k \leq N$, the monomial quasisymmetric polynomial $M_{\alpha}(x_1, \ldots, x_N)$ is the sum of all monomials $x_{i_1}^{\alpha_1}$ $\frac{\alpha_1}{i_1} x_{i_2}^{\alpha_2}$ $\frac{\alpha_2}{i_2}\cdots x_{i_k}^{\alpha_k}$ $\frac{\alpha_k}{i_k}$ for which $1 \leq i_1 < i_2 < \cdots < i_k \leq N$. For $N \geq n$, ${M_\alpha(x_1, \ldots, x_N): \alpha \in \text{Comp}_n}$ is readily seen to be a basis of QSym_N.

Next, for $\alpha \in \text{Comp}_n$ with length at most N, define Gessel's fundamental quasisymmetric polynomial [10] by

(2)
$$
F_{\alpha}(x_1,\ldots,x_N)=\sum x_{w_1}x_{w_2}\cdots x_{w_n},
$$

where we sum over all subscript sequences $w = w_1w_2\cdots w_n$ such that $1 \leq w_1 \leq w_2 \leq \cdots \leq w_n$ $w_n \leq N$ and for all $j \in sub(\alpha)$, $w_j < w_{j+1}$. In other words, strict increases in the subscripts are required in the "breaks" between parts of the composition α . Call sequences w satisfying these conditions $\text{sub}(\alpha)$ -compatible, and write $x^w = x_{w_1} \cdots x_{w_n}$. A routine inclusion-exclusion argument (cf. §3.8 below) shows that for $N \geq n$, $\{F_{\alpha}(x_1,\ldots,x_N): \alpha \in \text{Comp}_n\}$ is a basis of QSym^n_N . Note that some authors index fundamental quasisymmetric polynomials by pairs n, T where $T \subseteq [n-1]$. Additionally, various letters $(F, L, Q, \text{etc.})$ have been used to denote these polynomials.

As in the symmetric case, we would like to have quasisymmetric Hall-Littlewood polynomials (depending on a parameter t) that interpolate between F_{α} (when $t = 0$) and M_{α} (when $t = 1$). We sketch the definition of one such family of polynomials, introduced and studied by Hivert [15]. Quasisymmetric functions arise as the invariants of a certain action of \mathfrak{S}_N on polynomials. From this action, one can define divided difference operators in a degenerate Hecke algebra $H_N(0)$ which can then be lifted to $H_N(q)$. Hivert's quasisymmetric Hall-Littlewood polynomials thereby arise from a corresponding t-analogue \Box_{ω} of the Weyl symmetrizer. For a composition α of length $k \leq N$, define

$$
G_{\alpha}(x_1,\ldots,x_N;t) = \frac{1}{[k]!_t[N-k]!_t} \boxdot_{\omega} (x_1^{\alpha_1} \cdots x_k^{\alpha_k}).
$$

As in the case of symmetric Hall-Littlewood polynomials, there is a more concrete combinatorial definition of G_{α} giving its expansion into monomials. We discuss this definition in §3.10.

Hivert's $G_{\alpha}(t)$ are quasisymmetric versions of the Hall-Littlewood polynomials $P_{\lambda}(t)$; when $t = 0$, the latter specialize to Schur polynomials. In light of these relationships, the $G_{\alpha}(t)$ can be viewed as a quasisymmetric t-analogue of Schur functions. However, a more direct "quasisymmetric Schur function" has been introduced by Haglund, Luoto, Mason, and van Willigenburg [13] via specializations of nonsymmetric Macdonald polynomials to Demazure atoms. These have a combinatorial expansion, which we now describe.

Given a composition α , one forms its *diagram* by placing α_i boxes, or cells, in the *i*-th row from top to bottom, and left-justifying the rows. The entries in the diagram of α are given matrix coordinates (i, j) . A filling T of the diagram of α is a semistandard composition tableau (SSCT) if the following three conditions hold.

- (C1) The entries in each row are weakly decreasing when read from left to right.
- $(C2)$ Entries in the leftmost column of T are strictly increasing when read from top to bottom.
- (C3) Entries satisfy the *triple rule*, namely, if (i, k) and (j, k) are two cells in the same column, with $i < j$, then:
	- if $\alpha_i \ge \alpha_j$, then either $T(j,k) < T(i,k)$ or $T(i,k-1) < T(j,k)$;
	- if $\alpha_i < \alpha_j$, then either $T(j,k) < T(i,k)$ or $T(i,k) < T(j,k+1)$.

It follows from these conditions that all entries in the same column of an SSCT must be distinct.

The content of an SSCT T is cont $(T) = \alpha = (\alpha_1, \alpha_2, \ldots)$, where α_i is the number of times i appears in T. The corresponding monomial is $x^T = x^{\alpha}$. For example, the following picture shows an SSCT of shape 5264, content 32313212, and monomial $x_1^3x_2^2x_3^3x_4x_5^3x_6^2x_7x_8^2$.

A quasisymmetric Schur function S_{α} is defined as the generating series of semistandard composition tableaux, i.e., $\mathcal{S}_{\alpha} = \sum x^T$, where the sum runs over all SSCT of shape α . Part of their importance stems from the fact that they provide refinements of Schur functions, as in the formula

$$
s_{\lambda} = \sum_{\text{sort}(\alpha) = \lambda} S_{\alpha},
$$

where sort(α) is the partition obtained by organizing the parts of α from largest to smallest.

As discussed, the Hall-Littlewood functions specialize to Schur functions and monomial symmetric functions when t equals 0 and 1, respectively. The Schur P-functions are the $t = -1$ specializations. They are indexed by strict partitions, namely, partitions into distinct parts. The Schur P-functions are another example of a family of symmetric functions with combinatorial expansions in terms of both a collection of semistandard objects and a collection of standard objects. The first one is the classical expansion of the Schur P-functions as a generating series for shifted semistandard Young tableaux, where each tableau is weighted by its corresponding monomial (see, e.g., [14, 25, 27]). The second one is the expansion given by Stembridge [28] of the Schur P-functions as a sum over shifted standard Young tableaux of the corresponding peak quasisymmetric functions, denoted K_{α} . The K_{α} span an important subalgebra of QSym called the peak quasisymmetric algebra. While we will not be concerned with Schur P-functions in this paper, transition matrices involving the peak quasisymmetric functions will be presented.

Let Comp_n' be the set of compositions of n with no parts of length 1, except perhaps the last one. Note that $\alpha \in \text{Comp}'_n$ if and only if $\text{sub}(\alpha)$ is a subset of $[n-1]$ with no consecutive elements. For each $\alpha \in \text{Comp}'_n$, Stembridge [28] defines the peak quasisymmetric function K_{α} as the generating series of certain enriched P-partitions. We direct the reader to [28] for this definition and motivation. In this paper, however, we will use as a definition their expansion into fundamental quasisymmetric functions.

For $B \subseteq [n-1]$, let $B + 1 = \{b+1 : b \in B\} \setminus \{n\}$. Furthermore, we write \triangle for the symmetric difference between two sets, namely, $A \triangle B = (A \cup B) \setminus (A \cap B)$. Then, as in [28, Proposition 3.5] for $\alpha \in \text{Comp}'_n$, let

$$
K_{\alpha} = \sum_{\beta:\ \text{sub}(\alpha) \subseteq \text{sub}(\beta) \triangle(\text{sub}(\beta)+1)} F_{\beta}.
$$

3. Review of Known Transition Matrices

In the theory of symmetric and quasisymmetric polynomials, much combinatorial information is encoded in the transition matrices between various bases. Given two bases $B = \{B_\lambda : \lambda \in$ Par_n and $C = \{C_\lambda : \lambda \in \text{Par}_n\}$ of Sym_N^n , the transition matrix $\mathcal{M}(B, C)$ is the unique matrix (with entries in $\mathbb F$ and rows and columns indexed by partitions of n) such that

$$
B_{\lambda} = \sum_{\mu \in \text{Par}_n} \mathcal{M}(B, C)_{\lambda, \mu} C_{\mu}.
$$

Given a third basis D, it follows readily that $\mathcal{M}(B, D) = \mathcal{M}(B, C)\mathcal{M}(C, D)$ and $\mathcal{M}(C, B) =$ $\mathcal{M}(B,C)^{-1}$. We define $\mathcal{M}(B,C)$ similarly if B and C are bases of QSym_N^n , but here the rows and columns of the matrix are indexed by compositions of n. Finally, if B is a basis of Sym_N^n and C is a basis of QSymⁿ_N, then $\mathcal{M}(B, C)$ is a rectangular matrix expressing each B_λ as an **F**-linear combination of the C_{α} 's.

This section gives combinatorial formulas for previously known transition matrices associated to some of the edges in Figure 1. Transitions to the monomial bases offer alternate explicit definitions for Schur polynomials and the various forms of Hall-Littlewood polynomials. Specific examples of these transition matrices appear in Appendix 7. SAGE code for computing these transition matrices is available on the third author's website [32].

3.1. $\mathcal{M}(s, m)$. The expansion of Schur polynomials into monomials uses semistandard tableaux. For later work, we will also need tableaux of skew shape. Suppose $\lambda, \nu \in \text{Par}$ satisfy $\nu \subseteq \lambda$, i.e., $\nu_i \leq \lambda_i$ for all *i*. Define the *skew diagram*

$$
\lambda/\nu = \{(i,j) \in \mathbb{N}^+ \times \mathbb{N}^+ : 1 \le i \le \ell(\lambda), \nu_i < j \le \lambda_i\}.
$$

We will draw skew diagrams using the English convention where the longest rows are at the top. For $N \in \mathbb{N}^+$, a semistandard tableau of shape λ/ν with entries in $[N] = \{1, 2, ..., N\}$ is a function $T : \lambda/\nu \to [N]$ that is weakly increasing along rows and strictly increasing down columns. Writing $n = |\lambda/\nu|$, a standard tableau of shape λ/ν is a bijection $S : \lambda/\nu \to [n]$ that is also a semistandard tableau. Let $SSYT_N(\lambda/\nu)$ be the set of all semistandard tableaux of shape λ/ν with entries in [N], and let $SYT(\lambda/\nu)$ be the set of all standard tableaux of shape λ/ν . For any $T \in \text{SSYT}_N(\lambda/\nu)$, the *content of* T is the composition cont $(T) = (\alpha_1, \dots, \alpha_N)$, where α_i is the number of times i appears in a cell of T. The content monomial x^T is $x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_N^{\alpha_N}$.

The *skew Schur polynomial* in N variables can now be defined as

$$
s_{\lambda/\nu}(x_1,\ldots,x_N)=\sum_{T\in\text{SSYT}_N(\lambda/\nu)}x^T.
$$

The ordinary Schur polynomial s_λ is obtained by taking $\nu = (0)$ here. Skew Schur polynomials are symmetric, although this is not obvious from the combinatorial definition. Consequently, we have the following expansion of Schur polynomials in terms of the monomial symmetric polynomials.

Theorem 2. For all $\lambda, \mu \in \mathrm{Par}_n$, $\mathcal{M}(m, s)_{\lambda, \mu}$ is the number of semistandard tableaux of shape $λ$ and content $μ$. (This number is also called the Kostka number $K_{λ,μ}$.)

3.2. $\mathcal{M}(m, s)$. Eğecioğlu and Remmel [7] found the following combinatorial formula for the inverse Kostka matrix $\mathcal{M}(m, s)$. Fix $\mu \in \text{Par. A special rim-hook}$ is a sequence of cells in the diagram of μ that begins in the leftmost column and moves up and right through the diagram. The sign of a rim-hook occupying r rows is $(-1)^{r-1}$. A special rim-hook tableau S of shape μ is a dissection of the diagram of μ into a disjoint union of special rim-hooks. The sign of S, 7

 $sgn(S)$, is the product of the signs of the rim-hooks in it. The type of S is the integer partition obtained by listing the lengths of the rim-hooks in S in decreasing order.

Theorem 3. [7, Theorem 1] For all $\lambda, \mu \in \text{Par}_n$, $\mathcal{M}(m, s)_{\lambda, \mu} = \sum_{S} \text{sgn}(S)$ summed over all special rim-hook tableaux S of shape μ and type λ .

3.3. $\mathcal{M}(s, P)$. Lascoux and Schützenberger [18] first discovered a combinatorial formula for the "t-Kostka matrix" $\mathcal{M}(s, P)$ involving the famous "charge" statistic; many details were subsequently supplied by Butler [3]. Given a permutation w of [n], let $\text{IDes}(w)$ be the set of $k < n$ such that $k + 1$ appears to the left of k in w, and let $\text{chg}(w) = \sum_{k \in \text{IDes}(w)} (n - k)$.

Next, let v be a word of partition content (i.e., for all $k \geq 1$, the number of $(k + 1)$'s in v is no greater than the number of k 's). Extract one or more permutations from v as follows. Scan v from left to right marking the first 1, then the first 2 after that, etc., returning to the beginning of v when the right end is reached. Do this until the largest symbol has been marked. Remove the marked symbols from v (in the order they appear) to get the first permutation. Continue to extract permutations in this way until all symbols of v have been used, and let $\text{chg}(v)$ be the sum of the charges of the associated permutations. Finally, given a semistandard tableau T of partition content, let $w(T)$ be the word obtained by reading symbols row by row from top to bottom (i.e., longest row first), reading each row from right to left. Then define $chg(T) = chg(w(T)).$

Theorem 4. [3, 18] For all $\lambda, \mu \in \text{Par}_n$, $\mathcal{M}(s, P)_{\lambda, \mu} = \sum_T t^{\text{chg}(T)}$ summed over all semistandard tableaux T of shape λ and content μ .

3.4. $\mathcal{M}(P, s)$. Carbonara [4] gave a combinatorial reformulation of (1) that describes entries of the "inverse t-Kostka matrix" $\mathcal{M}(P, s)$ in terms of special tournament matrices. An $n \times n$ tournament matrix is a matrix B with entries in $\{0,1\}$ such that $B_{i,i} = 0$ for all i and, for all $i \neq j$, exactly one of $B_{i,j}$ and $B_{j,i}$ equals 1. Given $\lambda, \mu \in \text{Par}$ having length at most n, we say the tournament matrix B has type λ and shape μ iff the sequence $(\lambda_i + \sum_{j=1}^n B_{i,j} : 1 \leq i \leq n)$ is a rearrangement of the sequence $(\mu_i + n - i : 1 \leq i \leq n)$. Such a matrix is special (for λ) iff for all $i < j$ with $\lambda_i = \lambda_j$, $B_{ij} = 1$. Let $\Gamma_{\lambda,\mu}^*$ be the set of all $n \times n$ special tournament matrices of type λ and shape μ , where $n = \max(\ell(\lambda), \ell(\mu)).$

We define signs and weights for $B \in \Gamma_{\lambda,\mu}^*$ as follows. Since the entries of $(\mu_i + n - i : 1 \leq$ $i \leq n$) are distinct, there is a unique $w \in \mathfrak{S}_n$ such that $\lambda_i + \sum_{j=1}^n B_{i,j} = \mu_{w(i)} + n - w(i)$ for $1 \leq i \leq n$. Define sgn(B) = sgn(w), where sgn(w) is the usual sign of the permutation w. Define $wt(B) = \sum_{i>j} B_{i,j}$, which is the number of nonzero entries of B below the diagonal.

Theorem 5. [4, Theorem 2] For all $\lambda, \mu \in \text{Par}_n$, $\mathcal{M}(P, s)_{\lambda, \mu} = \sum_{B \in \Gamma_{\lambda, \mu}^*} \text{sgn}(B)(-t)^{\text{wt}(B)}$.

3.5. $\mathcal{M}(P,m)$. Macdonald [23, §III.5, p. 229] gives a formula for the monomial expansion of skew Hall-Littlewood polynomials $P_{\lambda/\nu}(x_1,\ldots,x_N;t)$ which yields $\mathcal{M}(P,m)$ by taking $\nu = (0)$. We introduce the following combinatorial model for Macdonald's formula.

Assume λ/ν is a skew shape with $N \geq \ell(\lambda)$. Given $T \in \text{SSYT}_{N}(\lambda/\nu)$, define a cell c with entry $v = T(c)$ to be special for T iff c is not in column 1 and there are no v's in the column of T just left of c's column. In this case, the weight of c is the number of cells weakly below c in the column just left of c that either have entries less than v or are part of the diagram for ν . Formally, the set of special cells for T is

$$
Sp(T) = \{(i, j) \in \lambda/\nu : j > 1 \text{ and for all } u \text{ with } (u, j - 1) \in \lambda/\nu, T(u, j - 1) \neq T(i, j)\}.
$$

The *weight* of a special cell (i, j) is

$$
\text{wt}(i,j) = |\{(u,j-1) \in \lambda/\nu : u \ge i \text{ and } T(u,j-1) < T(i,j)\}| + |\{(u,j-1) \in \nu/(0) : u \ge i\}|.
$$

Now define the set of starred semistandard tableaux

$$
SSYT_N^*(\lambda/\nu) = \{ (T, E) : T \in SSYT_N(\lambda/\nu) \text{ and } E \subseteq Sp(T) \}.
$$

A starred tableau $T^* = (T, E)$ has sign $sgn(T^*) = (-1)^{|E|}$, t-weight tstat $(T^*) = \sum_{c \in E} \text{wt}(c)$, content cont(T^{*}) = cont(T), x-weight $x^{T^*} = x^T$, and overall weight sgn(T^{*})t^{tstat(T*)} x^{T^*} .

For $T \in \text{SSYT}_N(\lambda/\nu)$, Macdonald defines $\psi_T(t) = \prod_{c \in \text{Sp}(T)} (1 - t^{\text{wt}(c)})$. Then Macdonald's monomial expansion of the skew Hall-Littlewood polynomials is

$$
P_{\lambda/\nu}(x_1,\ldots,x_N;t) = \sum_{T \in \text{SSYT}_N(\lambda/\nu)} \psi_T(t)x^T.
$$

Expanding the product in $\psi_T(t)$ using the distributive law, we get $\sum_{E \subseteq Sp(T)} \prod_{c \in E} (-t^{\text{wt}(c)})$. Comparing to the overall weight of starred tableaux, we find that

(3)
$$
P_{\lambda/\nu}(x_1,\ldots,x_N;t) = \sum_{T^* \in \text{SSYT}_N^*(\lambda/\nu)} \text{sgn}(T^*) t^{\text{tstat}(T^*)} x^{T^*}.
$$

Since $P_{\lambda/\nu}$ is known to be a symmetric polynomial, we deduce the following formula.

Theorem 6. For all $\lambda, \mu \in \text{Par}_n$, $\mathcal{M}(P,m)_{\lambda,\mu} = \sum_{T^*} \text{sgn}(T^*) t^{\text{stat}(T^*)}$ summed over all starred semistandard tableaux T^* of shape λ and content μ .

Example 7. Let $\lambda = 8654$, $\nu = 0$, $N \ge 8$, and

(4)
$$
T = \frac{\frac{1111222455}{2233668}}{\frac{33447}{55555}}
$$

In (4), the special cells are indicated by the underlined entries. Specifically,

$$
Sp(T) = \{(1,4), (1,6), (1,7), (2,5), (2,6), (3,3), (3,5)\}.
$$

These special cells have respective weights $1, 1, 1, 1, 3, 2, 1, 2$. So T contributes the term $(1 (t)^{4}(1-t^{2})^{2}(1-t^{3})x^{T}$ to P_{λ} . A typical starred tableau is $T^{*} = (T, \{(1,4), (1,6), (2,6)\})$. It can be pictured as follows:

$$
T^* = \frac{\frac{1}{2} \frac{1}{2} \frac{1}{3} \frac{1}{3} \frac{2}{4} \frac{4^*}{5} \frac{5}{5}}{\frac{3}{5} \frac{3}{5} \frac{4}{5} \frac{4}{5} \frac{7}{5}}.
$$

The overall weight of this object is $(-1)^3 t^{1+1+2} x_1^3 x_2^4 x_3^4 x_4^3 x_5^6 x_6 x_7 x_8 = -t^4 x^T$.

3.6. $\mathcal{M}(s, F)$. The fundamental quasisymmetric expansion of Schur polynomials is a sum over standard tableaux, rather than semistandard tableaux. Given $\lambda \in \text{Par}_n$ and $S \in \text{SYT}(\lambda)$, define the descent set $Des(S)$ to be the set of $k < n$ such that $k + 1$ appears in a lower row of S than k. Define the descent composition $Des'(S) = comp(Des(S))$ to be the composition associated to this subset of $[n-1]$. Gessel [10] first proved that

$$
s_{\lambda}(x_1,\ldots,x_N)=\sum_{S\in \mathrm{SYT}(\lambda)}F_{\mathrm{Des}'(S)}(x_1,\ldots,x_N).
$$

9

This formula can be proved bijectively by identifying the individual monomials in $F_{\text{Des}'(S)}$ as the content monomials of semistandard tableaux of shape λ that "standardize" to S. In terms of transition matrices, we can state Gessel's result as follows.

Theorem 8. For all $\lambda \in \text{Par}_n$ and $\alpha \in \text{Comp}_n$, $\mathcal{M}(s, F)_{\lambda, \alpha}$ is the number of standard tableaux with shape λ and descent set sub(α).

3.7. $\mathcal{M}(m, M)$. For $\lambda \in \text{Par}$, it is immediate that $m_{\lambda} = \sum_{\alpha} M_{\alpha}$ summed over all compositions α whose parts can be sorted to give the parts of λ. Therefore:

Theorem 9. For all $\lambda \in \text{Par}_n$ and $\alpha \in \text{Comp}_n$, $\mathcal{M}(m, M)_{\lambda, \alpha}$ is 1 if sort $(\alpha) = \lambda$ and 0 otherwise.

3.8. $\mathcal{M}(F,M)$ and $\mathcal{M}(M,F)$.

Theorem 10. For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(F, M)_{\alpha, \beta}$ is 1 if $\beta \succeq \alpha$ and 0 otherwise; whereas $\mathcal{M}(M,F)_{\alpha,\beta}$ is $(-1)^{\ell(\beta)-\ell(\alpha)}$ if $\beta \succeq \alpha$ and 0 otherwise.

Proof. (sketch) Using (2), one may verify that $F_{\alpha} = \sum_{\beta \succeq \alpha} M_{\beta}$, which is equivalent to the stated formula for $\mathcal{M}(F,M)_{\alpha,\beta}$. One obtains the formula for $\mathcal{M}(M,F)=\mathcal{M}(F,M)^{-1}$ using the Möbius inversion theorem on the poset of subsets of $[n-1]$ ordered by set inclusion. \Box

3.9. $\mathcal{M}(G, F)$. Let $\alpha, \beta \in \text{Comp}_n$ with β finer than α . Say $\ell(\alpha) = k$ and $\ell(\beta) = m$. By definition, there exist indices $0 = i_0 < i_1 < \cdots < i_k = m$ such that $\alpha_j = \beta_{i_{j-1}+1} + \cdots + \beta_{i_j}$ for $1 \leq j \leq k$. The *refining composition* $\text{Bre}(\beta, \alpha) = (i_1 - i_0, i_2 - i_1, \dots, i_k - i_{k-1})$ records the number of parts of β derived from each part of α . Define $s(\alpha, \beta) = \sum_{j=1}^{k} j(i_j - i_{j-1} - 1)$. Note that in the notation $\text{Bre}(\beta, \alpha)$ from [15], the finer composition is listed first, but in the function s (and g, ξ defined in §5.1), we list the finer composition second. This ordering is more convenient when working with transition matrices. In [15, Theorem 6.6], Hivert showed that for all $N \geq n$ and $\alpha \in \mathrm{Comp}_n$,

$$
G_{\alpha}(x_1,\ldots,x_N;t)=\sum_{\beta:\beta\succeq\alpha}(-1)^{\ell(\beta)-\ell(\alpha)}t^{s(\alpha,\beta)}F_{\beta}(x_1,\ldots,x_N).
$$

In other words:

Theorem 11. For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(G, F)_{\alpha, \beta}$ is $(-1)^{\ell(\beta)-\ell(\alpha)}t^{s(\alpha,\beta)}$ if $\beta \succeq \alpha$ and 0 otherwise.

Example 12. Take $\beta = 1221431211$ and $\alpha = 55314$. Then $\text{Bre}(\beta, \alpha) = 32113$ and

$$
s(\alpha, \beta) = 1 \cdot 2 + 2 \cdot 1 + 3 \cdot 0 + 4 \cdot 0 + 5 \cdot 2 = 14.
$$

So $\mathcal{M}(G,F)_{\alpha,\beta} = (-1)^5 t^{14}.$

3.10. $\mathcal{M}(G, M)$. In [15, eq. (105)], Hivert showed that for all $\alpha \in \text{Comp}_n$ and $N \geq n$,

$$
G_{\alpha}(x_1,\ldots,x_N;t) = \sum_{\beta:\beta\succeq\alpha} M_{\beta}(x_1,\ldots,x_N;t) \prod_{i=1}^{\ell(\mathrm{Bre}(\beta,\alpha))} (1-t^i)^{\mathrm{Bre}(\beta,\alpha)_i-1}.
$$

Hence:

Theorem 13. For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(G, M)_{\alpha, \beta}$ is $\prod_{i=1}^{\ell(\text{Bre}(\beta, \alpha))} (1-t^i)^{\text{Bre}(\beta, \alpha)_i-1}$ if $\beta \succeq \alpha$ and 0 otherwise.

3.11. $\mathcal{M}(\mathcal{S}, M)$. As defined in [13], $\mathcal{S}_{\alpha} = \sum_{T} x^{T}$ summed over all SSCT of shape α . So:

Theorem 14. Let $SSCT(\alpha, \beta)$ denote the set of all semistandard composition tableaux of shape α and content β . For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(\mathcal{S}, M)_{\alpha, \beta} = |\text{SSCT}(\alpha, \beta)|$.

3.12. $\mathcal{M}(\mathcal{S}, F)$. A standard composition tableau (SCT) T is a SSCT with content $(1, 1, \ldots, 1)$. The descent set $Des(T)$ is the set of those i that lie weakly to the left of $i + 1$. The descent *composition* for T is $comp(Des(T))$. For example, given

 $Des(T) = \{3, 5, 8, 9, 12, 15\}$ and $comp(Des(T)) = (3, 2, 3, 1, 3, 3, 2).$

Theorem 15. [13, Theorem 6.2] Let $SCT(\alpha, \beta)$ denote the set of all standard composition tableaux of shape α and descent composition β . For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(\mathcal{S}, F)_{\alpha,\beta} = |\text{SCT}(\alpha, \beta)|$.

Example 16. Consider the F-expansion of S_{13} . The only two SCT of shape 13 are

with descent compositions 13 and 22. So $S_{13} = F_{13} + F_{22}$.

3.13. $\mathcal{M}(K, F)$ and $\mathcal{M}(K, M)$. Stembridge [28] expressed the peak quasisymmetric functions as sums of both fundamental and monomial quasisymmetric functions.

Theorem 17. [28, Proposition 3.5] For all $\alpha \in \text{Comp}_n'$ and $\beta \in \text{Comp}_n$,

(5)
$$
\mathcal{M}(K, F)_{\alpha, \beta} = \begin{cases} 1, & \text{if sub}(\alpha) \subseteq \text{sub}(\beta) \triangle (\text{sub}(\beta) + 1); \\ 0, & \text{otherwise.} \end{cases}
$$

(6)
$$
\mathcal{M}(K,M)_{\alpha,\beta} = \begin{cases} 2^{\ell(\beta)-\ell(\alpha)}, & \text{if sub}(\alpha) \subseteq \text{sub}(\beta) \cup (\text{sub}(\beta)+1); \\ 0, & \text{otherwise.} \end{cases}
$$

Example 18.

$$
K_{31} = F_{31} + F_{22} + F_{121} + F_{112} = M_{31} + M_{22} + 2M_{211} + 2M_{121} + 2M_{112} + 4M_{1111}.
$$

4. F-expansion of Skew Hall-Littlewood Polynomials

4.1. **Expansion of** $P_{\lambda/\mu}$. Recall from §3.5 the combinatorial formula (3) for the monomial expansion of the skew Hall-Littlewood polynomials $P_{\lambda/\mu}(x_1,\ldots,x_N;t)$. This section converts this formula to an expansion of these polynomials in terms of the fundamental quasisymmetric basis. In particular, this provides a combinatorial interpretation for the entries of $\mathcal{M}(P, F)$. We remark that one can also obtain $\mathcal{M}(P, F)$ by multiplying the known matrices $\mathcal{M}(P, s)$ and $\mathcal{M}(s, F)$. However, this produces a quite complicated interpretation for the coefficients in $\mathcal{M}(P, F)$ as signed combinations of standard tableaux and special tournaments. The new interpretation developed below is much simpler.

To state our result, we need a few more definitions. Given a skew diagram λ/μ with n cells, let SYT^{*}(λ/μ) be the set of starred tableaux $S^* = (S, E)$ such that S is a standard tableau of shape λ/μ . In this case, observe that Sp(S) consists of all cells in the diagram not in column 1. So E can be an arbitrary subset of cells of λ/μ not in column 1. Define the ascent set of S^* , denoted Asc(S^*), to be the set of all $k < n$ such that either (a) $k+1$ appears in S in a lower row than k, or (b) there exist u, i, j with $S(u, j - 1) = k$, $S(i, j) = k + 1$, and $(i, j) \in E$. The second alternative says that $k + 1$ appears in a cell of E located in the next column after the column containing k. Define $\text{Asc}'(S^*) = \text{comp}(\text{Asc}(S^*))$ to be the associated composition.

11

Theorem 19. For all skew shapes λ/μ .

$$
P_{\lambda/\mu}(x_1,\ldots,x_N;t) = \sum_{S^* \in \text{SYT}^*(\lambda/\mu)} \text{sgn}(S^*) t^{\text{tstat}(S^*)} F_{\text{Asc}'(S^*)}(x_1,\ldots,x_N).
$$

In particular, for all $\lambda \in \text{Par}_n$ and $\alpha \in \text{Comp}_n$, $\mathcal{M}(P, F)_{\lambda,\alpha} = \sum_{S^*} \text{sgn}(S^*) t^{\text{tstat}(S^*)}$ summed over all starred standard tableaux S^* having shape λ and ascent set $\text{sub}(\alpha)$.

Proof. Let Y be the set of pairs (S^*, w) where $S^* = (S, E) \in SYT^*(\lambda/\mu)$ and w is an Asc (S^*) compatible word of length n (see §2.3). The overall weight of $(S^*, w) \in Y$ is $sgn(S^*)t^{tstat(S^*)}x^w$. Keeping in mind the definition of F_{α} , we see that the generating function for the weighted set Y is

(7)
$$
\sum_{y \in Y} \mathrm{wt}(y) = \sum_{S^* \in \mathrm{SYT}^*(\lambda/\mu)} \mathrm{sgn}(S^*) t^{\mathrm{tstat}(S^*)} F_{\mathrm{Asc}'(S^*)}(x_1, \dots, x_N).
$$

Comparing (3) and (7), the theorem will be proved if we can construct a sign-preserving, weight-preserving bijection $f : SSYT^*(\lambda/\mu) \to Y$.

Example 20. Let $\lambda = 65211$, and consider the starred standard tableau

S [∗] = 1 2 3 8 [∗] 9 15[∗] 4 6 7 [∗] 13[∗] 14 5 12 10 11

.

For this object, $E = \{(1,4), (1,6), (2,3), (2,4)\}\text{, } \text{Asc}(S^*) = \{3, 4, 6, 7, 9, 10, 14\}\text{, and } \text{Asc}'(S^*) =$ 31212141. A typical object in Y is (S^*, w) where

$$
w = 111 < 2 < 33 < 4 < 55 < 6 < 7888 < 9.
$$

The marked ascents in w are mandated for w to be $Asc(S^*)$ -compatible, and there is an extra ascent between $w_{11} = 7$ and $w_{12} = 8$. The overall weight of (S^*, w) is

$$
(-1)^4 t^{2+2+1+1} x_1^3 x_2 x_3^2 x_4 x_5^2 x_6 x_7 x_8^3 x_9 = t^6 x^w.
$$

Continuing the proof, we must define a "standardization map" $f : SSYT^*(\lambda/\mu) \to Y$ and an "unstandardization map" $g: Y \to \text{SSYT}^*(\lambda/\mu)$ such that f preserves signs and weights, $f \circ g = \text{id}_Y$, and $g \circ f = \text{id}_{\text{SSYT}^*(\lambda/\mu)}$. Given a semistandard tableau $T \in \text{SSYT}_N(\lambda/\mu)$, recall that there is a standard tableau stdz (T) , called the *standardization of* T, defined as follows. Suppose the entries in T consist of m_1 ones, m_2 twos, etc. Define $M_0 = 0$ and $M_i = m_1 + m_2 + \cdots + m_i$ for $1 \leq i \leq N$. We obtain stdz(T) from T by replacing the m_i occurrences of i in T, from left to right, by the integers $M_{i-1} + 1, M_{i-1} + 2, \ldots, M_i$. Now, given $T^* = (T, E) \in \text{SSYT}^*(\lambda/\mu)$, define $f(T^*) = (S^*, w) = ((\text{stdz}(T), E), w)$, where w consists of the symbols in T in increasing order.

We must check that $f(T^*) \in Y$ for all $T^* = (T, E) \in \text{SSYT}^*(\lambda/\mu)$. First, $S = \text{stdz}(T)$ is a standard tableau of shape λ/μ . Second, since $E \subseteq Sp(T)$, every cell of E is not in column 1, and so $E \subseteq Sp(S)$. Third, we claim w is an Asc (S^*) -compatible sequence. To check this, assume $k < n$ and $w_k = w_{k+1} = v$. By definition of standardization, the unique occurrences of k and $k + 1$ in S were used to relabel cells that both originally contained occurrences of v in T. Since the horizontal strip of cells in T containing v gets relabeled from left to right, $k + 1$ cannot appear in S in a lower row than k. So condition (a) in the definition of $Asc(S^*)$ does not hold for k. Can condition (b) hold for k? If so, there are u, i, j with $S(u, j - 1) = k$ and $S(i, j) = k + 1$ and $(i, j) \in E$. But then $T(u, j - 1) = v = T(i, j)$ means that (i, j) is not a

special cell for T, which contradicts $E \subseteq Sp(T)$. So we conclude that $k \notin \text{Asc}(S^*)$, proving the claim.

With notation as above, observe that $x^{T^*} = x^T = x^w$, so that f preserves the x-weight. Since applying f does not change E, f preserves signs. Finally, suppose $(i, j) \in E$ with $T(i, j) = v$. Since v cannot appear in column $j-1$ of T, it follows from the definition of standardization that the cells contributing to $wt(i, j)$ in the computation of tstat (T^*) are precisely the cells contributing to $wt(i, j)$ in the computation of tstat(S^*). So, f preserves the t-weight.

All that remains is to define the two-sided inverse g for f. Given $(S^*, w) = ((S, E), w) \in Y$, suppose w consists of m_1 ones followed by m_2 twos, etc. With M_i defined as before, let T : $\lambda/\mu \to [N]$ be obtained from S by replacing all symbols $M_{i-1} + 1, \ldots, M_i$ by i's, for $1 \le i \le N$. Then set $g(S^*, w) = (T, E)$. We must check that (T, E) lies in the codomain X. Note that w is Asc(S^*)-compatible, so $w_k = w_{k+1}$ implies that conditions (a) and (b) in the definition of ascent set are both false for this k . The falsehood of condition (a) ensures that T will be a semistandard tableau. On the other hand, the falsehood of condition (b) guarantees that every cell of E is special for T (not just special for S). So $(T, E) \in X$ as needed.

Knowing that f and g do map into their stated codomains, it is now immediate that $f \circ q = id_V$ and $g \circ f = \text{id}_{SSYT^*(\lambda/\mu)}$. (This verification does not involve the sets E or the ascent sets; one only needs the fact that the usual standardization of a semistandard tableau is reversible if the content word w is known). So the proof is complete. \Box

Example 21. Applying f to the starred semistandard tableau T^* from Example 7 gives $f(T^*)$ = (S^*, w) , where

$$
S^* = \frac{\begin{array}{|c|c|c|c|c|}\n\hline\n1 & 2 & 3 & 6 & 7 & 14 & 19 & 20 \\
\hline\n4 & 5 & 10 & 11 & 21 & 23 & \\
\hline\n8 & 9 & 12 & 13 & 22 & \\
\hline\n15 & 16 & 17 & 18 & & \\
\hline\n\end{array}
$$

and $w = 11122223333444555555678$. Observe that w is Asc(S^{*})-compatible, since Asc(S^{*}) = $\{3, 7, 11, 14, 20, 21, 22\}$. Moreover, $g(S^*, w) = T^*$.

Example 22. Applying g to the object $(S^*, w) \in Y$ from Example 20 gives the starred semistandard tableau

$$
T^* = \frac{\frac{1}{2} \frac{1}{3} \frac{1}{4} \frac{1}{8} \frac{5}{8}}{\frac{3}{6} \frac{8}{7}}.
$$

Note that, as required, T is semistandard and E does consist of special cells for T . Moreover, $f(T^*) = (S^*, w).$

Example 23. Using Theorem 19, we can make the following calculation. Each term corresponds to the starred standard tableau shown below it:

$$
P_{21}(t) = F_{21} - tF_{111} + F_{12} - t^2F_{111}.
$$

$$
\frac{1}{3} \qquad \frac{1}{3} \qquad \frac{1}{2} \qquad \frac{1}{2} \qquad \frac{1}{2} \qquad \frac{1}{3}.
$$

Remark 24 (Alternate Formula for $\mathcal{M}(P, s)$). As described in §3.4, Carbonara [4] expresses the entries of the inverse t-Kostka matrix $\mathcal{M}(P, s)$ as signed, weighted sums of special tournament matrices. An alternative description can be obtained by following $\mathcal{M}(P, F)$ by the projection from QSym to Sym given in [6]. The entry of $\mathcal{M}(P, s)_{\lambda,\mu}$ is again described as a sum of signed, 13

weighted objects. However, in this description the objects are pairs (S^*,T) where $S^* \in \text{SYT}^*(\lambda)$ and T is a "flat special rim-hook tableau" of shape μ and content Asc'(S^{*}).

In addition to working for skew Hall-Littlewood polynomials, this new description may have computational advantages. For $n = 4$, there are 37 special tournament matrices that contribute to the calculation of $\mathcal{M}(P, s)$. However, only 23 pairs (S^*, T) are now needed. We note that these pairs do not correspond to a subclass of special tournament matrices in any simple way. Carbonara's description computes the value $\mathcal{M}(P, s)_{4,22} = 0$ via the fact that there are no special tournament matrices with parameters $\lambda = 4$ and $\mu = 22$. There are two such pairs (S^*,T) , albeit of opposite sign and equal weight.

4.2. **Expansion of** $Q_{\lambda/\mu}$. We recall the definition of the skew Hall-Littlewood polynomials $Q_{\lambda/\mu}(x_1,\ldots,x_N;t)$ [23, §III.5]. For $r \in \mathbb{N}^+$, let $\phi_r(t) = (1-t)(1-t^2)\cdots(1-t^r)$. For a partition λ with $m_i(\lambda)$ parts equal to i for each i, define $b_{\lambda}(t) = \prod_{i \geq 1} \phi_{m_i(\lambda)}(t)$. For partitions $\mu \subseteq \lambda$, define $b_{\lambda/\mu}(t) = b_{\lambda}(t)/b_{\mu}(t)$. Finally, define

$$
Q_{\lambda/\mu}(x_1,\ldots,x_N;t) = b_{\lambda/\mu}(t) P_{\lambda/\mu}(x_1,\ldots,x_N;t).
$$

From this, Theorem 19 immediately gives

$$
Q_{\lambda/\mu}(x_1,\ldots,x_N;t) = \sum_{S^* \in \text{SYT}^*(\lambda/\mu)} \text{sgn}(S^*)t^{\text{tstat}(S^*)} b_{\lambda/\mu}(t) F_{\text{Asc}'(S^*)}(x_1,\ldots,x_N).
$$

On the other hand, Macdonald [23, §III.5, pp. 227–229] gives the following monomial expansion of $Q_{\lambda/\mu}$. Suppose $\nu \subseteq \rho$ are partitions such that $\theta = \rho/\nu$ is a horizontal strip. Let θ_i' be the number of cells of θ in column i, so $\theta'_{i} \in \{0,1\}$ since θ is a horizontal strip. Define

$$
\phi_{\rho/\nu}(t) = \prod_{i \in I} (1 - t^{m_i(\rho)})
$$

where $I = \{i \geq 1 : \theta'_i = 1 \text{ and } \theta'_{i+1} = 0\}$. For $T \in \text{SSYT}_N(\lambda/\mu)$, view T as a nested sequence of partitions $\mu = \lambda^{(0)} \subseteq \lambda^{(1)} \subseteq \cdots \subseteq \lambda^{(N)} = \lambda$, where $\lambda^{(i)}/\lambda^{(i-1)}$ is a horizontal strip consisting of the cells in T with entry i, and let $\phi_T(t) = \prod_{i=1}^N \phi_{\lambda^{(i)}/\lambda^{(i-1)}}(t)$. Macdonald's expansion for $Q_{\lambda/\mu}$ is

(8)
$$
Q_{\lambda/\mu}(x_1,\ldots,x_N;t) = \sum_{T \in \text{SSYT}(\lambda/\mu)} \phi_T(t)x^T.
$$

By imitating the proof we gave in $\S 4.1$, we can use (8) to derive an alternative fundamental quasisymmetric expansion for $Q_{\lambda/\mu}$ (and hence also for $P_{\lambda/\mu}$).

We merely sketch the necessary changes in the first formula and its proof. For a tableau $T \in \text{SSYT}_N(\lambda/\mu)$, define the set of Q-special cells for T by

$$
\text{QSp}(T) = \{(i,j) \in \lambda/\mu : \text{ for all } u \in \lambda/\mu, T(u,j+1) \neq T(i,j)\}.
$$

So, a cell c is Q-special for T iff the entry of T in c does not also appear in the column immediately to the right of c. The weight of a Q -special cell (i, j) is defined to be

$$
wt_Q(i,j) = m_j(\lambda^{(T(i,j))}) = i - |\{u : (u,j+1) \in \mu/(0) \text{ or } T(u,j+1) \le T(i,j)\}|.
$$

Now let $SSYT_N^{*Q}(\lambda/\mu) = \{(T, E) : T \in SSYT_N(\lambda/\mu) \text{ and } E \subseteq QSp(T)\}\$ be the set of Q starred semistandard tableaux. Since $\phi_T(t) = \prod_{c \in \text{QSp}(T)} (1 - t^{\text{wt}_Q(c)})$, it follows as before that

$$
Q_{\lambda/\mu}(x_1,\ldots,x_N;t) = \sum_{\substack{T^* \in \text{SSYT}_N^{*Q}(\lambda/\mu) \\ 14}} \text{sgn}(T^*)t^{\text{tstat}_Q(T^*)}x^{T^*}
$$

,

where $sgn(T^*) = (-1)^{|E|}$, $tstat_Q(T^*) = \sum_{c \in E} wt_Q(c)$, and $x^{T^*} = x^T$. Define Q-starred standard tableaux in the obvious way. For $S \in \text{SYT}(\lambda/\mu)$, observe that *every* cell in S is Q-special for S. Given $S^* = (S, E)$ with $E \subseteq \lambda/\mu$ and $k < n = |\lambda/\mu|$, define $k \in \text{QAsc}(S^*)$ iff either (a) $k+1$ appears in S in a lower row than k; or (b) $k \in E$ and $k + 1$ appears in S in the next column after k's column. Let $QAsc'(S^*) = \text{comp}(QAsc(S^*))$. Define Y as before, replacing Asc (S^*) by $QAsc(S^*)$. One should now check that the proof given above adapts to the present situation without difficulty. We therefore have the following result.

Theorem 25. For all skew shapes λ/μ ,

$$
Q_{\lambda/\mu}(x_1,\ldots,x_N;t) = \sum_{S^* \in \text{SYT}^{*Q}(\lambda/\mu)} \text{sgn}(S^*)t^{\text{tstate}_Q(S^*)} F_{\text{QAsc}'(S^*)}(x_1,\ldots,x_N).
$$

In particular, for all $\lambda \in \text{Par}_n$ and $\alpha \in \text{Comp}_n$, $\mathcal{M}(Q,F)_{\lambda,\alpha} = \sum_{S^*} \text{sgn}(S^*) t^{\text{stata}_Q(S^*)}$ summed over all Q-starred standard tableaux having shape λ and Q-ascent set sub(α).

Dividing through by $b_{\lambda/\mu}(t)$ gives an analogous F-expansion for $P_{\lambda/\mu}$.

5. Transition Matrices Giving G-Expansions

This section discusses combinatorial formulas for the transition matrices $\mathcal{M}(F, G)$, $\mathcal{M}(M, G)$, $\mathcal{M}(P,G)$, $\mathcal{M}(\mathcal{S},G)$ and $\mathcal{M}(K,G)$.

5.1. $\mathcal{M}(F, G)$. Let $\alpha, \beta \in \text{Comp}_n$ with β finer than α . Define $\xi_{\alpha,\beta}(j)$ to be j if β_j and β_{j+1} are formed from the same part of α and 0 otherwise. Set $g(\alpha, \beta) = \sum_{j=1}^{\ell(\beta)-1} \xi_{\alpha,\beta}(j)$.

Theorem 26. For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(F, G)_{\alpha, \beta}$ is $t^{g(\alpha, \beta)}$ if $\beta \succeq \alpha$ and 0 otherwise.

Proof. Since $\mathcal{M}(F, G)$ is the unique matrix such that $\mathcal{M}(G, F)\mathcal{M}(F, G) = I$, it is enough (by Theorem 11) to show that for all compositions $\beta \succeq \alpha$,

(9)
$$
\sum_{\gamma:\,\beta\geq\gamma\geq\alpha}(-1)^{\ell(\gamma)-\ell(\alpha)}t^{s(\alpha,\gamma)}t^{g(\gamma,\beta)}=\delta_{\alpha,\beta}.
$$

Recall that $s(\alpha, \gamma) = \sum_{j=1}^{\ell(\alpha)} j(\text{Bre}(\gamma, \alpha)_j - 1)$. If $\alpha = \beta$ then there is a single term that is readily seen to equal 1. So suppose $\beta \succ \alpha$. We define a sign-reversing involution $\gamma \mapsto \gamma'$ on the set of compositions γ with $\beta \succeq \gamma \succeq \alpha$, as follows. Let j be as small as possible such that $\text{Bre}(\beta, \gamma)_j > 1$ or Bre $(\gamma, \alpha)_j > 1$. Equivalently, this is the smallest j such that not all of α_j , β_j and γ_j are equal. We know that such a j must exist since $\alpha \neq \beta$.

If $\gamma_j = \beta_j$ (and hence $\beta_j < \alpha_j$), then set

(10)
$$
\gamma' = (\gamma_1, \dots, \gamma_{j-1}, \gamma_j + \gamma_{j+1}, \gamma_{j+2}, \dots, \gamma_{\ell(\gamma)}).
$$

Otherwise, let

(11)
$$
\gamma' = (\gamma_1, \ldots, \gamma_{j-1}, \beta_j, \gamma_j - \beta_j, \gamma_{j+1}, \ldots, \gamma_{\ell(\gamma)}).
$$

If γ' is defined according to (10), then $s(\alpha, \gamma') = s(\alpha, \gamma) - j$ and $g(\gamma', \beta) = g(\gamma, \beta) + j$. On the other hand, if γ' is defined according to (11), then $s(\alpha, \gamma') = s(\alpha, \gamma) + j$ and $g(\gamma', \beta) = g(\gamma, \beta) - j$. It follows immediately that the map $\gamma \mapsto \gamma'$ is a sign-reversing, t-weight preserving involution on the set of γ for which $\beta \succeq \gamma \succeq \alpha$. Hence, the sum in (9) is zero as desired.

Example 27. Let $\alpha = 212135$, $\gamma = 2121323$ and $\beta = 212112212$. Note that $\text{Bre}(\gamma, \alpha) = 111112$ and $\text{Bre}(\beta, \gamma) = 1111212$. It follows that $s(\alpha, \gamma) = 6 \cdot (2 - 1) = 6$ and $g(\gamma, \beta) = 5 + 8 = 13$. The smallest j for which the parts are not all equal is $j = 5$. The parts γ_5 and β_5 are not 15 equal, so γ' gets defined according to (11): $\gamma' = 21211223$. Hence $\text{Bre}(\gamma', \alpha) = 111122$ and Bre(β , γ') = 11111112. So $s(\alpha, \gamma') = 5 \cdot 1 + 6 \cdot 1 = 11$ and $g(\gamma', \beta) = 8$.

Example 28. Using Theorem 26, we calculate $F_3 = G_3 + tG_{21} + tG_{12} + t^3G_{111}$, $F_{21} = G_{21} + tG_{111}$, $F_{12} = G_{12} + t^2 G_{111}$, and $F_{111} = G_{111}$.

$$
5.2. \mathcal{M}(M, G).
$$

Theorem 29. For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(M, G)_{\alpha, \beta}$ is $(-1)^{\ell(\beta)-\ell(\alpha)} \prod_{j:\xi_{\alpha,\beta}(j)=j} (1-t^j)$ if $\beta \succeq \alpha$ and 0 otherwise.

Proof. Fix $\alpha, \beta \in \text{Comp}_n$. Since $\mathcal{M}(M, G) = \mathcal{M}(M, F) \mathcal{M}(F, G)$,

$$
\mathcal{M}(M, G)_{\alpha, \beta} = \sum_{\gamma \in \text{Comp}_n} \mathcal{M}(M, F)_{\alpha, \gamma} \mathcal{M}(F, G)_{\gamma, \beta}
$$

=
$$
\sum_{\gamma: \beta \succeq \gamma \succeq \alpha} (-1)^{\ell(\gamma) - \ell(\alpha)} t^{g(\gamma, \beta)}
$$

=
$$
(-1)^{\ell(\beta) - \ell(\alpha)} \sum_{\gamma: \beta \succeq \gamma \succeq \alpha} (-1)^{\ell(\gamma) - \ell(\beta)} t^{g(\gamma, \beta)}.
$$

Any composition γ in the above sum is completely determined by specifying the values of $\xi_{\gamma,\beta}(j)$ for those j such that $\xi_{\alpha,\beta}(j) = j$. This follows since $\beta \succeq \gamma \succeq \alpha$ and $\xi_{\alpha,\beta}(j) = 0$ force $\xi_{\gamma,\beta}(j) = 0$. On the other hand, for each j with $\xi_{\alpha,\beta}(j) = j$, we can always choose $\xi_{\gamma,\beta}(j)$ to be either 0 or j when building γ . Every time we choose to set $\xi_{\gamma,\beta}(j) = j$, we are increasing the length difference between β and γ by 1. Additionally, we are increasing the t-weight by j. Since all of these choices are independent, we can write

$$
\sum_{\gamma:\beta\geq\gamma\geq\alpha}(-1)^{\ell(\gamma)-\ell(\beta)}t^{g(\gamma,\beta)}=\prod_{j:\xi_{\alpha,\beta}(j)=j}(1-t^j).
$$

The theorem follows. \Box

Example 30. Consider $\alpha = 22$ and $\beta = 1111$. Then $\xi_{\alpha,\beta}(1) = 1$, $\xi_{\alpha,\beta}(2) = 0$ and $\xi_{\alpha,\beta}(3) = 3$. So $\mathcal{M}(M, G)_{\alpha, \beta} = (-1)^2 (1-t)(1-t^3)$.

Example 31. We calculate $M_3 = G_3 - (1-t)G_{21} - (1-t)G_{12} + (1-t)(1-t^2)G_{111}$, $M_{21} =$ $G_{21} - (1 - t)G_{111}$, $M_{12} = G_{12} - (1 - t^2)G_{111}$, and $M_{111} = G_{111}$.

5.3.
$$
\mathcal{M}(P, G)
$$
. By multiplying $\mathcal{M}(P, F)$ and $\mathcal{M}(F, G)$, we obtain the formula

(12)
$$
\mathcal{M}(P,G)_{\lambda,\beta} = \sum_{\substack{S^*=(S,E)\in \text{SYT}^*(\lambda)\\ \text{Asc}'(S^*)\preceq\beta}} (-1)^{|E|} t^{\text{tstat}(S^*)+g(\text{Asc}'(S^*),\beta)}.
$$

As described in Theorem 32 below, the contribution of each standard Young tableau S to (12) can be expressed as a product. One advantage of this latter form is that a frequent group of cancellations can be identified; see Corollary 33. In order to present these results, we introduce some new notation.

For $S \in \text{SYT}(\lambda)$, define $\text{Sp}(S)$ and wt(c) as in §4.1. We define the following subset of $\text{Sp}(S)$:

$$
Esp(S) = \{c \in Sp(S) : \text{Asc}(S, \{c\}) \neq \text{Asc}(S, \emptyset)\}.
$$

So, $c \in \text{Esp}(S)$ if and only if $S(c)-1$ appears in the column of S just left of the column containing $S(c)$ and $S(c)$ −1 appears weakly lower than $S(c)$ does. The subset Esp(S) keeps track of which cells actually affect Asc' (S^*) for starred tableaux with underlying tableau S. As we are only concerned here with standard tableaux S, we will let $c_j = c_j(S)$ denote the cell of S in which

j appears. Finally, note that the descent set of S, Des(S), is contained in Asc(S^{*}) for any S^* with underlying tableau S.

Our intent is to derive a simplified version of (12) in which the main sum extends over standard tableaux $S \in \text{SYT}(\lambda)$ rather than starred standard tableaux. To obtain $S^* = (S, E)$ from S, we will build the subset E by choosing to include or exclude various cells $c \in Sp(S)$. The final object S^* is required to satisfy $\text{Asc}'(S^*) \leq \beta$, or equivalently, $\text{Asc}(S^*) \subseteq \text{sub}(\beta)$. By the remark at the end of the last paragraph, this requirement will be met only if $Des(S) \subseteq sub(\beta)$. So we need only consider standard tableau S satisfying this condition.

The choices of which $c = c_{j+1}(S) \in Sp(S)$ to include in E can be made independently. We consider each such c according to whether $j \in sub(\beta)$ and whether $c \in Esp(S)$. We first note that if $j \notin \text{sub}(\beta)$ and $c \in \text{Esp}(S)$, then inclusion of c in E would cause Asc' $(S, E) \npreceq \beta$. Hence, this case need not be considered further. For each of the remaining possibilities for j and c , we consider the net effect on the signed t-weight caused by the decision to include or exclude c from E. For each $j \in sub(\beta)$, let $m_j = m_j(\beta)$ be the number of elements in sub(β) that are at most j.

- (1) $j \notin \text{sub}(\beta)$ and $c \notin \text{Esp}(S)$. If c is included in E, the t-weight contribution will be $-t^{\text{wt}(c)}$; otherwise the contribution will be 1.
- (2) $j \in sub(\beta)$ and $c \in Esp(S)$. If c is included in E, the t-weight contribution will be $-t^{\text{wt}(c)}$; otherwise it will be t^{m_j} .
- (3) $j \in sub(\beta)$ and $c \notin Esp(S)$. Since $c \notin Esp(S)$, the value of $\xi_{Asc'(S^*),\beta}(j)$ depends only on whether or not $j \in Des(S)$. Hence, there is a corresponding t-weight contribution of t^{m_j} if and only if $j \notin \text{Des}(S)$. As in Case 1, there is an additional contribution to the t-weight (coming from the tstat function) of $-t^{wt(c)}$ or 1 according to whether or not c is in E .

Observing that the factor $1-t^{wt(c)}$ occurs in both Case 1 and Case 3, we reorganize the cases slightly to obtain the following.

Theorem 32. For all $\lambda \in \mathrm{Par}_n$ and $\beta \in \mathrm{Comp}_n$,

(13)
$$
\mathcal{M}(P,G)_{\lambda,\beta} = \sum_{\substack{S \in \text{SYT}(\lambda) \\ \text{Des}(S) \subseteq \text{sub}(\beta) \\ c_{j+1} \in \text{Esp}(S)}} \prod_{\substack{f \in \text{sub}(\beta) : \\ (f^{m_j} - t^{\text{wt}(c_{j+1})})}} (t^{m_j} - t^{\text{wt}(c_{j+1})}) \prod_{\substack{j \in [n-1] : \\ c_{j+1} \in \text{Sp}(S) \setminus \text{Esp}(S)}} t^{m'_j} (1 - t^{\text{wt}(c_{j+1})}),
$$

where $m'_j = m_j$ if $j \in sub(\beta) \setminus Des(S)$ and 0 otherwise.

Corollary 33. If $m_j = \text{wt}(c_{j+1})$ for some $j \in \text{sub}(\beta)$ with $c_{j+1} \in \text{Esp}(S)$, then S can be omitted from the sum in (13).

Example 34. Let $\lambda = 32$ and $\beta = 1211$. Note that $sub(\beta) = \{1, 3, 4\}$ and so $m_1 = 1$, $m_3 = 2$, and $m_4 = 3$. In Table 1 we list the five elements of SYT(32) (referred to from left to right as S_1, \ldots, S_5 along with pertinent data. The row labeled \prod_1 (resp. \prod_2) gives the contributions from the first (resp. second) product in (13). Since $\text{Des}(S_2), \text{Des}(S_3) \not\subseteq \text{sub}(\beta), \prod_1 \text{ and } \prod_2 \text{ have}$ been left blank for these two tableaux. (For reference, the corresponding products for these tableaux are $(t-t) \cdot t^2(1-t)(1-t)$ and $(t-t)(t^2-t)(t^3-t^2) \cdot 1$, respectively.) Note that Corollary 33 applies to S_1 with $j = m_j = 1$. So the only contributions are from the last two columns and we find that

$$
\mathcal{M}(P,G)_{32,1211} = (t^2 - t)(1 - t) + (t^3 - t^2)(1 - t) = -t^4 + t^3 + t^2 - t.
$$

17

	5	3	$\ddot{ }$	٠Î 4 5 $\mathcal{D}_{\mathcal{L}}$	Ð റ 4
Des(S)	$\{3\}$	$\{2,4\}$	$2\}$	$\{1,4\}$	$\{1,3\}$
Sp(S)	$\{c_2,c_3,c_5\}$	$\{c_2, c_4, c_5\}$	$\{c_2, c_4, c_5\}$	$\{c_3, c_4, c_5\}$	$\{c_3, c_4, c_5\}$
$\mathrm{Esp}(S)$	$\{c_2, c_3, c_5\}$	$\{c_2\}$	$\{c_2, c_4, c_5\}$	$\{c_3, c_4\}$	$\{c_3, c_5\}$
\prod_1	$(t-t)(t^3-t)$			τ ²	t^3 . $-t^2$
\mathcal{L}_{2}					$-t)$

5.4. $\mathcal{M}(\mathcal{S}, G)$. For an SSCT T, define the *ascent set*, denoted Asc(T), to be the set of values i for which the leftmost occurrence of i appears strictly to the right of the rightmost occurrence of $i + 1$. If Asc $(T) = \{a_1, a_2, \ldots\}$, we let $SumAsc(T) = a_1 + a_2 + \cdots$. For example, the SSCT

$$
T = \frac{\begin{array}{|c|c|c|c|c|}\n\hline\n2 & 2 & 2 & 1 & 1 \\
\hline\n4 & 3 & & & \\
\hline\n7 & 5 & 5 & 5 & 3 & 3 & 1 \\
\hline\n8 & 8 & 6 & 6 & & \\
\hline\n\end{array}
$$

has ascent set $Asc(T) = \{1, 3, 6\}$ and $SumAsc(T) = 1 + 3 + 6 = 10$.

Theorem 35. For all $\alpha, \beta \in \text{Comp}_n$, $\mathcal{M}(\mathcal{S}, G)_{\alpha, \beta} = \sum_{T \in \text{SSCT}(\alpha, \beta)} t^{\text{SumAsc}(T)}$.

Proof. Fix $\alpha, \beta \in \text{Comp}_n$. We proceed to justify the following string of equalities:

$$
\mathcal{M}(\mathcal{S}, G)_{\alpha, \beta} = \sum_{\gamma \in \text{Comp}_n} \mathcal{M}(\mathcal{S}, F)_{\alpha, \gamma} \mathcal{M}(F, G)_{\gamma, \beta}
$$

$$
= \sum_{\gamma: \gamma \preceq \beta} |\text{SCT}(\alpha, \gamma)| t^{g(\gamma, \beta)}
$$

$$
= \sum_{T' \in \text{SSCT}(\alpha, \beta)} t^{\text{SumAsc}(T')}.
$$

The first equality follows from the fact that $\mathcal{M}(\mathcal{S}, G) = \mathcal{M}(\mathcal{S}, F)\mathcal{M}(F, G)$. The second follows from Theorems (15) and (26). To prove the third, we construct a bijection

$$
f = f_{\alpha,\beta} : \bigcup_{\gamma:\gamma\preceq\beta} \mathrm{SCT}(\alpha,\gamma) \to \mathrm{SSCT}(\alpha,\beta)
$$

such that for all $T \in \mathrm{SCT}(\alpha, \gamma)$ with $\gamma \preceq \beta$, SumAsc $(f(T)) = g(\gamma, \beta)$. Fix T in the domain of f, say $T \in \text{SCT}(\alpha, \gamma)$ with $\gamma \leq \beta$. For $0 \leq i \leq \ell(\gamma)$, let $d_i = \gamma_1 + \cdots + \gamma_i$, so $\{d_0, \ldots, d_{\ell(\gamma)}\}$ $Des(T) \cup \{0,n\}$. Then for $1 \leq i \leq \ell(\gamma)$, the entries $d_{i-1}+1,\ldots,d_i$, when read from right to left in T, appear in increasing order. Let A_i be the set of cells occupied by these values. Since β is finer than γ , there exist indices $0 = i_0 < i_1 < \cdots < i_{\ell(\gamma)} = \ell(\beta)$ such that $\gamma_j = \beta_{i_{j-1}+1} + \cdots + \beta_{i_j}$ for all j. Construct $T' = f(T)$ by labeling the cells in each set A_j from right to left by the sequence

$$
\underbrace{(i_{j-1}+1)\cdots(i_{j-1}+1)}_{\beta_{i_{j-1}+1}}\underbrace{(i_{j-1}+2)\cdots(i_{j-1}+2)}_{\beta_{i_{j-1}+2}}\cdots\underbrace{i_{j}\cdots i_{j}}_{\beta_{i_{j}}};
$$

A simple check confirms that T' satisfies (C1), (C2), and (C3) in the definition of an SSCT and has content β , so f does map into the codomain SSCT(α , β). Furthermore, note that the ascents in T' occur only inside the blocks A_j . Each such place where the value changes from k to $k+1$

corresponds to the inclusion of k in $Asc(T')$ and hence the contribution of k to SumAsc (T') . Furthermore, such a change only happens when the parts β_k and β_{k+1} belong to the same part of γ . In this case, we have $\xi_{\gamma,\beta}(k) = k$ and hence there will be a contribution of k to $g(\gamma,\beta)$. Thus SumAsc(T') = $g(\gamma, \beta)$. Since this relabeling is reversible, T can be reconstructed from $T' = f(T)$, so f is a bijection. See Example 36 for an illustration of the bijection.

Example 36. Let $\alpha = 5274$, $\beta = 33313212$, $\gamma = 64332$, and

$$
T = \frac{\frac{6}{10} \frac{5}{9}}{\frac{16}{18} \frac{13}{12} \frac{11}{18} \frac{8}{7} \frac{7}{1}}.
$$

First note that $g(\gamma, \beta) = 1 + 3 + 6 = 10$, since the i for which β_i and β_{i+1} came from the same part of γ are precisely 1, 3 and 6. In order to construct T', note that the first part of γ is a 6, which is refined by the first two parts of β (both equal to 3). Thus, we first need to look at the horizontal strip of cells containing the letters $1, 2, 3, 4, 5, 6$ and change these letters to 1, 1, 1, 2, 2, 2, respectively. The other conversions, based on the parts of β that fit into each part of γ , are as follows:

$$
7, 8, 9, 10 \mapsto 3, 3, 3, 4
$$

$$
11, 12, 13 \mapsto 5, 5, 5
$$

$$
14, 15, 16 \mapsto 6, 6, 7
$$

$$
17, 18 \mapsto 8, 8.
$$

Thus, the resulting T' with content β is

$$
T' = \frac{\begin{array}{|c|c|c|c|c|} \hline 2 & 2 & 2 & 1 & 1 \\\hline 4 & 3 & & & \\ \hline 7 & 5 & 5 & 5 & 3 & 3 & 1 \\\hline 8 & 8 & 6 & 6 & & \\ \hline \end{array}}.
$$

Note that $\text{Asc}(T') = \{1, 3, 6\}$, so $\text{SumAsc}(T') = 1 + 3 + 6 = 10 = g(\gamma, \beta)$.

Example 37. Using Theorem 35, the G-expansion of $S_{13}(t)$ is

$$
G_{13} + G_{22} + t^2 G_{211} + t^2 G_{121} + t G_{112} + t^2 G_{112} + t^4 G_{1111} + t^5 G_{1111}.
$$
\n
$$
\boxed{1}
$$
\n
$$
\boxed{1}
$$
\n
$$
\boxed{1}
$$
\n
$$
\boxed{1}
$$
\n
$$
\boxed{2}
$$
\n
$$
\boxed{2}
$$
\n
$$
\boxed{1}
$$
\n
$$
\boxed{3}
$$
\n
$$
\boxed{2}
$$
\n
$$
\boxed{1}
$$
\n
$$
\boxed{3}
$$
\n
$$
\boxed{4}
$$
\n
$$
\boxed{3}
$$
\n
$$
\boxed{4}
$$

5.5. M(K, G). Consider compositions $\alpha \in \text{Comp}'_n$ and $\beta \in \text{Comp}_n$ such that $\text{sub}(\alpha) \subseteq$ $\text{sub}(\beta) \cup (\text{sub}(\beta) + 1)$. Let $\text{sub}(\alpha) = \{a_1 < \cdots < a_m\}$ and $\text{sub}(\beta) = \{b_1 < \cdots < b_p\}$. For $A \subseteq [n-1]$, let $A-1$ denote the set obtained by subtracting one from every element in A and removing 0 if it appears. Define the polynomial $k(\alpha, \beta)$ as follows. For all $b_i \in sub(\beta)$, let

(14)
$$
k(b_i) = \begin{cases} 1+t^i, & \text{if } b_i \notin \text{sub}(\alpha) \cup (\text{sub}(\alpha)-1); \\ t^{i-1}+t^i, & \text{if } b_i \in \text{sub}(\alpha) \text{ and } b_i-1=b_{i-1}; \\ 1, & \text{otherwise.} \end{cases}
$$

Finally, define $k(\alpha, \beta) = \prod_{i=1}^{p} k(b_i)$.

Example 38. If $\alpha = 4253$ and $\beta = 12213221$, we have sub(α) = {4,6,11}, sub(α) – 1 = $\{3, 5, 10\}$, and $\text{sub}(\beta) = \{1, 3, 5, 6, 9, 11, 13\}$. Thus, $k(\alpha, \beta) = (1 + t) \cdot (t^3 + t^4) \cdot (1 + t^5) \cdot (1 + t^7)$.

Here is a graphical way of looking at this example. Depict a composition $\alpha = (\alpha_1, \ldots, \alpha_m)$ of n on the interval $[0, n]$ by highlighting the points in sub(α). As shown in Figure 2, highlight the 19

points in [0, n] between an element $a-1 \in sub(\alpha)-1$ and $a \in sub(\alpha)$. No two of the highlighted subintervals share an endpoint, because of the assumption that $\alpha \in \text{Comp}'_n$. Now, the three possible cases in (14) turn into the following:

- (1) b_i does not touch any of the highlighted intervals (e.g., $b_1 = 1$, $b_5 = 9$, and $b_7 = 13$);
- (2) b_{i-1} and b_i are the two ends of a highlighted interval (e.g., $b_3 = 5$ and $b_4 = 6$); or
- (3) b_i is an end of a highlighted interval, but b_{i-1} is not at the other end of the highlighted interval (e.g., $b_2 = 3$ and $b_6 = 11$).

$\mathrm{sub}(\alpha)$					14
$\mathrm{sub}(\alpha) - 1$ o		3	h,	10	14
		3 4	-6 5	10 -11	14
$sub(\beta)$	0	3	6 5	9 11	13 14
	$1+t$		$t^3 + t^4$	$1 + t^5$	$+ t^7$

FIGURE 2. For $sub(\alpha) = \{4, 6, 11\}$ and $sub(\beta) = \{1, 3, 5, 6, 9, 11, 13\}$, we have $k(\alpha, \beta) = (1+t) \cdot (t^3 + t^4) \cdot (1+t^5) \cdot (1+t^7).$

Theorem 39. For $\alpha \in \text{Comp}_n'$ and $\beta \in \text{Comp}_n$, $\mathcal{M}(K, G)_{\alpha, \beta}$ is $k(\alpha, \beta)$ if $\text{sub}(\alpha) \subseteq \text{sub}(\beta) \cup$ $(sub(\beta) + 1)$ and 0 otherwise.

Proof. Fix $\alpha \in \text{Comp}_n'$ and $\beta \in \text{Comp}_n$. Since $\mathcal{M}(K, G) = \mathcal{M}(K, F)\mathcal{M}(F, G)$,

$$
\mathcal{M}(K, G)_{\alpha, \beta} = \sum_{\gamma \in \text{Comp}_n} \mathcal{M}(K, F)_{\alpha, \gamma} \mathcal{M}(F, G)_{\gamma, \beta}
$$

$$
= \sum_{\gamma : \text{ sub}(\alpha) \subseteq \text{sub}(\gamma) \triangle (\text{sub}(\gamma) + 1)} t^{g(\gamma, \beta)}.
$$

We now describe a sequence of choices that will construct exactly the compositions γ indexing the sum above. By examining how each choice contributes to $g(\gamma, \beta) = \sum_j \xi_{\gamma, \beta}(j)$ in the power of t, we will obtain the product $k(\alpha, \beta)$. For convenience of notation, let $A = sub(\alpha) =$ ${a_1 < \cdots < a_m}$ and $B = sub(\beta) = {b_1 < \cdots < b_l}$ as above. Rather than constructing γ itself, we make choices concerning the members of the set $C = sub(\gamma)$. The requirements $\operatorname{sub}(\alpha) \subseteq \operatorname{sub}(\gamma) \triangle (\operatorname{sub}(\gamma) + 1)$ and $\beta \succeq \gamma$ translate into the requirements $A \subseteq C \triangle (C + 1)$ and $C \subseteq B$. Since we must have $C \subseteq B$, we need only decide, for each $b_i \in B$, whether to include or exclude b_i from the set C. When making these decisions, it is required that each $a_j \in A$ belong to exactly one of the sets C or $(C+1)$ being built. Equivalently, for each $a_j \in A$, exactly one of the numbers a_j or $a_j - 1$ must be chosen for inclusion in C. Since $C \subseteq B$, this requirement on A will only be possible if $A \subseteq B \cup (B+1)$. Thus if $A \nsubseteq B \cup (B+1)$, we must have $\mathcal{M}(K, G)_{\alpha, \beta} = 0$. From now on, assume $A \subseteq B \cup (B + 1)$. As in (14) and Example 38, there are three cases to consider.

• Case 1: $b_i \notin A \cup (A-1)$ (i.e., b_i does not touch any of the highlighted intervals in the picture). In this case, we may freely choose to include b_i as a member of C or not, which will not affect the validity of the condition $A \subseteq C \bigtriangleup (C + 1)$. Including b_i in $C = sub(\gamma)$ will cause β_i and β_{i+1} to come from different parts of γ , so that $\xi_{\gamma,\beta}(i) = 0$ for this choice. Excluding b_i from C will cause β_i and β_{i+1} to come from the same part of γ ,

so that $\xi_{\gamma,\beta}(i) = i$ for this choice. The net contribution to $t^{g(\gamma,\beta)}$ based on the choice involving this b_i is therefore $(1 + t^i) = k(b_i)$.

- Case 2: $b_i \in A$ and $b_{i-1} = b_i 1$ (i.e., b_{i-1} and b_i are the two ends of a highlighted interval in the picture). In this case, note that b_i equals some a_j , and $b_{i-1} = a_j - 1$, so the requirement on A forces us to include exactly one of b_{i-1} or b_i in C. If we decide that $b_{i-1} \notin C$ and $b_i \in C$, we see (as in Case 1) that $\xi_{\gamma,\beta}(i-1) = i-1$ and $\xi_{\gamma,\beta}(i) = 0$. If, instead, we decide that $b_{i-1} \in C$ and $b_i \notin C$, we have $\xi_{\gamma,\beta}(i-1) = 0$ and $\xi_{\gamma,\beta}(i) = i$. The net contribution to $t^{g(\gamma,\beta)}$ based on the choice involving both b_{i-1} and b_i is therefore $(t^{i-1} + t^i) = k(b_i).$
- Case 3: The first two cases do not occur (i.e., b_i is one end of a highlighted interval in the picture, but b_{i-1} is not at the other end of this interval). Consider several subcases. First, suppose $b_i = a_j - 1$ for some $a_j \in A$, and $a_j \notin B$ (i.e., b_i is the left end of an interval, and the right end of this interval is not in B). The requirement on A forces us to include b_i in C, giving $\xi_{\gamma,\beta}(i) = 0$ for this i and a contribution of $k(b_i) = 1$ to $t^{g(\gamma,\beta)}$. Second, suppose $b_i = a_j - 1$ for some $a_j \in A$, and $a_j = b_{i+1}$ is in B as well (i.e., b_i is the left end of an interval, and the right end $b_i + 1 = b_{i+1}$ is also in B). Then we will decide the membership of b_i in C as part of the Case 2 analysis for index $i+1$. Since the contribution of this b_i to $t^{g(\gamma,\beta)}$ has already been accounted for elsewhere, we multiply by $k(b_i) = 1$ in this subcase. Third, suppose $b_i = a_j$ for some $a_j \in A$, and $a_j - 1$ is not in B (i.e., b_i is the right end of an interval whose left end is not in B). The requirement on A forces us to include b_i in C, giving a contribution of $k(b_i) = 1$ to $t^{g(\gamma,\beta)}$.

Combining all the choices using the product rule for weighted sets, we see that

$$
\mathcal{M}(K, G)_{\alpha, \beta} = \sum_{C: A \subseteq C \triangle (C+1), C \subseteq B} t^{g(\text{comp}(C), \text{comp}(B))} = \prod_{i=1}^{p} k(b_i) = k(\alpha, \beta)
$$

when $A \subseteq B \cup (B + 1)$.

Example 40.

$$
K_{22} = G_{22} + G_{13} + (1+t^2)G_{211} + (1+t^2)G_{121} + (t+t^2)G_{112} + (t+t^2)(1+t^3)G_{1111}.
$$

6. Transformed Hall-Littlewood Polynomials

We conclude the paper with a brief discussion of transition matrices involving plethystically transformed Hall-Littlewood polynomials.

6.1. **Definition of** Q'_λ . For any $\lambda \in \text{Par}$, recall that $Q_\lambda(x;t)$ is the Hall-Littlewood polynomial obtained from $P_{\lambda}(x;t)$ by the formula $Q_{\lambda}(x;t) = b_{\lambda}(t)P_{\lambda}(x;t)$ (see §4.2). Define the transformed Hall-Littlewood symmetric function by setting

$$
Q'_{\lambda}(x;t) = Q_{\lambda}\left[\frac{x}{1-t};t\right] = Q_{\lambda}(x_i t^j : i \ge 1, j \ge 0);
$$

in other words, we replace variables x_1, x_2, \ldots in Q_λ by the monomials $x_i t^j$ for all $i \geq 1$ and $j \geq 0$. We can also define Q'_{λ} using plethystic substitution. Recall that the power-sums p_k are algebraically independent generators of the algebra Sym. Then $Q'_{\lambda} = \phi(Q_{\lambda})$, where ϕ is the unique F-algebra homomorphism on Sym sending each p_k to $p_k/(1-t^k)$. More information on the transformed Hall-Littlewood polynomials can be found in Macdonald's book [23] and in the papers [5, 8, 16, 31].

6.2. The Transformed Macdonald Polynomials H_{μ} . One can also obtain Q'_{λ} as the $q =$ 0 specialization of the transformed Macdonald polynomials $H_{\lambda}(x;q,t)$. For a partition $\lambda =$ $(\lambda_1, \lambda_2, \ldots, \lambda_k)$, set $n(\lambda) = \sum_{i=0}^k (i-1)\lambda_i$. Haglund [11] found an explicit combinatorial formula for the fundamental quasisymmetric expansion of the modified Macdonald polynomials $\tilde{H}_{\lambda}(x;q,t) = t^{n(\lambda)}H_{\lambda}(x;q,1/t)$, which was proved by Haglund, Haiman, and Loehr in [12]. This formula is readily adjusted to yield a formula for H_{λ} of the form

$$
H_{\lambda}(x;q,t) = \sum_{w \in S_n} t^{\text{comaj}_{\lambda}(w)} q^{\text{inv}_{\lambda}(w)} F_{\text{comp(IDEs}(w))},
$$

where $\lambda \in \text{Par}_n$, IDes (w) is the sum of all $i < n$ such that $i + 1$ appears to the left of i in $w_1 \cdots w_n$, and comally and inv_{λ} are certain permutation statistics depending on λ .

We define the statistics $\text{comaj}_{\lambda}(w)$ and $\text{inv}_{\lambda}(w)$ using the specific example $n = 9, \lambda = (4, 3, 2),$ and $w = 428165793$. Fill the cells in the diagram of λ with the letters in w, working from the shortest row up to the longest row, filling each row from left to right, as shown here:

This filled diagram determines a list of *column words*, obtained by reading the entries in each column from bottom to top. For a word $v = v_1 \cdots v_k$, let comaj(v) be the sum of all $i < k$ such that $v_i < v_{i+1}$. Let $\text{comaj}_{\lambda}(w)$ be the sum of $\text{comaj}(v)$ over all column words v in the filled diagram for w. In our example,

 $\text{comaj}_{\lambda}(w) = \text{comaj}(485) + \text{comaj}(217) + \text{comaj}(69) + \text{comaj}(3) = 1 + 2 + 1 + 0 = 4.$

Next, $inv_{\lambda}(w)$ counts the *inversion triples* in the filled diagram for w, which are defined as follows. Consider three cells in the diagram positioned as shown here:

.

So b is directly above a and c occurs somewhere to the right of a in the same row. We also allow a and c to be in the top row of the diagram, in which case we take $b = \infty$. This triple of cells is an inversion triple iff $a < b < c$ or $b < c < a$ or $c < a < b$. In our example, the inversion triples (a, b, c) are $(5, \infty, 3)$, $(7, \infty, 3)$, $(9, \infty, 3)$, $(8, 5, 6)$, and $(4, 8, 2)$, so $inv_{\lambda}(w) = 5$. Since IDes $(w) = \{1, 3, 5, 7\}$, w contributes a monomial t^4q^5 to the coefficient of F_{12222} in $H_{432}(x; q, t)$.

Theorem 41 ([12]). For all $\lambda \in \mathrm{Par}_n$ and $\alpha \in \mathrm{Comp}_n$, $\mathcal{M}(H, F)_{\lambda,\alpha}$ is $\sum_w t^{\mathrm{comaj}_{\lambda}(w)}q^{\mathrm{inv}_{\lambda}(w)}$ summed over all $w \in S_n$ with $\text{IDes}(w) = \text{sub}(\alpha)$.

6.3. $\mathcal{M}(Q', F)$. The transformed Hall-Littlewood polynomials are obtained from the transformed Macdonald polynomials by setting $q = 0$, i.e., $Q'_{\lambda}(x;t) = H_{\lambda}(x;0,t)$. Using Theorem 41, we therefore obtain the following combinatorial description of the fundamental quasisymmetric expansion of Q' .

Theorem 42. For all $\lambda \in \text{Par}_{n}$ and $\alpha \in \text{Comp}_{n}$, $\mathcal{M}(Q', F)_{\lambda,\alpha}$ is $\sum_{w} t^{\text{comaj}_{\lambda}(w)}$ summed over all $w \in S_n$ with $\text{IDes}(w) = \text{sub}(\alpha)$ and $\text{inv}_{\lambda}(w) = 0$.

On the other hand, it follows from [23, p. 241] that $\mathcal{M}(Q', s) = M(s, P)^T$, the transpose of the t-Kostka matrix. Accordingly, $\mathcal{M}(Q', F) = \mathcal{M}(s, P)^T \mathcal{M}(s, F)$. Carrying out the matrix multiplication and using the interpretations of the latter matrices from §3.3 and §3.6, we obtain a second combinatorial interpretation of $\mathcal{M}(Q',F)$:

Theorem 43. For all $\lambda \in \text{Par}_n$ and $\alpha \in \text{Comp}_n$, $\mathcal{M}(Q', F)_{\lambda,\alpha}$ is $\sum_{S,T} t^{\text{chg}(T)}$ summed over all pairs (S,T) where S is a standard tableau with $Des(S) = sub(\alpha)$, T is a semistandard tableau with content λ , and S and T have the same shape $\mu \in \text{Par}_n$.

The objects appearing in this theorem may remind the reader of the Robinson-Schensted-Knuth (RSK) algorithm. Indeed, the connection between Theorem 42 and Theorem 43 is explained in [12, §7] by a careful analysis involving the RSK algorithm.

6.4. Open Problems. By matrix multiplication, we obtain a formula for $\mathcal{M}(Q',G)_{\lambda,\alpha} =$ $\sum_{\beta \in \text{Comp}_n} \mathcal{M}(Q', F)_{\lambda, \beta} \mathcal{M}(F, G)_{\beta, \alpha}$. We leave it as an open problem to find a more direct combinatorial interpretation of this transition matrix.

Another open question is to define an appropriate notion of the "plethystically transformed" Hivert quasisymmetric function $G'_{\alpha}(x;t)$, which should play a role in QSym analogous to the role of Q'_{λ} in Sym. It would be interesting to study the combinatorics of transition matrices to and from such a basis.

Finally, one could develop transition matrices between Hivert's H-basis (see Remark 1) and other known bases of NSym, for example, those constructed in [1, 2, 9, 13, 29].

7. Appendix: Examples of Transition Matrices

This appendix lists specific examples of transition matrices (old and new) discussed in this paper. In each case, we give the relevant matrix for $n = 4$ (many computed using SAGE [26, 32]).

$$
\mathcal{M}(s,m) = \begin{pmatrix}\n4 & 31 & 22 & 211 & 1111 \\
31 & 1 & 1 & 1 & 1 \\
0 & 1 & 1 & 2 & 3 \\
0 & 0 & 1 & 1 & 2\n\end{pmatrix}.
$$
\n
$$
\mathcal{M}(s,m) = \begin{pmatrix}\n4 & 31 & 22 & 211 & 1111 \\
0 & 0 & 0 & 1 & 3 \\
1111 & 0 & 0 & 0 & 1\n\end{pmatrix}.
$$
\n
$$
\mathcal{M}(m,s) = \begin{pmatrix}\n4 & 31 & 22 & 211 & 1111 \\
22 & 0 & 1 & -1 & 1 \\
211 & 0 & 0 & 0 & 1 & -3 \\
0 & 0 & 1 & -1 & 1\n\end{pmatrix}.
$$
\n
$$
\mathcal{M}(s,P) = \begin{pmatrix}\n4 & 31 & 22 & 211 & 1111 \\
21 & 0 & 0 & 0 & 1 & -3 \\
1111 & 0 & 0 & 0 & 0 & 1\n\end{pmatrix}.
$$
\n
$$
\mathcal{M}(s,P) = \begin{pmatrix}\n4 & 31 & 22 & 211 & 1111 \\
0 & 1 & t & t+t^2 & t^3 + t^4 + t^5 \\
211 & 0 & 0 & 1 & t & t^2 + t^4 \\
0 & 0 & 0 & 1 & t + t^2 + t^3 \\
1111 & 0 & 0 & 0 & 0 & 1\n\end{pmatrix} = \mathcal{M}(Q',s)^T.
$$
\n
$$
\mathcal{M}(P,s) = \begin{pmatrix}\n4 & 31 & 22 & 211 & 1111 \\
0 & 1 & -t & -t & t^2 + t^3 \\
0 & 1 & -t & -t & t^2 + t^3 \\
0 & 0 & 1 & -t & t^3 \\
1111 & 0 & 0 & 0 & 1\n\end{pmatrix} = \mathcal{M}(s,Q')^T.
$$

.

$$
\begin{array}{cccccccccccc} 4 & 31 & 22 & 211 & 13 & 121 & 112 & 1111 \\ 3 & 1 & 1 & 1 & -t & 1 & -t & 0 & 1-t & 0 & (1-t)^3 \\ 31 & 0 & 1 & 0 & 1-t & 0 & 1-t & 0 & (1-t)^3 \\ 22 & 0 & 0 & 1 & 1-t^2 & 0 & 0 & 1-t & (1-t)(1-t^2) \\ 23 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1-t & (1-t)^2 - t^2 \\ 121 & 0 & 0 & 0 & 0 & 1 & 1-t^2 & 1-t^2 & (1-t^2)^2 \\ 122 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1-t^2 \\ 123 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1-t^3 \\ 124 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1-t^3 \\ 125 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1-t^3 \\ 4 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 31 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 4 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 31 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 22 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 1 \\ 23 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 14 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 111 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 31 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 31 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 32 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 33 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 34 & 0 & 0 &
$$

 $\mathcal{M}(Q,F)=D\cdot\mathcal{M}(P,F)$ where $D_{5\times 5}$ has diagonal entries

 $1-t$, $(1-t)^2$, $(1-t)(1-t^2)$, $(1-t)^2(1-t^2)$, $(1-t)(1-t^2)(1-t^3)(1-t^4)$.

25

$$
\mathcal{M}(F,G) = \begin{pmatrix} 4 & 31 & 22 & 211 & 13 & 121 & 112 & 1111 \\ 4 & t & t & t^3 & t & t^3 & t^3 & t^6 \\ 31 & 0 & 1 & 0 & t & 0 & t & 0 & t^3 \\ 22 & 0 & 0 & 1 & t^2 & 0 & 0 & t & t^4 \\ 0 & 0 & 1 & 0 & 0 & 0 & t & t^4 \\ 13 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & t \\ 121 & 0 & 0 & 0 & 0 & 1 & 0 & t^2 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & t^2 \\ 111 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.
$$

 $\mathcal{M}(M,G) =$

$$
\mathcal{M}(P,G) = \begin{pmatrix} 4 & 31 & 22 & 211 & 13 & 121 & 112 & 1111 \\ 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 31 & 0 & 1 & -t+1 & -t^3+t^2 & 1 & t^2-t & 0 & 0 \\ 0 & 0 & 1 & t^2-t & 0 & -t+1 & 0 & 0 \\ 211 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 1111 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.
$$

$$
\mathcal{M}(\mathcal{S}, G) = \begin{pmatrix} 4 & 31 & 22 & 211 & 13 & 121 & 112 & 1111 \\ 4 & t & t & t^3 & t & t^3 & t^3 & t^6 \\ 31 & 0 & 1 & 0 & t & 0 & t & 0 & t^3 \\ 22 & 0 & 0 & 1 & t^2 & 0 & 1 & t & t^2 + t^4 \\ 0 & 0 & 1 & t^2 & 0 & 1 & t & t^2 + t^4 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & t \\ 121 & 0 & 0 & 1 & t^2 & 1 & t^2 & t + t^2 & t^4 + t^5 \\ 112 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & t^2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & t^3 \\ 1111 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.
$$

 $\mathcal{M}(K, G) =$

 $\sqrt{ }$ \mathcal{L} 4 31 22 211 13 121 112 1111 4 $(1 + t)(1 + t^2)(1 + t^2)$ $1 + t (1 + t)(1 + t^2)$ $(1 + t)(1 + t^2) (1 + t^2)(1 + t^3)$ $\begin{array}{cccc} 4 & 1 & +t & 1+t & (1+t)(1+t^2) & 1+t & (1+t)(1+t^2) & (1+t)(1+t^2) & (1+t)(1+t^2)(1+t^3) \\ 31 & 0 & 1 & t(1+t) & 0 & 1+t & 1+t & t^2(1+t)^2 \end{array}$ $22 \begin{bmatrix} 0 & 0 & 1 \end{bmatrix}$ $1+t^{2}$ 1 $1 + t^2$ $t(1+t)$ $t(1+t)(1+t^3)$ \cdot $\mathcal{M}(H, F) =$

 $\sqrt{ }$ $\overline{}$ 4 31 22 211 13 121 112 1111 $4 \qquad \qquad \notag \qquad \qquad 4 + q^2 + q^3 \qquad q + 2 q^2 + q^3 + q^4 \qquad q^3 + q^4 + q^5 \quad q + q^2 + q^3 \qquad q^2 + q^3 + 2 q^4 + q^5$ $q^3+q^4+q^5$ q^6 31 $\left[t \quad 1+qt+q^2t \quad 1+q+qt+2q^2t \quad q+q^2+q^3t \quad 1+qt+q^2t \quad 2q+q^2+q^2t+q^3t \right]$ $t \qquad q + q^2 + q^3$ t q^3 $22 \int_0^2 t^2 + q t + q t^2 + q^2 t^2 + q^3 t^3 + q^2 t^2 + q^4 t + q^2 t^2 + q^3 t^3 + q^2 t^4 + q^2 t^2 + q^2 t^4$ $q+qt+q^2$ t q^2 211 $\begin{vmatrix} t^3 & t + t^2 + q t^3 & 2t + t^2 + q t^2 + q t^3 & 1 + q t + q t^2 & t + t^2 + q t^3 & 1 + t + q t + 2 q t^2 \end{vmatrix}$ $1 + qt + qt^2$ q $1111 \begin{array}{c} 1 \ t^6 \ \ t^3 + t^4 + t^5 \ \end{array} \qquad t^2 + t^3 + 2 t^4 + t^5 \qquad \qquad t + t^2 + t^3 \qquad t^3 + t^4 + t^5 \qquad \qquad t + 2 t^2 + t^3 + t^4$ $t + t^2 + t^3$ 1 \setminus $\begin{matrix} \end{matrix}$ $\mathcal{M}(Q',F) =$

.

8. Acknowledgments

We thank the anonymous referees for very helpful comments.

REFERENCES

- [1] C. Berg, N. Bergeron, F. Saliola, Serrano L., and M. Zabrocki. A lift of the Schur and Hall-Littlewood bases to non-commutative symmetric functions. Canadian J. Math, To appear.
- [2] N. Bergeron and M. Zabrocki. q and q, t-analogs of non-commutative symmetric functions. Discrete Math, 298:79–103, 2005.
- [3] Lynne Butler. Subgroup lattices and symmetric functions. Mem. Amer. Math. Soc., 112(539), 1994.
- [4] Joaquin O. Carbonara. A combinatorial interpretation of the inverse t-Kostka matrix. Discrete Math., 193(1- 3):117–145, 1998. Selected papers in honor of Adriano Garsia (Taormina, 1994).
- [5] J. Désarménien, B. Leclerc, and J.-Y. Thibon. Hall-Littlewood functions and Kostka-Foulkes polynomials in representation theory. Sém. Lothar. Combin., 32:Art. B32c, approx. 38 pages, 1994.
- [6] E. Egge, N. Loehr, and G. Warrington. From quasisymmetric expansions to Schur expansions via a modified inverse Kostka matrix. European J. Combin., 31(8):2014–2027, 2010.
- [7] O. Eğecioğlu and J. Remmel. A combinatorial interpretation of the inverse Kostka matrix. Linear Multilinear Algebra, 26:59–84, 1990.
- [8] Adriano Garsia. Orthogonality of Milne's polynomials and raising operators. Discrete Math., 99:247–264, 1992.
- [9] Israel M. Gelfand, Daniel Krob, Alain Lascoux, Bernard Leclerc, Vladimir S. Retakh, and Jean-Yves Thibon. Noncommutative symmetric functions. Adv. Math., 112(2):218–348, 1995.
- [10] Ira M. Gessel. Multipartite P-partitions and inner products of skew Schur functions. In Combinatorics and algebra (Boulder, Colo., 1983), volume 34 of Contemp. Math., pages 289–317. Amer. Math. Soc., Providence, RI, 1984.
- [11] J. Haglund. A combinatorial model for the Macdonald polynomials. Proc. Natl. Acad. Sci. USA, 101(46):16127–16131 (electronic), 2004.
- [12] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for Macdonald polynomials. J. Amer. Math. Soc., 102:2690-2696, 2005.
- [13] J. Haglund, K. Luoto, S. Mason, and S. van Willigenburg. Quasisymmetric Schur functions. J. Combin. Theory Ser. A, 118(2):463–490, 2011.
- [14] M. D. Haiman. On mixed insertion, symmetry, and shifted Young tableaux. J. Combin. Theory Ser. A, 50:196–225, 1989.
- [15] Florent Hivert. Hecke algebras, difference operators, and quasi-symmetric functions. Adv. Math., 155(2):181– 238, 2000.
- [16] Anatol Kirillov. New combinatorial formula for modified Hall-Littlewood polynomials (in q-series from a contemporary perspective). Contemp. Math., 254:283–333, 2000.
- [17] A. Lascoux, J.-C. Novelli, and J.-Y. Thibon. Noncommutative symmetric functions with matrix parameters. J. Algebraic Combin., 37:621–642, 2013.
- [18] A. Lascoux and M.-P. Schützenberger. Sur une conjecture de H. O. Foulkes. C. R. Acad. Sci. Paris Sér. A-B, 286A:323–A324, 1978.
- [19] Alain Lascoux, Bernard Leclerc, and Jean-Yves Thibon. Ribbon tableaux, Hall-Littlewood functions and unipotent varieties. Sém. Lothar. Combin., 34:Art. B34g, approx. 23 pp. (electronic), 1995.
- [20] D. E. Littlewood. On certain symmetric functions. Proc. London Math. Soc. (3), 11:485–498, 1961.
- [21] N. Loehr and G. Warrington. Nested quantum Dyck paths and $\nabla(s_\lambda)$. Intl. Math. Research Notices, 2008(5):article ID rnm157, 29 pages, 2008.
- [22] N. Loehr and G. Warrington. Quasisymmetric expansions of Schur-function plethysms. Proc. of Amer. Math. Soc., 140:1159–1171, 2012.
- [23] I. G. Macdonald. Symmetric functions and Hall polynomials. Oxford Mathematical Monographs. Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [24] J.-C. Novelli, J.-Y. Thibon, and L. K. Williams. Combinatorial Hopf algebras, noncommutative Hall-Littlewood functions, and permutation tableaux. Adv. Math., 224(4):1311–1348, 2010.
- [25] B. E. Sagan. Shifted tableaux, Schur Q-functions, and a conjecture of R. P. stanley. J. Combin. Theory Ser. A, 45:62–103, 1987.
- [26] W. A. Stein et al. Sage Mathematics Software (Version 4.7). The Sage Development Team, 2009. http://www.sagemath.org.
- [27] J. R. Stembridge. Shifted tableaux and the projective representations of symmetric groups. Adv. Math., 74:87–134, 1989.
- [28] J. R. Stembridge. Enriched P-partitions. Trans. Amer. Math. Soc., 349:763–788, 1997.
- [29] L. Tevlin. Noncommutative analogs of monomial symmetric functions, Cauchy identity, and Hall scalar product. arXiv:0712.2201, 2007.
- [30] Lenny Tevlin. Noncommutative symmetric Hall-Littlewood polynomials. In 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), Discrete Math. Theor. Comput. Sci. Proc., AO, pages 915–925. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2011.
- [31] S. Ole Warnaar and Wadim Zudilin. Dedekind's η-function and Rogers-Ramanujan identities. Bull. Lond. Math. Soc., 44:1–11, 2012.
- [32] Gregory S. Warrington. Sage worksheet for Hall-Littlewood transition matrices (available online), 2013.

Dept. of Mathematics, Virginia Tech, Blacksburg, VA 24061-0123, and Mathematics Department, United States Naval Academy, Annapolis, MD 21402-5002

E-mail address: nloehr@vt.edu

LABORATOIRE DE COMBINATOIRE ET D'INFORMATIQUE MATHÉMATIQUE (LACIM), UNIVERSITÉ DU QUÉBEC À MONTRÉAL, MONTRÉAL, QC, CANADA

 $E\text{-}mail$ $address:$ serrano@lacim.ca

Dept. of Mathematics and Statistics, University of Vermont, Burlington, VT 05401 E-mail address: gregory.warrington@uvm.edu