# ASYMPTOTIC PROPERTIES OF GROUND STATES OF SCALAR FIELD EQUATIONS WITH A VANISHING PARAMETER

#### VITALY MOROZ AND CYRILL B. MURATOV

ABSTRACT. We study the leading order behavior of positive solutions of the equation

$$-\Delta u + \varepsilon u - |u|^{p-2}u + |u|^{q-2}u = 0, \qquad x \in \mathbb{R}^N,$$

where  $N \geq 3$ , q > p > 2 and when  $\varepsilon > 0$  is a small parameter. We give a complete characterization of all possible asymptotic regimes as a function of p, q and N. The behavior of solutions depends sensitively on whether p is less, equal or bigger than the critical Sobolev exponent  $p^* = \frac{2N}{N-2}$ . For  $p < p^*$  the solution asymptotically coincides with the solution of the equation in which the last term is absent. For  $p > p^*$  the solution asymptotically coincides with the solution asymptotically coincides with the solution of the equation in which the solution of the equation with  $\varepsilon = 0$ . In the most delicate case  $p = p^*$  the asymptotic behavior of the solutions is given by a particular solution of the critical Emden–Fowler equation, whose choice depends on  $\varepsilon$  in a nontrivial way.

## 1. INTRODUCTION.

1.1. Setting of the problem. This paper deals with the analysis of positive solutions of the scalar field equation

$$(P_{\varepsilon}) \qquad -\Delta u + \varepsilon u - |u|^{p-2}u + |u|^{q-2}u = 0 \quad \text{in } \mathbb{R}^N,$$

where  $N \geq 3$ , q > p > 2 and  $\varepsilon > 0$ . Specifically, we are interested in the case where  $\varepsilon$  is a small parameter, with all other parameters fixed. Our goal is to understand the behavior of ground state solutions of  $(P_{\varepsilon})$  for  $\varepsilon \ll 1$ . By a ground state solution of  $(P_{\varepsilon})$  we understand a positive weak solution  $u_{\varepsilon} \in H^1(\mathbb{R}^N) \cap L^p(\mathbb{R}^N) \cap L^q(\mathbb{R}^N)$  of  $(P_{\varepsilon})$ . These solutions are critical points (saddles) of the energy

(1.1) 
$$\mathcal{E}_{\varepsilon}(u) := \int_{\mathbb{R}^N} \frac{1}{2} |\nabla u|^2 + \frac{\varepsilon}{2} |u|^2 \, dx - \frac{1}{p} |u|^p + \frac{1}{q} |u|^q \, dx.$$

The existence and uniqueness of ground state solutions of  $(P_{\varepsilon})$  with  $\varepsilon > 0$  is well known. The existence goes back to Strauss [26, Example 2] and Berestycki and Lions [5, Example 2]. Note that by strict convexity of the integrand in  $\mathcal{E}_{\varepsilon}(u)$  for large |u| every weak solution of  $(P_{\varepsilon})$  is essentially bounded, and so by elliptic regularity these are classical solutions of  $(P_{\varepsilon})$  that decay uniformly to zero as  $|x| \to \infty$ . Then the classical Gidas-Ni-Nirenberg symmetry result [14, Theorem 2] implies that every ground state solution of  $(P_{\varepsilon})$  is spherically symmetric about some point. The uniqueness of a spherically symmetric ground state is rather delicate and was proved only quite recently by Serrin and Tang [28, Theorem 4 (ii)]. The following theorem summarizes all the above results.

Date: September 20, 2018.

<sup>1991</sup> Mathematics Subject Classification. 35J61; 49J40; 35B40.

Key words and phrases. Pokhozhaev identity; Concentration compactness; Asymptotics.

The work of C. B. M. was supported, in part, by NSF via grants DMS-0718027, DMS-0908279 and DMS-1119724.

**Theorem A** ([26, 5, 14, 28]). Let  $N \geq 3$  and q > p > 2. There exists  $\varepsilon_* > 0$ such that  $(P_{\varepsilon})$  has no ground state solutions for  $\varepsilon \geq \varepsilon_*$ , while for every  $\varepsilon \in (0, \varepsilon_*)$ equation  $(P_{\varepsilon})$  admits a unique ground state solution  $u_{\varepsilon} \in C^{\infty}(\mathbb{R}^N)$  such that  $u_{\varepsilon}(x)$ is a monotone decreasing function of |x| and there exists  $C_{\varepsilon} > 0$  such that

(1.2) 
$$\lim_{|x|\to\infty} |x|^{\frac{N-1}{2}} e^{\sqrt{\varepsilon}|x|} u_{\varepsilon}(x) = C_{\varepsilon} > 0.$$

Furthermore, every ground state solution of  $(P_{\varepsilon})$  is a translate of  $u_{\varepsilon}$ .

We note that the threshold value  $\varepsilon_*$  in Theorem 1.1 is simply the smallest value of  $\varepsilon > 0$  for which the energy  $\mathcal{E}_{\varepsilon}$  is non-negative and can be easily computed explicitly.

We are interested in the asymptotic behavior of the ground states  $u_{\varepsilon}$  as  $\varepsilon \to 0$ . This question naturally arises in the studies of various bifurcation problems, for which  $(P_{\varepsilon})$  can be considered as a canonical normal form (see e.g. [9, 30]). Problem  $(P_{\varepsilon})$  itself may also be considered as a prototypical example of a bifurcation problem for elliptic equations. In fact, our results are expected to remain valid for a broader class of scalar field equations whose nonlinearity has the leading terms in the expansion around zero which coincide with the ones in  $(P_{\varepsilon})$ . Let us also mention that problem  $(P_{\varepsilon})$  appears in the studies of non-classical nucleation near spinodal in mesoscopic models of phase transitions [7, 22, 29], as well as in the studies of the decay of false vacuum in quantum field theories [8].

In order to understand the asymptotic behavior of  $u_{\varepsilon}$  as  $\varepsilon \to 0$ , we again note that for  $u \ge 1$  the energy density in  $\mathcal{E}_{\varepsilon}(u)$  is strictly convex. Hence we may conclude that the ground state solution  $u_{\varepsilon}$  in Theorem 1.1 satisfies a uniform upper bound

(1.3) 
$$u_{\varepsilon}(0) \leq 1 \quad \text{for all } \varepsilon \in (0, \varepsilon_*).$$

Elliptic regularity then implies that locally over compact sets the solution  $u_{\varepsilon}$  converges as  $\varepsilon \to 0$  to a radial solution of the limit equation

(P<sub>0</sub>) 
$$-\Delta u - |u|^{p-2}u + |u|^{q-2}u = 0 \quad \text{in } \mathbb{R}^{N}.$$

It is known that (here and everywhere below  $p^* := \frac{2N}{N-2}$ ):

- for 2 \*</sup> equation (P<sub>0</sub>) has no finite energy solutions, which is a direct consequence of Pokhozhaev's identity (see Remark 5.1);
- for  $p > p^*$  equation ( $P_0$ ) admits a unique radial ground state solution. The existence goes back to [5, Theorem 4], see also [20, 21], while the uniqueness was proved in [21, 18].

Note that the natural energy space for equation  $(P_{\varepsilon})$  is the usual Sobolev space  $H^1(\mathbb{R}^N) = \{u \in L^2(\mathbb{R}^N) : \|\nabla u\|_{L^2} < \infty\}$ , while for  $p \ge p^*$  the limit equation  $(P_0)$  is variationally well-posed in the homogeneous Sobolev space  $D^1(\mathbb{R}^N)$ , defined as the completion of  $C_0^{\infty}(\mathbb{R}^N)$  with respect to the Dirichlet norm  $\|\nabla u\|_{L^2}$ . Clearly,  $H^1(\mathbb{R}^N) \subsetneq D^1(\mathbb{R}^N)$  and as a consequence, no natural perturbation setting (in the spirit of the implicit function theorem) is available to analyze the family of equations  $(P_{\varepsilon})$  as  $\varepsilon \to 0$ . In fact, a linearization of  $(P_0)$  around the ground state solution is not a Fredholm operator and has zero as the bottom of the essential spectrum in  $L^2(\mathbb{R}^N)$ . As a consequence, advanced Lyapunov–Schmidt type reduction methods of Ambrosetti and Malchiodi [3] are not applicable to the family of equations  $(P_{\varepsilon})$ .

If we introduce the *canonical* rescaling associated with the lowest order nonlinear term in  $(P_{\varepsilon})$ :

(1.4) 
$$v(x) = \varepsilon^{-\frac{1}{p-2}} u\left(\frac{x}{\sqrt{\varepsilon}}\right),$$

then  $(P_{\varepsilon})$  transforms into the equation

$$(R_{\varepsilon}) \qquad -\Delta v + v = |v|^{p-2}v - \varepsilon^{\frac{q-p}{p-2}}|v|^{q-2}v \quad \text{in } \mathbb{R}^{N}.$$

The limit problem associated to  $(R_{\varepsilon})$  as  $\varepsilon \to 0$  has the form

$$(R_0) \qquad -\Delta v + v = |v|^{p-2}v \quad \text{in } \mathbb{R}^N$$

It is well-known that:

- for  $p \ge p^*$  equation  $(R_0)$  has no finite energy solutions, which is a direct consequence of Pokhozhaev's identity [23, 5]);
- for  $2 equation (<math>R_0$ ) admits a unique radial ground state solution. The existence goes back at least to [26], the uniqueness was proved in [17].

The advantage of the rescaling (1.4) is that at least in the range  $2 both <math>(R_{\varepsilon})$  and the limit problem  $(R_0)$  are variationally well-posed in the same Sobolev space  $H^1(\mathbb{R}^N)$ . Then the rescaled problem  $(R_{\varepsilon})$  could be naturally seen as a small perturbation of the limit problem  $(R_0)$  and the family of ground states  $(v_{\varepsilon})$  of problem  $(R_{\varepsilon})$  could be rigorously interpreted as a perturbation of the ground state solution of the limit problem  $(R_0)$ . This could be done e.g. by using a combination of the variational and Lyapunov–Schmidt perturbation techniques as developed by Ambrosetti, Malchiodi et al., see [3] and further references therein.

The distinction between the asymptotic behaviors of the solutions of problem  $(P_{\varepsilon})$  as  $\varepsilon \to 0$  depending on the value of p as compared to  $p^*$  was first pointed out in [22]. There it was also observed that the asymptotic behavior of the ground states  $u_{\varepsilon}$  for  $p = p^*$  is not controlled by the solution set structure of either  $(P_0)$  or  $(R_0)$ . Formal asymptotic analysis of [22] explains that, in fact, three different asymptotic regimes have to be distinguished in  $(P_{\varepsilon})$ : the subcritical case  $2 , the supercritical case <math>p > p^*$  and the most delicate critical case  $p = p^*$ .

It this work, using an adaptation of the constrained minimization techniques developed by H. Berestycki and P.-L. Lions in [5], combined with the Pokhozhaev identities associated with  $(P_{\varepsilon})$  and relevant limit problems, we provide a complete analysis of these three asymptotic regimes. The analysis confirms and extends the ideas introduced in [22] and gives a full characterization of the asymptotic behavior of ground state solutions of  $(P_{\varepsilon})$  for  $\varepsilon \to 0$ .

**Notations.** For  $\varepsilon \ll 1$  and  $f(\varepsilon), g(\varepsilon) \ge 0$ , we write  $f(\varepsilon) \le g(\varepsilon), f(\varepsilon) \sim g(\varepsilon)$  and  $f(\varepsilon) \simeq g(\varepsilon)$ , implying that there exists  $\varepsilon_0 > 0$  such that for every  $0 < \varepsilon \le \varepsilon_0$ :

- $f(\varepsilon) \lesssim g(\varepsilon)$  if there exists C > 0 independent of  $\varepsilon$  such that  $f(\varepsilon) \leq Cg(\varepsilon)$ ;
- $f(\varepsilon) \sim g(\varepsilon)$  if  $f(\varepsilon) \lesssim g(\varepsilon)$  and  $g(\varepsilon) \lesssim f(\varepsilon)$ ;
- $f(\varepsilon) \simeq g(\varepsilon) \text{ if } f(\varepsilon) \sim g(\varepsilon) \text{ and } \lim_{\varepsilon \to 0} \frac{f(\varepsilon)}{g(\varepsilon)} = 1.$

We also use the standard notations f = O(g) and f = o(g), bearing in mind that  $f \ge 0$  and  $g \ge 0$ . As usual,  $C, c, c_1$ , etc., denote generic positive constants independent of  $\varepsilon$ .

# 2. Main results.

2.1. Subcritical case  $2 . Since in the subcritical case the limit equation <math>(P_0)$  has no ground state solutions, in view of (1.3) the family of ground states  $u_{\varepsilon}$  must converge to zero, locally over compact subsets of  $\mathbb{R}^N$ . To describe the asymptotic behavior of  $u_{\varepsilon}$  we use the rescaling (1.4) which transforms  $(P_{\varepsilon})$  into equation  $(R_{\varepsilon})$ . For  $2 , let <math>v_0(x)$  denote the unique radial ground state

solution of the limit equation  $(R_0)$ . It is well-known that  $v_0 \in C^{\infty}(\mathbb{R}^N)$ ,  $v_0(x)$  is a monotone decreasing function of |x| and that

(2.1) 
$$\lim_{|x|\to\infty} |x|^{\frac{N-1}{2}} e^{|x|} u_{\varepsilon}(x) = C_0 > 0,$$

cf. [5]. The advantage of the rescaling (1.4) is that both  $(R_{\varepsilon})$  and the limit problems  $(R_0)$  are variationally well-posed in the Sobolev space  $H^1(\mathbb{R}^N)$ . Note however that  $(R_0)$  is translationally invariant and hence the radial ground state  $v_0(x)$  is not an isolated solution. As a consequence, an Implicit Function Theorem argument is not directly applicable to  $(R_{\varepsilon})$ . Nevertheless, it is known that the linearization operator  $-\Delta + 1 - (p-1)v_0^{p-2}$  of  $(R_0)$  around the ground state  $v_0$  is a Fredholm operator in  $H^1(\mathbb{R}^N)$ , see [3, Lemma 4.1]. Then perturbation techniques in [3] could be easily adapted in order to show that for all sufficiently small  $\varepsilon > 0$  equation  $(R_{\varepsilon})$  admits a radial ground state  $v_{\varepsilon}(x)$  which converges to  $v_0(x)$  as  $\varepsilon \to 0$ . Rescaling back to the original variable and taking into account the uniqueness of the radial ground state of  $(P_{\varepsilon})$  we arrive at the following (folklore) result.

**Theorem 2.1.** Let  $2 . As <math>\varepsilon \to 0$ , the rescaled family of ground states

(2.2) 
$$v_{\varepsilon}(x) := \varepsilon^{-\frac{1}{p-2}} u_{\varepsilon}\left(\frac{x}{\sqrt{\varepsilon}}\right)$$

converges to  $v_0(x)$  in  $H^1(\mathbb{R}^N)$ ,  $L^q(\mathbb{R}^N)$  and  $C^2(\mathbb{R}^N)$ . In particular,

(2.3) 
$$u_{\varepsilon}(0) \simeq \varepsilon^{\frac{1}{p-2}} v_0(0).$$

In the last section of this work we provide a short alternative proof of this result based only upon variational methods which are developed in the main part of this paper and without explicit references to perturbation techniques.

Remark 2.2. For  $p \ge p^*$  Pokhozhaev's identity implies that  $(R_0)$  has no nontrivial solutions in  $H^1(\mathbb{R}^N)$ . In fact, it is known that  $v_0(0) \to \infty$  as  $p \uparrow p^*$ . Note that a complete asymptotic characterization of the ground states of equations  $(R_0)$  as  $p \uparrow p^*$  (and more general *m*-Laplace equations of type  $(R_0)$ ) was given in [12, 13, 11]. More specifically (see [13, Corollary 1]), if  $\delta := p^* - p$ , then for  $\delta \downarrow 0$  it holds

(2.4) 
$$v_0(0) \simeq \beta_N \begin{cases} \delta^{-\frac{N-2}{4}}, & N \ge 5, \\ \delta^{-\frac{1}{2}} |\log \delta|, & N = 4, \\ \delta^{-\frac{1}{2}}, & N = 3, \end{cases}$$

for some explicit constants  $\beta_N > 0$ . This suggests that for  $p = p^*$  rescaling (1.4) fails to capture the behavior of the ground states  $u_{\varepsilon}$  and a different approach is needed to handle the critical and supercritical case. Note also that the asymptotic behavior of ground states of "slightly" subcritical elliptic problems in the context of bounded domains was studied in [4, 6, 16, 24].

2.2. Supercritical case  $p > p^*$ . In contrast to the subcritical case, for  $p > p^*$  the limit equation  $(P_0)$  admits a unique radial ground state solution  $u_0(x) > 0$ . It is known that  $u_0 \in C^2(\mathbb{R}^N)$ ,  $u_0(x)$  is a monotone decreasing function of |x| and that

(2.5) 
$$\lim_{|x| \to \infty} |x|^{N-2} u_0(x) = C_0 > 0$$

see [5, Theorem 4] or [20, 21] for the existence, and [21, 18] for the uniqueness proofs. However, as was already mentioned, the linearization operator  $-\Delta - (p-1)u_0^{p-2}$  of  $(P_0)$  around the ground state  $u_0$  is not Fredholm and has zero as the bottom of the essential spectrum in  $L^2(\mathbb{R}^N)$ . As a consequence, standard perturbation methods are not applicable to  $(P_0)$ . Using a direct analysis of the family of constrained minimizations problem associated to  $(P_{\epsilon})$ , we prove the following. **Theorem 2.3.** Let  $p > p^*$ . As  $\varepsilon \to 0$ , the family of ground states  $u_{\varepsilon}$  converges to  $u_0$  in  $D^1(\mathbb{R}^N)$ ,  $L^q(\mathbb{R}^N)$  and  $C^2(\mathbb{R}^N)$ . In particular,

(2.6) 
$$u_{\varepsilon}(0) \simeq u_0(0).$$

In addition,  $\varepsilon ||u_{\varepsilon}||_2^2 \to 0$ .

Remark 2.4. For  $p = p^*$  Pokhozhaev's identity implies that  $(P_0)$  has no nontrivial solutions in  $D^1(\mathbb{R}^N)$ . In fact, it is not difficult to show that  $u_0(0) \to 0$  as  $p \downarrow p^*$ . Moreover, if  $\delta := p - p^*$ , then for  $\delta \downarrow 0$  we prove

(2.7) 
$$\delta^{\frac{1}{q-p^*}} \lesssim u_0(0) \lesssim \delta^{\frac{1}{q+N}},$$

and, provided that  $q > \frac{N(N+2)}{2(N-2)}$ ,

(2.8) 
$$u_0(0) \sim \delta^{\frac{1}{q-p^*}}.$$

See Section 5.4 for further details and full statements. Note that related estimates for the asymptotics of ground states of  $(P_0)$  with fixed  $q > p > p^*$  on a sequence of expanding domains were studied in [20, 21].

2.3. Critical case  $p = p^*$ . In the critical case both the unrescaled limit equation  $(P_0)$  and the "canonically" rescaled equation  $(R_0)$  have no nontrivial finite energy solutions. We are going to show that after a suitable rescaling the correct limit equation for  $(P_{\varepsilon})$  is in fact given by the critical Emden–Fowler equation

$$(R_*) \qquad -\Delta U = U^{p^*-1} \quad \text{in } \mathbb{R}^N.$$

It is well-known that the radial ground states of  $(R_*)$  are given by the function

(2.9) 
$$U_1(x) := \left(1 + \frac{|x|^2}{N(N-2)}\right)^{-\frac{N-2}{2}}$$

and the family of its rescalings

(2.10) 
$$U_{\lambda}(x) := \lambda^{-\frac{N-2}{2}} U_1\left(\frac{x}{\lambda}\right), \qquad \lambda > 0.$$

Our main result in this work is the following.

**Theorem 2.5.** Let  $p = p^*$ . There exists a rescaling  $\lambda_{\varepsilon} : (0, \varepsilon_*) \to (0, \infty)$  such that as  $\varepsilon \to 0$ , the rescaled family of ground states

(2.11) 
$$v_{\varepsilon}(x) := \lambda_{\varepsilon}^{\frac{1}{p-2}} u_{\varepsilon}\left(\lambda_{\varepsilon}x\right)$$

converges to  $U_1(x)$  in  $D^1(\mathbb{R}^N)$ ,  $L^q(\mathbb{R}^N)$  and  $C^2(\mathbb{R}^N)$ . Moreover,

(2.12) 
$$\lambda_{\varepsilon} \sim \begin{cases} \varepsilon^{-\frac{p-2}{2q-4}}, & N \ge 5, \\ (\varepsilon \log \frac{1}{\varepsilon})^{-\frac{1}{q-2}}, & N = 4, \\ \varepsilon^{-\frac{1}{q-4}}, & N = 3. \end{cases}$$

and

(2.13) 
$$u_{\varepsilon}(0) \sim \begin{cases} \varepsilon^{\frac{1}{q-2}}, & N \ge 5, \\ (\varepsilon \log \frac{1}{\varepsilon})^{\frac{1}{q-2}}, & N = 4, \\ \varepsilon^{\frac{1}{2q-8}}, & N = 3. \end{cases}$$

*Remark* 2.6. Asymptotics (2.12) and (2.13) were first derived in [22] using methods of formal asymptotic expansions. Theorem 2.5, in particular, justifies the values of precise asymptotic constants found in [22].

2.4. **Outline.** The rest of the paper is organized as follows. In Section 3 we introduce a variational characterization of the ground states  $u_{\varepsilon}$  of the problem  $(P_{\varepsilon})$  as well as some other preliminary results. In Section 4 we study the critical case  $p = p^*$  and prove Theorem 2.5. In Section 5 we consider the supercritical case  $p > p^*$  and prove Theorem 2.3. Finally, in Section 6 we will revisit the subcritical case 2 and sketch a simple variational proof of Theorem 2.1, in the spirit of our previous arguments.

## 3. VARIATIONAL CHARACTERIZATION OF THE GROUND STATES.

The existence and properties of the ground state  $u_{\varepsilon}$  of equation  $(P_{\varepsilon})$ , as summarized in Theorem A, could be established in several different ways, e.g. by means of ODE techniques. Here we shall utilize a variational characterization of the ground states  $u_{\varepsilon}$  developed by Berestycki and Lions in [5].

Given q > p > 2 and  $\varepsilon \ge 0$  set

(3.1) 
$$f_{\varepsilon}(u) := \begin{cases} 0, & u < 0, \\ u^{p-1} - u^{q-1} - \varepsilon u, & u \in [0, 1], \\ -\varepsilon, & u > 1, \end{cases} \qquad F_{\varepsilon}(u) := \int_{0}^{u} f_{\varepsilon}(s) ds.$$

In view of (1.3) and since we are interested only in positive solutions of  $(P_{\varepsilon})$ , the nonlinearity in  $(P_{\varepsilon})$  may be always replaced by its bounded truncation  $f_{\varepsilon}(u)$  from (3.1).

For  $\varepsilon > 0$ , consider the constrained minimization problem

$$(S_{\varepsilon}) \qquad S_{\varepsilon} := \inf\left\{ \int_{\mathbb{R}^N} |\nabla w|^2 \, dx \right| \, w \in H^1(\mathbb{R}^N), \ p^* \int_{\mathbb{R}^N} F_{\varepsilon}(w) \, dx \ = 1 \right\}.$$

As was proved in [5, Theorem 2], there exists  $\varepsilon_* > 0$  depending only on p and q such that for all  $\varepsilon \in (0, \varepsilon_*)$  minimization problem  $(S_{\varepsilon})$  admits a positive radially symmetric minimizer  $w_{\varepsilon}(x)$ . Further, there exists a Lagrange multiplier  $\theta_{\varepsilon} > 0$  such that

(3.2) 
$$-\Delta w_{\varepsilon} = \theta_{\varepsilon} f_{\varepsilon}(w_{\varepsilon}) \quad \text{in } \mathbb{R}^{N}.$$

In particular, the minimizer  $w_{\varepsilon}$  satisfies Nehari's identity

(3.3) 
$$\int_{\mathbb{R}^N} |\nabla w_{\varepsilon}|^2 \, dx = \theta_{\varepsilon} \int f_{\varepsilon}(w_{\varepsilon}) w_{\varepsilon} \, dx$$

and Pokhozhaev's identity (see e.g. [5, Proposition 1])

(3.4) 
$$\int_{\mathbb{R}^N} |\nabla w_{\varepsilon}|^2 \, dx = \theta_{\varepsilon} p^* \int F_{\varepsilon}(w_{\varepsilon}) \, dx$$

The latter immediately implies that

(3.5) 
$$\theta_{\varepsilon} = S_{\varepsilon}.$$

Then a direct calculation involving (3.5) shows that the rescaled function

(3.6) 
$$u_{\varepsilon}(x) := w_{\varepsilon}\left(\frac{x}{\sqrt{S_{\varepsilon}}}\right)$$

is the radial ground state of  $(P_{\varepsilon})$ , described in Theorem A. Another simple consequence of (3.4) is that  $(P_{\varepsilon})$  has no nontrivial finite energy solutions for  $\varepsilon \geq \varepsilon_*$ .

Equivalently to  $(S_{\varepsilon})$ , we may seek to minimize the quotient

(3.7) 
$$\mathcal{S}_{\varepsilon}(w) := \frac{\|\nabla w\|_2^2}{\left(p^* \int_{\mathbb{R}^N} F_{\varepsilon}(w) \, dx\right)^{\frac{N-2}{N}}}, \qquad w \in \mathcal{M}_{\varepsilon},$$

where

(3.8) 
$$\mathcal{M}_{\varepsilon} := \Big\{ 0 \le u \in D^1(\mathbb{R}^N), \ \int_{\mathbb{R}^N} F_{\varepsilon}(w) \, dx > 0 \Big\}.$$

Clearly, if we set  $w_{\lambda}(x) := w(\lambda x)$  then  $S_{\varepsilon}(w_{\lambda}) = S_{\varepsilon}(w)$  for all  $\lambda > 0$ , that is  $S_{\varepsilon}$  is invariant with respect to dilations. This implies that

(3.9) 
$$S_{\varepsilon} = \inf_{w \in \mathcal{M}_{\varepsilon}} \mathcal{S}_{\varepsilon}(w).$$

In addition, since clearly  $\mathcal{M}_{\varepsilon_2} \subset \mathcal{M}_{\varepsilon_1}$  for  $\varepsilon_2 > \varepsilon_1 > 0$ , (3.9) shows that  $S_{\varepsilon}$  is a monotone nondecreasing function of  $\varepsilon \in (0, \varepsilon_*)$ .

One of the consequences of Pokhozhaev's identity (3.4) is an expression for the total energy of the solution

(3.10) 
$$\mathcal{E}_{\varepsilon}(u_{\varepsilon}) = \left(\frac{1}{2} - \frac{1}{p^*}\right) S_{\varepsilon}^{\frac{N}{2}},$$

see [5, Corollary 2], which shows that  $u_{\varepsilon}$  is indeed a ground state, i.e. a nontrivial solution with the least energy.

We will be frequently using the following well known decay and compactness properties of radial functions on  $\mathbb{R}^N$ .

Lemma 3.1. [5, Lemma A.IV, Theorem A.I'].

(1) Let  $s \ge 1$  and let  $u \in L^s(\mathbb{R}^N)$  be a radial non-increasing function. Then for every  $x \ne 0$  it holds

(3.11) 
$$u(x) \le C_{s,N} |x|^{-\frac{N}{s}} ||u||_s,$$

where  $C_{s,N} = |B_1(0)|^{-\frac{1}{s}}$ .

- (2) Let  $u_n \in H^1(\mathbb{R}^N)$  be a sequence of radial non-decreasing functions such that  $u_n \rightharpoonup u$  in  $H^1(\mathbb{R}^N)$ . Then upon extraction of a subsequence
- (3.12)  $u_n \to u \text{ in } L^{\infty}(\mathbb{R}^N \setminus B_r(0)) \text{ and } L^s(\mathbb{R}^N \setminus B_r(0)) \quad \forall r > 0 \text{ and } \forall s > p^*.$

4. Critical case  $p = p^*$ .

Throughout this section we always assume that  $p = p^*$ . In this critical case Pokhozhaev's identity implies that both the limit equation  $(P_0)$  and the canonically rescaled limit equation  $(R_0)$  have no positive finite energy solutions. We are going to show that after a suitably chosen rescaling, the limit equation for  $(P_{\varepsilon})$  is in fact given by the critical Emden–Fowler equation.

#### 4.1. Critical Emden–Fowler equation. Let

$$(S_*) \qquad S_* := \inf\left\{\int_{\mathbb{R}^N} |\nabla w|^2 \, dx \ \Big| \ w \in D^1(\mathbb{R}^N), \ \int_{\mathbb{R}^N} |w|^p \, dx = 1\right\}$$

be the optimal constant in the Sobolev inequality

(4.1) 
$$\int_{\mathbb{R}^N} |\nabla w|^2 \, dx \ge S_* \left( \int_{\mathbb{R}^N} |w|^p \, dx \right)^{2/p}, \qquad \forall w \in D^1(\mathbb{R}^N).$$

It is easy to see that  $S_*$  is achieved by translations of the rescaled family

(4.2) 
$$W_{\lambda}(x) := U_{\lambda}\left(\sqrt{S_*}x\right),$$

where  $U_{\lambda}(x)$  are the ground states of the critical Emden–Fowler equation  $(R_*)$ , explicitly defined by (2.9). Clearly,

(4.3) 
$$||W_{\lambda}||_{p} = 1, \qquad ||\nabla W_{\lambda}||_{2}^{2} = S_{*}.$$

A straightforward computation leads to the explicit expression

(4.4) 
$$\|\nabla U_{\lambda}\|_{2}^{2} = \|U_{\lambda}\|_{p}^{p} = S_{*}^{\frac{N}{2}}$$

Note that the family of minimizers  $W_{\lambda}$  solves the Euler–Lagrange equation

(4.5) 
$$-\Delta W = S_* W^{p-1} \quad \text{in } \mathbb{R}^N.$$

4.2. Variational estimates of  $S_{\varepsilon}$ . For our purposes it is convenient to consider the dilation invariant Sobolev quotient

(4.6) 
$$\mathcal{S}_*(w) := \frac{\int_{\mathbb{R}^N} |\nabla w|^2 \, dx}{\left(\int_{\mathbb{R}^N} |w|^p \, dx\right)^{\frac{N-2}{N}}}, \qquad w \in D^1(\mathbb{R}^N), \quad w \neq 0,$$

so that

(4.7) 
$$S_* = \inf_{0 \neq w \in D^1(\mathbb{R}^N)} \mathcal{S}_*(w).$$

Denote

(4.8) 
$$\sigma_{\varepsilon} := S_{\varepsilon} - S_*$$

In order to control  $\sigma_{\varepsilon}$  in terms of  $\varepsilon$ , we shall use Sobolev's minimizers  $W_{\lambda}$  as a family of test functions for  $S_{\varepsilon}$ . Note that since  $W_{\lambda} \in L^2(\mathbb{R}^N)$  only if  $N \geq 5$ , we shall consider the higher and lower dimensions separately. Straightforward calculations show that  $W_{\lambda} \in L^s(\mathbb{R}^N)$  for all  $s > \frac{N}{N-2}$ , with

(4.9) 
$$\|W_{\lambda}\|_{s}^{s} = \lambda^{N-\frac{2s}{p-2}} \|W_{1}\|_{s}^{s} = \lambda^{-\frac{N-2}{2}(s-p)} \|W_{1}\|_{s}^{s}.$$

In particular, if  $N \geq 5$  then  $W_{\lambda} \in L^2(\mathbb{R}^N)$  and

(4.10) 
$$||W_{\lambda}||_{2}^{2} = \lambda^{2} ||W_{1}||_{2}^{2}$$

To consider dimensions N = 3, 4, given  $R \gg \lambda$ , we introduce a cut off function  $\eta_R \in C_c^{\infty}(\mathbb{R})$  such that  $\eta_R(r) = 1$  for  $|r| < R, 0 < \eta(r) < 1$  for R < |r| < 2R,  $\eta_R(r) = 0$  for |r| > 2R and  $|\eta'(r)| \le 2/R$ . We then compute as in, e.g., [27, Chapter III, proof of Theorem 2.1]<sup>1</sup>

(4.13) 
$$\int |\nabla(\eta_R W_\lambda)|^2 = S_* + O\left(\left(\frac{R}{\lambda}\right)^{-(N-2)}\right).$$

(4.14) 
$$\int |\eta_R W_\lambda|^p = 1 - O\left(\left(\frac{R}{\lambda}\right)^{-N}\right)$$

(4.15) 
$$\int |\eta_R W_{\lambda}|^q = \lambda^{-\frac{N-2}{2}(q-p)} ||W_1||_q^q \Big( 1 - O\Big( \Big(\frac{R}{\lambda}\Big)^{-(N-2)\Big(q-\frac{N}{N-2}\Big)} \Big) \Big),$$

(4.16) 
$$\int |\eta_R W_\lambda|^2 = \lambda^2 ||\eta_{R/\lambda}^2 W_1||_2^2 = \begin{cases} O(\lambda^2 \log \frac{R}{\lambda}), & N = 4, \\ O(\lambda R), & N = 3. \end{cases}$$

Using the above calculations we obtain an upper estimate of  $\sigma_{\varepsilon}$  which is essential for further considerations.

<sup>1</sup>Note that if  $0 < U \in H^1_{loc}(\mathbb{R}^N)$  solves

(4.11) 
$$-\Delta U = kU^{p-1}, \qquad x \in \mathbb{R}^N,$$

for some  $k \neq 0$ , then

(4.12) 
$$\int |\nabla(\eta U)|^2 dx = k \int \eta^2 |U|^p dx + \int |\nabla \eta|^2 U^2 dx \quad \text{for all } \eta \in C_c^\infty(\mathbb{R}^N).$$

See also [27, Chapter III, proof of Theorem 2.1].

Lemma 4.1. We have

(4.17) 
$$0 < \sigma_{\varepsilon} \lesssim \begin{cases} \varepsilon^{\frac{q-p}{q-2}}, & N \ge 5, \\ (\varepsilon \log \frac{1}{\varepsilon})^{\frac{q-4}{q-2}}, & N = 4, \\ \varepsilon^{\frac{q-6}{2q-8}}, & N = 3. \end{cases}$$

In particular,  $\sigma_{\varepsilon} \to 0$  as  $\varepsilon \to 0$ .

*Proof.* To prove that  $\sigma_{\varepsilon} > 0$  simply note that

(4.18) 
$$S_* \leq \mathcal{S}_*(w_{\varepsilon}) < \mathcal{S}_{\varepsilon}(w_{\varepsilon}) = S_{\varepsilon}.$$

We shall now establish the upper bound on  $\sigma_{\varepsilon}$ , which clearly tends to zero as  $\varepsilon \to 0$ .

CASE  $N \geq 5$ . Using  $W_{\lambda}$  as a family of test functions, we obtain that  $W_{\lambda} \in \mathcal{M}_{\varepsilon}$  for sufficiently small  $\varepsilon$  and sufficiently large  $\lambda$ , and we have

(4.19) 
$$S_{\varepsilon}(W_{\lambda}) \leq \frac{S_{*}}{\left(1 - \beta_{2}\varepsilon\lambda^{2} - \beta_{q}\lambda^{-\frac{N-2}{2}(q-p)}\right)^{\frac{N-2}{N}}},$$

where

(4.20) 
$$\beta_2 := \frac{p}{2} \|W_1\|_2^2, \qquad \beta_q := \frac{p}{q} \|W_1\|_q^q.$$

To minimize the right hand side of (4.19), we have to minimize the scalar function

(4.21) 
$$\psi(\lambda) := \beta_2 \varepsilon \lambda^2 + \beta_q \lambda^{-\frac{N-2}{2}(q-p)}$$

It is easy to see that  $\psi$  achieves its minimum in scaling at

(4.22) 
$$\lambda_{\varepsilon} = \varepsilon^{-\frac{2}{(N-2)(q-2)}}$$

and

(4.23) 
$$\min_{\lambda>0} \psi \sim \psi(\lambda_{\varepsilon}) \sim \varepsilon^{\frac{q-p}{q-2}}.$$

For  $N \geq 5$ , we conclude that

(4.24) 
$$\mathcal{S}_{\varepsilon}(W_{\lambda}) \leq \frac{S_{*}}{\left(1 - \psi(\lambda_{\varepsilon})\right)^{\frac{N-2}{N}}} = S_{*}\left(1 + O(\psi(\lambda_{\varepsilon}))\right) = S_{*} + O\left(\varepsilon^{\frac{q-p}{q-2}}\right),$$

and the bound (4.17) is achieved on the function  $W_{\lambda_{\varepsilon}}$ , where  $\lambda_{\varepsilon}$  is given by (4.22).

CASE N = 4. Assume  $R \gg \lambda$ . Testing against  $\eta_R W_{\lambda}$  and using calculations in (4.13)–(4.16) with p = 4, we obtain

(4.25) 
$$S_{\varepsilon}(\eta_R W_{\lambda}) \leq \left(S_* + O\left(\left(\frac{R}{\lambda}\right)^{-2}\right)\right) \\ \times \left(\left[1 - O\left(\left(\frac{R}{\lambda}\right)^{-4}\right)\right] - \varepsilon \lambda^2 O\left(\log\frac{R}{\lambda}\right) - \beta_q \lambda^{-(q-4)} \left[1 - O\left(\left(\frac{R}{\lambda}\right)^{-2(q-2)}\right)\right]\right)^{-\frac{1}{2}} \\ \leq S_* \left(1 + O(\psi(\lambda, R))\right),$$

where

(4.26) 
$$\psi(\lambda, R) = \varepsilon \lambda^2 O\left(\log \frac{R}{\lambda}\right) + O\left(\left(\frac{R}{\lambda}\right)^{-2}\right) + \beta_q \lambda^{-(q-4)} \left[1 - o(1)\right].$$

Choose

(4.27) 
$$\lambda_{\varepsilon} = \left(\varepsilon \log \frac{1}{\varepsilon}\right)^{-\frac{1}{q-2}}, \qquad R_{\varepsilon} = \varepsilon^{-\frac{1}{2}}.$$

A routine calculation shows that as  $\varepsilon \to 0$ ,

(4.28) 
$$\log \frac{R_{\varepsilon}}{\lambda_{\varepsilon}} \sim \log \frac{1}{\varepsilon},$$

and hence

(4.29) 
$$\psi(\lambda_{\varepsilon}, R_{\varepsilon}) \sim \left(\varepsilon \log \frac{1}{\varepsilon}\right)^{\frac{q-4}{q-2}}.$$

Thus bound (4.17) is achieved by the test function  $\eta_{R_{\varepsilon}}W_{\lambda_{\varepsilon}}$ , where  $\lambda_{\varepsilon}$  and  $R_{\varepsilon}$  are given by (4.27).

CASE N = 3. Assume  $R \gg \lambda$ . Testing against  $\eta_R W_{\lambda}$  and using calculations in (4.13)–(4.16) with p = 6, we obtain

$$(4.30) \quad \mathcal{S}_{\varepsilon}(\eta_R W_{\lambda}) \leq \left(S_* + O\left(\left(\frac{R}{\lambda}\right)^{-1}\right)\right) \\ \times \left(\left[1 - O\left(\left(\frac{R}{\lambda}\right)^{-3}\right)\right] - \varepsilon \lambda O(R) - \beta_q \lambda^{-\frac{1}{2}(q-6)} \left[1 - O\left(\left(\frac{R}{\lambda}\right)^{-(q-3)}\right)\right]\right)^{-\frac{1}{3}} \\ \leq S_* \left(1 + O(\psi(\lambda, R))\right),$$

where

(4.31) 
$$\psi(\lambda, R) = \varepsilon \lambda O(R) + O\left(\left(\frac{R}{\lambda}\right)^{-1}\right) + \beta_q \lambda^{-\frac{1}{2}(q-6)} \left[1 - o(1)\right].$$

Choosing

(4.32) 
$$\lambda_{\varepsilon} = \varepsilon^{-\frac{1}{q-4}}, \qquad R_{\varepsilon} = \varepsilon^{-\frac{1}{2}},$$

we then find that

(4.33) 
$$\psi(\lambda_{\varepsilon}, R_{\varepsilon}) \sim \varepsilon^{\frac{q-6}{2q-8}}$$

and the bound (4.17) is achieved on the test function  $\eta_{R_{\varepsilon}}W_{\lambda_{\varepsilon}}$ , where  $\lambda_{\varepsilon}$  and  $R_{\varepsilon}$  is given by (4.32).

4.3. **Pokhozhaev estimates.** Nehari identity (3.3) combined with Pokhozhaev's identity (3.4) lead to the following important relations.

**Lemma 4.2.** Set  $\kappa := \frac{q(p-2)}{2(q-p)} > 0$ . Then

(4.34) 
$$\|w_{\varepsilon}\|_{q}^{q} = \kappa \varepsilon \|w_{\varepsilon}\|_{2}^{2},$$

(4.35) 
$$\|w_{\varepsilon}\|_{p}^{p} = 1 + (\kappa + 1)\varepsilon\|w_{\varepsilon}\|_{2}^{2}$$

*Proof.* Since  $w_{\varepsilon}$  is a minimizer of  $(S_{\varepsilon})$ , identities (3.3)–(3.5) read as

(4.36) 
$$1 = \|w_{\varepsilon}\|_{p}^{p} - \|w_{\varepsilon}\|_{q}^{q} - \varepsilon \|w_{\varepsilon}\|_{2}^{2},$$

(4.37) 
$$1 = \|w_{\varepsilon}\|_{p}^{p} - \frac{p}{q}\|w_{\varepsilon}\|_{q}^{q} - \frac{p}{2}\varepsilon\|w_{\varepsilon}\|_{2}^{2}$$

Then the conclusion follows by a direct algebraic computation.

Lemma 4.3. 
$$\varepsilon(\kappa+1) \|w_{\varepsilon}\|_{2}^{2} \leq \frac{N}{N-2} S_{*}^{-1} \sigma_{\varepsilon} (1+o(1))$$

*Proof.* Since  $w_{\varepsilon}$  is a minimizer of  $(S_{\varepsilon})$ , with the help of Lemma 4.2 we obtain

(4.38) 
$$S_* \leq \mathcal{S}_*(w_{\varepsilon}) = \frac{\|\nabla w_{\varepsilon}\|_2^2}{\|w_{\varepsilon}\|_p^2} = \frac{S_{\varepsilon}}{\left(1 + (\kappa + 1)\varepsilon \|w_{\varepsilon}\|_2^2\right)^{\frac{N-2}{N}}},$$

or, equivalently,

(4.39) 
$$S_*^{\frac{N}{N-2}} \left( 1 + (\kappa+1)\varepsilon \|w_{\varepsilon}\|_2^2 \right) \le S_{\varepsilon}^{\frac{N}{N-2}}.$$

Since  $\sigma_{\varepsilon} := S_{\varepsilon} - S_*$ , rearranging and differentiating, for  $\varepsilon \to 0$  we obtain

(4.40) 
$$S_*^{\frac{N}{N-2}}(\kappa+1)\varepsilon \|w_{\varepsilon}\|_2^2 \le S_{\varepsilon}^{\frac{N}{N-2}} - S_*^{\frac{N}{N-2}} = \frac{N}{N-2}S_*^{\frac{2}{N-2}}\sigma_{\varepsilon} + o(\sigma_{\varepsilon}),$$

so the conclusion follows.

Combining the results of the three lemmas just proved, we obtain the following result concerning the asymptotic behavior of different norms associated with the minimizer  $w_{\varepsilon}$  of  $(S_{\varepsilon})$ .

**Corollary 4.4.** As  $\varepsilon \to 0$ , we have

(4.41) 
$$||w_{\varepsilon}||_{p}^{p} \to 1, \quad ||w_{\varepsilon}||_{q}^{q} \to 0, \quad \varepsilon ||w_{\varepsilon}||_{2}^{2} \to 0.$$

4.4. Optimal rescaling. Following [19], consider the concentration function

(4.42) 
$$Q_{\varepsilon}(\lambda) = \int_{B_{\lambda}} |w_{\varepsilon}|^p \, dx,$$

where here and everywhere below  $B_{\lambda}$  is the ball of radius  $\lambda$  centered at the origin. Clearly,  $Q_{\varepsilon}(\cdot)$  is strictly monotone increasing, with  $\lim_{\lambda\to 0} Q_{\varepsilon}(\lambda) = 0$  and  $\lim_{\lambda\to\infty} Q_{\varepsilon}(\lambda) = ||w_{\varepsilon}||_{p}^{p} \to 1$  as  $\varepsilon \to 0$  in view of Corollary 4.4. Therefore, the equation  $Q_{\varepsilon}(\lambda) = Q_{*}$  with

(4.43) 
$$Q_* := \int_{B_1} |W_1(x)|^p \, dx < 1,$$

has a unique solution  $\lambda = \lambda_{\varepsilon} > 0$  whenever  $\varepsilon \ll 1$ :

(4.44) 
$$Q_{\varepsilon}(\lambda_{\varepsilon}) = Q_*.$$

Similarly, since the function

(4.45) 
$$Q_0(\lambda) := \int_{B_{\lambda^{-1}}} |W_1(x)|^p \, dx = \int_{B_1} |W_\lambda(x)|^p \, dx$$

is strictly monotone decreasing, with  $\lim_{\lambda\to 0} Q_0(\lambda) = 1$  and  $\lim_{\lambda\to\infty} Q_0(\lambda) = 0$ , there is a unique solution to the equation  $Q_0(\lambda) = Q_*$ . In fact, by the definition of  $Q_*$  this equation is satisfied if and only if  $\lambda = 1$ .

Using the value of  $\lambda_{\varepsilon}$  implicitly determined by (4.44), we define the rescaled family

(4.46) 
$$v_{\varepsilon}(x) := \lambda_{\varepsilon}^{\frac{N-2}{2}} w_{\varepsilon}(\lambda_{\varepsilon} x)$$

Note that

(4.47) 
$$||v_{\varepsilon}||_{p} = ||w_{\varepsilon}||_{p} = 1 + o(1), \qquad ||\nabla v_{\varepsilon}||_{2}^{2} = ||\nabla w_{\varepsilon}||_{2}^{2} = S_{*} + o(1),$$

i.e.  $(v_{\varepsilon})$  is a minimizing family for  $S_*$ . Note also that

(4.48) 
$$\int_{B_1} |v_{\varepsilon}(x)|^p \, dx = Q_*$$

The next statement is a direct consequence of the Concentration–Compactness Principle of P.L.Lions, cf. [27, Chapter I, Theorem 4.9].

Lemma 4.5. 
$$\|\nabla (v_{\varepsilon} - W_1)\|_2 \to 0$$
 and  $\|v_{\varepsilon} - W_1\|_p \to 0$  as  $\varepsilon \to 0$ .

Proof. By (4.47), for any sequence  $\varepsilon_n \to 0$  there exist a subsequence  $(\varepsilon_{n'})$  such that  $(v_{\varepsilon_{n'}})$  converges weakly in  $D^1(\mathbb{R}^N)$  to some radial function  $w_0 \in D^1(\mathbb{R}^N)$ . Applying the Concentration–Compactness Principle (cf. [27, Chapter I, Theorem 4.9] or [31, Theorem 1.41]) to  $\|v_{\varepsilon}\|_p^{-1}v_{\varepsilon}$ , we further conclude that in fact  $(v_{\varepsilon_{n'}})$  converges to

 $w_0$  strongly in  $D^1(\mathbb{R}^N)$  and  $L^p(\mathbb{R}^N)$ . As a consequence,  $||w_0||_p = 1$  and hence  $w_0$  is a radial minimizer of  $(S_*)$ , that is  $w_0 \in \{W_\lambda\}_{\lambda>0}$ . Furthermore,

(4.49) 
$$\int_{B_1} |w_0(x)|^p \, dx = Q_*$$

We therefore conclude that  $w_0 = W_1$ . Finally, by uniqueness of the limit the full sequence  $(v_n)$  converges to  $W_1$  strongly in  $D^1(\mathbb{R}^N)$  and  $L^p(\mathbb{R}^N)$ .

4.5. Rescaled equation estimates. The rescaled minimizer  $v_{\varepsilon}$  defined in (4.46) solves the equation

$$(R_{\varepsilon}^{*}) \qquad -\Delta v_{\varepsilon} + S_{\varepsilon} \varepsilon \lambda_{\varepsilon}^{2} v_{\varepsilon} = S_{\varepsilon} \left( |v_{\varepsilon}|^{p-2} v_{\varepsilon} - \lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}} |v_{\varepsilon}|^{q-2} v_{\varepsilon} \right),$$

obtained from the Euler–Lagrange equation (3.2) for  $(S_{\varepsilon})$ . From the definition of  $v_{\varepsilon}$  we obtain

(4.50) 
$$\|v_{\varepsilon}\|_{q}^{q} = \lambda_{\varepsilon}^{\frac{2(q-p)}{p-2}} \|w_{\varepsilon}\|_{q}^{q}, \qquad \|v_{\varepsilon}\|_{2}^{2} = \lambda_{\varepsilon}^{-2} \|w_{\varepsilon}\|_{2}^{2}.$$

From Lemma 4.2 and Lemma 4.3 we then derive the essential relation

(4.51) 
$$\lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}} \|v_{\varepsilon}\|_{q}^{q} = \kappa \varepsilon \lambda_{\varepsilon}^{2} \|v_{\varepsilon}\|_{2}^{2} \lesssim \sigma_{\varepsilon}$$

which leads to the following two–sided estimate.

Lemma 4.6.  $\sigma_{\varepsilon}^{-\frac{p-2}{2(q-p)}} \lesssim \lambda_{\varepsilon} \lesssim \varepsilon^{-\frac{1}{2}} \sigma_{\varepsilon}^{\frac{\varepsilon}{2}}$ .

*Proof.* Follows directly from (4.51) by observing that

(4.52) 
$$\liminf_{\varepsilon \to 0} \|v_{\varepsilon}\|_q > 0, \qquad \liminf_{\varepsilon \to 0} \|v_{\varepsilon}\|_2 > 0$$

To prove the latter, we note that by Lemma 4.5 and in view of the embedding  $L^q(B_1) \subset L^p(B_1)$  we have

$$(4.53) \ c \|v_{\varepsilon}\chi_{B_1}\|_q \ge \|v_{\varepsilon}\chi_{B_1}\|_p \ge \|W_1\chi_{B_1}\|_p - \|(W_1 - v_{\varepsilon})\chi_{B_1}\|_p = \|W_1\chi_{B_1}\|_p - o(1),$$

where here and below  $\chi_{B_R}$  is the characteristic function of  $B_R$ . Similarly, in view of the embedding  $L^p(B_1) \subset L^2(B_1)$  we obtain

(4.54) 
$$\|v_{\varepsilon}\chi_{B_1}\|_2 \ge \|W_1\chi_{B_1}\|_2 - \|(W_1 - v_{\varepsilon})\chi_{B_1}\|_2 = \|W_1\chi_{B_1}\|_2 - o(1),$$

so the assertion follows.

Using estimate (4.17), we extract from Lemma 4.6 a lower bound

(4.55) 
$$\lambda_{\varepsilon} \gtrsim \sigma_{\varepsilon}^{-\frac{1}{2}\frac{p-2}{q-p}} \gtrsim \begin{cases} \varepsilon^{-\frac{p-2}{2q-4}}, & N \ge 5, \\ (\varepsilon \log \frac{1}{\varepsilon})^{-\frac{1}{q-2}}, & N = 4, \\ \varepsilon^{-\frac{1}{q-4}}, & N = 3, \end{cases}$$

and an upper bound

(4.56) 
$$\lambda_{\varepsilon} \lesssim \begin{cases} \varepsilon^{-\frac{1}{2}\frac{p-2}{q-2}}, & N \ge 5, \\ \varepsilon^{-\frac{1}{q-2}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{q-4}{2q-4}}, & N = 4, \\ \varepsilon^{-\frac{q-2}{4(q-4)}}, & N = 3. \end{cases}$$

Note that for  $N \ge 5$  the above lower and upper estimates are equivalent, and as a consequence we obtain the following.

**Corollary 4.7.** Assume  $N \geq 5$ . Then  $||v_{\varepsilon}||_q$  and  $||v_{\varepsilon}||_2$  are bounded.

*Proof.* Follows from (4.51), (4.55) and (4.56).

In the lower dimensions the growth of  $||v_{\varepsilon}||_2$  is to be taken into account to obtain matching bounds, so instead of (4.56) we shall use a more explicit upper bound

(4.57) 
$$\lambda_{\varepsilon} \lesssim \frac{\varepsilon^{-1/2} \sigma_{\varepsilon}^{\frac{1}{2}}}{\|v_{\varepsilon}\|_{2}} \lesssim \|v_{\varepsilon}\|_{2}^{-1} \begin{cases} \varepsilon^{-\frac{1}{q-2}} \left(\log \frac{1}{\varepsilon}\right)^{\frac{q-4}{2q-4}}, & N=4, \\ \varepsilon^{-\frac{q-2}{4(q-4)}}, & N=3, \end{cases}$$

which is also a combination of (4.51) and (4.17).

4.6. A lower barrier. To control the norm  $||v_{\varepsilon}||_2$ , we note that

$$(4.58) \quad -\Delta v_{\varepsilon} + S_{\varepsilon} \varepsilon \lambda_{\varepsilon}^{2} v_{\varepsilon} = S_{\varepsilon} \left( v_{\varepsilon}^{p-1} - \lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}} v_{\varepsilon}^{q-1} \right) \ge -V_{\varepsilon}(x) v_{\varepsilon}, \qquad x \in \mathbb{R}^{N},$$
  
where

(4.59) 
$$V_{\varepsilon}(x) := S_{\varepsilon} \lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}} v_{\varepsilon}^{q-2}(x).$$

According to the radial estimate (3.11),

(4.60) 
$$u_{\varepsilon}(x) \le C_p |x|^{-\frac{2}{p-2}} ||u_{\varepsilon}||_p,$$

Using (4.47) and the fact that  $\lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}} \lesssim \sigma_{\varepsilon} \to 0$  by Lemmas 4.1 and 4.6, for sufficiently small  $\varepsilon > 0$  we obtain (4.61)

$$V_{\varepsilon}(x) = S_{\varepsilon}\lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}}v_{\varepsilon}^{q-2}(x) \le S_{\varepsilon}\lambda_{\varepsilon}^{-\frac{2(q-p)}{p-2}}C_{p}^{q-2}\|v_{\varepsilon}\|_{p}^{q-2}|x|^{-\frac{2(q-2)}{p-2}} \le C|x|^{-\frac{2(q-2)}{p-2}},$$

where the constant C > 0 does not depend on  $\varepsilon$  or x. Therefore, for small  $\varepsilon > 0$  solutions  $v_{\varepsilon} > 0$  satisfy the linear inequality

(4.62) 
$$-\Delta v_{\varepsilon} + V_0(x)v_{\varepsilon} + S_{\varepsilon}\varepsilon\lambda_{\varepsilon}^2 v_{\varepsilon} \ge 0, \qquad x \in \mathbb{R}^N,$$

where  $V_0(x) := C|x|^{-\frac{2(q-2)}{p-2}}$ .

**Lemma 4.8.** There exists R > 0 and c > 0 such that for all small  $\varepsilon > 0$ 

(4.63) 
$$v_{\varepsilon}(x) \ge c|x|^{-(N-2)}e^{-\sqrt{\varepsilon}S_{\varepsilon}\lambda_{\varepsilon}|x|} \qquad (|x| > R).$$

*Proof.* Define the barrier

(4.64) 
$$h_{\varepsilon}(x) := \left(|x|^{-(N-2)} + |x|^{\beta}\right) e^{-\sqrt{\varepsilon S_{\varepsilon}} \lambda_{\varepsilon} |x|},$$

where  $\beta~<0$  is fixed in such a way that

(4.65) 
$$-(N-2) - \frac{2(q-p)}{p-2} < \beta < -(N-2)$$

and the value of c>0 will be specified later. A direct computation then shows that for some  $R\gg 1$  one get

$$(4.66) \quad -\Delta h_{\varepsilon} + V_{0}(x)h_{\varepsilon} + \varepsilon\lambda_{\varepsilon}^{2}h_{\varepsilon} \\ = \left\{ -\beta(\beta + N - 2)|x|^{\beta - 2} + C\left(|x|^{-(N-2)} + |x|^{\beta}\right)|x|^{-2\frac{q-2}{p-2}} + \sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon}\left((2\beta + N - 1)|x|^{\beta - 1} + (3 - N)|x|^{-(N-1)}\right) \right\} e^{-\sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon}|x|} \\ \le \left\{ -\beta(\beta + N - 2)|x|^{\beta - 2} + 2C|x|^{-(N-2)-2\frac{q-2}{p-2}} \right\} e^{-\sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon}|x|} \le 0,$$

for all |x| > R, where  $R \gg 1$  can be chosen independent of  $\varepsilon > 0$ .

Note that Lemmas 4.5 and 3.1 imply

(4.67) 
$$\| (v_{\varepsilon} - W_1) \chi_{B_{2R} \setminus B_{R/2}} \|_{\infty} \to 0,$$

and hence

(4.68) 
$$v_{\varepsilon}(R) \ge \frac{1}{2}W_1(R)$$

for all sufficiently small  $\varepsilon > 0$ . Choose c > 0 so that

(4.69) 
$$c(R^{-(N-2)} + R^{\beta}) \le \frac{1}{2}W_1(R).$$

Then

(4.70) 
$$v_{\varepsilon} \ge ch_{\varepsilon} \quad \text{for } |x| > R,$$

by the comparison principle for the operator  $-\Delta + V_0 + \varepsilon \lambda_{\varepsilon}^2$ , (see, e.g., [2, Theorem 2.7]).

4.7. Case N = 3 and N = 4 completed. We shall apply Lemma 4.8 to obtain matching estimates on the blow-up of  $||v_{\varepsilon}||_2$  in low dimensions.

**Lemma 4.9.** If N = 3 then  $||v_{\varepsilon}||_2^2 \gtrsim \frac{1}{\sqrt{\varepsilon}\lambda_{\varepsilon}}$ .

*Proof.* Assuming N = 3 we directly calculate from Lemma 4.8,

(4.71) 
$$\|v_{\varepsilon}\|_{2}^{2} \ge \int_{\mathbb{R}^{3} \setminus B_{R}} |v_{\varepsilon}|^{2} dx \ge \int_{R}^{\infty} c^{2} e^{-2\sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon}r} dr \ge \frac{C}{\sqrt{\varepsilon}\lambda_{\varepsilon}},$$

which is what is required.

As an immediate corollary, using (4.57), we obtain an upper estimate of  $\lambda_{\varepsilon}$  which matches the lower bound of (4.55) in the case N = 3.

**Corollary 4.10.** If N = 3 then  $\lambda_{\varepsilon} \lesssim \varepsilon^{-\frac{1}{q-4}}$ .

Next we consider the case N = 4.

**Lemma 4.11.** If N = 4 then  $||v_{\varepsilon}||_2^2 \gtrsim \log \frac{1}{\sqrt{\varepsilon}\lambda_{\varepsilon}}$ .

*Proof.* Assuming N = 4 we directly calculate using Lemma 4.8,

$$(4.72) \quad \|v_{\varepsilon}\|_{2}^{2} \ge \int_{\mathbb{R}^{4} \setminus B_{R}} |v_{\varepsilon}|^{2} \, dx \ge \int_{R}^{\infty} c^{2} r^{-1} e^{-2\sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon} r} \, dr = c^{2} \Gamma(0, 2\sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon} R),$$

where

(4.73) 
$$\Gamma(0,t) = -\log(t) - \gamma + O(t), \qquad t \searrow 0,$$

is the incomplete Gamma function and  $\gamma \approx 0.5772$  is the Euler constant [1]. Hence we obtain for sufficiently small  $\varepsilon$ :

(4.74) 
$$\|v_{\varepsilon}\|_{2}^{2} \ge c^{2}(-\log(2\sqrt{\varepsilon S_{\varepsilon}}\lambda_{\varepsilon}R) - \gamma) \ge C\log\left(\frac{1}{\sqrt{\varepsilon}\lambda_{\varepsilon}}\right),$$

which is what is required.

**Corollary 4.12.** If 
$$N = 4$$
 then  $\lambda_{\varepsilon} \lesssim \left(\varepsilon \log \frac{1}{\varepsilon}\right)^{-\frac{1}{q-2}}$ .

*Proof.* An immediate corollary of (4.51) and (4.17) is the relation

(4.75) 
$$C\varepsilon\lambda_{\varepsilon}^{2}\log\frac{1}{\sqrt{\varepsilon}\lambda_{\varepsilon}} \le \left(\varepsilon\log\frac{1}{\varepsilon}\right)^{\frac{q-4}{q-2}}$$

Note that  $\varepsilon^{\delta_1} \leq \sqrt{\varepsilon}\lambda_{\varepsilon} \leq \varepsilon^{\delta_2}$  for some  $\delta_{1,2} \geq 0$  and  $\varepsilon$  small enough, which is a consequence of (4.56) and (4.55). Therefore,

(4.76) 
$$\log \frac{1}{\sqrt{\varepsilon}\lambda_{\varepsilon}} \sim \log \frac{1}{\varepsilon},$$

and the conclusion follows.

4.8. Further estimates. The results in the previous section could be used in a standard way to improve upon some earlier estimates.

An immediate consequence of the sharp upper estimates of  $\lambda_{\varepsilon}$  is the following.

# **Corollary 4.13.** $||v_{\varepsilon}||_q = O(1).$

The boundedness of the  $L^q$  norm also allows to reverse estimates of  $||v_{\varepsilon}||_2$  via (4.51).

### Corollary 4.14.

(4.77) 
$$\|v_{\varepsilon}\|_{2}^{2} = \begin{cases} O(1), & N \ge 5, \\ O(\log \frac{1}{\varepsilon}), & N = 4, \\ O(\varepsilon^{-\frac{1}{2}\frac{q-6}{q-4}}), & N = 3. \end{cases}$$

We now prove that the  $L^q$  bound also implies an  $L^{\infty}$  bound.

**Lemma 4.15.**  $||v_{\varepsilon}||_{\infty} = O(1).$ 

*Proof.* Note that by  $(R_{\varepsilon}^*)$  the function  $v_{\varepsilon}$  is a positive solution of the linear inequality

(4.78)  $-\Delta v_{\varepsilon} - V_{\varepsilon}(x)v_{\varepsilon} \le 0, \qquad x \in \mathbb{R}^{N},$ 

where

(4.79) 
$$V_{\varepsilon}(x) := S_{\varepsilon} v_{\varepsilon}^{p-2}(x).$$

From the radial estimate (3.11) we obtain

(4.80) 
$$v_{\varepsilon}(x) \le C_q \|v_{\varepsilon}\|_q |x|^{-\frac{\gamma}{q}}.$$

Hence, using Corollary 4.13 we obtain

(4.81) 
$$V_{\varepsilon}(x) \le S_{\varepsilon} C_q^{p-2} ||v_{\varepsilon}||_q^{p-2} |x|^{-\frac{N(p-2)}{q}} \le C_* |x|^{-\frac{2p}{q}},$$

for some constant  $C_* > 0$  which does not depend on  $\varepsilon$  or x. As a consequence,  $v_{\varepsilon}$  is a positive solution of the linear inequality

(4.82) 
$$-\Delta v_{\varepsilon} - V_*(x)v_{\varepsilon} \le 0, \qquad x \in \mathbb{R}^N,$$

where  $V_*(x) = C_*|x|^{-\frac{2p}{q}} \in L^s_{loc}(\mathbb{R}^N)$ , for some s > N/2. The result can then be concluded by the weak Harnack inequality for subsolutions of (4.82) (cf. [25, Remark 5.1 on p. 226]). Here we give an elementary proof that also works in the present context. Integrating the inequality in (4.82) over a ball and applying divergence theorem, by monotonic decrease of  $v_{\varepsilon}(x)$  in |x| we have

(4.83) 
$$|\nabla v_{\varepsilon}(x)| \leq \frac{C}{|x|^{N-1}} \int_{B_{|x|}(0)} V_{*}(y) v_{\varepsilon}(y) \, dy \leq C' v_{\varepsilon}(0) |x|^{1-\frac{2p}{q}},$$

for some C, C' > 0 independent of  $\varepsilon$  or x. Integrating again along the straight line from 0 to  $x_0$ , we obtain

(4.84) 
$$v_{\varepsilon}(0) \le v_{\varepsilon}(x_0) + C'' v_{\varepsilon}(0) |x_0|^{\frac{2(q-p)}{q}},$$

for some C'' > 0 independent of  $\varepsilon$  or x. We then conclude by choosing  $|x_0|$  sufficiently small independently of  $\varepsilon$ , using (4.80) and Corollary 4.13.

A standard consequence of the  $L^\infty$  bound and elliptic regularity theory is the following convergence statement.

**Corollary 4.16.**  $v_{\varepsilon} \to U_1$  in  $C^2(\mathbb{R}^N)$  and  $L^s(\mathbb{R}^N)$  for any  $s \ge p$ . In particular, (4.85)  $v_{\varepsilon}(0) \simeq W_1(0).$  Proof. Indeed, a consequence of the  $L^{\infty}$  bound of Lemma 4.15 and convergence in  $D^1(\mathbb{R}^N)$  via compactness result for monotone radial functions in Lemma 3.1 is convergence in  $L^s(\mathbb{R}^N)$  for any  $s \geq p$ . Then Calderón–Zygmund estimate [15, Theorem 9.11] implies convergence in  $W^{2,s}_{loc}(\mathbb{R}^N)$  and, hence, by Sobolev embedding also in  $C^{1,\alpha}_{loc}(\mathbb{R}^N)$ . Since the nonlinearity in  $(R^*_{\varepsilon})$  is smooth, using Schauder's estimates [15, Theorem 6.2, 6.6] we conclude convergence in  $C^2_{loc}(\mathbb{R}^N)$ . Finally, taking into account that the constants in Schauder estimates are uniform with respect to translations, we deduce convergence in  $C^2(\mathbb{R}^N)$ .

Taking into account that

(4.86) 
$$u_{\varepsilon}(0) \sim \lambda_{\varepsilon}^{-\frac{\tau}{p-2}} v_{\varepsilon}(0)$$

we can use (4.85) to estimate the amplitude of  $u_{\varepsilon}(0)$  to derive (2.13), which completes the proof of Theorem 2.5.

5. Supercritical case  $p > p^*$ .

5.1. The limit equation. For  $p > p^*$  the limit equation

(P<sub>0</sub>) 
$$-\Delta u - |u|^{p-2}u + |u|^{q-2}u = 0 \quad \text{in } \mathbb{R}^{N},$$

admits a unique positive radial ground state solution  $u_0 \in D^1(\mathbb{R}^N)$ . Further, it is known that  $u_0 \in C^2(\mathbb{R}^N)$ ,  $u_0(x)$  is monotone decreasing function of |x|, and there exists  $C_0 > 0$  such that

(5.1) 
$$\lim_{|x| \to \infty} |x|^{N-2} u_0(x) = C_0 > 0,$$

see [5, Theorem 4] for the existence, or [20, 21] for the existence and asymptotic decay, and [21, 18] for the uniqueness proofs.

Similarly to (3.6), the ground state  $u_0$  admits a variational characterization in the Sobolev space  $D^1(\mathbb{R}^N)$  via the rescaling

(5.2) 
$$u_0(x) := w_0\left(\frac{x}{\sqrt{S_0}}\right),$$

where  $w_0 > 0$  is the radial (i.e., depending only on |x|) minimizer of the constrained minimization problem

(S<sub>0</sub>) 
$$S_0 := \inf \left\{ \int_{\mathbb{R}^N} |\nabla w|^2 \, dx \, \middle| \, w \in D^1(\mathbb{R}^N), \, p^* \int_{\mathbb{R}^N} F_0(w) \, dx \, = 1 \right\}$$

where  $F_0$  is defined by (3.1) (see [5, Section 5]). Similarly to (3.2)–(3.5), one concludes that the minimizer  $w_0$  solves the Euler–Lagrange equation

(5.3) 
$$-\Delta w_0 = S_0 (|w_0|^{p-2} w_0 - |w_0|^{q-2} w_0) \quad \text{in } \mathbb{R}^N$$

Further,  $w_0$  satisfies Nehari's identity

(5.4) 
$$\int_{\mathbb{R}^N} |\nabla w_0|^2 \, dx = S_0 \int \left( |w_0|^p - |w_0|^q \right) \, dx$$

and Pokhozhaev's identity (see e.g. [5, Proposition 1])

(5.5) 
$$\int_{\mathbb{R}^N} |\nabla w_0|^2 \, dx = S_0 p^* \int \left(\frac{|w_0|^p}{p} - \frac{|w_0|^q}{q}\right) \, dx.$$

Taking into account that  $\|\nabla w_0\|_2^2 = S_0$ , we then derive from Nehari and Pokhozhaev's identities the relation

(5.6) 
$$\|w_0\|_p^p - \|w_0\|_q^q = \frac{p^*}{p} \|w_0\|_p^p - \frac{p^*}{q} \|w_0\|_q^q = 1,$$

which leads to the explicit expressions

(5.7) 
$$\|w_0\|_p^p = \frac{(q-p^*)p}{(q-p)p^*}, \qquad \|w_0\|_q^q = \frac{(p-p^*)q}{(q-p)p^*}$$

Remark 5.1. Note that the arguments leading to (5.7) also give non-existence of non-trivial weak solutions  $u \in D^1(\mathbb{R}^N) \cap L^p(\mathbb{R}^N) \cap L^q(\mathbb{R}^N)$  of problem  $(P_0)$  in the case 2 and <math>q > p.

5.2. Energy and norms estimates. To control the relations between  $S_{\varepsilon}$  and  $S_0$  it is convenient to consider the equivalent to  $(S_0)$  scaling invariant quotient

(5.8) 
$$S_0(w) := \frac{\int_{\mathbb{R}^N} |\nabla w|^2 \, dx}{\left(p^* \int_{\mathbb{R}^N} F_0(w) \, dx\right)^{\frac{N-2}{N}}}, \qquad w \in D^1(\mathbb{R}^N), \quad \int_{\mathbb{R}^N} F_0(w) \, dx > 0.$$

Then

(5.9) 
$$S_0 = \inf_{\substack{w \in D^1(\mathbb{R}^N) \\ F_0(w) > 0}} \mathcal{S}_0(w)$$

**Lemma 5.2.**  $0 < S_{\varepsilon} - S_0 \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .

*Proof.* To show that  $S_0 < S_{\varepsilon}$  simply note that

(5.10) 
$$S_0 \leq \mathcal{S}_0(w_{\varepsilon}) < \mathcal{S}_{\varepsilon}(w_{\varepsilon}) = S_{\varepsilon}.$$

To control  $S_{\varepsilon}$  from above we will use the minimizer  $w_0$  as a test function for  $(S_{\varepsilon})$ . In view of (5.1), we have  $w_0 \in L^2(\mathbb{R}^N)$  if and only if  $N \geq 5$ . Therefore we shall consider the higher and lower dimensions separately.

CASE  $N \geq 5$ . Testing  $(S_{\varepsilon})$  against  $w_0$ , we obtain

(5.11) 
$$S_{\varepsilon} \leq S_{\varepsilon}(w_0) \leq \frac{S_0}{\left(1 - \varepsilon \|w_0\|_{L^2(\mathbb{R}^N)}^2\right)^{\frac{N-2}{N}}} \leq S_0 + O(\varepsilon),$$

which proves the claim for  $N \geq 5$ .

To consider the lower dimensions, given R > 1 we introduce a cutoff function  $\eta_R \in C_c^{\infty}(\mathbb{R})$  such that  $\eta_R(r) = 1$  for |r| < R,  $0 < \eta(r) < 1$  for R < |r| < 2R,  $\eta_R(r) = 0$  for |r| > 2R and  $|\eta'(r)| \le 2/R$ . Then taking into account (5.1), for  $s > \frac{N}{N-2}$  we compute

(5.12) 
$$\int_{\mathbb{R}^N} |\nabla(\eta_R w_0)|^2 = S_0 + O(R^{-(N-2)}),$$

(5.13) 
$$\int_{\mathbb{R}^N} |\eta_R w_0|^s \, dx = \|w_0\|_{L^s(\mathbb{R}^N)}^s \left(1 - O(R^{N-s(N-2)})\right),$$

(5.14) 
$$\int_{\mathbb{R}^N} |\eta_R w_0|^2 = \begin{cases} O(\log(R)), & N = 4, \\ O(R), & N = 3. \end{cases}$$

CASE N = 4. Let  $R = \varepsilon^{-1}$ . Testing  $(S_{\varepsilon})$  against  $\eta_R w_0$  and using the fact that p > 4, we obtain

(5.15) 
$$S_{\varepsilon} \leq S_{\varepsilon}(w_{0}) \leq \frac{S_{0} + O(R^{-2})}{\left(1 - O(R^{-4}) - \varepsilon O(\log R)\right)^{1/2}} \\ \leq \frac{S_{0} + O(\varepsilon^{2})}{\left(1 - O(\varepsilon^{4}) - O(\varepsilon \log \frac{1}{\varepsilon})\right)^{\frac{1}{2}}} \leq S_{0} + O(\varepsilon \log \frac{1}{\varepsilon}),$$

which proves the claim.

CASE N = 3. Let  $R = \varepsilon^{-1/2}$ . Testing  $(S_{\varepsilon})$  against  $\eta_R w_0$  and using the fact that p > 6, we obtain

(5.16) 
$$S_{\varepsilon} \leq S_{\varepsilon}(w_{0}) \leq \frac{S_{0} + O(R^{-1})}{\left(1 - O(R^{-6}) - \varepsilon O(R)\right)^{1/3}}$$
$$\leq \frac{S_{0} + O(\varepsilon^{1/2})}{\left(1 - O(\varepsilon^{3/2}) - O(\varepsilon^{1/2})\right)^{1/3}} \leq S_{0} + O(\varepsilon^{1/2}),$$
which completes the proof.

which completes the proof.

**Lemma 5.3.**  $||w_{\varepsilon}||_{\infty} \leq 1$  and  $||w_{\varepsilon}||_{s} \leq 1$  for all  $s > p^{*}$ .

*Proof.* In view of 
$$(1.3)$$
 and  $(3.6)$  we have

(5.17) 
$$\|w_{\varepsilon}\|_{\infty} = \|u_{\varepsilon}\|_{\infty} \le 1.$$

Using Sobolev's inequality and Lemma 5.2 we also obtain

(5.18) 
$$\|w_{\varepsilon}\|_{p^*}^2 \leq S_*^{-1} \|\nabla w_{\varepsilon}\|_2^2 = S_*^{-1} S_{\varepsilon} = S_*^{-1} S_0 (1+o(1)).$$

Then for every  $s > p^*$ 

$$\|w_{\varepsilon}\|_{s}^{s} \leq \|w_{\varepsilon}\|_{p^{*}}^{p^{*}},$$

so the assertion follows.

Lemma 5.4.  $\varepsilon \|w_{\varepsilon}\|_2^2 \to 0.$ 

*Proof.* Since  $w_{\varepsilon}$  is a minimizer of  $(S_{\varepsilon})$ , we have

(5.20) 
$$1 = p^* \int_{\mathbb{R}^N} F_{\varepsilon}(w_{\varepsilon}) dx = p^* \int_{\mathbb{R}^N} F_0(w_{\varepsilon}) dx - p^* \frac{\varepsilon}{2} ||w_{\varepsilon}||_2^2.$$

Therefore

(5.21) 
$$S_0(w_{\varepsilon}) = \frac{\|\nabla w_{\varepsilon}\|_2^2}{\left(p^* \int_{\mathbb{R}^N} F_0(w) \, dx\right)^{\frac{N-2}{N}}} = \frac{S_{\varepsilon}}{\left(1 + \frac{p^*}{2}\varepsilon \|w_{\varepsilon}\|_2^2\right)^{\frac{N-2}{N}}}.$$

Assume to the contrary of the statement of the Lemma that  $\limsup_{\varepsilon \to 0} \varepsilon \|w_{\varepsilon}\|_{2}^{2} =$ m > 0. Then by Lemma 5.2 for any sequence  $\varepsilon_n \to 0$  we obtain

(5.22) 
$$S_0 \leq S_0(w_{\varepsilon_n}) = \frac{S_{\varepsilon_n}}{\left(1 + \frac{p^*}{2}\varepsilon_n \|w_{\varepsilon_n}\|_2^2\right)^{\frac{N-2}{N}}} \leq \frac{S_0(1+o(1))}{1 + \frac{p^*}{2}m} < S_0,$$
  
a contradiction.

a contradiction.

5.3. **Proof of Theorem 2.3.** Consider a sequence of  $\varepsilon_n \to 0$ . Since  $\|\nabla w_{\varepsilon_n}\|_2^2 =$  $S_{\varepsilon_n} \to S_0$ , the sequence  $(\varepsilon_n)$  contains a subsequence, still denoted  $(\varepsilon_n)$ , such that  $w_{\varepsilon_n} \to \bar{w}$  in  $D^1(\mathbb{R}^N)$  and  $w_{\varepsilon_n} \to \bar{w}$  a.e. in  $\mathbb{R}^N$ , (5.23)

where  $\bar{w} \in D^1(\mathbb{R}^N)$  is a radial function. By Lemma 5.3, the sequence  $(w_{\varepsilon_n})$  is bounded in  $L^{p^*}(\mathbb{R}^N)$  and  $L^{\infty}(\mathbb{R}^N)$ . Using Lemma 3.1 and Sobolev inequality, we also obtain a uniform bound

(5.24) 
$$w_{\varepsilon}(x) \le C|x|^{-\frac{N-2}{2}} \|\nabla w_{\varepsilon}\|_{2} \le 2C|x|^{-\frac{N-2}{2}} S_{0},$$

for  $\varepsilon$  sufficiently small. Using Lemma 3.1 we conclude that

(5.25) 
$$w_{\varepsilon_n} \to \bar{w} \text{ in } L^s(\mathbb{R}^N) \text{ for any } s \in (p^*, \infty).$$

18

Taking into account Lemma 5.4 and (5.20) we also obtain

(5.26) 
$$\int_{\mathbb{R}^N} F_0(\bar{w}) dx = \lim_{n \to \infty} \int_{\mathbb{R}^N} F_0(w_{\varepsilon_n}) dx = \lim_{n \to \infty} \left( 1 + p^* \frac{\varepsilon_n}{2} \|w_{\varepsilon_n}\|_2^2 \right) = 1.$$

By the weak lower semicontinuity we also conclude that

(5.27) 
$$\|\nabla \bar{w}\|_2^2 \le \liminf_{n \to \infty} \|\nabla w_{\varepsilon_n}\|_2^2 = S_0,$$

that is  $\bar{w}$  is a minimizer for  $(S_0)$ . By the uniqueness of the radial minimizer of  $(S_0)$  we conclude that  $\bar{w} = w_0$ .

We now claim that 
$$(w_{\varepsilon_n})$$
 converges strongly to  $w_0$  in  $D^1(\mathbb{R}^N)$ . Indeed, we have

(5.28) 
$$\|\nabla(w_{\varepsilon_n} - w_0)\|_2^2 = \|\nabla w_{\varepsilon_n}\|_2^2 + \|\nabla w_0\|_2^2 - 2\int_{\mathbb{R}^N} \nabla w_{\varepsilon_n} \cdot \nabla w_0 \, dx$$
$$= S_{\varepsilon_n} + S_0 - 2\int_{\mathbb{R}^N} \nabla w_{\varepsilon_n} \cdot \nabla w_0 \, dx.$$

Estimating the last term and taking into account (5.3), (5.6) and the fact that by Hölder inequality

(5.29) 
$$\left| \int_{\mathbb{R}^N} f_0(w_0)(w_{\varepsilon} - w_0) dx \right| \leq \|f_0(w_0)\|_{\frac{p}{p-1}} \|w_{\varepsilon} - w_0\|_p$$
$$\leq C \|w_0\|_p^{p-1} \|w_{\varepsilon} - w_0\|_p \to 0,$$

we obtain

(5.30) 
$$\int_{\mathbb{R}^N} \nabla w_{\varepsilon_n} \cdot \nabla w_0 \, dx = S_0 \int_{\mathbb{R}^N} f_0(w_0) w_{\varepsilon_n} \, dx$$
$$= S_0 \int_{\mathbb{R}^N} f_0(w_0) w_0 \, dx + S_0 \int_{\mathbb{R}^N} f_0(w_0) (w_{\varepsilon_n} - w_0) dx$$
$$= S_0 (1 + o(1)),$$

which proves the claim.

Since  $(w_{\varepsilon_n})$  converges to  $w_0$  in  $D^1(\mathbb{R}^N)$  and in  $L^s(\mathbb{R}^N)$  for any  $s \ge p^*$ , similarly to the proof of Corollary 4.16 by the standard elliptic regularity we conclude that  $(w_{\varepsilon_n})$  converges to  $w_0$  in  $C^2(\mathbb{R}^N)$ . The proof of of Theorem 2.3 is then completed by taking into account the uniqueness of  $w_0$ .

5.4. Remarks on a slightly supercritical limit problem. Here we discuss the asymptotic behavior as  $p \downarrow p^*$  of the minimizer  $w_0$  of the limit variational problem  $(S_0)$ . For convenience, set  $\delta := p - p^* > 0$ . To highlight the dependance on  $\delta$ , in this section we denote the ground state energy in (5.9) by  $S_0^{\delta}$ , while  $w_0^{\delta}$  will be used to denote the corresponding minimizer. Also, in this section the asymptotic notation such as  $\leq$ , etc., is in terms of  $\delta \to 0$ .

The following summarizes our results regarding the asymptotic behavior of  $w_0^{\delta}$  as  $\delta \downarrow 0$ .

**Proposition 5.5.**  $0 < S_0^{\delta} - S_* \to 0$  for  $\delta \downarrow 0$ . In addition, it holds

(5.31) 
$$\delta^{\frac{1}{q-p^*}} \lesssim w_0^{\delta}(0) \lesssim \delta^{\frac{1}{q+N}},$$

and, provided that  $q > \frac{N(N+2)}{2(N-2)}$ ,

(5.32) 
$$w_0^{\delta}(0) \sim \delta^{\frac{1}{q-p^*}}.$$

Let us note, however, that the asymptotic of  $w_0^{\delta}(0)$  for general values of q is open, and numerical evidence suggests that the conclusion of (5.32) is *false* for q sufficiently close to  $p^*$ .

To prove Proposition 5.5, we first establish a few basic estimates for the behavior of the minimizer of the quotient in (5.8) as  $\delta \to 0$ .

Lemma 5.6.  $||w_0^{\delta}||_{\infty} \leq 1$ ,  $||\nabla w_0^{\delta}||_2 \lesssim 1$ ,  $||w_0^{\delta}||_p \lesssim 1$  and  $||w_0^{\delta}||_q \lesssim \delta$ .

Proof. The first inequality is an immediate consequence of  $||u_0||_{\infty} \leq 1$  and (5.2). To prove the second estimate, consider a suitable fixed test function  $w \in C_0^{\infty}(\mathbb{R}^N)$  satisfying  $0 \leq w \leq 1$ . Then from (5.8) and (5.9) we can conclude that  $S_0^{\delta} \leq 1$  as  $\delta \to 0$ , implying the result. The last two inequalities are immediate consequences of (5.7).

We now establish a rough upper bound on the amplitude of  $w_0^{\delta}$ .

Lemma 5.7.  $||w_0^{\delta}||_{\infty} \lesssim \delta^{\frac{1}{q+N}}$ .

*Proof.* In view of the gradient estimate of Lemma 5.6, by Calderón–Zygmund inequality [15, Theorem 9.11] applied to  $w_0^{\delta}$  solving (5.3) we conclude that  $\|w_0^{\delta}\|_{W^{2,p}_{loc}(\mathbb{R}^N)}$ is uniformly bounded and, hence,  $\|\nabla w_0^{\delta}\|_{\infty} \leq C$  for some C > 0 independent of  $\delta$  for sufficiently small  $\delta > 0$ . This yields the following estimate for some c > 0independent of  $\delta$ :

(5.33) 
$$c \|w_0^{\delta}\|_{\infty}^{q+N} \le \frac{1}{2^q} \|w_0^{\delta}\|_{\infty}^q |B_R(0)| \le \int_{B_R(0)} |w_0^{\delta}|^q dx \le \|w_0^{\delta}\|_q^q,$$

where  $R = \|w_0^{\delta}\|_{\infty}/(2C)$ , and we used monotonicity of  $w_0^{\delta}(x)$  in |x|. The result then follows from the fact that  $\|w_0^{\delta}\|_q^q \sim \delta$  by (5.7).

The relations in (5.7) immediately lead to the following lower bound on  $w_0^{\delta}(0)$ . Lemma 5.8.  $\|w_0^{\delta}\|_{\infty} \gtrsim \delta^{\frac{1}{q-p^*}}$ .

*Proof.* Indeed, by (5.7) we have

(5.34) 
$$\delta \|w_0^{\delta}\|_p^p \le \frac{p}{q}(q-p^*)\|w_0^{\delta}\|_{\infty}^{q-p^*-\delta}\|w_0^{\delta}\|_p^p$$

and the result follows from  $||w_0^{\delta}||_p^p > 0$  and smallness of  $\delta$ .

Importantly, for sufficiently large q we can prove a matching upper bound, yielding the precise asymptotic behavior of the minimizer's amplitude as  $\delta \to 0$ .

**Lemma 5.9.** If 
$$q > \frac{N(N+2)}{2(N-2)}$$
 then  $||w_0^{\delta}||_{\infty} \lesssim \delta^{\frac{1}{q-p^*}}$ .

Proof. In view of Lemmas 3.1 and 5.6 and Sobolev inequality, we have

(5.35) 
$$w_0^{\delta}(x) \leq \min \left\{ C_{p^*} |x|^{-\frac{N}{p^*}} \|w_0^{\delta}\|_{p^*}, C_q |x|^{-\frac{N}{q}} \|w_0^{\delta}\|_q \right\},$$
$$\lesssim \min \left\{ |x|^{-\frac{N-2}{2}}, \delta^{\frac{1}{q}} |x|^{-\frac{N}{q}} \right\}.$$

In view of (5.1), (5.2) and Lemma 5.3, we can apply Newtonian kernel to  $(P_0^{\delta})$ . We obtain

(5.36) 
$$w_0^{\delta}(x) = S_0^{\delta} A_N \int_{\mathbb{R}^N} \frac{(w_0^{\delta}(y))^{p-1} - (w_0^{\delta}(y))^{q-1}}{|x-y|^{N-2}} dy,$$

where  $A_N = \frac{\Gamma((N-2)/2)}{4\pi^{N/2}}$ . In particular, for  $q > \frac{N(N+2)}{2(N-2)}$  we have (with a slight abuse of notation)

$$(5.37) w_0^{\delta}(0) = \frac{S_0^{\delta}}{N-2} \int_0^{\infty} \left( (w_0^{\delta}(r))^{p-1} - (w_0^{\delta}(r))^{q-1} \right) r \, dr$$

$$\leq S_*(1+o(1)) \int_0^{\infty} (w_0^{\delta}(r))^{p^*-1} r \, dr$$

$$\lesssim \int_0^{\infty} \min\left\{ r^{-\frac{N-2}{2}}, \delta^{\frac{1}{q}} r^{-\frac{N}{q}} \right\}^{\frac{N+2}{N-2}} r \, dr,$$

$$\lesssim \int_R^{\infty} r^{-\frac{N}{2}} \, dr + \delta^{\frac{N+2}{q(N-2)}} \int_0^R r^{1-\frac{N(N+2)}{q(N-2)}} \, dr,$$

$$\lesssim R^{-\frac{N-2}{2}} + \delta^{\frac{N+2}{q(N-2)}} R^{2-\frac{N(N+2)}{q(N-2)}}.$$

Minimizing for  $q > \frac{N(N+2)}{2(N-2)}$  the function

(5.38) 
$$\psi_{\delta}(R) := R^{-\frac{N-2}{2}} + \delta^{\frac{N+2}{q(N-2)}} R^{2-\frac{N(N+2)}{q(N-2)}}$$

we obtain

(5.39) 
$$\min_{R>0} \psi_{\delta}(R) = \psi_{\delta}(R_*) \sim \delta^{\frac{1}{q-p^*}},$$

where  $R_* \sim \delta^{-\frac{2}{(N-2)(q-p^*)}}$ .

We also establish the energy convergence estimate.

**Lemma 5.10.**  $0 < S_0^{\delta} - S_* \to 0 \text{ as } \delta \to 0.$ 

*Proof.* Taking into account (5.7) we obtain

(5.40) 
$$S_* \leq S_*(w_0^{\delta}) = \frac{\|\nabla w_0^{\delta}\|_2^2}{\|w_0^{\delta}\|_p^2} = \left(\frac{p^*(q-p)}{p(q-p^*)}\right)^{2/p} S_0^{\delta} < S_0^{\delta}.$$

To control  $S_0^{\delta}$  from above we will use the Sobolev minimizers  $(W_{\lambda})_{\lambda>0}$  as a family of test function for  $(S_0^{\delta})$ . Using (4.3) we obtain

(5.41) 
$$S_0^{\delta}(W_{\lambda}) = \frac{S_*}{\left(\frac{p^*}{p}\lambda^{-\frac{N-2}{2}\delta} \|W_1\|_p^p - \frac{p^*}{q}\lambda^{-\frac{N-2}{2}(q-p^*)} \|W_1\|_q^q\right)^{\frac{N-2}{N}}}$$

To minimize the right hand side of (5.41), we need to maximize for  $\lambda > 0$  the scalar function

(5.42) 
$$\psi(\lambda) := \frac{p^*}{p} \|W_1\|_p^p \lambda^{-\frac{1}{2}(N-2)(p-p^*)} - \frac{p^*}{q} \|W_1\|_q^q \lambda^{-\frac{1}{2}(N-2)(q-p^*)}.$$

It is easy to see that  $\psi$  achieves its maximum at

(5.43) 
$$\lambda_* := \left(\frac{p(q-p^*) \|W_1\|_q^q}{q(p-p^*) \|W_1\|_p^p}\right)^{\frac{2}{N-2}\frac{1}{q-p}}$$

and

(5.44) 
$$\psi(\lambda_*) = A(p, p^*, q) \|W_1\|_p^{\frac{p(q-p^*)}{q-p}} \|W_1\|_q^{-\frac{q(p-p^*)}{q-p}},$$

where

(5.45) 
$$A(p,p^*,q) := p^* p^{-\frac{q-p^*}{q-p}} q^{\frac{p-p^*}{q-p}} \left\{ \left(\frac{q-p^*}{p-p^*}\right)^{-\frac{p-p^*}{q-p}} - \left(\frac{q-p^*}{p-p^*}\right)^{-\frac{q-p^*}{q-p}} \right\} > 0.$$

In particular, when  $\delta = p - p^* \to 0$  we have  $A(p, p^*, q) \simeq 1$  and  $\psi(\lambda_*) \simeq 1$ , so

(5.46) 
$$S_0^{\delta} \le S_0^{\delta}(W_{\lambda_*}) = (\psi(\lambda_*))^{-\frac{N-2}{N}} S_* = S_* (1+o(1)),$$

which completes the proof.

Remark 5.11. Instead of  $W_{\lambda}$  we can use rescalings of an arbitrary function  $w \in D^1(\mathbb{R}^N) \cap L^q(\mathbb{R}^N)$  as a family of test function in (5.41). Then, taking into account that by Sobolev imbedding  $w \in L^{p^*}(\mathbb{R}^N)$  and, hence, by interpolation we have  $w \in L^p(\mathbb{R}^N)$  as well, the above argument with generic  $q > p > p^*$  leads to

(5.47) 
$$S_0(p, p^*, q) A^{\frac{2}{p^*}}(p, p^*, q) \|w\|_p^{\frac{2p(q-p^*)}{p^*(q-p)}} \le \|\nabla w\|_2^2 \|w\|_q^{\frac{2q(p-p^*)}{p^*(q-p)}},$$

which could be interpreted as a supercritical Gagliardo–Nirenberg type inequality. Similar ideas where used in [10] to establish sharp constants in the classical Gagliardo–Nirenberg inequality, which *formally* coincides with (5.47) when  $1 < q < p < p^*$ .

## 6. Subcritical case 2 revisited: proof of Theorem 2.1.

In the subcritical case Pokhozhaev's identity implies that the limit equation  $(P_0)$  has no positive finite energy solutions. As discussed in the Introduction, to understand the asymptotic behavior of the ground states  $u_{\varepsilon}$  we consider the rescaling in (1.4), which transforms  $(P_{\varepsilon})$  into  $(R_{\varepsilon})$ , with the associated limit problem as  $\varepsilon \to 0$  given by  $(R_0)$  (see Sec. 1).

Let  $G_{\varepsilon}: \mathbb{R} \to \mathbb{R}$  be a bounded  $C^2$ -function such that

(6.1) 
$$G_{\varepsilon}(w) := \frac{1}{p} |w|^p - \frac{1}{2} |w|^2 - \frac{\varepsilon^{\frac{q-p}{p-2}}}{q} |w|^q$$

for  $0 \le w \le \varepsilon^{-\frac{1}{p-2}}$ ,  $G_{\varepsilon}(w) \le 0$  for  $w > \varepsilon^{-\frac{1}{p-2}}$ , and  $G_{\varepsilon}(w) = 0$  for  $w \le 0$ . For  $\varepsilon \in [0, \varepsilon^*)$ , consider a family of the constrained minimization problems

$$(S'_{\varepsilon}) \qquad S'_{\varepsilon} := \inf\left\{ \int_{\mathbb{R}^N} |\nabla w|^2 \, dx \right| \, w \in H^1(\mathbb{R}^N), \ p^* \int_{\mathbb{R}^N} G_{\varepsilon}(w) \, dx \ = 1 \right\}$$

Note that all the problems  $(S'_{\varepsilon})$ , including the limit problem  $(S'_0)$ , are well posed in the same energy space  $H^1(\mathbb{R}^N)$ . According to [5, Theorem 2],  $(S_{\varepsilon})$  admits a radial positive minimizer  $w_{\varepsilon}$  for every  $\varepsilon \in [0, \varepsilon^*)$ . In view of its uniqueness [17], the rescaled function

(6.2) 
$$v_{\varepsilon}(x) := w_{\varepsilon}\left(\frac{x}{\sqrt{S'_{\varepsilon}}}\right),$$

coincides with the radial ground state of  $(R_{\varepsilon})$ .

In order to estimate  $S'_{\varepsilon}$ , consider the associated dilation invariant representation

(6.3) 
$$\mathcal{S}'_{\varepsilon}(w) := \frac{\int_{\mathbb{R}^N} |\nabla w|^2 \, dx}{\left(p^* \int_{\mathbb{R}^N} G_{\varepsilon}(w) \, dx\right)^{\frac{N-2}{N}}}, \qquad w \in \mathcal{M}'_{\varepsilon},$$

where  $\mathcal{M}_{\varepsilon}' := \{ 0 \le u \in H^1(\mathbb{R}^N), \int_{\mathbb{R}^N} G_{\varepsilon}(w) \, dx > 0 \}.$  Clearly (6.4)  $S_{\varepsilon}' = \inf_{w \in \mathcal{M}'_{\varepsilon}} S_{\varepsilon}'(w)$ 

and for sufficiently small  $\varepsilon$  we have

(6.5) 
$$S'_0 \leq \mathcal{S}'_0(w_{\varepsilon}) < \mathcal{S}'_{\varepsilon}(w_{\varepsilon}) = S'_{\varepsilon}.$$

Indeed, since by definition  $p^* \int_{\mathbb{R}^N} G_{\varepsilon}(w_{\varepsilon}) dx = 1$  and  $G_{\varepsilon}(s)$  is a decreasing function of  $\varepsilon$  for each s > 0, we have  $w_{\varepsilon} \in \mathcal{M}'_0$ , and the second inequality again follows

by monotonicity of  $G_{\varepsilon}(s)$  in  $\varepsilon$ . At the same time, by continuity  $w_0 \in \mathcal{M}'_{\varepsilon}$  for sufficiently small  $\varepsilon$ . Therefore, using  $w_0$  as a test function for  $(S'_{\varepsilon})$ , we obtain for sufficiently small  $\varepsilon$ 

(6.6) 
$$S'_{\varepsilon} \leq S'_{\varepsilon}(w_0) = \frac{S'_0}{\left(1 - \frac{p^*}{q} \varepsilon^{\frac{q-p}{p-2}} \|w_0\|_q^q\right)^{\frac{N-2}{N}}} \leq S'_0 + O\left(\varepsilon^{\frac{q-p}{p-2}}\right).$$

Therefore,  $S'_{\varepsilon} \to S'_0$ .

Arguing as in the proof of Lemma 4.2, we may conclude that

(6.7) 
$$\|w_{\varepsilon}\|_{p}^{p} = \frac{(q-p^{*})p}{(q-p)p^{*}} + \frac{p(q-2)}{2(q-p)}\|w_{\varepsilon}\|_{2}^{2}.$$

Then, using this identity to compute  $S'_0(w_{\varepsilon})$  and the convergence of  $S'_{\varepsilon}$  to  $S'_0$ , after some tedious algebra we obtain

(6.8) 
$$\lim_{\varepsilon \to 0} \|w_{\varepsilon}\|_{2}^{2} = \frac{2(p^{*}-p)}{p^{*}(p-2)}, \qquad \lim_{\varepsilon \to 0} \|w_{\varepsilon}\|_{p}^{p} = \frac{(p^{*}-2)p}{(p-2)p^{*}}$$

In particular, this implies that  $p^* \int_{\mathbb{R}^N} G_0(w_{\varepsilon}) dx \to 1$  as  $\varepsilon \to 0$ . Hence, there exists a rescaling  $\lambda_{\varepsilon} \to 1$  such that  $p^* \int_{\mathbb{R}^N} G_0(\tilde{w}_{\varepsilon}) dx = 1$  and  $S'_{\varepsilon}(\tilde{w}_{\varepsilon}) \to S'_0$  for  $\tilde{w}_{\varepsilon}(x) := w_{\varepsilon}(\lambda_{\varepsilon}x)$ . This implies that  $(\tilde{w}_{\varepsilon})$  is a minimizing family for  $(S'_0)$  that satisfies the constraint used in the analysis of [5]. Then, applying [5, Theorem 2] we conclude that for a sequence  $\varepsilon_n \to 0$  we have  $\tilde{w}_{\varepsilon_n} \to \bar{w}$  strongly in  $H^1(\mathbb{R}^N)$ , and in view of the convergence of  $(\lambda_{\varepsilon})$  we have  $w_{\varepsilon_n} \to \bar{w}$  as well, where  $\bar{w}$  is the minimizer of  $(S'_0)$  satisfying the constraint. Therefore, by uniqueness of minimizers of  $(R_0)$  [17], we have  $\bar{w} = w_0$  and the limit is a full limit.

Finally, arguing as in the proof of Lemma 4.15, using  $||w_{\varepsilon}||_{p^*}$  instead of the  $L^q$  norm to control the growth of  $w_{\varepsilon}$  at the origin, we also conclude that  $||w_{\varepsilon}||_{\infty} \leq 1$  as  $\varepsilon \to 0$ . Then by standard elliptic regularity, similarly to the proof of Corollary 4.16, we conclude that  $w_{\varepsilon}$  converges to  $w_0$  in  $L^s(\mathbb{R}^N)$  for any  $s \geq 2$  and in  $C^2(\mathbb{R}^N)$ , which completes the proof of Theorem 2.1.

## References

- M. ABRAMOWITZ, I. A. STEGUN, Handbook of Mathematical Functions. Courier Dover Publications, 1964. 1046 pp.
- [2] S. AGMON, Bounds on exponential decay of eigenfunctions of Schrödinger operators. In: Schrödinger operators. Lecture Notes in Mathematics, 1159, pp.1–38. Springer Verlag, Berlin, 1985.
- [3] A. AMBROSETTI, A. MALCHIODI, Perturbation methods and semilinear elliptic problems on R<sup>n</sup>. Progress in Mathematics, 240. Birkhäuser Verlag, Basel, 2006. xii+183 pp.
- [4] F.V. ATKINSON, L.A. PELETIER, Elliptic equations with nearly critical growth. J. Differential Equations 70 (1987), 349–365.
- [5] H. BERESTYCKI, P.-L. LIONS, Nonlinear scalar field equations. I: Existence of a ground state. Arch. Rational Mech. Anal. 82 (1983), 313–345.
- [6] H. BREZIS, L.A. PELETIER, Asymptotics for elliptic equations involving critical exponents. In: Partial Differential Equations and Calculus of Variations, Birkhäuser, 1989, 149–192.
- [7] J. W. CAHN AND J. E. HILLIARD, Free energy of a nonuniform system .3. nucleation in a 2-component incompressible fluid. J. Chem. Phys., 31 (1959), 688–699.
- [8] S. COLEMAN, Fate of the false vacuum: Semiclassical theory. Phys. Rev. D, 15 (1977), 2929– 2936.
- [9] M. CROSS AND P. C. HOHENBERG, Pattern formation outside of equilibrium. Rev. Mod. Phys., 65 (1993), 851–1112.
- [10] M. DEL PINO AND J. DOLBEAULT, Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions. J. Math. Pures Appl. 81 (9) (2002), 847–875.
- [11] A. FERRERO, F. GAZZOLA, Asymptotic behavior of ground states of quasilinear elliptic problems with two vanishing parameters, Part III. J. Diff. Eq. 198 2004, 53–90.

- [12] F. GAZZOLA, J. SERRIN, Asymptotic behavior of ground states of quasilinear elliptic problems with two vanishing parameters. Ann. Inst. Henri Poincaré, Analyse non Linéaire 19 (2002), 477–504.
- [13] F. GAZZOLA, L. PELETIER, P. PUCCI, J. SERRIN, Asymptotic behavior of ground states of quasilinear elliptic problems with two vanishing parameters, Part II. Ann. Inst. Henri Poincaré, Analyse non Linéaire 20 (2003), 947–974.
- [14] B. GIDAS, W.M NI, L. NIRENBERG, Symmetry of positive solutions of nonlinear elliptic equations in R<sup>n</sup>. Adv. Math., Suppl. Stud. 7A (1981), 369–403.
- [15] D. GILBARG AND N. S. TRUDINGER, Elliptic Partial Differential Equations of Second Order. Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [16] Z.-C. HAN, Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent. Ann. Inst. H. Poincaré Anal. Non Linéaire 8 (1991), 159–174.
- [17] M. K. KWONG, Uniqueness of positive solutions of  $\Delta u u + u^p = 0$  in  $\mathbb{R}^n$ . Arch. Rational Mech. Anal. **105** (1989), 243–266.
- [18] M. K. KWONG, J. B. MCLEOD, L. PELETIER, W. TROY, On ground state solutions of  $-\Delta u = u^p u^q$ . J. Differential Equations 95 (1992), 218-239.
- [19] P.-L. LIONS, The concentration-compactness principle in the calculus of variations. The locally compact case. I. Ann. Inst. H. Poincaré Anal. Non Linéaire 1 (1984), 109-145.
- [20] F. MERLE, L. PELETIER, Asymptotic behavior of positive solutions of elliptic equations with critical and supercritical growth. I. The radial case. Arch. Rational Mech. Anal. 112 (1990), no. 1, 1-19.
- [21] F. MERLE, L. PELETIER, Asymptotic behavior of positive solutions of elliptic equations with critical and supercritical growth. II. The non-radial case. J. Funct. Anal. 105 (1992), no. 1, 1-41.
- [22] C. MURATOV, E. VANDEN-EIJNDEN, Breakup of Universality in the Generalized Spinodal Nucleation Theory. J. Stat. Phys. 114 (2004), 605–623.
- [23] S. POHOŻAEV, Eigenfunctions of the equation  $\Delta u + \lambda f(u) = 0$ . Dokl. Akad. Nauk SSSR 165 (1965) 36–39.
- [24] O. REY, Proof of two conjectures of H. Brézis and L. A. Peletier. Manuscripta Math. 65 (1989), 19–37.
- [25] G. STAMPACCHIA, Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus. Ann. Inst. Fourier (Grenoble) 15 (1965) fasc. 1, 189–258.
- [26] W.A. STRAUSS, Existence of solitary waves in higher dimensions. Comm. Math. Phys. 55 (1977), 149–162.
- [27] M. STRUWE, Variational Methods. Springer-Verlag, 1990.
- [28] J. SERRIN, M. TANG, Uniqueness of ground states for quasilinear elliptic equations. Indiana Univ. Math. J. 49 (2000), 897–923.
- [29] C. UNGER AND W. KLEIN, Nucleation theory near classical spinodal. Phys. Rev. B, 29 (1984), 2698–2708.
- [30] W. VAN SAARLOOS AND P. C. HOHENBERG, Fronts, pulses, sources and sinks in generalized complex ginzburg-landau equations. Physica D, 56 (1992), 303–367.
- [31] M. WILLEM, Minimax theorems. Birkhäuser, 1996.

SWANSEA UNIVERSITY, DEPARTMENT OF MATHEMATICS, SINGLETON PARK, SWANSEA, SA2 8PP, WALES, UNITED KINGDOM

E-mail address: v.moroz@swansea.ac.uk

DEPARTMENT OF MATHEMATICAL SCIENCES, NEW JERSEY INSTITUTE OF TECHNOLOGY, UNIVERSITY HEIGHTS, NEWARK NJ 07102, USA

*E-mail address*: muratov@njit.edu