

On Lagrangian formulations for arbitrary bosonic HS fields on Minkowski backgrounds

ALEXANDER A. RESHETNYAK*

Laboratory of Non-Linear Media Physics, Institute of
Strength Physics and Materials Science, 634021 Tomsk, Russia

Abstract

We review the details of unconstrained Lagrangian formulations for Bose particles propagated on an arbitrary dimensional flat space-time and described by the unitary irreducible integer higher-spin representations of the Poincare group subject to Young tableaux $Y(s_1, \dots, s_k)$ with k rows. The procedure is based on the construction of Verma modules and finding auxiliary oscillator realizations for the symplectic $sp(2k)$ algebra which encodes the second-class operator constraints subsystem in the HS symmetry algebra. Application of an universal BRST approach reproduces gauge-invariant Lagrangians with reducible gauge symmetries describing the free dynamics of both massless and massive bosonic fields of any spin with appropriate number of auxiliary fields.

1 Introduction

Growth of the interest to higher-spin (HS) field theory is mainly stipulated by the hopes to reconsider the problems of a unique description of variety of elementary particles and all known interactions especially due to expected output of LHC on the planned capacity. Remind, that it suspects both the proof of supersymmetry display, the answer on the question on existence of Higgs boson, and probably a new insight on origin of Dark Matter ([1]). Because of HS field theory is closely related to superstring theory, which operates with an infinite tower of bosonic and fermionic HS fields it can be treated as an approach to study a structure of superstring theory from field-theoretic viewpoint. Some of the aspects of current state of HS field theory are discussed in the reviews [2]. The paper considers the last results of constructing Lagrangian formulations (LFs) for free integer HS fields on flat $\mathbb{R}^{1,d-1}$ -space-time subject to arbitrary Young tableaux $Y(s_1, \dots, s_k)$ in Fronsdal metric-like formalism within BFV-BRST approach [3], and based on the results presented in [4] (see Ref.[4], for detailed bibliography on various aspects).

It is known that for $d > 4$ space-time dimensions, there appear, besides totally symmetric irreducible representations of Poincare or (Anti)-de-Sitter ((A)dS) algebras the mixed-symmetry representations determined by more than one spin-like parameters [5], [6]. Whereas

*reshet@ispms.tsc.ru

for the former ones the LF's both for massless and massive free higher-spin fields is well enough developed [7], [8], [9], [10], [11] for the latter the problem of their field-theoretic description is not completely solved. So, the main result within the problem of unconstrained LF for arbitrary massless mixed-symmetry HS fields on a Minkowski space-time was obtained in [12] with use of unfolded form of equations of motion for the field in "frame-like" formulation. In the "metric-like" formulation corresponding Lagrangians were derived in closed manner for only reducible Poincare group $ISO(1, d-1)$ representations in [13]. In turn, the Labastida-like [5] constrained LF with off-shell traceless and mixed-symmetry holonomic constraints for arbitrary mixed-symmetric higher spin fields were recently studied in [14] and on a basis of detour complexes from the BRST quantization of worldline diffeomorphism invariant systems in [15].

The paper is devoted to the construction of unconstrained general gauge-invariant Lagrangians for both massless and massive mixed-symmetry tensor fields of rank $s_1 + s_2 + \dots + s_k$, with any integer numbers $s_1 \geq s_2 \geq \dots \geq s_k \geq 1$ for $k \leq [d/2]$ in a $\mathbb{R}^{1,d-1}$ space, the fields being elements of Poincare-group irreps with a Young tableaux (YT) having k rows. Our approach is based on the BFV-BRST construction [3] which in application to free HS field theory consists of three steps: transforming of conditions that determine the representations with a given spin into a topological gauge system of mixed-class operator constraints in an auxiliary Fock space. Then, the subsystem of the second-class constraints, is converted, with a preservation of the initial algebraic structure, into a system of first-class constraints alone in an enlarged Fock space with respect to which one constructs the BRST charge. Finally, the Lagrangian for a HS field is constructed in terms of the BRST charge in such a way that, first, the corresponding equations of motion reproduce the initial constraints, second, the LF contain all appropriate auxiliary and Stuckelberg fields. The BRST approach to LF of HS field theories has been developed for arbitrary massless and massive bosonic fields in Minkowski and AdS spaces in [16]–[20].

2 Integer HS Symmetry Algebra for Bosonic fields

A massless integer spin irreducible representation of Poincare group in $\mathbb{R}^{1,d-1}$ is described by a tensor field $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} \equiv \Phi_{\mu_1^1 \dots \mu_{s_1}^1, \mu_1^2 \dots \mu_{s_2}^2, \dots, \mu_1^k \dots \mu_{s_k}^k}(x)$ of rank $\sum_{i \geq 1}^k s_i$ and generalized spin $\mathbf{s} = (s_1, s_2, \dots, s_k)$, ($s_1 \geq s_2 \geq \dots \geq s_k > 0, k \leq [d/2]$) subject to a YT with k rows of respective length s_1, s_2, \dots, s_k

$$\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} \longleftrightarrow \begin{array}{|c|c|c|c|c|c|c|c|} \hline \mu_1^1 & \mu_2^1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \mu_{s_1}^1 \\ \hline \mu_1^2 & \mu_2^2 & \cdot & \cdot & \cdot & \cdot & \cdot & \mu_{s_2}^2 & \\ \hline \dots & & & & & & & & \\ \hline \mu_1^k & \mu_2^k & \cdot & \cdot & \cdot & \cdot & \mu_{s_k}^k & & \\ \hline \end{array}, \quad (1)$$

This field is symmetric with respect to the permutations of each type of Lorentz indices μ^i , (for the signature $\eta_{\mu\nu} = \text{diag}(+, -, \dots, -)$, $\mu, \nu = 0, 1, \dots, d-1$) and obeys to the Klein-Gordon, divergentless (2), traceless (3) and mixed-symmetry equations (4) [for $i, j = 1, \dots, k$; $l_i, m_i = 1, \dots, s_i$]:

$$\partial^\mu \partial_\mu \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = 0, \quad \partial^{\mu^i} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = 0, \quad (2)$$

$$\eta^{\mu^i \mu^j} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = \eta^{\mu^i \mu^j} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = 0, \quad l_i < m_i, \quad (3)$$

$[\downarrow, \rightarrow]$	$t^{i_1 j_1}$	$t_{i_1 j_1}^+$	l_0	l^i	l^{i+}	$l^{i_1 j_1}$	$l^{i_1 j_1+}$	g_0^i
$t^{i_2 j_2}$	$A^{i_2 j_2, i_1 j_1}$	$B^{i_2 j_2}_{i_1 j_1}$	0	$l_{j_2} \delta^{i_2 i}$	$-l^{i_2+} \delta^{j_2 i}$	$l^{\{j_1 j_2 \delta^{i_1 i_2}\}}$	$-l^{i_2} \{i_1 + \delta^{j_1 i_2}\}$	$F^{i_2 j_2, i}$
$t_{i_2 j_2}^+$	$-B^{i_1 j_1}_{i_2 j_2}$	$A_{i_1 j_1, i_2 j_2}^+$	0	$l_{i_2} \delta_{j_2}^i$	$-l_{j_2}^+ \delta_{i_2}^i$	$l_{i_2}^{\{j_1 \delta_{j_2}^{i_1}\}}$	$-l_{j_2} \{j_1 + \delta_{i_2}^{i_1}\}$	$-F_{i_2 j_2}^{i+}$
l_0	0	0	0	0	0	0	0	0
l^j	$-l^{j_1} \delta^{i_1 j}$	$-l_{i_1} \delta_{j_1}^j$	0	0	$l_0 \delta^{j i}$	0	$-\frac{1}{2} l^{\{i_1 + \delta^{j_1 i_2}\}}$	$l^j \delta^{i j}$
l^{j+}	$l^{i_1 + \delta^{j_1 j}}$	$l_{j_1}^+ \delta_{i_1}^j$	0	$-l_0 \delta^{j i}$	0	$\frac{1}{2} l^{\{i_1 \delta^{j_1 i_2}\}}$	0	$-l^{j+} \delta^{i j}$
$l^{i_2 j_2}$	$-l^{j_1} \{j_2 \delta^{i_2 i_1}\}$	$-l_{i_1}^{\{i_2 + \delta_{j_1}^{j_2}\}}$	0	0	$-\frac{1}{2} l^{\{i_2 \delta^{j_2 i_1}\}}$	0	$L^{i_2 j_2, i_1 j_1}$	$l^i \{i_2 \delta^{j_2 i}\}$
$l^{i_2 j_2+}$	$l^{i_1} \{i_2 + \delta^{j_2 i_1}\}$	$l_{j_1}^{\{j_2 + \delta_{i_1}^{i_2}\}}$	0	$\frac{1}{2} l^{\{i_2 + \delta^{j_2 i_1}\}}$	0	$-L^{i_1 j_1, i_2 j_2}$	0	$-l^i \{i_2 + \delta^{j_2 i}\}$
g_0^j	$-F^{i_1 j_1, j}$	$F_{i_1 j_1}^{j+}$	0	$-l^i \delta^{i j}$	$l^{i+} \delta^{i j}$	$-l^j \{i_1 \delta^{j_1 i_2}\}$	$l^j \{i_1 + \delta^{j_1 i_2}\}$	0

Table 1: HS symmetry algebra $\mathcal{A}(Y(k), \mathbb{R}^{1, d-1})$.

$$\Phi_{(\mu^1)_{s_1}, \dots, \{(\mu^i)_{s_i}, \dots, \underbrace{\mu_1^j \dots \mu_{l_j}^j}_{\dots}, \dots, \mu_{s_j}^j, \dots, (\mu^k)_{s_k}} = 0, \quad i < j, \quad 1 \leq l_j \leq s_j, \quad (4)$$

where the bracket below denote that the indices in it do not include in symmetrization, i.e. the symmetrization concerns only indices $(\mu^i)_{s_i}, \mu_{l_j}^j$ in $\{(\mu^i)_{s_i}, \dots, \underbrace{\mu_1^j \dots \mu_{l_j}^j}_{\dots}, \dots, \mu_{s_j}^j, \dots, (\mu^k)_{s_k}\}$.

Simultaneous description of all $ISO(1, d-1)$ group irreps maybe reformulated in a standard manner with an auxiliary Fock space \mathcal{H} , generated by k pairs of bosonic creation $a_{\mu^i}^i(x)$ and annihilation $a_{\nu^j}^{j+}(x)$ operators, $i, j = 1, \dots, k, \mu^i, \nu^j = 0, 1, \dots, d-1$: $[a_{\mu^i}^i, a_{\nu^j}^{j+}] = -\eta_{\mu^i \nu^j} \delta^{i j}$ and a set of constraints for an arbitrary string-like (so called *basic*) vector $|\Phi\rangle \in \mathcal{H}$,

$$|\Phi\rangle = \sum_{s_1=0}^{\infty} \sum_{s_2=0}^{s_1} \dots \sum_{s_k=0}^{s_{k-1}} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}}(x) \prod_{i=1}^k \prod_{l_i=1}^{s_i} a_i^{+\mu^i} |0\rangle, \quad (5)$$

$$(l_0, l^i, l^{ij}, t^{i_1 j_1}) |\Phi\rangle = (\partial^\mu \partial_\mu, -i a_\mu^i \partial^\mu, \frac{1}{2} a_\mu^i a^{j\mu}, a_\mu^{i_1+} a^{j_1 \mu}) |\Phi\rangle = 0, \quad i \leq j; \quad i_1 < j_1. \quad (6)$$

The set of $(k(k+1)+1)$ primary constraints (6), $\{o_\alpha\} = \{l_0, l^i, l^{ij}, t^{i_1 j_1}\}$, are equivalent to Eqs. (2)–(4) for all spins. In turn, additional condition, $g_0^i |\Phi\rangle = (s_i + \frac{d}{2}) |\Phi\rangle$ for number particles operators, $g_0^i = -a_\mu^{i+} a^{\mu i} + \frac{d}{2}$, makes (6) to be equivalent to Eqs. (2)–(4) for given spin $\mathbf{s} = (s_1, s_2, \dots, s_k)$.

The procedure of LF implies the Hermiticity of BFV-BRST operator Q , $Q = C^\alpha o_\alpha + \dots$, that means the extension of the set $\{o_\alpha\}$ up to one of $\{o_I\} = \{o_\alpha, o_\alpha^+, g_0^i\}$, which is closed with respect to hermitian conjugation related to standard scalar product on \mathcal{H} and commutator multiplication $[\ , \]$. Operators o_I satisfy to the Lie-algebra commutation relations, $[o_I, o_J] = f_{IJ}^K o_K$, for structure constants $f_{IJ}^K = -f_{JI}^K$, to be determined from the multiplication table 1.

The products $B_{i_1 j_1}^{i_2 j_2}, A^{i_2 j_2, i_1 j_1}, F^{i_1 j_1, i}, L^{i_2 j_2, i_1 j_1}$ in the table 1 are given by the relations,

$$B^{i_2 j_2}_{i_1 j_1} = (g_0^i - g_0^j) \delta_{i_1}^{i_2} \delta_{j_1}^{j_2} + (t_{j_1}^{j_2} \theta_{j_1}^{j_2} + t_{j_1}^{j_2+} \theta_{j_1}^{j_2}) \delta_{i_1}^{i_2} - (t_{i_1}^{i_2} \theta_{i_1}^{i_2} + t_{i_1}^{i_2} \theta_{i_1}^{i_2}) \delta_{j_1}^{j_2}, \quad (7)$$

$$A^{i_2 j_2, i_1 j_1} = t^{i_1 j_2} \delta^{i_2 j_1} - t^{i_2 j_1} \delta^{i_1 j_2}, \quad F^{i_2 j_2, i} = t^{i_2 j_2} (\delta^{j_2 i} - \delta^{i_2 i}), \quad (8)$$

$$L^{i_2 j_2, i_1 j_1} = \frac{1}{4} \left\{ \delta^{i_2 i_1} \delta^{j_2 j_1} \left[2g_0^{i_2} \delta^{i_2 j_2} + g_0^{i_2} + g_0^{j_2} \right] - \left(\delta^{j_2 \{i_1 [t_{j_1}^{i_2} \theta_{i_2 j_1} + t^{i_2 j_1}] + \theta_{j_1}^{i_2}\}} + (j_2 \leftrightarrow i_2) \right) \right\}, \quad (9)$$

with Heaviside θ -symbol θ^{ij} . From the Hamiltonian analysis of the dynamical systems the operators $\{o_I\}$ contain respective $2k^2$ second-class $\{o_a\} = \{l^{ij}, t^{i_1 j_1}, l_{ij}^+, t_{i_1 j_1}^+\}$, $(2k+1)$ first-class $\{l_0, l^i, l_j^+\}$ constraints subsystems and k elements g_0^i which forms the non-vanishing in \mathcal{H} matrix $\Delta_{ab}(g_0^i)$ in $[o_a, o_b] \sim \Delta_{ab}$ for topological gauge system.

We called in [4] the algebra of the operators O_I as *integer higher-spin symmetry algebra in Minkowski space with a Young tableaux having k rows* and denoted it as $\mathcal{A}(Y(k), \mathbb{R}^{1,d-1})$.

The subsystem of the second-class constraints $\{o_a\}$ together with $\{g_0^i\}$ forms the subalgebra in $\mathcal{A}(Y(k), \mathbb{R}^{1,d-1})$ which is isomorphic to the symplectic $sp(2k)$ algebra in turn to be Howe dual [21] to the Lorentz algebra $so(1, d-1)$ (see details in [4] and in [14] as well). Note, the elements g_0^i , form a basis in the Cartan subalgebra whereas l^{ij}, t_i^j are the basis of low-triangular subalgebra in $sp(2k)$.

Having constructed the HS symmetry algebra, we can not still construct BRST operator Q with respect to the elements o_I from $\mathcal{A}(Y(k), \mathbb{R}^{1,d-1})$ due to second-class constraints $\{o_a\}$ presence in it. One should to convert symplectic algebra $sp(2k)$ of $\{o_a, g_0^i\}$ into enlarged set of operators O_I with only first-class constraints.

3 Verma module and Oscillator realization for $sp(2k)$

We consider an additive conversion procedure developed within BRST approach, (see e.g. [17]), which implies the enlarging of o_I to $O_I = o_I + o'_I$, with additional parts o'_I to be given on a new Fock space \mathcal{H}' being independent on \mathcal{H} . In this case the elements O_I are given on $\mathcal{H} \otimes \mathcal{H}'$ so that the requirement for O_I to be in involution, i.e. $[O_I, O_J] \sim O_K$, leads to the series of the same algebraic relations, for O_I and o'_I as those for o_I .

Leaving aside the details of Verma module (special representation space [22]) construction for the symplectic algebra $sp(2k)$ of new operators o'_I considered in [4], we present here theirs explicit oscillator form in terms of new $2k^2$ creation and annihilation operators $(B^c; B_d^+) = (b_{ij}^+, d_{rs}^+; b_{ij}, d_{rs})$, $i, j, r, s = 1, \dots, k; i \leq j; r < s$ as follows,

$$g_0^i = \sum_{l \leq m} b_{lm}^+ b_{lm} (\delta^{il} + \delta^{im}) + \sum_{r < s} d_{rs}^+ d_{rs} (\delta^{is} - \delta^{ir}) + h^i, \quad (10)$$

$$l'_{ij} = b_{ij}^+ \quad t'_{lm} = d_{lm}^+ - \sum_{n=1}^{l-1} d_{nl} d_{nm}^+ - \sum_{n=1}^k (1 + \delta_{nl}) b_{nm}^+ b_{ln}, \quad (11)$$

$$t'_{lm} = - \sum_{n=1}^{l-1} d_{nl}^+ d_{nm} + \sum_{p=0}^{m-l-1} \sum_{k_1=l+1}^{m-1} \dots \sum_{k_p=l+p}^{m-1} C^{k_p m}(d^+, d) \prod_{j=1}^p d_{k_j-1 k_j} \quad (12)$$

$$- \sum_{n=1}^k (1 + \delta_{nm}) b_{nl}^+ b_{nm}, \quad k_0 \equiv l, \quad \text{where}$$

$$C^{lm}(d^+, d) \equiv \left(h^l - h^m - \sum_{n=m}^k (d_{ln}^+ d_{ln} + d_{mn}^+ d_{mn}) + \sum_{n=l+1}^{m-1} d_{nm}^+ d_{nm} - d_{lm}^+ d_{lm} \right) d_{lm} \quad (13)$$

$$- \sum_{n=l+1}^{m-1} d_{ln}^+ d_{nm} + \sum_{n=m+1}^k \left\{ d_{mn}^+ - \sum_{n'=1}^{m-1} d_{n'n}^+ d_{n'm} \right\} d_{ln}, \quad \text{for } l < m.$$

Note, first, that B_c, B_d^+ satisfy to the standard (only nonvanishing) commutation relations, $[B_c, B_d^+] = \delta_{cd}$, second, the arbitrary parameters h^i in (10) serve to reproduce correct LF fo HS field with given spin \mathbf{s} , whereas the form of the rest elements l'_{ij} , for $i \leq j$, to be expressed by means of $C^{lm}(d^+, d)$ may be found in [4]. At last, the operators l'_{ij}, t'_{lm} are not Hermitian conjugated respectively l'_{ij}, t'_{lm} in \mathcal{H}' with respect to standard scalar product as

in \mathcal{H} . To restore that property we introduce new scalar product,

$$\langle \Phi_1 | K' E^{-\alpha} | \Phi_2 \rangle = \langle \Phi_2 | K' E'^{\alpha} | \Phi_1 \rangle^*, \quad \langle \Phi_1 | K' g_0^i | \Phi_2 \rangle = \langle \Phi_2 | K' g_0^i | \Phi_1 \rangle^*, \quad (14)$$

for $(E'^{\alpha}; E^{-\alpha}) = (l'_{ij}, t'_{lm}; l'^{+}_{ij}, t'^{+}_{lm})$ with the operator K' , relating standard scalar products in Verma module and in \mathcal{H}' [4]¹.

4 BRST-BFV operator and Lagrangian formulations

Because of the algebra of O_I under consideration is a Lie algebra $\mathcal{A}(Y(k), \mathbb{R}^{1,d-1})$ the BFV-BRST operator Q' can be constructed in the standard way by the formula,

$$Q' = O_I \mathcal{C}^I + \frac{1}{2} \mathcal{C}^I \mathcal{C}^J f_{JI}^K \mathcal{P}_K \quad (15)$$

with the constants f_{JI}^K from the table 1, for the constraints $O_I = (L_0, L_i^+, L_i, L_{ij}, L_{ij}^+, T_{ij}, T_{ij}^+, G_0^i)$, fermionic ghost fields and conjugated to them momenta $(\mathcal{C}^I, \mathcal{P}_I) = ((\eta_0, \mathcal{P}_0); (\eta^i, \mathcal{P}_i^+); (\eta_i^+, \mathcal{P}_j); (\eta^{ij}, \mathcal{P}_{ij}^+); (\eta_{ij}^+, \mathcal{P}_{ij}); (\vartheta_{rs}, \lambda_{rs}^+); (\vartheta_{rs}^+, \lambda_{rs}); (\eta_G^i, \mathcal{P}_G))$ with the properties

$$\eta^{ij} = \eta^{ji}, \vartheta_{rs} = \vartheta_{rs} \theta^{sr}, \{\vartheta_{rs}, \lambda_{tu}^+\} = \delta_{rt} \delta_{su}, \{\mathcal{P}_j, \eta_i^+\} = \delta_{ij}, \{\eta_{lm}, \mathcal{P}_{ij}^+\} = \delta_{li} \delta_{jm} \quad (16)$$

and non-vanishing anticommutators $\{\eta_0, \mathcal{P}_0\} = \iota$, $\{\eta_G^i, \mathcal{P}_G^j\} = \iota \delta^{ij}$ for zero-mode ghosts². The property of the BRST operator to be Hermitian is defined by the rule with respect to the scalar product $\langle | \rangle$ in $\mathcal{H}_{tot} = \mathcal{H} \otimes \mathcal{H}' \otimes \mathcal{H}_{gh}$,

$$Q'^+ K = K Q', \quad \text{for } K = \hat{1} \otimes K' \otimes \hat{1}_{gh}. \quad (17)$$

To construct LF for bosonic HS fields in a $\mathbb{R}^{1,d-1}$ Minkowski space we partially follow the algorithm of [17], [19], which is a particular case of our construction, corresponding to $s_3 = 0$. First, we extract the dependence of Q' (15) on the ghosts $\eta_G^i, \mathcal{P}_G^i$, to obtain the BRST operator Q only for the system of converted first-class constraints $\{O_I\} \setminus \{G_0^i\}$ and generalized spin operator σ^i :

$$Q' = Q + \eta_G^i (\sigma^i + h^i) + \mathcal{A}^i \mathcal{P}_G^i, \quad \text{where} \quad (18)$$

$$\begin{aligned} Q &= \frac{1}{2} \eta_0 L_0 + \eta_i^+ L^i + \sum_{l \leq m} \eta_{lm}^+ L^{lm} + \sum_{l < m} \vartheta_{lm}^+ T^{lm} + \frac{\iota}{2} \sum_l \eta_l^+ \eta^l \mathcal{P}_0 - \sum_{l < n < m} \vartheta_{lm}^+ \vartheta^n \lambda^{nm} \\ &+ \sum_{n < l < m} \vartheta_{lm}^+ \vartheta_n^m \lambda^{nl} - \sum_{i < l < j} (\vartheta_{lj}^+ \vartheta_i^{+l} - \vartheta_{il}^+ \vartheta^{+lj}) \lambda^{ij} - \sum_{n, l < m} (1 + \delta_{ln}) \vartheta_{lm}^+ \eta^{+n} \mathcal{P}^{mn} \\ &+ \sum_{n, l < m} (1 + \delta_{mn}) \vartheta_{lm}^+ \eta^m \mathcal{P}^{+ln} + \frac{1}{2} \sum_{l < m, n \leq m} \eta_{nm}^+ \eta^n \lambda^{lm} - \left[\frac{1}{2} \sum_{l \leq m} (1 + \delta_{lm}) \eta^m \eta_{lm}^+ \right. \\ &\left. + \sum_{l < m} \vartheta_{lm} \eta^{+m} + \sum_{m < l} \vartheta_{ml}^+ \eta^{+m} \right] \mathcal{P}^l + h.c. \quad \text{and} \quad (19) \end{aligned}$$

$$\begin{aligned} \sigma^i &= G_0^i - h^i - \eta_i \mathcal{P}_i^+ + \eta_i^+ \mathcal{P}_i + \sum_m (1 + \delta_{im}) (\eta_{im}^+ \mathcal{P}^{im} - \eta_{im} \mathcal{P}_{im}^+) \\ &+ \sum_{l < i} [\vartheta_{li}^+ \lambda^{li} - \vartheta^{li} \lambda_{li}^+] - \sum_{i < l} [\vartheta_{il}^+ \lambda^{il} - \vartheta^{il} \lambda_{il}^+], \quad (20) \end{aligned}$$

¹The case of the massive bosonic HS fields whose system of second-class constraints contains additionally to elements of $sp(2k)$ algebra the constraints of isometry subalgebra of Minkowski space l^i, l_i^+, l_0 may be treated via procedure of dimensional reduction of the algebra $\mathcal{A}(Y(k), \mathbb{R}^{1,d})$ for massless HS fields to one $\mathcal{A}(Y(k), \mathbb{R}^{1,d-1})$ for massive HS fields, (see [4]). Now, the wave equation in (2) is changed on Klein-Gordon equation corresponding to the constraint l_0 ($l_0 = \partial^\mu \partial_\mu + m^2$) acting on the same *basic* vector $|\Phi\rangle$ (5).

²The ghosts possess the standard ghost number distribution, $gh(\mathcal{C}^I) = -gh(\mathcal{P}_I) = 1 \implies gh(Q') = 1$.

with some inessential for LF operatorial quantities \mathcal{A}^i . Then, we choose a representation of \mathcal{H}_{tot} as, $(\eta_i, \eta_{ij}, \vartheta_{rs}, \mathcal{P}_0, \mathcal{P}_i, \mathcal{P}_{ij}, \lambda_{rs}, \mathcal{P}_G^i)|0\rangle = 0$ and suppose that the field vectors $|\chi\rangle$ as well as the gauge parameters $|\Lambda\rangle$ do not depend on ghosts η_G^i

$$|\chi\rangle = \sum_n \prod_{i \leq j, r < s}^k (b_{ij}^+)^{n_{ij}} (d_{rs}^+)^{p_{rs}} (\eta_0^+)^{n_{f0}} \prod_{i, j, l \leq m, n \leq o} (\eta_i^+)^{n_{fi}} (\mathcal{P}_j^+)^{n_{pj}} (\eta_{lm}^+)^{n_{flm}} (\mathcal{P}_{no}^+)^{n_{pno}} \\ \times \prod_{r < s, t < u} (\vartheta_{rs}^+)^{n_{frs}} (\lambda_{tu}^+)^{n_{\lambda tu}} |\Phi(a_i^+)^{n_{f0}(n)_{fi}(n)_{pj}(n)_{flm}(n)_{pno}(n)_{frs}(n)_{\lambda tu}} \rangle_3. \quad (21)$$

We denote by $|\chi^k\rangle$ the state (21) satisfying to $gh(|\chi^k\rangle) = -k$. Thus, the physical state having the ghost number zero is $|\chi^0\rangle$, the gauge parameters $|\Lambda\rangle$ having the ghost number -1 is $|\chi^1\rangle$ and so on. The vector $|\chi^0\rangle$ must contain physical string-like vector $|\Phi\rangle = |\Phi(a_i^+)^{(0)_{f0}(0)_{fi}(0)_{pj}(0)_{flm}(0)_{pno}(0)_{frs}(0)_{\lambda tu}} \rangle_{(0)_{ij}(0)_{rs}}}$:

$$|\chi^0\rangle = |\Phi\rangle + |\Phi_A\rangle, \quad \text{where } |\Phi_A\rangle|_{[B_a^+ = C^I = \mathcal{P}_I = 0]} = 0 \quad (22)$$

Independence of the vectors (21) on η_G^i transforms the equation for the physical state $Q'|\chi^0\rangle = 0$ and the BRST complex of the reducible gauge transformations, $\delta|\chi\rangle = Q'|\chi^1\rangle$, $\delta|\chi^1\rangle = Q'|\chi^2\rangle, \dots, \delta|\chi^{(r-1)}\rangle = Q'|\chi^{(r)}\rangle$, to the relations:

$$1) Q|\chi^0\rangle = 0, \quad 2) \delta|\chi^0\rangle = Q|\chi^1\rangle, \quad \dots \quad r) \delta|\chi^{r-1}\rangle = Q|\chi^{(r)}\rangle, \quad (23)$$

$$1) (\sigma^i + h^i)|\chi^0\rangle = 0, \quad 2) (\sigma^i + h^i)|\chi^1\rangle = 0, \quad \dots \quad r) (\sigma^i + h^i)|\chi^r\rangle = 0, \quad (24)$$

Where $r - 1 = k(k + 1) - 1$ is the stage of reducibility both for massless and for the massive bosonic HS field. Resolving the spectral problem from the Eqs.(24) we determine the eigenvectors of the operators σ^i : $|\chi^0\rangle_{(n)_k}, |\chi^1\rangle_{(n)_k}, \dots, |\chi^s\rangle_{(n)_k}, n_1 \geq n_2 \geq \dots n_k \geq 0$ and corresponding eigenvalues of the parameters h^i (for massless HS fields),

$$-h^i = n_i + \frac{d-2-4i}{2}, \quad i = 1, \dots, k, \quad n_1, \dots, n_{k-1} \in \mathbb{Z}, n_k \in \mathbb{N}_0, \quad (25)$$

Let us fix some values of $n_i = s_i$. Then one should substitute h^i corresponding to the chosen s_i (25) into K (17), Q (19) and relations (23), (24). Thus, e.g., the equation of motion in (23) corresponding to the field with given spin (s_1, \dots, s_k) has the form, $Q_{(n)_k}|\chi^0\rangle_{(n)_k} = 0$, with nilpotent $Q_{(n)_k}$ and the same for the gauge transformations in (23).

Following to bosonic one- [16], [18] and two-row cases , [17], [19] one can show that last equation may be derived from the Lagrangian action for fixed spin $(n)_k = (s)_k$,

$$\mathcal{S}_{(s)_k} = \int d\eta_0 {}_{(s)_k} \langle \chi^0 | K_{(s)_k} Q_{(s)_k} | \chi^0 \rangle_{(s)_k}, \quad \left(\implies \frac{\delta \mathcal{S}_{(s)_k}}{\delta {}_{(s)_k} \langle \chi^0 |} = Q_{(s)_k} | \chi^0 \rangle_{(s)_k} = 0 \right), \quad (26)$$

where the standard scalar product for the creation and annihilation operators in \mathcal{H}_{tot} is assumed.

Concluding, one can prove the Lagrangian action (26) indeed reproduces the basic conditions (2)–(4) for massless (massive) HS fields. General action (26) gives, in principle, a straight receipt to obtain the Lagrangian for any component field from general vector $|\chi^0\rangle_{(s)_k}$.

³The brackets $(n)_{fi}, (n)_{pj}, (n)_{ij}$ in (21) means, e.g., for $(n)_{ij}$ the set of indices $(n_{11}, \dots, n_{1k}, \dots, n_{k1}, \dots, n_{kk})$. The sum above is taken over n_l, n_{ij}, p_{rs} and running from 0 to infinity, and over the rest n 's from 0 to 1.

Acknowledgements The author is thankful to the organizers of the Dubna International Workshop SQS'11 for the hospitality. A.R. is grateful to K. Alkalaev, M. Grigoriev, D. Francia, V. Gershun, E. Latini, P.M. Lavrov, Yu.M. Zinoviev for valuable discussions and comments and to I.L. Buchbinder for collaboration. The work was supported by the RFBR grant, project Nr. 12-02-00121 and by LRSS grant Nr.224.2012.2.

References

- [1] Feldman D., Perez P.F., Nath P. R-parity Conservation via the Stueckelberg Mechanism: LHC and Dark Matter Signals// JHEP. 2012. V. 1201. P. 038, [arXiv:1109.2901 [hep-ph]].
- [2] Vasiliev M. Higher spin gauge theories in various dimensions// Fortsch. Phys. 2004. V. 52 p. 702–717, [arXiv:hep-th/0401177]; Sorokin D. Introduction to the classical theory of higher spins// AIP Conf. Proc. 2005. V. 767. P. 172–202, [arXiv:hep-th/0405069]; Bouatta N., Compère G., Sagnotti A. An introduction to free higher-spin fields// Preprint [arXiv:hep-th/0409068]; Bekaert X., Cnockaert S., Iazeolla C., Vasiliev M.A. Nonlinear higher spin theories in various dimensions// Preprint [arXiv:hep-th/0503128]; Fotopoulos A., Tsuluaia M. Gauge Invariant Lagrangians for Free and Interacting Higher Spin Fields. A review of BRST formulation// Int.J.Mod.Phys. A. 2008. V. 24. P. 1–60, [arXiv:0805.1346[hep-th]].
- [3] Fradkin E.S., Vilkovisky G.A. Quantization of relativistic systems with constraints// Phys. Lett. B. 1975. V. 55. P. 224–226; Batalin I.A., Vilkovisky G.A. Relativistic S-matrix of dynamical systems with boson and fermion constraints// Phys. Lett. B. 1977. V. 69. P. 309–312; Batalin I.A., Fradkin E.S. Operator quantization of relativistic dynamical systems subject to first class constraints// Phys. Lett. B. 1983. V. 128. P. 303.
- [4] Buchbinder I.L., Reshetnyak A.A. General Lagrangian formulation for higher spin fields with arbitrary index symmetry. I. Bosonic fields
Preprint [arXiv:1110.5044[hep-th]].
- [5] Labastida J.M.F., Morris T.R. Massless mixed symmetry bosonic free fields// Phys. Lett. B. 1986. V. 180. P. 101–106; Labastida J.M.F. Massless fermionic free fields// Phys. Lett. B. 1987. V.186. P. 365–369; Massless bosonic free fields// Phys. Rev. Lett. 1987. V. 58. P. 531–534; Massless particles in arbitrary representations of the Lorentz group// Nucl. Phys. B. 1989. V. 322. P. 185–209.
- [6] Metsaev R.R. Massless mixed symmetry bosonic free fields in d-dimensional anti-de Sitter space-time// Phys. Lett. B. 1995. V. 354. P. 78-84.
- [7] Fronsdal C. Massless fields with integer spin// Phys. Rev. D. 1978. V.18. P. 3624–3629.
- [8] Fronsdal C. Singletons and massless, integer-spin fields on de Sitter space// Phys. Rev. D. 1979. V.20. P. 848–856; Vasiliev M.A. 'Gauge' Form Of Description Of Massless Fields With Arbitrary Spin (in Russian)// Yad.Fiz. 1980. V. 32. P. 855–861.
- [9] Vasiliev M.A. Free massless fermionic fields of arbitrary spin in D-dimensional anti-de Sitter space// Nucl. Phys. B. 1988. V.301. P. 26–51; Lopatin V.E., Vasiliev M.A. Free massless bosonic fields of arbitrary spin in D-dimensional de Sitter space// Mod. Phys. Lett. A. 1998. V. 3. P. 257–265.

- [10] Singh L.P.S., Hagen C.R. Lagrangian formulation for arbitrary spin. 1. The bosonic case// Phys. Rev. D. 1974. V. 9. P. 898–909; Lagrangian formulation for arbitrary spin. 2. The fermionic case// Phys. Rev. D. 1974. V. 9. P. 910–920; Ponomarev D.S., Vasiliev M.A. Frame-Like Action and Unfolded Formulation for Massive Higher-Spin Fields// Nucl. Phys. B. 2010. V. 839. P. 466–498, [arXiv:1001.0062[hep-th]].
- [11] Zinoviev Yu.M. On massive high spin particles in AdS// Preprint [arXiv:hep-th/0108192]; Metsaev R.R. Massive totally symmetric fields in AdS(d)// Phys.Lett. B. 2004. V. 590 P. 95–104, [arXiv:hep-th/0312297]; Fermionic fields in the d-dimensional anti-de Sitter spacetime// Phys. Lett. B. 1998. V.419. P. 49–56, [arXiv:hep-th/9802097]; Light-cone form of field dynamics in anti-de Sitter space-time and AdS/CFT correspondence// Nucl. Phys. B. 1999. V. 563. P. 295–348, [arXiv:hep-th/9906217]; Massless arbitrary spin fields in AdS(5)// Phys. Lett. B. 2002. V.531. P. 152–160, [arXiv:hep-th/0201226].
- [12] Skvortsov E.D. Frame-like Actions for Massless Mixed-Symmetry Fields in Minkowski space// Nucl. Phys. B. 2009. V. 808. P. 569, [arXiv:0807.0903[hep-th]].
- [13] Campoleoni A., Francia D., Mourad J., Sagnotti A. Unconstrained Higher Spins of Mixed Symmetry. I. Bose Fields// Nucl.Phys. B. 2009. V. 815. P. 289–357, [arXiv:0810.4350].
- [14] K.B. Alkalaev, M. Grigoriev, I.Y. Tipunin, Massless Poincare modules and gauge invariant equations// Nucl. Phys. B. 2009. V. 823 P. 509, [arXiv:0811.3999 [hep-th]].
- [15] D. Cherney, E. Latini, A. Waldron, Generalized Einstein Operator Generating Functions, Phys. Lett. B682 (2010) 472–475, [arXiv:0909.4578[hep-th]]; BRST Detour Quantization, J. Math. Phys. 51 (2010) 062302, [arXiv:0906.4814[hep-th]]; F. Bastianelli, O. Corradini, E. Latini, Spinning particles and higher spin fields on (A)dS backgrounds, JHEP 0811 (2008) 054, [arXiv:0810.0188[hep-th]].
- [16] Burdik C., Pashnev A., Tsulaia M. Auxiliary representations of Lie algebras and the BRST constructions// Mod. Phys. Lett. A. 2000. V. 15. P. 281–291, [arXiv:hep-th/0001195].
- [17] Burdik C., Pashnev A., Tsulaia M. On the mixed symmetry irreducible representations of the Poincare group in the BRST approach// Mod. Phys. Lett. A. 2001. V. 16. P. 731–746, [arXiv:hep-th/0101201]; The Lagrangian description of representations of the Poincare group// Nucl. Phys. Proc. Suppl. 2001. V. 102. P. 285–292, [arXiv:hep-th/0103143].
- [18] Buchbinder I.L., Krykhtin V.A. Gauge invariant Lagrangian construction for massive bosonic higher spin fields in D dimensions// Nucl. Phys. B. 2005. V.727 P. 536–563.
- [19] Buchbinder I.L., Krykhtin V.A., Takata H. Gauge invariant Lagrangian construction for massive bosonic mixed-symmetry higher spin fields// Phys. Lett. B. 2007. V. 656. P. 253, [arXiv:0707.2181[hep-th]]; Moshin P.Yu., Reshetnyak A.A., BRST Approach to Lagrangian Formulation for Mixed-Symmetry Fermionic Higher-Spin Fileds// JHEP. 2007. V. 0710 P. 040, [arXiv:0706.0386[hep-th]].

- [20] Buchbinder I.L., Krykhtin V.A., Lavrov P.M. Gauge invariant Lagrangian formulation of higher spin massive bosonic field theory in AdS space// Nucl. Phys. B. 2007. V. 762. P. 344–376, [arXiv:hep-th/0608005]; Buchbinder I.L., Krykhtin V.A., Reshetnyak A.A. BRST approach to Lagrangian construction for fermionic higher spin fields in AdS space// Nucl. Phys. B. 2007. V. 787. P. 211–240, [arXiv:hep-th/0703049].
- [21] Howe R. Transcending classical invariant theory// J. Amer. Math. Soc. 1989. V. 3. P. 2; Remarks on classical invariant theory// Trans. Amer. Math. Soc. 1989. V. 2. P. 313.
- [22] Dixmier J. Algebres enveloppantes. Paris: Gauthier-Villars, 1974.