

# From quadratic Hamiltonians of polymomenta to abstract geometrical Maxwell-like and Einstein-like equations

Alexandru Oană and Mircea Neagu

## Abstract

The aim of this paper is to create a large geometrical background on the dual 1-jet space  $J^{1*}(\mathcal{T}, M)$  for a multi-time Hamiltonian approach of the electromagnetic and gravitational physical fields. Our geometric-physical construction is achieved starting only from a given quadratic Hamiltonian function

$$H = h_{ab}(t)g^{ij}(t, x)p_i^a p_j^b + U_{(a)}^{(i)}(t, x)p_i^a + \mathcal{F}(t, x)$$

which naturally produces a canonical nonlinear connection  $N$ , a canonical Cartan  $N$ -linear connection  $C\Gamma(N)$  and their corresponding local distinguished (d-) torsions and curvatures. In such a context, we construct some geometrical electromagnetic-like and gravitational-like field theories which are characterized by some natural geometrical Maxwell-like and Einstein-like equations. Some abstract and geometrical conservation laws for the multi-time Hamiltonian gravitational physical field are also given.

*2010 Mathematics Subject Classification:* 70S05, 53C07, 53C80.

*Key words and phrases:* dual 1-jet spaces, nonlinear connections, canonical Cartan  $N$ -linear connection, d-torsions and d-curvatures, geometrical Maxwell-like and Einstein-like equations.

## 1 Distinguished Riemannian geometrization of metrical multi-time Hamilton spaces

Recently, the studies of Atanasiu and Neagu (see the papers [2], [3] and [4]) initiated the new way of distinguished Riemannian geometrization for Hamiltonians depending on polymomenta, which represents in fact a natural "multi-time" extension of the already classical Hamiltonian geometry on cotangent bundles (synthesized in the Miron et al.'s book [13]). In what follows, we expose the main geometrical ideas which characterize the distinguished Riemannian geometrical approach of Hamiltonians depending on polymomenta (see for details the Oană and Neagu's papers [15], [16]).

Let us consider that  $h = (h_{ab}(t))$  is a semi-Riemannian metric on the "multi-time" (*temporal*) manifold  $\mathcal{T}^m$ , where  $m = \dim \mathcal{T}$ . Let  $g = (g^{ij}(t^c, x^k, p_k^c))$  be a symmetric d-tensor on the dual 1-jet space  $E^* = J^{1*}(\mathcal{T}, M)$ , which has the rank  $n = \dim M$  and a constant signature. At the same time, let us consider a smooth multi-time Hamiltonian function

$$E^* \ni (t^a, x^i, p_i^a) \rightarrow H(t^a, x^i, p_i^a) \in \mathbb{R},$$

which yields the *fundamental vertical metrical d-tensor*

$$G_{(a)(b)}^{(i)(j)} = \frac{1}{2} \frac{\partial^2 H}{\partial p_i^a \partial p_j^b},$$

where  $a, b = 1, \dots, m$  and  $i, j = 1, \dots, n$ .

**Definition 1** A multi-time Hamiltonian function  $H : E^* \rightarrow \mathbb{R}$ , having the fundamental vertical metrical d-tensor of the form

$$G_{(a)(b)}^{(i)(j)}(t^c, x^k, p_k^c) = \frac{1}{2} \frac{\partial^2 H}{\partial p_i^a \partial p_j^b} = h_{ab}(t^c) g^{ij}(t^c, x^k, p_k^c),$$

is called a **Kronecker h-regular multi-time Hamiltonian function**.

In this context, we introduce the following important geometrical concept:

**Definition 2** A pair  $MH_m^n = (E^* = J^{1*}(\mathcal{T}, M), H)$ , where  $m = \dim \mathcal{T}$  and  $n = \dim M$ , consisting of the dual 1-jet space and a Kronecker h-regular multi-time Hamiltonian function  $H : E^* \rightarrow \mathbb{R}$ , is called a **multi-time Hamilton space**.

**Example 3** Let us consider the Kronecker h-regular multi-time Hamiltonian function  $H_1 : E^* \rightarrow \mathbb{R}$ , given by

$$H_1 = \frac{1}{4mc} h_{ab}(t) \varphi^{ij}(x) p_i^a p_j^b, \quad (1)$$

where  $h_{ab}(t)$  ( $\varphi_{ij}(x)$ , respectively) is a semi-Riemannian metric on the temporal (spatial, respectively) manifold  $\mathcal{T}$  ( $M$ , respectively) having the physical meaning of **gravitational potentials**, and  $m$  and  $c$  are the known constants from Theoretical Physics representing the **mass of the test body** and the **speed of light**. Then, the multi-time Hamilton space  $\mathcal{GMH}_m^n = (E^*, H_1)$  is called the **multi-time Hamilton space of the gravitational field**.

**Example 4** If we consider on  $E^*$  a symmetric d-tensor field  $g^{ij}(t, x)$ , having the rank  $n$  and a constant signature, we can define the Kronecker h-regular multi-time Hamiltonian function  $H_2 : E^* \rightarrow \mathbb{R}$ , by setting

$$H_2 = h_{ab}(t) g^{ij}(t, x) p_i^a p_j^b + U_{(a)}^{(i)}(t, x) p_i^a + \mathcal{F}(t, x), \quad (2)$$

where  $U_{(a)}^{(i)}(t, x)$  is a  $d$ -tensor field on  $E^*$ , and  $\mathcal{F}(t, x)$  is a function on  $E^*$ . Then, the multi-time Hamilton space  $\mathcal{NEDMH}_m^n = (E^*, H_2)$  is called the **non-autonomous multi-time Hamilton space of electrodynamics**. The dynamical character of the gravitational potentials  $g_{ij}(t, x)$  (i.e., the dependence on the temporal coordinates  $t^c$ ) motivated us to use the word **"non-autonomous"**.

An important role for the subsequent development of our distinguished Riemannian geometrical theory for multi-time Hamilton spaces is represented by the following result (proved in the paper [2]):

**Theorem 5** *If we have  $m = \dim \mathcal{T} \geq 2$ , then the following statements are equivalent:*

- (i)  $H$  is a Kronecker  $h$ -regular multi-time Hamiltonian function on  $E^*$ .
- (ii) The multi-time Hamiltonian function  $H$  reduces to a multi-time Hamiltonian function of non-autonomous electrodynamic type. In other words, we have

$$H = h_{ab}(t)g^{ij}(t, x)p_i^a p_j^b + U_{(a)}^{(i)}(t, x)p_i^a + \mathcal{F}(t, x). \quad (3)$$

**Corollary 6** *The fundamental vertical metrical  $d$ -tensor of a Kronecker  $h$ -regular multi-time Hamiltonian function  $H$  has the form*

$$G_{(a)(b)}^{(i)(j)} = \frac{1}{2} \frac{\partial^2 H}{\partial p_i^a \partial p_j^b} = \begin{cases} h_{11}(t)g^{ij}(t, x^k, p_k^1), & m = \dim \mathcal{T} = 1 \\ h_{ab}(t^c)g^{ij}(t^c, x^k), & m = \dim \mathcal{T} \geq 2. \end{cases} \quad (4)$$

Following now the Miron's geometrical ideas from [13], the paper [2] proves that any Kronecker  $h$ -regular multi-time Hamiltonian function  $H$  produces a natural nonlinear connection on the dual 1-jet space  $E^*$ , which depends only by the given Hamiltonian function  $H$ :

**Theorem 7** *The pair of local functions  $N = \left( N_{1(i)b}^{(a)}, N_{2(i)j}^{(a)} \right)$  on  $E^*$ , where  $(\chi_{bc}^a)$  are the Christoffel symbols of the semi-Riemannian temporal metric  $h_{ab}$ )*

$$N_{1(i)b}^{(a)} = \chi_{bc}^a p_i^c, \\ N_{2(i)j}^{(a)} = \frac{h^{ab}}{4} \left[ \frac{\partial g_{ij}}{\partial x^k} \frac{\partial H}{\partial p_k^b} - \frac{\partial g_{ij}}{\partial p_k^b} \frac{\partial H}{\partial x^k} + g_{ik} \frac{\partial^2 H}{\partial x^j \partial p_k^b} + g_{jk} \frac{\partial^2 H}{\partial x^i \partial p_k^b} \right],$$

represents a nonlinear connection on  $E^*$ . This is called the **canonical nonlinear connection of the multi-time Hamilton space**  $MH_m^n = (E^*, H)$ .

Taking into account the Theorem 5 and using the *generalized spatial Christoffel symbols* of the  $d$ -tensor  $g_{ij}$ , which are given by

$$\Gamma_{ij}^k = \frac{g^{kl}}{2} \left( \frac{\partial g_{li}}{\partial x^j} + \frac{\partial g_{lj}}{\partial x^i} - \frac{\partial g_{ij}}{\partial x^l} \right),$$

we immediately obtain the following geometrical result:

**Corollary 8** For  $m = \dim \mathcal{T} \geq 2$ , the canonical nonlinear connection  $N$  of a multi-time Hamilton space  $MH_m^n = (E^*, H)$ , whose Hamiltonian function is given by (3), has the components

$$N_{1(i)b}^{(a)} = \chi_{bc}^a p_i^c, \quad N_{2(i)j}^{(a)} = -\Gamma_{ij}^k p_k^a + T_{(i)j}^{(a)},$$

where

$$T_{(i)j}^{(a)} = \frac{h^{ab}}{4} (U_{ib\bullet j} + U_{jb\bullet i}), \quad (5)$$

and

$$U_{ib} = g_{ik} U_{(b)}^{(k)}, \quad U_{kb\bullet r} = \frac{\partial U_{kb}}{\partial x^r} - U_{sb} \Gamma_{kr}^s.$$

The canonical nonlinear connection  $N = \left( N_{1(i)b}^{(a)}, N_{2(i)j}^{(a)} \right)$  on  $E^*$  allows us the construction of the *adapted bases*

$$\left\{ \frac{\delta}{\delta t^a}, \frac{\delta}{\delta x^i}, \frac{\partial}{\partial p_i^a} \right\} \subset \chi(E^*), \quad \{ dt^a, dx^i, \delta p_i^a \} \subset \chi^*(E^*),$$

where

$$\begin{aligned} \frac{\delta}{\delta t^a} &= \frac{\partial}{\partial t^a} - N_{1(j)a}^{(b)} \frac{\partial}{\partial p_j^b}, & \frac{\delta}{\delta x^i} &= \frac{\partial}{\partial x^i} - N_{2(j)i}^{(b)} \frac{\partial}{\partial p_j^b}, \\ \delta p_i^a &= dp_i^a + N_{1(i)b}^{(a)} dt^b + N_{2(i)j}^{(a)} dx^j. \end{aligned} \quad (6)$$

The main result of the metrical multi-time Hamilton geometry is the Theorem of existence of the Cartan canonical  $h$ -normal  $N$ -linear connection  $C\Gamma(N)$  (see [16]) which allows the subsequent development of our metrical multi-time Hamilton theory of physical fields.

**Theorem 9 (the Cartan canonical  $N$ -linear connection)** *On the metrical multi-time Hamilton space  $MH_m^n = (J^{1*}(\mathcal{T}, M), H)$ , endowed with the canonical nonlinear connection  $N$ , there exists an unique  $h$ -normal  $N$ -linear connection*

$$C\Gamma(N) = \left( \chi_{bc}^a, A_{jc}^i, H_{jk}^i, C_{j(c)}^{i(k)} \right),$$

having the metrical properties:

- (i)  $g_{ij|k} = 0, \quad g^{ij}|_{(c)}^{(k)} = 0,$
- (ii)  $A_{jc}^i = \frac{g^{il}}{2} \frac{\delta g_{lj}}{\delta t^c}, \quad H_{jk}^i = H_{kj}^i, \quad C_{j(c)}^{i(k)} = C_{j(c)}^{k(i)},$

where " $_{/a}$ ", " $_{|k}$ " and " $_{(c)}^{(k)}$ " represent the local covariant derivatives of  $C\Gamma(N)$ . Moreover, the adapted local coefficients  $H_{jk}^i$  and  $C_{j(c)}^{i(k)}$  of the **Cartan canonical connection**  $C\Gamma(N)$  have the expressions

$$H_{jk}^i = \frac{g^{ir}}{2} \left( \frac{\delta g_{jr}}{\delta x^k} + \frac{\delta g_{kr}}{\delta x^j} - \frac{\delta g_{jk}}{\delta x^r} \right), \quad C_{i(c)}^{j(k)} = -\frac{g_{ir}}{2} \left( \frac{\partial g^{jr}}{\partial p_k^c} + \frac{\partial g^{kr}}{\partial p_j^c} - \frac{\partial g^{jk}}{\partial p_r^c} \right).$$

**Remark 10** (i) *The Cartan canonical connection  $CT(N)$  of the multi-time Hamilton space  $MH_m^n$  verifies also the metrical properties*

$$h_{ab/c} = h_{ab|k} = h_{ab|_{(c)}^{(k)}} = 0, \quad g_{ij/c} = 0.$$

(ii) *In the case  $m = \dim \mathcal{T} \geq 2$ , the adapted coefficients of the Cartan canonical connection  $CT(N)$  of the multi-time Hamilton space  $MH_m^n$  reduce to*

$$A_{bc}^a = \chi_{bc}^a, \quad A_{jc}^i = \frac{g^{il}}{2} \frac{\partial g_{lj}}{\partial t^c}, \quad H_{jk}^i = \Gamma_{jk}^i, \quad C_{j(c)}^{i(k)} = 0. \quad (7)$$

Applying the formulas that determine the local d-torsions and d-curvatures of an  $h$ -normal  $N$ -linear connection  $D\Gamma(N)$  (see [15]) to the Cartan canonical connection  $CT(N)$ , we obtain (see [16]):

**Theorem 11** *The torsion tensor  $\mathbb{T}$  of the Cartan canonical connection  $CT(N)$  of the multi-time Hamilton space  $MH_m^n$  is determined by the adapted local d-components*

	$h_{\mathcal{T}}$	$h_M$		$v$	
	$m \geq 1$	$m = 1$	$m \geq 2$	$m = 1$	$m \geq 2$
$h_{\mathcal{T}}h_{\mathcal{T}}$	0	0	0	0	$R_{(r)ab}^{(f)}$
$h_M h_{\mathcal{T}}$	0	$T_{1j}^r$	$T_{aj}^r$	$R_{(r)1j}^{(1)}$	$R_{(r)aj}^{(f)}$
$vh_{\mathcal{T}}$	0	0	0	$P_{(r)1(1)}^{(1)(j)}$	$P_{(r)a(b)}^{(f)(j)}$
$h_M h_M$	0	0	0	$R_{(r)ij}^{(1)}$	$R_{(r)ij}^{(f)}$
$vh_M$	0	$P_{i(1)}^{r(j)}$	0	$P_{(r)i(1)}^{(1)(j)}$	0
$vv$	0	0	0	0	0

where

(i) *for  $m = \dim \mathcal{T} = 1$ , we have*

$$T_{1j}^r = -A_{j1}^r, \quad P_{i(1)}^{r(j)} = C_{i(1)}^{r(j)}, \quad P_{(r)1(1)}^{(1)(j)} = \frac{\partial N_{(r)1}^{(1)}}{\partial p_j^1} + A_{r1}^j - \delta_r^j \chi_{11}^1,$$

$$P_{(r)i(1)}^{(1)(j)} = \frac{\partial N_{(r)i}^{(1)}}{\partial p_j^1} + H_{ri}^j, \quad R_{(r)1j}^{(1)} = \frac{\delta N_{(r)1}^{(1)}}{\delta x^j} - \frac{\delta N_{(r)j}^{(1)}}{\delta t},$$

$$R_{(r)ij}^{(1)} = \frac{\delta N_{(r)i}^{(1)}}{\delta x^j} - \frac{\delta N_{(r)j}^{(1)}}{\delta x^i};$$

(ii) *for  $m = \dim \mathcal{T} \geq 2$ , using the equality (5) and the notations*

$$\chi_{fab}^c = \frac{\partial \chi_{fa}^c}{\partial t^b} - \frac{\partial \chi_{fb}^c}{\partial t^a} + \chi_{fa}^d \chi_{db}^c - \chi_{fb}^d \chi_{da}^c,$$

$$\mathfrak{A}_{kij}^r = \frac{\partial \Gamma_{ki}^r}{\partial x^j} - \frac{\partial \Gamma_{kj}^r}{\partial x^i} + \Gamma_{ki}^p \Gamma_{pj}^r - \Gamma_{kj}^p \Gamma_{pi}^r,$$

we have

$$\begin{aligned} T_{aj}^r &= -A_{ja}^r, & P_{(r)a(b)}^{(f)(j)} &= \delta_b^f A_{ra}^j, & R_{(r)ab}^{(f)} &= \chi_{gab}^f p_r^g, \\ R_{(r)aj}^{(f)} &= -\frac{\partial N_{(r)j}^{(f)}}{\partial t^a} - \chi_{ca}^f T_{(r)j}^{(c)}, \\ R_{(r)ij}^{(f)} &= -\mathfrak{R}_{rij}^k p_k^f + \left[ T_{(r)i|j}^{(f)} - T_{(r)j|i}^{(f)} \right]. \end{aligned}$$

**Theorem 12** *The curvature tensor  $\mathbb{R}$  of the Cartan connection  $CT(N)$  of the multi-time Hamilton space  $MH_m^n$  is determined by the following adapted local curvature  $d$ -tensors:*

	$h_{\mathcal{T}}$	$h_M$		$v$	
	$m \geq 1$	$m = 1$	$m \geq 2$	$m = 1$	$m \geq 2$
$h_{\mathcal{T}}h_{\mathcal{T}}$	$\chi_{abc}^d$	0	$R_{ibc}^l$	0	$-R_{(l)(a)bc}^{(d)(i)}$
$h_M h_{\mathcal{T}}$	0	$R_{i1k}^l$	$R_{ibk}^l$	$-R_{(i)(1)1k}^{(1)(l)} = -R_{i1k}^l$	$-R_{(l)(a)bk}^{(d)(i)}$
$vh_{\mathcal{T}}$	0	$P_{i1(1)}^{l(k)}$	0	$-P_{(i)(1)1(1)}^{(1)(l)(k)} = -P_{i1(1)}^{l(k)}$	0
$h_M h_M$	0	$R_{ijk}^l$	$\mathfrak{R}_{ijk}^l$	$-R_{(i)(1)jk}^{(1)(l)} = -R_{ijk}^l$	$-R_{(l)(a)jk}^{(d)(i)}$
$vh_M$	0	$P_{ij(1)}^{l(k)}$	0	$-P_{(i)(1)j(1)}^{(1)(l)(k)} = -P_{ij(1)}^{l(k)}$	0
$vv$	0	$S_{i(1)(1)}^{l(j)(k)}$	0	$-S_{(i)(1)(1)(1)}^{(1)(l)(j)(k)} = -S_{i(1)(1)}^{l(j)(k)}$	0

where, for  $m = \dim \mathcal{T} \geq 2$ , we have the relations

$$-R_{(l)(a)bc}^{(d)(i)} = \delta_i^d \chi_{abc}^d - \delta_a^d R_{lbc}^i, \quad -R_{(l)(a)bk}^{(d)(i)} = -\delta_a^d R_{lbc}^i, \quad -R_{(i)(a)jk}^{(d)(l)} = -\delta_a^d \mathfrak{R}_{ijk}^l,$$

and, generally, the following formulas are true:

(i) for  $m = \dim \mathcal{T} = 1$ , we have

$$\begin{aligned} \chi_{111}^1 &= 0, \\ R_{i1k}^l &= \frac{\delta A_{i1}^l}{\delta x^k} - \frac{\delta H_{ik}^l}{\delta t} + A_{i1}^r H_{rk}^l - H_{ik}^r A_{r1}^l + C_{i(1)}^{l(r)} R_{(r)1k}^{(1)}, \\ R_{ijk}^l &= \frac{\delta H_{ij}^l}{\delta x^k} - \frac{\delta H_{ik}^l}{\delta x^j} + H_{ij}^r H_{rk}^l - H_{ik}^r H_{rj}^l + C_{i(1)}^{l(r)} R_{(r)jk}^{(1)}, \\ P_{i1(1)}^{l(k)} &= \frac{\partial A_{i1}^l}{\partial p_k^1} - C_{i(1)/1}^{l(k)} + C_{i(1)}^{l(r)} P_{(r)1(1)}^{(1)(k)}, \\ P_{ij(1)}^{l(k)} &= \frac{\partial H_{ij}^l}{\partial p_k^1} - C_{i(1)|j}^{l(k)} + C_{i(1)}^{l(r)} P_{(r)j(1)}^{(1)(k)}, \\ S_{i(1)(1)}^{l(j)(k)} &= \frac{\partial C_{i(1)}^{l(j)}}{\partial p_k^1} - \frac{\partial C_{i(1)}^{l(k)}}{\partial p_j^1} + C_{i(1)}^{r(j)} C_{r(1)}^{l(k)} - C_{i(1)}^{r(k)} C_{r(1)}^{l(j)}, \end{aligned}$$

(ii) for  $m = \dim \mathcal{T} \geq 2$ , we have

$$\begin{aligned}\chi_{abc}^d &= \frac{\partial \chi_{ab}^d}{\partial t^c} - \frac{\partial \chi_{ac}^d}{\partial t^b} + \chi_{ab}^f \chi_{fc}^d - \chi_{ac}^f \chi_{fb}^d, \\ R_{ibc}^l &= \frac{\partial A_{ib}^l}{\partial t^c} - \frac{\partial A_{ic}^l}{\partial t^b} + A_{ib}^r A_{rc}^l - A_{ic}^r A_{rb}^l, \\ R_{ibk}^l &= \frac{\partial A_{ib}^l}{\partial x^k} - \frac{\partial \Gamma_{ik}^l}{\partial t^b} + A_{ib}^r \Gamma_{rk}^l - \Gamma_{ik}^r A_{rb}^l, \\ \mathfrak{R}_{ijk}^l &= \frac{\partial \Gamma_{ij}^l}{\partial x^k} - \frac{\partial \Gamma_{ik}^l}{\partial x^j} + \Gamma_{ij}^r \Gamma_{rk}^l - \Gamma_{ik}^r \Gamma_{rj}^l.\end{aligned}$$

In the next Sections, following the physical and geometrical ideas of the already classical Lagrangian geometry of physical fields (see [12], [13] and [14]), we construct a possible multi-time Hamiltonian approach of the electromagnetic and gravitational physical fields, which is characterized by some natural geometrical Maxwell-like and Einstein-like equations. To reach this aim, we consider a multi-time Hamilton space  $MH_m^n = (J^{1*}(T, M), H)$  endowed with its canonical nonlinear connection

$$N = \left( N_{1(i)b}^{(a)}, N_{2(i)j}^{(a)} \right),$$

and we also consider the Cartan canonical connection of the space  $MH_m^n$ , which is locally expressed by

$$CT(N) = \left( \chi_{bc}^a, A_{jc}^i, H_{jk}^i, C_{j(c)}^{i(k)} \right).$$

## 2 Multi-time Hamilton electromagnetism. Geometrical Maxwell-like equations

Let us consider the *canonical Liouville-Hamilton d-tensor field of polymomenta*

$$\mathbb{C}^* = p_i^a \frac{\partial}{\partial p_i^a},$$

together with the fundamental vertical metrical d-tensor  $G_{(a)(b)}^{(i)(j)}$  of the multi-time Hamilton spaces  $MH_m^n$ . These geometrical objects allow us to construct the *metrical deflection d-tensors*

$$\begin{aligned}\Delta_{(a)b}^{(i)} &= G_{(a)(c)}^{(i)(k)} \Delta_{(k)b}^{(c)} = p_{(a)/b}^{(i)}, & \Delta_{(a)j}^{(i)} &= G_{(a)(c)}^{(i)(k)} \Delta_{(k)j}^{(c)} = p_{(a)|j}^{(i)}, \\ \vartheta_{(a)(b)}^{(i)(j)} &= G_{(a)(c)}^{(i)(k)} \vartheta_{(k)(b)}^{(c)(j)} = p_{(a)|b}^{(i)(j)},\end{aligned}$$

where  $p_{(a)}^{(i)} = G_{(a)(c)}^{(i)(k)} p_k^c$  and " /<sub>b</sub>", "|<sub>j</sub>" and "|<sub>(b)</sub>" are the local covariant derivatives induced by the Cartan connection  $CT(N)$ .

Taking into account the expressions of the local covariant derivatives of the Cartan connection  $CT(N)$  (see the paper [15]), by a direct calculation, we obtain

**Proposition 13** *The metrical deflection  $d$ -tensors of the multi-time Hamilton spaces  $MH_m^n$  have the expressions:*

(i) *for  $m = \dim \mathcal{T} = 1$ , we have*

$$\begin{aligned} \Delta_{(1)1}^{(i)} &= -h_{11}g^{ik}A_{k1}^r p_r^1, & \Delta_{(1)j}^{(i)} &= h_{11}g^{ik} \left[ -N_{2(k)j}^{(1)} - H_{kj}^r p_r^1 \right], \\ \vartheta_{(1)(1)}^{(i)(j)} &= h_{11}g^{ij} - h_{11}g^{ik}C_{k(1)}^r p_r^1; \end{aligned} \quad (8)$$

(ii) *for  $m = \dim \mathcal{T} \geq 2$ , we have*

$$\begin{aligned} \Delta_{(a)b}^{(i)} &= -h_{ac}g^{ik}A_{kb}^r p_r^c, & \Delta_{(a)j}^{(i)} &= -\frac{g^{ik}}{4}(U_{ka\bullet j} + U_{ja\bullet k}), \\ \vartheta_{(a)(b)}^{(i)(j)} &= h_{ab}g^{ij}. \end{aligned} \quad (9)$$

In order to construct our metrical multi-time Hamiltonian theory of electromagnetism, we introduce the following concept:

**Definition 14** *The distinguished 2-form on  $J^{1*}(\mathcal{T}, M)$ , locally defined by*

$$\mathbb{F} = F_{(a)j}^{(i)} \delta p_i^a \wedge dx^j + f_{(a)(b)}^{(i)(j)} \delta p_i^a \wedge \delta p_j^b, \quad (10)$$

where

$$F_{(a)j}^{(i)} = \frac{1}{2} \left[ \Delta_{(a)j}^{(i)} - \Delta_{(a)i}^{(j)} \right], \quad f_{(a)(b)}^{(i)(j)} = \frac{1}{2} \left[ \vartheta_{(a)(b)}^{(i)(j)} - \vartheta_{(a)(b)}^{(j)(i)} \right], \quad (11)$$

is called the **multi-time electromagnetic field of the metrical multi-time Hamilton space  $MH_m^n$** .

By a straightforward calculation, the Proposition 13 implies

**Proposition 15** *The components  $F_{(a)j}^{(i)}$  and  $f_{(a)(b)}^{(i)(j)}$  of the multi-time electromagnetic field  $\mathbb{F}$ , associated to the multi-time Hamilton space  $MH_m^n$ , have the following expressions:*

(i) *in the case  $m = \dim \mathcal{T} = 1$ , we have*

$$F_{(1)j}^{(i)} = \frac{h^{11}}{2} \left[ g^{jk} N_{2(k)i}^{(1)} - g^{ik} N_{2(k)j}^{(1)} + (g^{jk} H_{ki}^r - g^{ik} H_{kj}^r) p_r^1 \right], \quad f_{(1)(1)}^{(i)(j)} = 0;$$

(ii) *in the case  $m = \dim \mathcal{T} \geq 2$ , we have*

$$F_{(a)j}^{(i)} = \frac{1}{8} \left[ g^{jk} U_{ka\bullet i} - g^{ik} U_{ka\bullet j} + g^{jk} U_{ia\bullet k} - g^{ik} U_{ja\bullet k} \right], \quad f_{(a)(b)}^{(i)(j)} = 0.$$

The main result of our abstract geometrical Hamilton multi-time electromagnetism is given by



**Theorem 16** *The electromagnetic components  $F_{(a)j}^{(i)}$  of the multi-time Hamilton space  $MH_m^n$  are governed by the following **geometrical Maxwell-like equations**:*

(i) *for  $m = \dim \mathcal{T} = 1$ , we have*

$$\left\{ \begin{array}{l} F_{(1)k/1}^{(i)} = \frac{1}{2} \mathcal{A}_{\{i,k\}} \left\{ \Delta_{(1)1|k}^{(i)} + \Delta_{(1)r}^{(i)} T_{1k}^r + \vartheta_{(1)(1)}^{(i)(r)} R_{(r)1k}^{(1)} + R_{r1k}^i p_{(1)}^{(r)} \right\} \\ \sum_{\{i,j,k\}} F_{(1)j|k}^{(i)} = -\frac{1}{2} \sum_{\{i,j,k\}} \left\{ \vartheta_{(1)(1)}^{(i)(r)} R_{(r)jk}^{(1)} + R_{rjk}^i p_{(1)}^{(r)} \right\} \\ F_{(1)j|1}^{(i)(k)} = \frac{1}{2} \mathcal{A}_{\{i,j\}} \left\{ \vartheta_{(1)(1)|j}^{(i)(k)} - P_{rj(1)}^i p_{(1)}^{(r)(k)} - \Delta_{(1)r}^{(i)} C_{j(1)}^{r(k)} - \vartheta_{(1)(1)}^{(i)(r)} P_{(r)j(1)}^{(1)(k)} \right\}; \end{array} \right.$$

(ii) *for  $m = \dim \mathcal{T} \geq 2$ , we have*

$$\left\{ \begin{array}{l} F_{(a)k/b}^{(i)} = \frac{1}{2} \mathcal{A}_{\{i,k\}} \left\{ \Delta_{(a)b|k}^{(i)} + \Delta_{(a)r}^{(i)} T_{bk}^r + \vartheta_{(a)(f)}^{(i)(r)} R_{(r)bk}^{(f)} + R_{rbk}^i p_{(a)}^{(r)} \right\} \\ \sum_{\{i,j,k\}} F_{(a)j|k}^{(i)} = -\frac{1}{2} \sum_{\{i,j,k\}} \left\{ \vartheta_{(a)(f)}^{(i)(r)} R_{(r)jk}^{(f)} + \mathfrak{R}_{rjk}^i p_{(a)}^{(r)} \right\} \\ \sum_{\{i,j,k\}} F_{(a)j|c}^{(i)(k)} = 0, \end{array} \right.$$

where  $\mathcal{A}_{\{i,j\}}$  means an alternate sum,  $\sum_{\{i,j,k\}}$  means a cyclic sum, and we used the notations  $p_{(1)}^{(i)} = G_{(1)(1)}^{(i)(j)} p_j^1$  and  $p_{(a)}^{(i)} = G_{(a)(b)}^{(i)(j)} p_j^b$ .

**Proof.** The general Ricci identities (see [15] and [16]) applied to  $g^{ij}$  lead us to the equalities:

$$\begin{aligned} g^{ir} R_{rbk}^j + g^{jr} R_{rbk}^i &= 0, & g^{ir} R_{rkl}^j + g^{jr} R_{rkl}^i &= 0, \\ g^{ir} P_{rk(c)}^j + g^{jr} P_{rk(c)}^i &= 0. \end{aligned} \quad (12)$$

Let us consider the following non-metrical deflection d-tensor identities ([15]):

$$\begin{aligned} (d_1) \quad \Delta_{(p)b|k}^{(d)} - \Delta_{(p)k/b}^{(d)} &= p_r^d R_{pbk}^r - \Delta_{(p)r}^{(d)} T_{bk}^r - \vartheta_{(p)(f)}^{(d)(r)} R_{(r)bk}^{(f)}, \\ (d_2) \quad \Delta_{(p)j|k}^{(d)} - \Delta_{(p)k|j}^{(d)} &= p_r^d R_{pj k}^r - \vartheta_{(p)(f)}^{(d)(r)} R_{(r)jk}^{(f)}, \\ (d_3) \quad \Delta_{(p)j|c}^{(d)(k)} - \vartheta_{(p)(c)|j}^{(d)(k)} &= p_r^d P_{pj(c)}^r - \Delta_{(p)r}^{(d)} C_{j(c)}^{r(k)} - \vartheta_{(p)(f)}^{(d)(r)} P_{(r)j(c)}^{(f)(k)}, \end{aligned}$$

where  $\Delta_{(i)b}^{(a)} = p_{i/b}^a$ ,  $\Delta_{(i)j}^{(a)} = p_{i|j}^a$ ,  $\vartheta_{(i)(b)}^{(a)(j)} = p_i^a |_{(b)}^{(j)}$ .

Contracting the above deflection d-tensor identities with the fundamental vertical metrical d-tensor  $G_{(a)(d)}^{(i)(p)}$ , and using the equalities (12), we obtain the following *metrical deflection d-tensor identities*:

$$\begin{aligned} (d'_1) \quad \Delta_{(a)b|k}^{(i)} - \Delta_{(a)k/b}^{(i)} &= -p_{(a)}^{(r)} R_{rbk}^i - \Delta_{(a)r}^{(i)} T_{bk}^r - \vartheta_{(a)(f)}^{(i)(r)} R_{(r)bk}^{(f)}, \\ (d'_2) \quad \Delta_{(a)j|k}^{(i)} - \Delta_{(a)k|j}^{(i)} &= -p_{(a)}^{(r)} R_{rjk}^i - \vartheta_{(a)(f)}^{(i)(r)} R_{(r)jk}^{(f)}, \end{aligned}$$

$$(d'_3) \quad \Delta_{(a)j|^{(k)}_{(c)}}^{(i)} - \vartheta_{(a)(c)|j}^{(i)(k)} = -p_{(a)}^{(r)} P_{rj(c)}^{i(k)} - \Delta_{(a)r}^{(i)} C_{j(c)}^{r(k)} - \vartheta_{(a)(f)}^{(i)(r)} P_{(r)j(c)}^{(f)(k)}.$$

To obtain the first (respectively, the third) geometrical Maxwell-like equation, we permute the indices  $i$  and  $k$  in the identity  $(d'_1)$  (respectively, the indices  $i$  and  $j$  in the identity  $(d'_3)$ ), and we subtract this new identity from the initial one. For  $m = \dim \mathcal{T} \geq 2$  we find the following new identity:

$$F_{(a)j|^{(k)}_{(c)}}^{(i)} = \frac{1}{2} \left[ \vartheta_{(a)(c)|j}^{(i)(k)} - \vartheta_{(a)(c)|i}^{(j)(k)} \right].$$

Consequently, doing a cyclic sum by  $\{i, j, k\}$  for  $m \geq 2$ , we obtain what we were looking for.

Doing a cyclic sum after the indices  $\{i, j, k\}$  in the identity  $(d'_2)$ , it follows the second geometrical Maxwell-like equation. ■

### 3 Multi-time Hamilton gravitational field. Geometrical Einstein-like equations

Let us consider that  $h = (h_{ab}(t))$  is a fixed semi-Riemannian metric on the temporal manifold  $\mathcal{T}$  and let

$$N = \left( N_{1(i)b}^{(a)}, N_{2(i)j}^{(a)} \right)$$

be an "a priori" given nonlinear connection on the dual 1-jet space  $J^{1*}(\mathcal{T}, M)$ . Let

$$\delta p_i^a = dp_i^a + N_{1(i)b}^{(a)} dt^b + N_{2(i)j}^{(a)} dx^j$$

be the vertical distinguished 1-forms adapted to the nonlinear connection  $N$ .

An essential element in the development of our abstract geometrical multi-time Hamilton gravitational theory is given by the following definition:

**Definition 17** *From an abstract physical point of view, an adapted metrical d-tensor  $\mathbb{G}$  on the dual 1-jet space  $E^* = J^{1*}(\mathcal{T}, M)$ , locally expressed by*

$$\mathbb{G} = h_{ab} dt^a \otimes dt^b + g_{ij} dx^i \otimes dx^j + h_{ab} g^{ij} \delta p_i^a \otimes \delta p_j^b,$$

where  $g_{ij} = g_{ij}(t^c, x^k, x_c^k)$  is a symmetric d-tensor field of rank  $n = \dim M$  having a constant signature on  $E^* = J^{1*}(\mathcal{T}, M)$ , is called a **multi-time gravitational h-potential** on  $E^*$ .

Now, taking a multi-time Hamilton space  $MH_m^n = (E^*, H)$ , via its fundamental vertical metrical d-tensor  $G_{(a)(b)}^{(i)(j)}$  (which is given by (4)) and its canonical nonlinear connection  $N$ , we naturally construct a corresponding multi-time gravitational h-potential on  $E^*$ , setting

$$\mathbb{G} = h_{ab} dt^a \otimes dt^b + g_{ij} dx^i \otimes dx^j + h_{ab} g^{ij} \delta p_i^a \otimes \delta p_j^b.$$

At the same time, let us consider that

$$CT(N) = \left( \chi_{ab}^c, A_{jc}^k, H_{jk}^i, C_{j(c)}^{i(k)} \right)$$

is the Cartan canonical connection of the multi-time Hamilton space  $MH_m^n$ .

**Postulate.** *We postulate that the **geometrical Einstein-like equations**, which govern the multi-time gravitational h-potential  $\mathbb{G}$  of the multi-time Hamilton space  $MH_m^n$ , are the abstract geometrical Einstein equations attached to the Cartan canonical connection  $CT(N)$  and to the adapted metric  $\mathbb{G}$  on  $E^*$ , namely*

$$\text{Ric}(CT) - \frac{\text{Sc}(CT)}{2}\mathbb{G} = \mathcal{K}\mathbb{T}, \quad (13)$$

where  $\text{Ric}(CT)$  represents the **Ricci tensor** of the Cartan connection,  $\text{Sc}(CT)$  is the **scalar curvature**,  $\mathcal{K}$  is the **Einstein constant** and  $\mathbb{T}$  is an intrinsic d-tensor of matter, which is called the **stress-energy d-tensor of polynomials**.

In the adapted basis of vector fields

$$(X_A) = \left( \frac{\delta}{\delta t^a}, \frac{\delta}{\delta x^i}, \frac{\partial}{\partial p_i^a} \right),$$

which is produced by the canonical nonlinear connection  $N$  of the multi-time Hamilton space  $MH_m^n$ , the curvature tensor  $\mathbb{R}$  of the Cartan canonical connection  $CT(N)$  is locally expressed by

$$\mathbb{R}(X_C, X_B)X_A = \mathbf{R}_{ABC}^D X_D.$$

It follows that we have

$$R_{AB} = \text{Ric}(X_A, X_B) = \mathbf{R}_{ABD}^D, \quad \text{Sc}(CT) = G^{AB} R_{AB},$$

where

$$G^{AB} = \begin{cases} h^{ab}, & \text{for } A = a, B = b \\ g^{ij}, & \text{for } A = i, B = j \\ h^{ab}g_{ij}, & \text{for } A = \binom{a}{i}, B = \binom{b}{j} \\ 0, & \text{otherwise.} \end{cases} \quad (14)$$

Taking into account, on the one hand, the form of the metrical d-tensor  $\mathbb{G} = (G_{AB})$  of the multi-time Hamilton space  $MH_m^n$ , and, on the other hand, taking into account the expressions of the local curvature d-tensors attached to the Cartan canonical connection  $CT(N)$ , by direct computations, we find

**Proposition 18** *The Ricci tensor  $\text{Ric}(CT)$  of the Cartan canonical connection  $CT(N)$  of the multi-time Hamilton space  $MH_m^n$  is determined by the following adapted components:*

(i) for  $m = \dim \mathcal{T} = 1$ , we have

$$\begin{aligned}
R_{11} &:= \chi_{11} = 0, & R_{1i} &= R_{1i1}^1 = 0, \\
R_{1(1)}^{(i)} &= -P_{1(1)1}^{1(i)} = 0, & R_{i1} &= R_{i1r}^r, & R_{ij} &= R_{ijr}^r, \\
R_{(1)1}^{(i)} &:= -P_{(1)1}^{(i)} = -P_{r1(1)}^{i(r)}, & R_{i(1)}^{(j)} &:= -P_{i(1)}^{(j)} = -P_{ir(1)}^r, \\
R_{(1)(1)}^{(i)(j)} &:= -S_{(1)(1)}^{(i)(j)} = -S_{r(1)(1)}^{i(j)(r)}, & R_{(1)j}^{(i)} &:= -P_{(1)j}^{(i)} = -P_{rj(1)}^{i(r)};
\end{aligned}$$

(ii) for  $m = \dim \mathcal{T} \geq 2$ , we have

$$\begin{aligned}
R_{ab} &:= \chi_{ab} = \chi_{abf}^f, & R_{ai} &= R_{aif}^f = 0, \\
R_{a(b)}^{(j)} &= -P_{af(b)}^{f(j)} = 0, & R_{ia} &= R_{iar}^r, & R_{ij} &= R_{ijr}^r, \\
R_{(a)(b)}^{(i)(j)} &:= -S_{(a)(b)}^{(i)(j)} = -S_{r(a)(b)}^{i(j)(r)} = 0, & R_{i(b)}^{(j)} &:= -P_{i(b)}^{(j)} = -P_{ir(b)}^r = 0, \\
R_{(a)b}^{(i)} &:= -P_{(a)b}^{(i)} = -P_{rb(a)}^{i(r)} = 0, & R_{(a)j}^{(i)} &:= -P_{(a)j}^{(i)} = -P_{rj(a)}^{i(r)} = 0.
\end{aligned}$$

Using the notations  $\chi = h^{ab}\chi_{ab}$ ,  $R = g^{ij}R_{ij}$  and  $S = h^{11}g_{ij}S_{(1)(1)}^{(i)(j)}$ , we obtain

**Corollary 19** *The scalar curvature  $\text{Sc}(\text{CT})$  of the Cartan canonical connection  $\text{CT}(N)$  of the multi-time Hamilton space  $MH_m^n$  is given by the formulas:*

- (i) for  $m = \dim \mathcal{T} = 1$ , we have  $\text{Sc}(\text{CT}) = R - S$ ;
- (ii) for  $m = \dim \mathcal{T} \geq 2$ , we have  $\text{Sc}(\text{CT}) = \chi + R$ .

In this context, the main result of the Hamilton geometrical multi-time gravitational theory is offered by

**Theorem 20** *The **geometrical Einstein-like equations**, which govern the multi-time gravitational  $h$ -potential  $\mathbb{G}$  of the multi-time Hamilton space  $MH_m^n$ , have the following adapted local form:*

(i) for  $m = \dim \mathcal{T} = 1$ , we have

$$\left\{ \begin{array}{l}
-\frac{R-S}{2}h_{11} = \mathcal{K}\mathbb{T}_{11} \\
R_{ij} - \frac{R-S}{2}g_{ij} = \mathcal{K}\mathbb{T}_{ij} \\
-S_{(1)(1)}^{(i)(j)} - \frac{R-S}{2}h_{11}g^{ij} = \mathcal{K}\mathbb{T}_{(1)(1)}^{(i)(j)} \\
0 = \mathbb{T}_{1i}, \quad R_{i1} = \mathcal{K}\mathbb{T}_{i1} \\
0 = \mathbb{T}_{1(1)}^{(i)}, \quad -P_{i(1)}^{(j)} = \mathcal{K}\mathbb{T}_{i(1)}^{(j)} \\
-P_{(1)1}^{(i)} = \mathcal{K}\mathbb{T}_{(1)1}^{(i)}, \quad -P_{(1)j}^{(i)} = \mathcal{K}\mathbb{T}_{(1)j}^{(i)};
\end{array} \right. \quad (15)$$

(ii) for  $m = \dim \mathcal{T} \geq 2$ , we have

$$\left\{ \begin{array}{l} \chi_{ab} - \frac{\chi + R}{2} h_{ab} = \mathcal{K} \mathbb{T}_{ab} \\ R_{ij} - \frac{\chi + R}{2} g_{ij} = \mathcal{K} \mathbb{T}_{ij} \\ -\frac{\chi + R}{2} h_{ab} g^{ij} = \mathcal{K} \mathbb{T}_{(a)(b)}^{(i)(j)} \\ 0 = \mathbb{T}_{ai}, \quad R_{ia} = \mathcal{K} \mathbb{T}_{ia} \\ 0 = \mathbb{T}_{(a)b}^{(i)}, \quad 0 = \mathbb{T}_{a(b)}^{(j)} \\ 0 = \mathbb{T}_{i(b)}^{(j)}, \quad 0 = \mathbb{T}_{(a)j}^{(i)}, \end{array} \right. \quad (16)$$

where  $\mathbb{T}_{AB}$ ,  $A, B \in \left\{ a, i, \binom{a}{i} \right\}$  represent the adapted components of the stress-energy d-tensor of matter  $\mathbb{T}$ .

**Remark 21** In order to have the compatibility of the system of geometrical Einstein-like equations, it is necessary the "a priori" vanishing of certain adapted components of the stress-energy d-tensor of matter  $\mathbb{T}$ .

From a physical point of view, it is well known that in the classical Riemannian theory of gravity, the stress-energy d-tensor of matter have to verify some conservation laws. By a natural extension of the classical Riemannian conservation laws, in our geometrical Hamiltonian context, we postulate the following *generalized conservation laws* of the stress-energy d-tensor of polymomenta  $\mathbb{T}$ :

$$\mathbb{T}_{A|B}^B = 0, \quad \forall A \in \left\{ a, i, \binom{a}{i} \right\},$$

where  $\mathbb{T}_A^B = G^{BD} \mathbb{T}_{DA}$ . Consequently, by straightforward computations, we obtain

**Theorem 22** The *generalized conservation laws* of the Einstein-like equations of the multi-time Hamilton space  $MH_m^n$  are expressed by the following formulas:

(i) for  $m = \dim \mathcal{T} = 1$ , we have

$$\left\{ \begin{array}{l} \left[ \frac{R - S}{2} \right]_{/1} = R_{1|r}^r - P_{(r)1}^{(1)|r} \\ \left[ R_j^r - \frac{R - S}{2} \delta_j^r \right]_{|r} = P_{(r)j}^{(1)|r} \\ \left[ S_{(r)(1)}^{(1)(j)} + \frac{R - S}{2} \delta_r^j \right] \Big|_{(1)}^{(r)} = -P_{(1)|r}^{r(j)}, \end{array} \right. \quad (17)$$

where

$$\begin{aligned}
R_1^i &= g^{iq} R_{q1}, & P_{(i)1}^{(1)} &= h^{11} g_{iq} P_{(1)1}^{(q)}, & R_j^i &= g^{iq} R_{qj}, \\
P_{(i)j}^{(1)} &= h^{11} g_{iq} P_{(1)j}^{(q)}, & P_{(1)}^{i(j)} &= g^{iq} P_{q(1)}^{(j)} & S_{(i)(1)}^{(1)(j)} &= h^{11} g_{iq} S_{(1)(1)}^{(q)(j)};
\end{aligned}$$

(ii) for  $m = \dim \mathcal{T} \geq 2$ , we have

$$\left\{ \begin{array}{l} \left[ \chi_b^f - \frac{\chi + R}{2} \delta_b^f \right]_{/f} = -R_{b|r}^r \\ \left[ R_j^r - \frac{\chi + R}{2} \delta_j^r \right]_{|r} = 0, \end{array} \right. \quad (18)$$

where

$$\chi_b^f = h^{fc} \chi_{cb}, \quad R_j^i = g^{iq} R_{qj}, \quad R_b^i = g^{iq} R_{qb}.$$

**Open problem.** The authors of this paper consider that the finding of a possible real physical meaning of the present multi-time Hamiltonian geometrical theories of gravity and electromagnetism may be an open problem for physicists.

**Acknowledgements.** Many thanks go to Professor Gh. Atanasiu for his constant encouragements and useful suggestions concerning this research topic.

## References

- [1] G.S. Asanov, *Jet extension of Finslerian gauge approach*, Fortschritte der Physik, vol. **38**, no. **8** (1990), 571-610.
- [2] Gh. Atanasiu, M. Neagu, *Canonical nonlinear connections in the multi-time Hamilton geometry*, Balkan Journal of Geometry and Its Applications, vol. **14**, no. **2** (2009), 1-12.
- [3] Gh. Atanasiu, M. Neagu, *Distinguished tensors and Poisson brackets in the multi-time Hamilton geometry*, BSG Proceedings **16**, Geometry Balkan Press, Bucharest (2009), 12-27.
- [4] Gh. Atanasiu, M. Neagu, *Distinguished torsion, curvature and deflection tensors in the multi-time Hamilton geometry*, Electronic Journal "Differential Geometry - Dynamical Systems", vol. **11** (2009), 20-40.
- [5] I. Čomić, *Multi-time Lagrange spaces*, Electronic Journal "Differential Geometry - Dynamical Systems", vol. **7** (2005), 15-33.
- [6] G. Giachetta, L. Mangiarotti, G. Sardanashevily, *Covariant Hamiltonian field theory*, <http://arxiv.org/hep-th/9904062> (1999).
- [7] G. Giachetta, L. Mangiarotti, G. Sardanashevily, *Polysymplectic Hamiltonian formalism and some quantum outcomes*, <http://arxiv.org/hep-th/0411005> (2004).

- [8] M. Gotay, J. Isenberg, J.E. Marsden, R. Montgomery, *Momentum maps and classical fields. Part I. Covariant field theory*, <http://arxiv.org/physics/9801019> (2004).
- [9] M. Gotay, J. Isenberg, J.E. Marsden, *Momentum maps and classical fields. Part II. Canonical analysis of field theories*, <http://arxiv.org/math-ph/0411032> (2004).
- [10] I.V. Kanatchikov, *Basic structures of the covariant canonical formalism for fields based on the De Donder-Weyl theory*, <http://arxiv.org/hep-th/9410238> (1994).
- [11] I.V. Kanatchikov, *On quantization of field theories in polymomentum variables*, AIP Conf. Proc., vol. **453**, Issue **1** (1998), 356-367.
- [12] R. Miron, M. Anastasiei, *The Geometry of Lagrange Spaces: Theory and Applications*, Kluwer Academic Publishers, 1994.
- [13] R. Miron, D. Hrimiuc, H. Shimada, S.V. Sabău, *The Geometry of Hamilton and Lagrange Spaces*, Kluwer Academic Publishers, 2001.
- [14] M. Neagu, *Riemann-Lagrange Geometry on 1-Jet Spaces*, Matrix Rom, Bucharest, 2005.
- [15] A. Oană, M. Neagu, *The local description of the Ricci and Bianchi identities for an  $h$ -normal  $N$ -linear connection on the dual 1-jet space  $J^{1*}(\mathcal{T}, M)$* , <http://arxiv.org/math.DG/1111.4173> (2011).
- [16] A. Oană, M. Neagu, *A distinguished Riemannian geometrization for quadratic Hamiltonians of polymomenta*, <http://arxiv.org/math.DG/1112.5442> (2011).
- [17] D.J. Saunders, *The Geometry of Jet Bundles*, Cambridge University Press, New York, London, 1989.

Alexandru OANĂ and Mircea NEAGU  
 University Transilvania of Braşov,  
 Department of Mathematics - Informatics,  
 Blvd. Iuliu Maniu, no. 50, Braşov 500091, Romania.  
*E-mails:* alexandru.oana@unitbv.ro, mircea.neagu@unitbv.ro