BEC-BCS Crossover with Feshbach Resonance for a Three-Hyperfine-Species Model

Guojun Zhu^{*} and Anthony J. Leggett

Department of Physics, University of Illinois at Urbana-Champaign

We consider the behavior of an ultracold Fermi gas across a narrow Feshbach resonance, where the occupation of the closed channel may not be negligible. While the corrections to the singlechannel formulae associated with the nonzero chemical potential and with particle conservation have been considered in the existing literature, there is a further effect, namely the "inter-channel Pauli exclusion principle" associated with the fact that a single hyperfine species may be common to the two channels. We focus on this effect and show that, as intuitively expected, the resulting corrections are of order E_F/η , where E_F is the Fermi energy of the gas in the absence of interactions and η is the Zeeman energy difference between the two channels. We also consider the related corrections to the fermionic excitation spectrum, and briefly discuss the collective modes of the system.

I. INTRODUCTION

In a low-temperature dilute system, a short-range interaction can be characterized with a single parameter, a_s , a.k.a. the s-wave scattering length. A very desirable property of the Feshbach resonance in such a system is that the effective interaction is tunable experimentally through the Zeeman energy difference between channels which is in turn tunable through instruments such as a magnetic field [1–4].

$$a_s(B) = a_{bg} \left(1 + \frac{\Delta B}{B - B_0} \right) \tag{1}$$

where B_0 is the magnetic field at which the a_s diverges, i.e., the resonant point. This unique ability gives physicists a rare opportunity to study a many-body system under various interaction strengths, and thus connect different physics originally developed separately. Particularly for the fermionic gas, there are a series of theoretical works about uniform treatment of BEC and BCS since the 1960s [5–8], for which the dilute ultracold fermionic alkali gas with Feshbach resonances provides a perfect testing ground. Indeed, these theories work quite well qualitatively.

The two-body theory of the Feshbach resonance has a characteristic parameter, $\delta_c \sim a_{bg}^2 (\Delta B)^2$, defined as the detuning energy at which the weight of the bound state shifts from predominantly in the open channel to predominantly in the closed channel (see Fig. 1) [9]. Naïvely speaking, on the negative detuning side of any resonance (i.e. $\delta < 0$), the two particles should mostly stay in a (virtual) bound state of the closed channel (or "virtual state" in some other-type resonances). However, at the resonance point of a Feshbach resonance $(a_s \to \pm \infty)$, the atoms are mostly still in the open channel, and they do so down to a negative detuning $\delta \sim -\delta_c$. Only when the negative detuning from resonance is much more than δ_c , do atoms have the majority weight in the closed channel.




FIG. 1: Energy levels in a Feshbach resonance δ is the energy detuning from the resonance point. The horizontal line stands for the zero energy s-wave scattering state, $\psi \sim \frac{1}{r} - \frac{1}{a_s}$, which exists for any detuning. The lower solid line stands for the real bound state, which only exists for negative detuning $(\delta < 0, a_s > 0)$. The dashed line stands for the (uncoupled) closed-channel bound state. An interesting point to notice is that the real bound state appears earlier than the cross point of the (uncoupled) closed-channel bound state appears earlier than the cross point of the (uncoupled) closed-channel bound-state level and zero energy. Another important point to notice is the negative detuning $-\delta_c$. When the negative detuning is smaller than δ_c , this real bound state is composed mostly with atoms in the open channel and vice versa.

When considering a many-body system with a Feshbach resonance, an important question is how this energy scale, δ_c , compares to the many-body energy scale, namely, the Fermi energy of the free fermionic atoms, E_F . In the region not too far away from the resonance $(|\delta| \ll \delta_c)$, the closed-channel weight is negligible if the Fermi energy is much smaller than δ_c , (i.e., broad resonance). Crossover experiments are usually performed at detuning not too far from the resonance, and hence the closed channel can be safely ignored at the many-body level. Eventually, when the detuning is too far away, $|\delta| \gg \delta_c$, the bound state is almost like the uncoupled closed-channel bound state with a little dressing from the open channel. Nevertheless, such a large detuning in the broad resonance is not very interesting because

^{*} zhu.guojun2010@gmail.com

the resonance effect is very small and the s-wave scattering length, a_s , is close to its background value then. Given the above consideration, for most purposes, we can almost neglect the closed channel and the problem can be well-described as a two-species fermion system with a tunable interaction when it is not too far away from the resonance. The Feshbach resonance indeed serves as a simple "magic" knob to change the interaction strength. The original theories developed on single-channel models apply to this case directly. This is also the situation for two popular experimental systems (⁶Li atoms at 834G, 40 K atoms at 224G). Many theoretical works have been developed using either the single-channel model or the two-channel model with the broad resonance assumption (e.g. [10–12]). On the contrary, when the Fermi energy of the free fermionic atoms is comparable to or even larger than δ_c , the closed channel has to be included at the many-body level even for small detuning. Such a situation, previously considered in some works [13, 14], is the focus of the current work.

To complicate the problem further, configurations of Feshbach resonances often have one common hyperfine species between the two channels. There are three hyperfine species in the two channels instead of four species (two for each channel). Two most common systems (⁶Li at 834G. ⁴⁰K at 224G) both contain three species of fermions although they are broad resonances. The Pauli exclusion principle prevents atoms of both channels from occupying the same momentum level simultaneously because of this common species. This "inter-channel Pauli exclusion" has no counterpart in the two-body physics. This peculiar effect in many-body crossover problems has received little theoretical attention up to now. Nevertheless, narrow resonances do exist [3] and it is not inconceivable to perform many-body experiments using such resonances. The central concern of this paper is about these situations.

Roughly speaking, turning from two-body systems with Feshbach resonances to many-body systems brings three new effects into the original two-body problem. The first effect is closely associated with the Fermi energy: For a many-body fermionic system at low temperature, most fermions are inactive; only the fermions close to the Fermi surface participate in the interaction processes. Therefore, the energy often needs to be measured from the Fermi surface instead of from zero as in a two-body situation. The second effect relates to particle conservation. Unlike in the single-channel problem, there are two relevant densities in the two-channel problem: the density of atoms in the open channel, n_o , and the density of atoms in the closed channel, n_c . When the closed-channel weight is small (broad resonance), it is legitimate to treat the total density as the same as the open-channel density. However, in the narrow resonance, where the closed-channel weight is not negligible, counting becomes complicated. Extra care is required to specify which channel quantities such as "density" refer to. These two aspects have been extensively studied

previously [13, 14].

The last effect is unique to the three-species problem, where one common species is shared by both channels. The phase spaces of two channels overlap because of the common species, which prevents both channels have the occupation in the same level simultaneously. This effect is controlled by the wave-function overlap of the states in the two channels. A rough estimate of this overlap can be made: The uncoupled closed-channel bound state which is in resonance with the open-channel zero energy threshold has relatively small spatial extension, a_c . Its binding energy E_b is close to the Zeeman energy difference between two channels, η , when considering only situations not far from resonances. On the other hand, fermions in the open channel fill the lowest momentum states up to typically the Fermi energy, E_F . By a simple dimensional argument, the ratio E_F/η must control the overlap effect. How it modifies the many-body picture is the central topic of this paper.

The present paper is divided as follows: Section II defines the many-body model and introduces the appropriate notation for the 3-species case. Section III lists the mean-field result and its renormalization result. Section III A discusses the fermionic excitation within the meanfield level, and section IV discusses the bosonic modes that are beyond the mean-field level. We conclude and discuss our approach in Section V. A couple of detailed calculations are list in the Appendix and further details of the calculations may be found in ref. [15].

II. MODEL AND NOTATION

We denote the three hyperfine species as a, b and c, where the open channel contains the pair of species (a, b)and the closed channel contains the pair of species (a, c). We denote the Grassmann variable for species i by ψ_i and write the two channels in the notation:

$$(\bar{\psi}\bar{\psi}) = \begin{pmatrix} \bar{\psi}_a \bar{\psi}_b & \bar{\psi}_a \bar{\psi}_c \end{pmatrix} \qquad (\psi\psi) = \begin{pmatrix} \psi_b \psi_a \\ \psi_c \psi_a \end{pmatrix} \qquad (2)$$

The two-body interaction can then be written as a (2×2) hermitian matrix \tilde{U}

$$\tilde{U} \equiv \begin{pmatrix} U & Y \\ Y^* & V \end{pmatrix} \tag{3}$$

Note that we have restricted the Hilbert space to only include pairs of (a, b) and (a, c). When only close-toresonance region is considered, this is a good approximation instead of the most generic description, where all three species are treated equivalently. Within this approximation, species b and c share the same chemical potential, but not with a; in other words, $n_a = n_b + n_c$. We can now write the finite-temperature action as

$$S(\bar{\psi},\psi) = \int_0^\beta d\tau \int d^d r \left[\sum_j \bar{\psi}_j (\partial_\tau - \frac{1}{2m} \nabla^2 - \mu + \eta_j) \psi_j - (\bar{\psi}\bar{\psi}) \tilde{U}(\psi\psi) \right]$$
(4)

Here η_j is the Zeeman energy for hyperfine species j. We choose the zero so that $\eta_a = \eta_b = 0$, $\eta_c = \eta$.

Let us illustrate above discussion in one example. In a common experimental setup for ⁶Li, atoms are usually prepared in the two lowest hyperfine levels: described by the direct product of two states, $|F = \frac{1}{2}, F_z = -\frac{1}{2}\rangle \otimes$ $|F = \frac{1}{2}, F_z = +\frac{1}{2}\rangle$. This is a good approximation until the two atoms are very close. For the atom-atom interaction that conserves the z-component of the total angular momentum, $F_z^{(1)} + F_z^{(2)}$, this channel mixes with four other possible channels of the same total z-direction angular momentum, i.e. $F_z^{(1)} + F_z^{(2)} = 0$: $|\frac{1}{2}, -\frac{1}{2}\rangle \otimes |\frac{3}{2}, +\frac{1}{2}\rangle$, $|\frac{3}{2}, -\frac{1}{2}\rangle \otimes |\frac{1}{2}, +\frac{1}{2}\rangle, |\frac{3}{2}, +\frac{3}{2}\rangle \otimes |\frac{3}{2}, -\frac{3}{2}\rangle, |\frac{3}{2}, +\frac{1}{2}\rangle \otimes |\frac{3}{2}, -\frac{1}{2}\rangle$ (All states are labeled as $|F, F_z\rangle$). Various resonances can take place. Nevertheless, it is usually sufficient to consider only the one in resonance and neglect all others, when close to the resonance. The closed channel in the most studied resonance with a magnetic filed close to 834G, is approximately $|\frac{3}{2}, -\frac{1}{2}\rangle \otimes |\frac{1}{2}, +\frac{1}{2}\rangle$ and the resonance is a three-species resonance[3, 16]. In this case, *a* is $|\frac{1}{2}, +\frac{1}{2}\rangle$; *b* is $|-\frac{1}{2}, +\frac{1}{2}\rangle$; and *c* is $|\frac{3}{2}, -\frac{1}{2}\rangle$.

We perform the Hubbard-Stratonovich transformation on Eq.4. Introduce 2-component auxiliary fields (order parameters), (Δ_1, Δ_2) , coupled to the fermionic fields as

$$\Delta \longrightarrow \Delta - \tilde{U}(\psi\psi) \tag{5}$$

We can introduce a spinor representation

$$\bar{\Psi} = \begin{pmatrix} \bar{\psi}_a & \psi_b & \psi_c \end{pmatrix} \qquad \Psi = \begin{pmatrix} \psi_a \\ \bar{\psi}_b \\ \bar{\psi}_c \end{pmatrix} \tag{6}$$

The action can then be rewritten in a more compact form with respect to Ψ and $\bar{\Psi}$

$$S(\bar{\Delta}, \Delta, \bar{\psi}_i, \psi_i) = \int_0^\beta d\tau \int d^d r \left[\Delta^{\dagger} \tilde{U}^{-1} \Delta - \bar{\Psi} \mathcal{G}^{-1} \Psi \right]$$
(7)

where the fermionic correlation \mathcal{G}^{-1} in the momentumfrequency representation is

$$\mathcal{G}^{-1} = \begin{pmatrix} i\omega_n - \xi_{\mathbf{k}} & \Delta_1 & \Delta_2 \\ \bar{\Delta}_1 & i\omega_n + \xi_{\mathbf{k}} & 0 \\ \bar{\Delta}_2 & 0 & i\omega_n + \xi_{\mathbf{k}} + \eta \end{pmatrix}$$
(8)

where $\xi_{\mathbf{k}} = \hbar^2 k^2 / 2m - \mu$. The action in Eq. 7 is bilinear in the quantities Ψ , $\bar{\Psi}$ and we can formally integrate them out, with the result

$$S(\bar{\Delta}, \Delta) = \int dx \left(\bar{\Delta} \tilde{U}^{-1} \Delta - \operatorname{tr} \ln \hat{\mathcal{G}}^{-1} \right)$$
(9)

III. MEAN-FIELD RESULT AND RENORMALIZATION

Eq. (8) can be inverted to get G. The final meanfield equations are (for simplicity, both Δ_i 's are taken as real.[17])

$$\begin{pmatrix} \Delta_1 \\ \Delta_2 \end{pmatrix} = \begin{pmatrix} U & Y \\ Y^* & V \end{pmatrix} \sum_{\mathbf{k}} \begin{pmatrix} h_{1\mathbf{k}} \\ h_{2\mathbf{k}} \end{pmatrix}$$
(10)

where $h_{1\mathbf{k}}$ and $h_{2\mathbf{k}}$ are expectations of the abnormal Green's function.

$$h_{1\mathbf{k}} = \left\langle \psi_{a,-\mathbf{k}} \psi_{b,+\mathbf{k}} \right\rangle = \Delta_1 \frac{E_{1\mathbf{k}} + \xi_{\mathbf{k}} + \eta}{(E_{1\mathbf{k}} + E_{2\mathbf{k}})(E_{1\mathbf{k}} + E_{3\mathbf{k}})}$$
(11)
$$h_{2\mathbf{k}} = \left\langle \psi_{a,-\mathbf{k}} \psi_{c,+\mathbf{k}} \right\rangle = \Delta_2 \frac{E_{1\mathbf{k}} + \xi_{\mathbf{k}}}{(E_{1\mathbf{k}} + E_{2\mathbf{k}})(E_{1\mathbf{k}} + E_{3\mathbf{k}})}$$
(12)

where $E_{i\mathbf{k}}$'s are the eigenvalues of the fermionic correlation Eq. 8 (see details in Sec. III A).

There is one number equation for each channel,

$$\sum_{\omega_n, \mathbf{k}} G_{22} e^{(-i\omega_n \delta_-)} = N_{open}$$
$$\sum_{\omega_n, \mathbf{k}} G_{33} e^{(-i\omega_n \delta_-)} = N_{close}$$

The Matsubara summation can be performed by the normal trick of multiplying the summand by a Fermi function and deforming the contour [18]. For the summation at zero temperature, we just need to consider the positive roots, $E_{1\mathbf{k}}$. It is straightforward to find

$$N_{\rm open} = \sum_{\mathbf{k}} \frac{(E_{1\,\mathbf{k}} - \xi_{\mathbf{k}})(E_{1\,\mathbf{k}} + \xi_{\mathbf{k}} + \eta) - \Delta_2^2}{(E_{1\,\mathbf{k}} + E_{2\,\mathbf{k}})(E_{1\,\mathbf{k}} + E_{3\,\mathbf{k}})}$$
(13)

$$N_{\text{closed}} = \sum_{\mathbf{k}} \frac{(E_{1\,\mathbf{k}} - \xi_{\mathbf{k}})(E_{1\,\mathbf{k}} + \xi_{\mathbf{k}}) - \Delta_{1}^{2}}{(E_{1\,\mathbf{k}} + E_{2\,\mathbf{k}})(E_{1\,\mathbf{k}} + E_{3\,\mathbf{k}})}$$
(14)

A. The Bogoliubov canonical transformation and the fermionic excitation

The fermionic correlation (Eq. 8) can be diagonalized with a (unitary) Bogoliubov canonical transformation. We break down the transformation into two steps $T_{\bf k}$ and $L_{\bf k}$.

$$B_{\omega_n,\mathbf{k}} = L^{\dagger}_{\mathbf{k}} T^{\dagger}_{\mathbf{k}} G^{-1}_{\omega_n,\mathbf{k}} T_{\mathbf{k}} L_{\mathbf{k}}$$
(15)

Here $B_{\mathbf{k}}$ is the diagonal matrix; $T_{\mathbf{k}}$ and $L_{\mathbf{k}}$ are both unitary transformations. We take $T_{\mathbf{k}}$ as the canonical transformation at the broad resonance limit, i.e., when we can ignore the inter-channel Pauli exclusion.

$$T_{\mathbf{k}} = \begin{pmatrix} u_{\mathbf{k}} & v_{\mathbf{k}} & 0\\ -v_{\mathbf{k}} & u_{\mathbf{k}} & 0\\ 0 & 0 & 1 \end{pmatrix}$$
(16)

where u_k and v_k are defined in a similar fashion as in the single-channel BCS problem

$$v_{\mathbf{k}}^2 \equiv 1 - u_{\mathbf{k}}^2 \equiv \frac{1}{2} \left(1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right) \tag{17}$$

$$E_{\mathbf{k}} \equiv (\xi_{\mathbf{k}}^2 + \Delta_1^2)^{1/2}$$
(18)

Note that in the narrow resonance here, $v_{\mathbf{k}}^2$ does not carry the physical meaning of the occupation number of the (open-channel) atoms, and $E_{\mathbf{k}}$ does not represent the fermionic excitation spectrum. $T_{\mathbf{k}}$ can be taken as the direct sum of two parts, the first two rows/columns describe the open-channel excitations, while the third row/column describes the closed-channel ones. It nevertheless is not sufficient to diagonalize the fermionic correlation in the narrow resonance.

Introduce a dimensionless scale ζ , (Please refer to Appendix A and B for details.)

$$\zeta = \frac{\Delta_2^2}{\Delta_1 \eta} \sim \left(\frac{E_F}{\eta}\right)^{\frac{1}{2}} \ll 1 \tag{19}$$

Here both Δ_1 and Δ_2 are their mean-field (saddle point) values. It is not hard to find the additional unitary transformation $L_{\mathbf{k}}$ to the first order in ζ . (Please refer to Appendix A for details.)

$$L_{\mathbf{k}} \approx I + \begin{pmatrix} 0 & -\frac{\Delta_1 \Delta_2}{4E_{\mathbf{k}}^2} & u_{\mathbf{k}} \\ \frac{\Delta_1 \Delta_2}{4E_{\mathbf{k}}^2} & 0 & v_{\mathbf{k}} \\ -u_{\mathbf{k}} & -v_{\mathbf{k}} & 0 \end{pmatrix} \frac{\Delta_2}{\eta} \equiv I + \delta_k \qquad (20)$$
$$L_{\mathbf{k}}^{\dagger} = I - \delta_{\mathbf{k}}$$

So we finally arrive at the diagonal matrix $B_{\omega_n,\mathbf{k}}$ to first order in ζ

$$B_{\omega_n,\mathbf{k}} = i\omega_n I - \begin{pmatrix} E_{1\mathbf{k}} & 0 & 0\\ 0 & -E_{2\mathbf{k}} & 0\\ 0 & 0 & -E_{3\mathbf{k}} \end{pmatrix}$$
(21)

The eigenvalues of $\hat{\mathcal{G}}^{-1}$ in $B_{\omega_n,\mathbf{k}}$ describes the dispersion spectrum of the fermionic excitations

$$E_{1\mathbf{k}} \approx E_{\mathbf{k}} + \frac{\Delta_2^2 u_{\mathbf{k}}^2}{\xi_{\mathbf{k}} + \eta} \approx E_{\mathbf{k}} + u_{\mathbf{k}}^2 \Delta_1 \zeta \tag{22}$$

$$E_{2\mathbf{k}} \approx E_{\mathbf{k}} - \frac{\Delta_2^2 v_{\mathbf{k}}^2}{\xi_{\mathbf{k}} + \eta} \approx E_{\mathbf{k}} - v_{\mathbf{k}}^2 \Delta_1 \zeta \tag{23}$$

$$E_{3\mathbf{k}} \approx \xi_{\mathbf{k}} + \eta - \frac{\Delta_2^2}{2(\xi_{\mathbf{k}} + \eta)} \approx \epsilon_{\mathbf{k}} + \eta - \frac{\zeta}{2} \Delta_1 \qquad (24)$$

 $E_{1\mathbf{k}}$ and $E_{2\mathbf{k}}$ correspond to the traditional Bogoliubov quasi-particle modes; while $E_{3\mathbf{k}}$ describes the fermionic excitation mostly in the closed channel. The correction due to the inter-channel Pauli exclusion are all of order ζ .

B. Renormalization of the mean-field equation

The summations in the gap equations above (Eq. 10) diverge when they are converted into integrals at 3D because of the artificial assumption of contact interactions. We can nevertheless remove the singularity in two steps. First, we notice that the closed-channel bound state is much smaller than the interparticle distance. Therefore the two-body correlation within the closed channel is almost unchanged from its two-body value. Given this consideration, we project the closed-channel correlation $h_{2\mathbf{k}}$ onto the two-body bound-state wave function of the uncoupled closed channel, ϕ . Second, the bare interaction in the open channel is replaced with a more physically meaningful quantity, the effective s-wave scattering length, \tilde{a}_s , which already incorporates the singularity. We briefly illustrate the procedure as follows, and please refer to [15] for more details.

First, we project the closed-channel correlation $h_{2{\bf k}}$ into ϕ

$$h_{2\mathbf{k}} = \alpha \phi_{\mathbf{k}} u_{\mathbf{k}}^2 \tag{25}$$

This projection is only for the high momentum, for the low momentum, the available phase space for the closed channel is much more limited because the atoms in the open channel center in the low momentum. More specifically, this restriction is represented by the factor $u_{\mathbf{k}}^2$. We come to this conclusion because the closed-channel bound state ϕ is much smaller in size than the interparticle distance. This guarantees that the low momentum states are dominated by the open channel. Comparing this equation with the two-body Schrödinger equation for ϕ , we find

$$\alpha = \frac{\sum_{\mathbf{k}\mathbf{k}'} \phi_{\mathbf{k}}^* Y_{\mathbf{k}\mathbf{k}'} h_{1\mathbf{k}'}}{(-E_b + \eta - 2\mu - \lambda_1)}$$
(26)

where

$$\lambda_1(\eta) \equiv -\sum_{\mathbf{k}} \phi_{\mathbf{k}}^* (E_{\mathbf{k}} - \xi_{\mathbf{k}}) \phi_{\mathbf{k}} - \sum_{\mathbf{k}\mathbf{k}'} \phi_{\mathbf{k}}^* v_{\mathbf{k}'}^2 V_{\mathbf{k}\mathbf{k}'} \phi_{\mathbf{k}'} \quad (27)$$

Put all these together into the gap equation of the open channel, we find

$$\Delta_{1\mathbf{p}} = \sum_{\mathbf{k}} \left(U_{\mathbf{pk}} + \frac{\sum_{\mathbf{k'p'}} Y_{\mathbf{pp'}} \phi_{\mathbf{p'}} \phi_{\mathbf{k'}}^* Y_{\mathbf{kk'}}}{(-E_b + \eta - 2\mu - \lambda_1)} \right) h_{1\mathbf{k}}$$
$$- \frac{\sum_{\mathbf{kk'p'}} Y_{\mathbf{pp'}} \phi_{\mathbf{p'}} \phi_{\mathbf{k'}}^* Y_{\mathbf{kk'}} v_{\mathbf{p'}}^2 h_{1\mathbf{k}}}{(-E_b + \eta - 2\mu - \lambda_1)}$$

Comparing to the similar equations in the two-body problem, we find the detuning from the resonance here differs from that of the two-body physics by $2\mu + \lambda_1$. 2μ describes the many-body shift of the starting point from zero to the Fermi surface, while λ_1 describes the interchannel Pauli exclusion. Furthermore, the last term in the r.h.s. is also unique for the many-body problem. This term has no singularity for high-momentum though because of the extra $v_{\mathbf{p}'}^2$ term in the integral.

Now we handle the above equation with the similar strategy as in the single-channel problem: we integrate out the high-momentum part and replace the bare interaction with the more physically observable effective open-channel a_s . Multiply both side with (1 + TG), where T is the scattering matrix for the open channel, and $G = (\omega - H_0)^{-1}$ is the Green's function for a free pair in the open channel.[19]

$$(1+TG)\Delta_1 = -Th_1 - \lambda_2$$
$$\lambda_{2\mathbf{p}} \equiv \alpha \sum_{\tilde{\mathbf{p}}\,\mathbf{p}'} (1+TG)_{\mathbf{p}\tilde{\mathbf{p}}} Y_{\tilde{\mathbf{p}}\mathbf{p}'} \phi_{\mathbf{p}'} v_{\mathbf{p}'}^2 \tag{28}$$

Using the zero energy value of the free pair Green's function $G(\omega = 0) = (-2\epsilon_{\mathbf{k}})^{-1}$ and introducing the low-momentum s-wave scattering length with extra shift

$$\tilde{a}_s = a_{\rm bg} \left(1 + \frac{\mathcal{K}}{\delta - 2\mu - \lambda_1}\right) \tag{29}$$

where $\mathcal{K} = \Delta_{\mu} B$, Δ_{μ} is the effective difference of the pair magnetic moments of two channels. And δ is the energy detuning between two channels, we have the renormalized open-channel gap equation

$$1 = -\frac{\lambda_2}{\Delta_1} - \left[\frac{4\pi\tilde{a}_s(\mu,\lambda_1)}{m}\sum(\frac{1}{2E_{\mathbf{k}}} - \frac{1}{2\epsilon_{\mathbf{k}}} - \frac{\Delta_2^2\xi_{\mathbf{k}}}{4(\xi_{\mathbf{k}} + \eta)E_{\mathbf{k}}^3})\right]$$
(30)

Here the two factors λ_1 and λ_2 , given respectively by Eqs. (27) and (28), as well as the last term in the above equation, describe the inter-channel Pauli exclusion effect. All of them involve overlap integrals between the open-channel wave function and the closed-channel wave function. (The factor $v^2_{\bf k}$ or $u^2_{\bf k}$ describes mostly the open-channel wave function, while ϕ describes the closed-channel wave function). The larger the overlap of the two, the larger are λ_1 and λ_2 . This has a very intuitive interpretation: more overlap leads to more severe inter-channel Pauli exclusion, which in turn leads to larger (corrections) terms. In our model, the openchannel wave function is spread over a large region of real space, (even on the BEC side, the real bound-state is very loosely bound comparing to the closed-channel bound state), while the closed-channel wave function is very sensitive to the binding energy, $E_b \approx \eta$. A closedchannel bound state is more spread out in real space and has larger overlap with the open-channel wave function, if it is closer to the threshold, i.e., the binding energy is smaller. Consequently, the terms λ_1 and λ_2 are larger in such cases. Nevertheless, λ_1 is much smaller than the Fermi energy E_F , or the other shift, the chemical potential, 2μ . So it is still legitimate to treat this shift as a

perturbation. In addition, it can be shown that all these terms are linear in the density. (Please refer to [15] for details.)

We can also use the expansion on $E_{i\mathbf{k}}$ in Eqs. (22-24) to rewrite the two number equations Eqs. (13, 14) to the first order in ζ

$$N_{\text{closed}} \approx \sum_{\mathbf{k}} \frac{\Delta_2^2}{(\xi_{\mathbf{k}} + \eta)(2\xi_{\mathbf{k}} + \eta)}$$
(31)

$$N_{open} \approx \sum_{\mathbf{k}} \left[\frac{E_{\mathbf{k}} - \xi_{\mathbf{k}}}{2E_{\mathbf{k}}} (1 + \frac{\Delta_1}{\eta}\zeta) - \frac{\Delta_1^3}{4E_{\mathbf{k}}^3}\zeta \right]$$
(32)

Note that the closed-channel correlation $h_{2\mathbf{k}}$ is forced into a particular simple form Eq. 25, which does not lead to divergence in high momentum in integration and therefore has already been "renormalized". This is because the two-body closed-channel wave function ϕ is much smaller comparing to the inter-particle distance and therefore deformed little in many-body case to the lowest order of ζ . This equation implicitly includes Δ_2 through $h_{2\mathbf{k}}$ and therefore we do not need any equation explicitly about Δ_2 .

In the summary, Eqs. (25, 30, 31, 32) together become the renormalized set of equations that determine the state of the system at the mean-field level.

C. Discussion of the mean-field solution

As discussed before, the correction of the narrow Feshbach resonance can be taken into account in two steps. First, omitting the inter-channel Pauli exclusion, we only consider the chemical potential μ in the shift and the extra counting due to the closed channel. Then in the second step, we can correct the previous result with quantities originated from the inter-channel Pauli exclusion unique to the three-species problem.

In the first step, the gap equation and the (openchannel) number equation are simplified to

$$1 = -\left[\frac{4\pi\tilde{a}_s(\mu)}{m}\sum(\frac{1}{2E_{\mathbf{k}}} - \frac{1}{2\epsilon_{\mathbf{k}}})\right]$$
(33)

$$N_{\rm open} = \sum_{\mathbf{k}} \frac{E_{\mathbf{k}} - \xi_{\mathbf{k}}}{2E_{\mathbf{k}}} \tag{34}$$

Here we only consider the shift of the chemical potential 2μ in \tilde{a}_s (Eq. 29),

$$\tilde{a}_s = a_{\rm bg} \left(1 + \frac{\mathcal{K}}{\delta - 2\mu}\right) \approx \frac{a_{\rm bg}\mathcal{K}}{\delta - 2\mu} \tag{35}$$

The above equations need to be solved self-consistently.

An interesting point to notice is that the gap Δ_1 saturates in the BEC side. This is because the effective attraction in the open channel where the pairing happens is limited by the real attractive strength in the closed channel. It can not become infinitely strong as in an ideal single-channel model. Mathematically, this can be seen from the gap equation, Eqs. (10-12). The closed-channel correlation is limited by the total density, and therefore the gap has a maximum.

Once this step is finished, we can look for the correction due to the inter-channel Pauli exclusion. Both λ_1 and λ_2 can be shown to be linear with the density, so can be calibrated by experiments of different densities. With the proper value of these two parameters, we can find the correction numerically. It is nevertheless not hard to show that the correction is of order ζ , so that we are warranted to treat this effect as a perturbation [15].

IV. BEYOND THE MEAN-FIELD: THE COLLECTIVE MODES

The order parameters (Δ_1, Δ_2) are defined in terms of the collective behaviors of many fermion atoms. Fluctuations of the order parameters thus signal the collective excitation modes of the system. Here with a twocomponent complex order parameter, four independent modes exist: two for the magnitude variation of each Δ_i , the internal phase between Δ_1 and Δ_2 , and the overall local phase $\theta(x)$ of Δ_1 and Δ_2 . A similar result has been obtained in the BCS limit by Catelani and Yuzbashyan,[20]. Two modes of the magnitudefluctuation are gapped and massive as expected. The modes of the phase-fluctuation are of more interest. We summarize our finding in the following. Please refer to [15] for more details.

A. The in-sync phase mode

The in-sync phase mode is the counterpart of the Anderson-Bogoliubov modes in the single-channel problem[8, 21]. In this mode, $\Delta_{1\mathbf{k}}$ and $\Delta_{2\mathbf{k}}$ rotate simultaneously, and the action $S(\bar{\Delta}_i, \Delta_i)$ (Eq. 9), is invariant for such fluctuation. We therefore conclude that there exists a massless (Goldstone) mode corresponding to the local phase invariance. Introduce the phase fluctuation θ ,

$$\Delta_i(x) \to \Delta_i e^{i2\theta(x)} \qquad \bar{\Delta}_i(x) \to \bar{\Delta}_i e^{-i2\theta(x)}$$

In order to focus on this particular mode, we replace every degrees of freedom except θ with their equilibrium values. For small perturbation around the equilibrium, we only retain the first non-trivial order of θ (the second order). It can then be shown that the fluctuation has a linear dispersion relation and therefore is a sound-like mode.

$$S[\theta] = \sum_{q} \theta(q)\theta(-q) \left[\frac{1}{2}\pi^{(0)}(0)\omega_{m}^{2} - \frac{n}{2m}\mathbf{q}^{2}\right]$$
(36)

where n is the density of the pairs and

$$\pi^{(0)}(0) \equiv \sum_{k} \operatorname{tr} \left(G_k \sigma_3 G_k \sigma_3 \right) \tag{37}$$

 G_k is the mean-field fermionic correlation in Eq. 8 and σ_3 is defined as

$$\sigma_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$
(38)

Following the same approach as the last section, G_k can be expanded with respect to ζ , after some algebra, we can write down the $\pi^{(0)}(0)$ to the first non-trivial order of ζ .

$$\pi^{(0)}(0) \approx \sum_{\mathbf{k}} \frac{\Delta_1^2}{E_{\mathbf{k}}^3} - \sum_{\mathbf{k}} \frac{\Delta_1^2 \Delta_2^2 \xi_{\mathbf{k}}}{2E_{\mathbf{k}}^5 (\xi_{\mathbf{k}} + \eta)}$$
(39)

 Δ_1 and Δ_2 are the mean-field values of two order parameters. The first term in the above formula is the same as that of the single-channel crossover[8]. It then becomes clear that the sound velocity follows the same structure as that for the single-channel crossover problem with a correction of order ζ .

B. The out-of-sync phase mode

The out-of-sync phase mode on the other hand is a novel mode associated with the two-channel problem, with no direct counterpart in the single-channel problem. When the phase fluctuation of two channels are out of sync, the inter-channel coupling strength changes. Thus, this mode is expected to be a gapped (massive) mode. Similar as Sec. IV A, we narrow down to the mode that the phases of two atoms (ψ_b and ψ_c) are opposite and leave all other degrees of freedom constant.

$$\begin{pmatrix} \psi_a(x)\\ \psi_b(x)\\ \psi_c(x) \end{pmatrix} \to \begin{pmatrix} \psi_a(x)\\ \psi_b(x)e^{+i\theta(x)}\\ \psi_c(x)e^{-i\theta(x)} \end{pmatrix}$$

The order parameters do not have a simple transformation because they are connected to two channels via 2×2 interaction matrix \tilde{U} , which mixes two channels (Eq. 5).

$$\begin{pmatrix} \Delta_1(x) \\ \Delta_2(x) \end{pmatrix} \to \begin{pmatrix} U & Y \\ Y^* & V \end{pmatrix} \begin{pmatrix} \psi_b \psi_a(x) e^{+i\theta(x)} \\ \psi_c \psi_a(x) e^{-i\theta(x)} \end{pmatrix}$$

This term cannot be easily written in terms of meanfield value Δ_i . On the other hand, as mentioned before, we freeze all other modes to their mean-field value. We therefore use another two-component of atom pairs ($\psi_b \psi_a, \psi_c \psi_a$), which is the linear recombination of (Δ_1, Δ_2). After some lengthy algebra, we can show that the action is

$$S[\theta] = \sum_{q} \theta(q)\theta(-q) \left[\frac{1}{4}\pi^{(0)}(0)(\omega_m^2 - \omega_0^2) - \frac{n}{4m}\mathbf{q}^2\right] (40)$$

where

$$\omega_0^2 = -\frac{16Yh_1^*h_2}{\pi^{(0)}(0)}$$
$$\tilde{h}_1 = \left\langle \psi_b \psi_a \right\rangle = \sum_{\mathbf{k}} h_{1\mathbf{k}} \qquad \tilde{h}_2 = \left\langle \psi_c \psi_a \right\rangle = \sum_{\mathbf{k}} h_{2\mathbf{k}}$$

Here we can see that ω_m has a finite value ω_0 at the zero momentum, which indicates a gapped mode. It can be shown that ω_0 is in the order of the ionization (pair breaking) energy, which is around Δ_1 on the BCS side, while around $|\mu|$ on the BEC side.

Again, please refer to [15] for more details.

V. CONCLUSION AND DISCUSSION

In this paper, we have studied the narrow Feshbach resonance in the three-species case where two channels share the same species.

In general, for the narrow resonance without a shared species, the main correction to the single-channel result comes from the extra counting of the atoms in the open channel, which leads to the extra shift 2μ in \tilde{a}_s , and the closed channel, which leads to the extra number equation. Two number equations exist, one for each channel. The open-channel number equation resembles the number equation of the single-channel model.

When there is a common species, however, the Pauli exclusion between two channels due to the common species in the three-species narrow resonance, calls for careful consideration.

Our treatment follows the the idea of "universality" [22, 23]. For a dilute system with a short-range potential, such as the dilute ultracold alkali gas, the shortrange part of a two-body correlation does not significantly change from two-body to many-body. This particular feature justifies using the two-body quantities as the boundary condition for the many-body correlations. In non-resonant situations, the s-wave scattering length, a_s , is a good candidate for such a purpose. However, in a Feshbach resonance, the open-channel scattering length becomes enormous, comparable to or even larger than the many-body length scale; furthermore, its value is very sensitive to the ratio of weights of atoms in the two channels. Therefore it is no longer suitable as the two-body characteristic quantity used in the many-body problem. A better candidate of two-body quantities as boundary condition here is the short-range part of the two-body wave function itself. In other word, we expect the twobody correlation of the many-body system matches the shape of the two-body wave function at short distances.

When the spatial extension a_c of the closed-channel bound state is of the order of the inter-particle distance, a_0 , or even larger, the Feshbach resonance in the manybody context is a genuine three-species many-body problem and no simple solution is available to our knowledge. By contrast, when the bound-state's spatial extension is much smaller than the interparticle distance,

the two-body correlation in the closed channel, which is expected to almost entirely concentrated in the shortrange part, is proportional to the two-body bound-state wave function. The ratio of the the bound-state size and the interparticle distance, $a_c/a_0 \sim \zeta$, serves as the expansion parameter and we can extract the effect of the inter-channel Pauli exclusion perturbatively. In essence, we can then ignore the many-body effects within closedchannel bound states, while only taking into consideration of the Pauli exclusion between channels and within the open channels. A few new parameters need to be introduced and can be calibrated from experiments, such as λ_1, λ_2 . Mean field properties can still be determined through gap equations and number equations similar to the single-channel case. The excitation modes are also close to the original single-channel result with correction of the order of ζ .

In our approach, we take the broad resonance result (or the single-channel crossover) as our zeroth order solution, upon which the expansion is performed. It is however known that the simple BCS-type pairing treatment is not adequate to quantitatively describe the whole BEC-BCS crossover region. Therefore the zeroth order solution used here can be improved through further theoretical development. Nevertheless, we expect the perturbation approach used here to build the narrow resonance result from the single-channel crossover result to remain valid. Once the zeroth order solution (for a broad resonance or a single channel BEC-BCS crossover model) is appropriately improved, the correction of the narrow resonance in such a parameter regime, can still be obtained by a procedure similar to the one described in this paper.

ACKNOWLEDGMENTS

We thank Professor Monique Combescot, Dr. Shizhong Zhang and Dr. Wei-Cheng Lee for many inspiring discussions. Part of this research is supported by the National Science Foundation under grant No. DMR 09-06921.

Appendix A: Diagonalization of the mean-field Green's function Eq. (8)

Apply T onto G^{-1} (Eq. 8), we have

$$T_{\mathbf{k}}^{\dagger}G_{\omega_{n},\mathbf{k}}^{-1}T_{\mathbf{k}} = i\omega_{n}I + \begin{pmatrix} -E_{\mathbf{k}} & 0 & u_{\mathbf{k}}\Delta_{2} \\ 0 & +E_{\mathbf{k}} & v_{\mathbf{k}}\Delta_{2} \\ u_{\mathbf{k}}\Delta_{2} & v_{\mathbf{k}}\Delta_{2} & +\xi_{\mathbf{k}} + \eta \end{pmatrix}$$
(A1)

We drop all the k subscripts in the rest of this section because matrices in this section are decoupled in momentum and we only deal with one particular momentum \mathbf{k} a time. The off-diagonal elements in the above matrix is regarded as perturbation because we only seek the solution around the BCS wave function (*T* transformation). We need to find a unitary transformation *L* to diagonalize this matrix. We notice that the first term is proportional to an identity matrix and does not change by unitary transformation, we only need to concentrate on the second term. We rescale all elements with $E_{\mathbf{k}}$ for simplicity in the following of this section.

$$y = \frac{\Delta_2}{E_{\mathbf{k}}}, \qquad t = \frac{\xi_{\mathbf{k}} + \eta}{E_{\mathbf{k}}},$$

And the second term in r.h.s. of Eq. A1 is

$$R = \begin{pmatrix} -1 & 0 & uy \\ 0 & 1 & vy \\ uy & vy & t \end{pmatrix}$$

The secular equation of R is (|xI - R| = 0)

$$(x^{2} - 1)(x - t) - y^{2}x + (u^{2} - v^{2})y^{2} = 0$$
 (A2)

We use $u^2 + v^2 = 1$ here. We assume at the zeroth order, the three eigenvalues are -1, 1 and t. (t has weak dependency on energy as $(\xi_k + \eta)/E_k$, however, at the low energy region of interest, we ignore ξ_k .) Both y and t are larger than 1, however, we will verify that given condition $y^2 \ll t$, the correction is indeed small and the expansion is reasonable (See Appendix B). Indeed, closechannel component can still be smaller than the openchannel component at low-k (in the order of k_F) due to the close-channel bound state is much smaller than the interparticle distance even when the total close-channel atom number is more than that of the open channel. And here all the quantities are about low-k unless specifically noticed. We expand the system to the first order of the dimensionless parameter $\zeta = y^2/t$ (19), and find

$$\begin{array}{rcrr} x^{(0)} & x^{(1)} & Eigenvector \\ -1 & -u^2 \tilde{\zeta} & \left(1 & \frac{uvy^2}{2t} & -\frac{uy}{t}\right) \\ 1 & -v^2 \tilde{\zeta} & \left(-\frac{uvy^2}{2t} & 1 & -\frac{vy}{t}\right) \\ t & \frac{1}{2} \tilde{\zeta} & \left(\frac{uy}{t} & \frac{vy}{t} & 1\right) \end{array}$$

Now it is easy to write down the corresponding diagonal matrix and the unitary transformation

$$B = i\omega_n I + E \begin{pmatrix} -1 - u^2 \tilde{\zeta} & 0 & 0\\ 0 & 1 - v^2 \tilde{\zeta} & 0\\ 0 & 0 & t + \frac{1}{2} \tilde{\zeta} \end{pmatrix}$$
(A3)

$$L = \begin{pmatrix} 1 & -\frac{uvy^2}{2t} & \frac{uy}{t} \\ \frac{uvy^2}{2t} & 1 & \frac{vy}{t} \\ -\frac{uy}{t} & -\frac{vy}{t} & 1 \end{pmatrix}$$
(A4)

Here L is not exactly unitary transformation, it is only unitary in the first order of $\tilde{\zeta}$. We have

$$B = i\omega_n I + E\left(L^{\dagger}RL\right) + o(\tilde{\zeta})$$

Restore the factor $E_{\mathbf{k}}$ and we can obtain Eq. (20) and Eq. (21). (Here we use $uv = \Delta_1/2E$.)

In the above treatment, the small parameter ζ is momentum dependent. If we restore the subscript **k** and scale it back with $E_{\mathbf{k}}$

$$\tilde{\zeta} = \frac{\Delta_2^2}{E_{\mathbf{k}}(\xi_{\mathbf{k}} + \eta)} \tag{A5}$$

A momentum-dependent small parameter is not very convenient to work with, so we take its maximum value in low momentum ($\leq E_F$). In the BCS-like states ($\mu > 0$), min $E_k = \Delta_1$, min $\xi_{\mathbf{k}} = 0$; in the BEC-like states ($\mu < 0$), min $E_k = \sqrt{\Delta_1^2 + \mu^2}$ and min $\xi_{\mathbf{k}} = |\mu|$. We take the smaller values and have our expanding small parameter $\zeta(19)$

$$\zeta = \max \tilde{\zeta} = \frac{\Delta_2^2}{\Delta_1 \eta} \tag{A6}$$

Appendix B: Smallness of the expansion factor ζ

Here we check the smallness of our expansion factor $\zeta(19)$. We have the closed-channel gap equation (10)

$$\Delta_2 = \sum Y h_{1\mathbf{k}} + \sum V h_{2\mathbf{k}} \tag{B1}$$

The first term on the right is relatively small comparing to the second term. Therefore, we drop the first term in estimation. Furthermore, we assume $h_{2\mathbf{k}} = \sqrt{N_c}\phi_{\mathbf{k}}$, where N_c is the total number of closed-channel pairs, and $\phi_{\mathbf{k}}$ is the normalized wave function of the isolated closedchannel potential satisfying the two-body Schrödinger equation

$$-E_b^{(0)}\phi_{\mathbf{p}} = 2\epsilon_{\mathbf{p}}\phi_{\mathbf{p}} - \sum_{\mathbf{k}} V\phi_{\mathbf{k}}$$
(B2)

Rearranging it, we have (especially at low momentum)

$$\sum_{\mathbf{k}} V \phi_{\mathbf{k}} = (2\epsilon_{\mathbf{p}} + E_b) \phi_{\mathbf{p}} \approx \eta \phi_{\mathbf{p}}$$

Here E_b is the binding energy of the closed-channel bound state, which is close to the Zeeman energy difference, η , around the Feshbach resonance. The second approximation is correct at low momentum not too far away from the resonance (smaller or in the same order of the Fermi momentum), i.e. $\epsilon_{\mathbf{p}} \ll E_b \approx \eta$. Put all these together, we have

$$\Delta_2 \approx \alpha E_b \phi_{k=0}$$

If we assume a simple exponentially decayed wave function:

$$\phi_{\mathbf{k}} = \sqrt{\frac{8\pi\kappa}{\mathcal{V}_0}} \frac{1}{k^2 + \kappa^2} \approx \sqrt{\frac{8\pi\kappa}{\mathcal{V}_0}} \frac{1}{\kappa^2} \tag{B3}$$

Here \mathcal{V}_0 is the total volume and κ is the characteristic momentum of the closed-channel bound state, $\eta \approx E_b =$ $\hbar^2 \kappa^2 / 2m$. The second approximation above is only for low momentum. Collect all these together, we have

$$\Delta_2 \approx \sqrt{N_c} \eta \sqrt{\frac{8\pi\kappa}{\mathcal{V}_0}} \frac{1}{\kappa^2} \sim \eta \left(\frac{k_{Fc}}{\kappa}\right)^{\frac{3}{2}} \tag{B4}$$

 $k_{Fc} \sim (N_c/\mathcal{V}_0)^{\frac{1}{3}}$ is the Fermi momentum corresponding to the density of the closed-channel pairs, which is much smaller than the characteristic momentum for the bound-state, κ .

Now let us get back to
$$\zeta$$
 (19)

$$\zeta = \frac{\Delta_2^2}{\Delta_1 \eta} \tag{19}$$

At the BCS limit, the closed-channel density is small, k_{Fc} is small and that makes ζ small; Moving toward the (narrow) resonance, where the closed-channel density is comparable to to the total density, at low energy, Δ_1 is in the order of the Fermi energy. We have (we no longer distinguish k_{Fc} with k_F)

$$\zeta = \frac{\Delta_2^2}{\Delta_1 \eta} \sim \frac{\eta^2 \frac{k_{Fc}^2}{\kappa^3}}{k_F^2 \eta} \sim \frac{k_F}{\kappa} \ll 1 \tag{B5}$$

which is also very small.

- [1] U. Fano, Phys. Rev. **124**, 1866 (1961).
- [2] H. Feshbach, Annals of Physics **19**, 287 (1962).
- [3] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Rev. Mod. Phys. 82, 1225 (2010).
- [4] C. J. Pethick and H. Smith, Bose-Einstein Condensation in Dilute Gases (Cambridge University Press, 2001).
- [5] D. M. Eagles, Phys. Rev. **186**, 456 (1969).
- [6] A. J. Leggett, in Proceedings of the XVIth Karpacz Winter School of Theoretical Physics, Karpacz, Poland (Springer-Verlag, 1980) pp. 13–27.
- [7] P. Nozières and S. Schmitt-Rink,
 J. of Low T. Phys. 59, 195 (1985),
 10.1007/BF00683774.
- [8] M. Randeria, in *Bose-Einstein Condensation*, edited by A. Griffin, D. Snoke, and S. Stringari (Cambridge University Press, 1995) pp. 355–392.
- [9] A. J. Leggett, *Quantum Liquids* (Oxford University Press, 2006) appendix 4A.
- [10] M. Holland, S. J. J. M. F. Kokkelmans, M. L. Chiofalo, and R. Walser, Phys. Rev. Lett. 87, 120406 (2001).
- [11] T.-L. Ho, Phys. Rev. Lett. 92, 090402 (2004).
- [12] J. N. Fuchs, A. Recati, and W. Zwerger, Phys. Rev. Lett. 93, 090408 (2004).
- [13] G. M. Bruun, A. D. Jackson, and E. E. Kolomeitsev, Phys. Rev. A 71, 052713 (2005).
- [14] V. Gurarie and L. Radzihovsky, Annals of Physics **322**, 2 (2007), january Special Issue 2007.
- [15] G. Zhu, BEC-BCS Crossover with Feshbach Resonance for Three-Hyperfine-Species Model, Ph.D. thesis, University of Illinois at Urbana-Champaign (arXiv:1212.6691) (2012), arXiv:1212.6691.

- [16] S. Zhang, Strongly interacting Fermi gases, radio frequency spectroscopy and universality, Ph.D. thesis, University of Illinois at Urbana-Champaign (2009).
- [17] When Y is not real, Δ_1 and Δ_2 cannot be both real even at the mean field level. Nevertheless, we can require one real, then the other will have a phase just to compensate the phase in Y. The final conclusion can be verified to remain valid.
- [18] See sec. 4.2.1 in [24], sec. 25 in [25].
- [19] Here we use the relation between the scattering *T*-matrix, the free pair Green's function $G = (\omega - H_0)^{-1}$ and the bare interaction $V = -U_{\text{eff}}$.

$$T = V + TGV = -(1 + TG)U_{\text{eff}}$$

- [20] G. Catelani and E. A. Yuzbashyan, Phys. Rev. A 78, 033615 (2008).
- [21] N. Nagaosa, Quantum Field Theory in Condensed Matter Physics (Springer, 1999) Original Japanese edition title: Quantum Field Theory in Strongly Correlated Systems; Published in Japan by Iwanami Shoten, Publisher, Tokyo in 1995; Translated by Stefan Heusler;.
- [22] S. Tan, Annals of Physics **323**, 2952 (2008).
- [23] S. Zhang and A. J. Leggett, Phys. Rev. A 79, 023601 (2009).
- [24] A. Altland and B. Simons, Condensed Matter Field Theory (Cambridge University Press, 2010).
- [25] A. L. Fetter and J. D. Walecka, Quantum Theory of Many-Particle Systems, dover ed. (Dover Publications, Inc. (Original Publisher: McGraw-Hill), 1971).